

Does women's place of birth affect their opportunity for an informed choice about Down syndrome screening? A population-based study in France.

Olivia Anselem (✉ olivia.anselem@aphp.fr)

Maternité Port-Royal <https://orcid.org/0000-0002-1516-3203>

Marie-Josèphe Saurel-Cubizolles

Université de Paris, CRESS, Obstetrical Perinatal and Pediatric Resarch Team, EPOPé, INSERM, INRA,
Paris, France

Babak Khoshnood

INSERM

Béatrice Blondel

INSERM

Priscille Sauvegrain

INSERM

Nathalie Bertille

INSERM

Elie Azria

Groupe hospitalier Paris Saint-Joseph

Research article

Keywords: Down syndrome, prenatal diagnosis, social disparities, implicit bias

Posted Date: December 3rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-33585/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background To examine disparities in the opportunity to make an informed choice about Down syndrome screening by maternal place of birth, in France, where the French national authority for health (HAS) recommends that physicians offer it to all. We used population-based data from the nationally representative French Perinatal Surveys in 2010 and 2016 (N=24,644 women) to analyze the opportunity for an informed choice for prenatal screening, measured by a composite indicator.

Results Among the 24 644 women in the study, 20 612 (83.6%) were born in France, 861 (3.5%) elsewhere in Europe, 1550 (6.3%) in North Africa, and 960 (3.9%) in Sub-Saharan Africa. The probability of screening was lower for women born outside France. After adjustment for survey year, maternal age, parity, education level, and level of perinatal care of the maternity, women born outside France had less often the opportunity to make an informed choice as compared with women born in France. This association remained essentially the same even after excluding women without adequate prenatal care.

Conclusions Women born outside France, including those with adequate prenatal care, had less opportunity than women born in France to make an informed choice about prenatal screening for Down syndrome.

Introduction

Down syndrome (trisomy 21) is the most common genetic cause of intellectual disability (1). Prenatal screening techniques for this disease have advanced considerably in recent decades. In France, the French national authority for health (HAS) coordinates the regularly updated strategy for offering screening in optimal conditions. HAS currently recommends that physicians inform all women that nuchal translucency can be measured at the first ultrasound examination, between 11 weeks and 13 weeks +6 days, and that they can have blood tests. They enable a risk calculation that integrates the maternal serum markers and can be assayed during the first or second trimester (2). The methods of Down syndrome screening in France are legally regulated by decrees published successively in 1997, 2009, and 2018 (3), which authorize both physicians and midwives to offer these screening tests. Until 2017, women for whom this screening showed an increased risk of Down syndrome were offered a diagnostic examination — amniocentesis or chorionic villus sampling — to determine fetal karyotype. Only that year did HAS include the study of fetal DNA in its strategy for detecting Down syndrome among women at increased risk (2).

Since 1997, the National Health Insurance Fund provides full coverage of screening for Down syndrome by the first-trimester ultrasound and the serum marker assays for all women with health insurance. Despite hopes that the disparities in prenatal diagnosis of Down syndrome observed before 1997 would disappear with its full financial coverage, they have persisted, as Khoshnood et al. showed in an analysis of data from the Paris Registry of Congenital Malformations covering the period from 1983 to 2003 (4). This analysis showed progress over time in use of this screening, but also persistence in social

disparities, with a screening rate that varied by socioeconomic status and by the mother's place of birth. In 2009 a new decree of the HAS recommended the generalization of Down syndrome screening at the first trimester, resulting in an integrated estimation of the risk of Down screening, possibly easier to implement than the previous screening. This important change in the screening strategy may have affected access to screening for certain social groups, especially foreign born women and changed the magnitudes of inequalities previously observed.

Work in other settings has shown similar results with important disparities in screening and in diagnosis. This is the case for example in the Netherlands, where women born in Turkey, North Africa, and the West Indies had rates of screening and diagnosis lower than those of native-born Dutch women (5). Studies showing disparities in the performance of screening do not, however, allow us to differentiate disparities in access to screening from differences in their use that result from women's deliberate choices. As some authors suggest, social groups may differ in their willingness to undergo an examination so closely related to a strong parental attitude, one that may be culturally or socially determined (6).

In England, Rowe et al. sought to avoid the question of individual choice by focusing on the offer of screening — asking women if the test was offered to them; they observed no geographic inequalities between the more and less advantaged areas (7). They also found that the test was offered to Asian women less often than to white women. In France, the existence of disparities in access to this screening between women according to their nationality has been demonstrated by analysis of the opportunities they had to make a choice about Down syndrome screening (8). Disparities in access fell between 1998 and 2003, but in 2003 91% of the women born in France had an opportunity to make a free and informed choice, compared with only 62% of those born in Sub-Saharan Africa.

The objective of this study is to analyze the opportunity to make an informed choice about Down syndrome screening and the rates of this screening in 2010s on a national scale in France, according to the mother's place of birth. Moreover, we also aimed to analyze the access to Down syndrome screening specifically for women with adequate prenatal care, in order to assess whether there are any disparities which are not linked to inequalities in prenatal follow-up but that could be linked to differences in terms of tests proposal by the professionals.

Methods

Data for this analysis come from the National Perinatal Surveys conducted in 2010 and 2016. These cross-sectional surveys collected data from all births at a gestational age of at least 22 weeks or with a birth weight of at least 500 grams during a single week, in all maternity units in metropolitan France and the overseas territories (9).

The data came both from the medical records, mainly on delivery, and the child's condition at birth, and a postpartum face-to-face questionnaires with parturients before their discharge. This interview focused on their prenatal care and their socioeconomic characteristics.

Women who did not speak French, were younger than age 18 and those who experienced stillbirth or termination of pregnancy were excluded from the study.

In 2010, 13 894 women with 14 142 babies were included, and in 2016, the sample comprised 15 187 women with 15 418 babies. The two survey samples were first compared to check the absence of major heterogeneity and to analyze the evolution in Down syndrome screening uptake. The two samples were then merged and analyzed together to improve the statistical power of the analysis.

We selected women who had lived in metropolitan France for at least one year when they gave birth and whose place of birth was known: 13 507 women in 2010 and 11 137 women in 2016.

Maternal characteristics studied

The mother's place of birth was classified in five categories: France, other European countries, North Africa, Sub-Saharan Africa, and the rest of the world.

In both surveys, the questionnaire used as the basis for the face-to-face interview included the following questions about Down syndrome screening: "Did you have fetal nuchal translucency measured at your first-trimester ultrasound to learn your risk of Down syndrome?" and "Did you have a blood test to learn your risk for Down syndrome (serum markers)?" The possible responses were "Yes," "No," or "I don't know". If the answer to the latter question was negative, the reasons could be: refusal; amniocentesis or chorionic villus sampling from the start; tests were not offered; late initiation of prenatal care or no prenatal care or prenatal care abroad; another reason, or "don't know".

To analyze access to Down syndrome blood screening test, we used a composite indicator built from these answers defined by Grupposo et al. (8). Accordingly, we considered that women had an opportunity to make an informed choice about Down syndrome screening if they had undergone or refused a serum marker assay or had had an initial amniocentesis or chorionic villus sampling. We considered that women had not had this opportunity when they did not know if they had had the assay or if it had not been performed for one of the following reasons: not offered, because of late antenatal care, for another reason, or for an unknown reason (Figure 1).

Adequate prenatal care was defined in accordance with French clinical practice guidelines (10) by at least 8 antenatal visits and at least 3 prenatal ultrasounds for term births, and according to gestational age at birth for preterm births (9).

The women's characteristics were previously described in the National Perinatal Survey report (11). The other covariables considered were age, parity, and educational level.

Statistical analyses

Pearson's Chi-square tests were used to compare rates of Down syndrome screening and the opportunity to make an informed choice between 2010 and 2016, according to the mother's country of birth.

Differences were considered significant when p was less than 0.05.

To study the association between the mothers' country of birth and rates of the different prenatal screening tests for Down syndrome, as well as the opportunity to make an informed choice about it, we used multivariate multinomial logistic regression models adjusted for the survey year, the individual factors (maternal age, parity, educational level, and prenatal care), and for the maternity ward characteristics (level of perinatal care: I, IIA, IIB, and III). The analysis was performed after fusion of the databases for 2010 and 2016.

For the analyzed sample, the rate of missing data per variable ranged only from 0.3% to 0.8% and thus did not justify an analysis with multiple imputations. The results presented are therefore those for the complete cases only.

The statistical analyses were performed with Stata 13 software.

Results

Among the 24 644 women included, 20 612 (83.6%) were born in France, 861 (3.5%) in another European country, 1550 (6.3%) in North Africa, 960 (3.9%) in Sub-Saharan Africa, and 661 (2.7%) in the rest of the world.

Prenatal screening for Down syndrome, whether by nuchal translucency measurements or maternal serum marker assays, rose significantly between 2010 and 2016 ($p<0.001$) (Table 1). The proportion of women reporting that they were not offered serum marker testing dropped from 12.7% in 2010 to 9.5% in 2016. Initial use of amniocentesis or chorionic villus sampling also regressed, from 9.7% in 2010 to 2.2% in 2016. The proportion of women who did not know why their serum markers were not tested fell from 2.6% in 2010 to 2.1% in 2016. On the other hand, refusal to have these markers assay rose, accounting for 59.1% of the reasons for non-testing in 2016, up from 44.0% in 2010.

Between these two surveys, the rate of women who did not have an "opportunity to make an informed choice" dropped significantly, from 8.4% in 2010 to 5.4% in 2016 ($p<0.001$).

Cumulatively in the two surveys, women born outside of France underwent Down syndrome screenings less often than women born in France (Table 2): 76.3% of the women born in other European countries, 62.6% of those born in North Africa, and 62.4% born in Sub-Saharan Africa reported fetal nuchal measurement, compared with 90.3% of the native Frenchwomen ($p<0.001$). Similarly, 77.9% of the women born in other European countries, 62.7% of those from North Africa, and 69.0% from Sub-Saharan African countries reported serum marker assays, compared with 89.5% of the French-born women ($p<0.001$).

The rates of women who did not know if they had had nuchal translucency measurements or serum marker assays were significantly higher among the women born abroad than among those born in France.

The analysis showed that the women born outside France had the "opportunity to make an informed choice" less often than the natives. Overall, 12.8% of the women born elsewhere in Europe, 20.1% of those born in North Africa, and 21.6% of those born in Sub-Saharan Africa did not have this opportunity, compared with 4.9% of the native Frenchwomen ($p<0.001$).

After adjustment for the survey year, maternal age, parity, educational level, and the maternity ward's level of perinatal care, birth abroad was associated with higher risk of not having the opportunity to make an informed choice (Table 3). Specifically, this was the case for the women born in other European countries (aOR 2.3, 95% CI 1.8-2.9), in North Africa (aOR 3.8, 3.3-4.5) and in Sub-Saharan Africa (aOR 3.3, 2.7-3.9).

After exclusion of the women without adequate prenatal care, these risks remained high: OR_a 2.4 (1.8-3.1) for the women born in other European countries, OR_a 4.0 (3.3-4.7) in North Africa and OR_a 3.2 (2.5-4.0) in Sub-Saharan Africa.

Discussion

Our study shows that women born abroad and living in France when they gave birth were less likely to be screened for Down syndrome – either or both of nuchal translucency measurements and serum marker assays – than women born in France. It also shows the rate of women who did not have the opportunity to choose the serum screening, for which consent is required, was higher for foreign born women than among women born in France.

The principal strength of our study is the good quality of the data from representative nationwide populations. Moreover, a face-to-face questionnaire used during an interview within days of delivery asked about the details of the different stages of screening and the reasons why they were not performed. The pooling of data from two different survey years has been taken into account by the statistical analysis. Moreover, a recall bias cannot be ruled out, to the extent that the information was collected during the postpartum period, but there it could exist equally in all groups. Women who did not speak French well enough to respond to the questionnaire were not questioned and are not included in this analysis. This might create a selection bias attenuating the strength of the association between the mother's place of birth and the availability of an informed choice. Moreover, the composite criterion of "an opportunity to make an informed choice" combines in a single category the women who had screening, those who refused it, and those who had an invasive examination. It is possible that some women had the screening without having understood it or without having been clearly informed about its purpose and its implications, because of language barriers.

Other authors have reported disparities in Down syndrome screening according to women's origin. In the Netherlands, Fransen et al. found a lower rate of participation in Down syndrome screening among women with at least one parent born abroad (North Africa, Turkey, Surinam, West Indies, other Western or non-Western countries) compared with women born to parents both born in the Netherlands (5). In Australia, Maxwell et al. showed that screening for Down syndrome took place less often for Aboriginal Australian women than for their non-Aboriginal counterparts (12). Other authors have studied the use of

diagnostic examinations such as amniocentesis or chorionic villus sampling and found the same disparities according to women's geographic origin (13,14).

Nonetheless, these studies cannot differentiate problems in access to these tests from the choice to not undergo screening. That is, the lower frequency of Down syndrome screening among women born abroad may result from their attitude towards this test, determined in part by social or cultural factors (6, 15–17). Therefore, these studies did not inform about the ability of the healthcare system to respond to the needs of different groups.

In our study, the use of the composite indicator "opportunity to make an informed choice" makes it possible to study the offer of screening and its disparities by maternal place of birth. Among the women who did not have the opportunity to make a choice, some lacked access to screening because of barriers to this access, which we know are numerous among migrant women (18). Unlike the previous study using this indicator, we also excluded women who arrived in France less than a year before giving birth and women whose prenatal care was considered suboptimal. The hypothesis that these disparities in access to prenatal care determine the disparities in access to an informed choice about this screening, is nonetheless largely questioned by the nearly identical results we found after those exclusions. These disparities in access to an informed choice are therefore not linked only to the late start of prenatal care. Moreover, women with problems speaking French were not included, these disparities cannot be attributed only to language barriers either.

Our study thus allows us to show that women born abroad, including those receiving adequate prenatal care, had less often the opportunity to make an informed choice about Down syndrome screening than those born in France; that is, differential care exists in this context. Several hypotheses might explain why some physicians and midwives responsible for telling women about the existence of and the reasons and procedures for these screening tests failed to do so. It is possible that some physicians have prejudices about the attitudes of women born abroad toward this screening or toward terminations of pregnancy after a positive diagnosis. A Delphi consensus process in France thus showed that experts considered that women born in North Africa and Muslim women were less likely to request terminations of pregnancy (19).

These prejudices or implicit biases concerning some ethnic or cultural groups can thus lead to differential care, manifested by the disparities observed here. While the existence of implicit bias has been demonstrated repeatedly by physicians in different specialties (20–26), including in perinatal medicine (27), very little proof exists of its manifestations in terms of differential care (20, 28–30). Other work nonetheless converges with ours to support this hypothesis. Rowe et al. also distinguished the offer of screening and its performance in England and showed that women born in Asia were offered screening less often (7). They also showed that when it is proposed, it is chosen less often. Some authors have used scales to study women's attitude toward screening by their socio-occupational category or their geographic origin (17, 31–35). The prospective study by Dormandy et al. of 1499 women in England

showed that women born abroad were no more unfavorable to screening than those born in England; nonetheless the women born abroad knew less about the screening (17).

Screening for Down syndrome is a complex process, becoming even more complex with technical progress, in particular the examination of fetal DNA in maternal blood. The level of knowledge of the women who accept it is clearly suboptimal, as demonstrated in France (36). Our study shows that inadequate information is still more marked among women born abroad, probably because their level of health literacy is inadequate and lower than that of women born in France, but possibly also because some physicians provide differential care related to implicit biases.

Conclusion

Women born outside France, including those with adequate prenatal care, had a lesser opportunity to make an informed choice about prenatal screening for Down syndrome as compared with those born in France.

The rapid technical advances in screening for fetal abnormalities have led to increasingly more effective tests, in particular examination of fetal DNA in maternal blood. At the same time, these advances have made the process of prenatal screening for Down syndrome increasingly complex. Hence, there is a greater risk of an increase in disparities in effective and informed access to various prenatal screening techniques. One result of these processes is an increase in the probability and degree of implicit bias in provision of screening tests.

Abbreviations

HAS: Haute Autorité de Santé, French national authority for health.

Declarations

Ethics approval and consent to participate

The surveys were approved by the National Council on Statistical Information (Comité du Label), the French Data Protection Authority (CNIL) and the INSERM ethics committee.

Consent for publication

Not applicable.

Availability of data and materials

The data supporting the conclusions of this article are available in the ADISP website
https://clicktime.symantec.com/3LxK1SeTZznFcNij7TwrWhE6H2?u=www.progedo-adisp.fr%2Fenquetes_sante.php

Competing interest

The authors declare they have no competing interest.

Funding

The national perinatal surveys are supported by the French ministry of health [Direction de la Recherche, des Études de l'Évaluation et des Statistiques (DREES), Direction Générale de la Santé (DGS) and Direction Générale de l'Organisation des Soins (DGOS)], and by Santé publique France.

Author's contributions

OA, EA, BB, PS and BK conceived the study idea. MJSC and NB performed the statistical analyses. OA and EA drafted the first version of the manuscript. All authors discussed the results and revised the manuscript. All authors have read and approved the final version of the manuscript.

Acknowledgments: The authors thank the Maternal and Child Health services in each district, the department heads in each maternity unit and the investigators who allowed data collection, and the women who agreed to be interviewed.

References

1. Roizen NJ, Patterson D. Down ' s syndrome. *The Lancet*. 2003 Apr 12;361(9365):1281-9;
2. Recommandation HAS Dépistage de la trisomie 21 2018. https://www.has-sante.fr/jcms/c_2899277/fr/depistage-de-la-trisomie-21;
3. Légifrance. Arrêté HAS dépistage de la trisomie 21. <https://www.legifrance.gouv.fr/affichTexte.do?cidTexte=JORFTEXT000020814373>
4. Khoshnood B, De Vigan C, Blondel B, Vodovar V, Cadio E, Goffinet F. Long-term trends for socio-economic differences in prenatal diagnosis of Down syndrome: Diffusion of services or persistence of disparities? *BJOG An Int J Obstet Gynaecol*. 2008;

5. Fransen MP, Schoonen MHHJD, Mackenbach JP, Steegers EAP, De Koning HJ, Laudy JAM, et al. Ethnic differences in participation in prenatal screening for Down syndrome: A register-based study. *Prenat Diagn.* 2010;
6. Learman LA, Kuppermann M, Gates E, Nease RF, Gildengorin V, Washington AE. Social and familial context of prenatal genetic testing decisions: Are there racial/ethnic differences? *Am J Med Genet.* 2003;
7. Rowe R, Puddicombe D, Hockley C, Redshaw M. Offer and uptake of prenatal screening for Down syndrome in women from different social and ethnic backgrounds. *Prenat Diagn.* 2008;
8. Grupposo MC, Khoshnood B, Supernant K, Blondel B. [Socioeconomic disparities for maternal serum screening of Down's syndrome: trends between 1998 and 2003 in France]. *J Gynecol Obstet Biol Reprod (Paris).* 2008 May;37(3):246-55.
9. Blondel B, Coulm B, Bonnet C, Goffinet F, Le Ray C; National Coordination Group of the National Perinatal Surveys. Trends in perinatal health in metropolitan France from 1995 to 2016: Results from the French National Perinatal Surveys. *J Gynecol Obstet Hum Reprod.* 2017 Dec;46(10):701-713.
10. Recommandation HAS Suivi de grossesse 2016. Available from: https://www.has-sante.fr/upload/docs/application/pdf/suivi_orientation_femmes_enceintes_synthese.pdf
11. Enquête nationale Périnatale 2016. Available from: http://www.xn-epop-inserm-ebb.fr/wp-content/uploads/2017/10/ENP2016_rapport_complet.pdf
12. Maxwell S, Brameld K, Bower C, Dickinson E, et al. Socio-demographic disparities in the uptake of prenatal screening and diagnosis in Western Australia. *Australian and New Zealand Journal of Obstetrics and Gynaecology.* 2011.
13. Khoshnood B, Pryde P, Blondel B, Lee KS. Socioeconomic and state-level differences in prenatal diagnosis and live birth prevalence of Down's syndrome in the United States. *Rev Epidemiol Sante Publique.* 2003;
14. Fransen MP, Wildschut HIJ, Mackenbach JP, Steegers EAP, Galjaard RJ, Essink-Bot ML. Ethnic and socio-economic differences in uptake of prenatal diagnostic tests for Down's syndrome. *Eur J Obstet Gynecol Reprod Biol.* 2010;
15. Kuppermann M, Learman LA, Gates E, Gregorich SE, Nease RF, Lewis J, et al. Beyond race or ethnicity and socioeconomic status: Predictors of prenatal testing for Down syndrome. *Obstet Gynecol.* 2006;
16. Kuppermann M, Nease RF, Gates E, Learman LA, Blumberg B, Gildengorin V, et al. How do women of diverse backgrounds value prenatal testing outcomes? *Prenat Diagn.* 2004;
17. Dormandy E, Michie S, Hooper R, Marteau TM. Low uptake of prenatal screening for Down syndrome in minority ethnic groups and socially deprived groups: A reflection of women's attitudes or a failure to facilitate informed choices? *Int J Epidemiol.* 2005;
18. Heaman M, Bayrampour H, Kingston D, Blondel B, Gissler M, Roth C, et al. Migrant women's utilization of prenatal care: A systematic review. *Maternal and Child Health Journal.* 2013.
19. Sauvageain P, Zeitlin J, Rico-Berrocal R. Why is perinatal and infant mortality high in the Seine-Saint-Denis district? A consultation with healthcare providers using a Delphi process. *J Gynecol Obstet Biol Reprod (Paris).* 2017 May;38(5):449-55.

Ia Reprod. 2016;

20. Van Ryn M, Burke J. The effect of patient race and socio-economic status on physicians' perceptions of patients. *Soc Sci Med.* 2000;
21. Haider AH, Schneider EB, Sriram N, Dossick DS, Scott VK, Swoboda SM, et al. Unconscious race and social class bias among acute care surgical clinicians and clinical treatment decisions. *JAMA Surg.* 2015;
22. Hausmann LRM, Myaskovsky L, Niyonkuru C, Oyster ML, Switzer GE, Burkitt KH, et al. Examining implicit bias of physicians who care for individuals with spinal cord injury: A pilot study and future directions. In: *Journal of Spinal Cord Medicine.* 2015.
23. Miner J, Biros MH, Trainor A, Hubbard D, Beltram M. Patient and physician perceptions as risk factors for oligoanalgesia: A prospective observational study of the relief of pain in the emergency department. *Acad Emerg Med.* 2006;
24. Todd KH, Samaroo N, Hoffman JR. Ethnicity as a Risk Factor for Inadequate Emergency Department Analgesia. *JAMA J Am Med Assoc.* 1993;
25. Wright JG, Coyte P, Hawker G, Bombardier C, Cooke D, Heck D, et al. Variation in orthopedic surgeons' perceptions of the indications for and outcomes of knee replacement. *CMAJ.* 1995;
26. Youm J, Chan V, Belkora J, Bozic KJ. Impact of socioeconomic factors on informed decision making and treatment choice in patients with hip and knee OA. *J Arthroplasty.* 2015;
27. Shapiro N, Wachtel E V., Bailey SM, Espiritu MM. Implicit Physician Biases in Periviability Counseling. *J Pediatr.* 2018;
28. Hirsh AT, Hollingshead NA, Ashburn-Nardo L, Kroenke K. The interaction of patient race, provider bias, and clinical ambiguity on pain management decisions. *J Pain.* 2015;
29. Sabin JA, Greenwald AG. The influence of implicit bias on treatment recommendations for 4 common pediatric conditions: Pain, urinary tract infection, attention deficit hyperactivity disorder, and asthma. *American Journal of Public Health.* 2012.
30. Chapman EN, Kaatz A, Carnes M. Physicians and implicit bias: How doctors may unwittingly perpetuate health care disparities. *Journal of General Internal Medicine.* 2013.
31. Bangsgaard L, Tabor A. Do pregnant women and their partners make an informed choice about first trimester risk assessment for Down syndrome, and are they satisfied with the choice? *Prenat Diagn.* 2013;
32. Gourounti K, Sandall J. Do pregnant women in Greece make informed choices about antenatal screening for Down's syndrome? A questionnaire survey. *Midwifery.* 2008;
33. Nagle C, Hodges R, Wolfe R, Wallace E. Reporting Down syndrome screening results: Women's understanding of risk. *Prenat Diagn.* 2009;
34. Schoonen M, Wildschut H, Essink-Bot ML, Peters I, Steegers E, De Koning H. The provision of information and informed decision-making on prenatal screening for Down syndrome: A questionnaire- and register-based survey in a non-selected population. *Patient Educ Couns.* 2012;

35. Pruksanusak N, Suwanrath C, Kor-Anantakul O, Prasartwanakit V, Leetanaporn R, Suntharasaj T, et al. A survey of the knowledge and attitudes of pregnant Thai women towards Down syndrome screening. *J Obstet Gynaecol Res*. 2009;
36. Favre R, Moutel G, Duchange N, Vayssi  re C, Kohler M, Bouffet N, et al. What about informed consent in first-trimester ultrasound screening for Down syndrome? *Fetal Diagn Ther*. 2008;

Tables

Table 1: Trends in Down syndrome screening between 2010 and 2016

	2010 %	2016 %	2010 + (n) % 2016	p
Did you have the fetal nuchal translucency measured at your first-trimester ultrasound to learn your risk of Down syndrome?			(24 592)	
No	5.6	5.7	5.6	
Yes	85.1	88.2	86.5	
I don't know	9.3	6.2	7.9	<0.001
Did you have a blood test to learn your risk for Down syndrome (serum markers)?			(24 484)	
No	12.7	9.5	11.3	
Yes	84.7	88.4	86.4	
I don't know	2.6	2.1	2.4	<0.001
If not, why didn't you have this assay?			(2 543)	
not offered	14.2	6.8	11.6	
I refused	44.0	59.1	49.2	
prenatal care was: too late/or none/	21.4	22.9	21.9	
or abroad				
amniocentesis or chorionic villus sampling	9.7	2.2	7.0	
other reason	5.6	5.2	5.5	
I don't know	5.2	3.8	4.7	
Opportunity to make an informed choice†			(24 265)	
Yes	91.7	94.7	93.0	
No	8.4	5.3	7.0	<0.001

† Answer "yes" if the serum marker assay was performed or refused, or if amniocentesis or a chorionic villus sampling was performed first. Answer "no" if the answer to the question "did you have a blood test to learn your risk of Down syndrome (serum markers)?" was "I don't know" or if the assay was not performed for one of the following reasons: not offered, because of late prenatal care, for another reason, or for an unknown reason (see Fig 1).

Table 2: Down syndrome screening according to mother's place of birth (years: 2010+2016)

	Total n=24 644	France n=20 612	Europe n=861	North Africa n=1550	Sub- Saharan Africa n=960	Other countries n=661	p
	%	%	%	%	%	%	
Did you have the fetal nuchal translucency measured at your first-trimester ultrasound to learn your risk of Down syndrome ?							
No	5.6	4.6	6.4	13.6	11.5	9.3	
Yes	86.5	90.3	76.3	62.6	62.4	72.3	
I don't know	7.9	5.2	17.3	23.8	26.1	18.4	<0.001
Did you have a blood test to learn your risk for Down syndrome?							
No	11.3	9.5	14.8	28.0	20.3	11.1	
Yes	86.4	89.5	77.9	62.7	69.0	79.8	
I don't know	2.4	1.1	7.2	9.3	10.7	9.1	<0.001
If not, why didn't you have this blood test?							
not offered	11.6	11.4	12.5	12.5	12.4	8.3	
I refused	49.2	47.9	50.0	58.3	43.2	46.7	
prenatal care was: too late/or none/or abroad	21.9	22.4	15.2	18.3	27.0	28.3	
amniocentesis or chorionic villus sampling first	7.0	8.8	9.8	1.5	0.60	3.3	
other reason	5.5	6.5	2.7	1.8	6.2	1.7	
I don't know	4.7	2.9	9.8	7.6	10.7	11.7	<0.001
Opportunity to make an informed choice†							
Yes	93.0	95.1	87.2	79.9	78.4	86.0	
No	7.0	4.9	12.8	20.1	21.6	14.0	<0.001

†Answer "yes" if the serum marker assay was performed or refused, or if amniocentesis or a chorionic villus sampling was performed first. Answer "no" if the answer to the question "did you have a blood test to learn your risk of Down syndrome (serum markers)?" was "I don't know" or if the assay was not performed for one of the following reasons: not offered, because of late prenatal care, for another reason, or for an unknown reason (see Fig 1).

Table 3: Association between mother's place of birth and Down syndrome screening (years: 2010+2016)

	France	Europe	North Africa	Sub-Saharan Africa	Other countries	p
n=20 612	n=861	n=1550	n=960	n=661		
aOR(95% CI)§	aOR (95% CI)§	aOR (95% CI)§	aOR (95% CI)§	aOR (95% CI)§		
No	1	1.55 (1.16-2.09)	3.94 (3.30-4.70)	2.58 (2.05-3.24)	3.10 (2.32-4.15)	
Yes	1	1	1	1	1	
I don't know	1	3.36 (2.73-4.13)	5.67 (4.88-6.59)	5.02 (4.20-6.00)	5.57 (4.43-6.99)	<0.001
Did you have the fetal nuchal translucency measured at your first-trimester ultrasound to learn your risk of Down syndrome?						
No	1	1.67 (1.36-2.05)	3.57 (3.13-4.08)	1.91 (1.60-2.29)	1.31 (1.01-1.70)	
Yes	1	1	1	1	1	
I don't know	1	5.24 (3.81-7.21)	8.30 (6.49-10.62)	6.65 (5.04-8.78)	9.14 (6.58-12.68)	<0.001
Did you have a blood test to learn your risk for Down syndrome?						
All women	1	2.27 (1.80-2.86)	3.84 (3.28-4.49)	3.27 (2.71-3.94)	3.28 (2.55-4.23)	<0.001
Women who had adequate prenatal care ‡	1	2.42 (1.85-3.15)	3.96 (3.31-4.75)	3.20 (2.55-4.00)	3.75 (2.84-4.96)	<0.001

• Odds ratio adjusted for survey year, parity, age, mother's educational level and maternity ward level with confidence interval

†Answer "yes" if the serum marker assay was performed or refused, or if amniocentesis or a chorionic villus sampling was performed first. Answer "no" if the answer to the question "did you have a blood test to learn your risk of Down syndrome (serum markers)?" was "I don't know" or if the assay was not performed for one of the following reasons: not offered, because of late prenatal care, for another reason, or for an unknown reason (see Fig 1).

#at least 8 prenatal visits and 3 ultrasounds

Figures




Figure 1

Construction of the indicator: opportunity to make an informed choice about Down syndrome screening




Figure 1

Construction of the indicator: opportunity to make an informed choice about Down syndrome screening