Fat mass prevalence and influence on nutritional status in ICU COVID-19 patients

Antonino De Lorenzo

Section of Clinical Nutrition and Nutrigenomics, Department of Biomedicine and Prevention, University of Rome Tor Vergata, Rome, 00133, Italy

Maria Grazia Tarsitano

Department of Experimental Medicine, University of Rome Sapienza, Rome, 00161, Italy

Carmela Falcone

Unit of Radiology, Grande Ospedale Metropolitano Bianchi Melacrino Morelli, Reggio Calabria, 89124, Italy

Laura Di Renzo

Section of Clinical Nutrition and Nutrigenomics, Department of Biomedicine and Prevention, University of Rome Tor Vergata, Rome, 00133, Italy

Lorenzo Romano

School of Specialization in Food Science, University of Rome Tor Vergata, Rome, 00133, Italy

Sebastiano Macheda

Unit of Intensive Care Medicine and Anesthesia, Grande Ospedale Metropolitano Bianchi Melacrino Morelli, Reggio Calabria, 89124

Anna Ferrarelli

Unit of Radiology, Grande Ospedale Metropolitano Bianchi Melacrino Morelli, Reggio Calabria, 89124, Italy

Demetrio Labate

Unit of Intensive Care Medicine and Anesthesia, Grande Ospedale Metropolitano Bianchi Melacrino Morelli, Reggio Calabria, 89124

Marco Tescione

Unit of Intensive Care Medicine and Anesthesia, Grande Ospedale Metropolitano Bianchi Melacrino Morelli, Reggio Calabria, 89124

Federico Bilotta

Department of Anesthesiology and Critical Care, University of Rome Sapienza, Rome, 00161, Italy

Paola Gualtieri (✉ paola.gualtieri@uniroma2.it)

Section of Clinical Nutrition and Nutrigenomics, Department of Biomedicine and Prevention, University of Rome Tor Vergata, Rome, 00133, Italy

Keywords: fat mass, Covid-19, BMI, obesity, ICU, PNI, steatosis

Posted Date: June 8th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-33974/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Fat mass prevalence and influence on nutritional status in ICU COVID-19 patients

Antonino De Lorenzo ^{1§}, Maria Grazia Tarsitano ^{2§}, Carmela Falcone ³, Laura Di Renzo ¹, Lorenzo Romano ⁴, Sebastiano Macheda⁵, Anna Ferrarelli ³, Demetrio Labate ⁵, Marco Tescione ⁵, Federico Bilotta ⁶ and Paola Gualtieri ^{1,*}

¹ Section of Clinical Nutrition and Nutrigenomics, Department of Biomedicine and Prevention, University of Rome Tor Vergata, Rome, 00133, Italy; delorenzo@uniroma2.it (A.D.L.); laura.di.renzo@uniroma2.it (L.D.R.); paola.gualtieri@uniroma2.it (P.G.);

² Department of Experimental Medicine, University of Rome Sapienza, Rome, 00161, Italy; mariagrazia.tarsitano@uniroma1.it (M.G.T.);

³ Unit of Radiology, Grande Ospedale Metropolitano Bianchi Melacrino Morelli, Reggio Calabria, 89124, Italy; milafalcone@gmail.com (C.F.); anferrarelli@yahoo.it (A.F.);

⁴ School of Specialization in Food Science, University of Rome Tor Vergata, Rome, 00133, Italy; lorenzo.romano@alumni.uniroma2.eu (L.R.);

⁵ Unit of Intensive Care Medicine and Anesthesia, Grande Ospedale Metropolitano Bianchi Melacrino Morelli, Reggio Calabria, 89124, Italy; nucciomacheda58@gmail.com (S.M.); labatedemetrio84@gmail.com (D.L.); marcotescione.mt@gmail.com (M.T.);

⁶ Department of Anesthesiology and Critical Care, University of Rome Sapienza, Rome, 00161, Italy; bilotta@tiscali.it (F.B.)

§ equal contribution

* Correspondence to Paola Gualtieri, Via Montpellier, +39-0672596856, paola.gualtieri@uniroma2.it

1 Abstract

2 Background: Obesity and steatosis are associated with COVID-19 severe pneumonia.
3 Elevated levels of pro-inflammatory cytokines and reduced immune response are typical of
4 these patients. In particular, adipose tissue is the organ playing the crucial role. So, it is
5 necessary to evaluate fat mass and not simpler body mass index (BMI), because BMI leaves
6 a portion of the obese population unrecognized. The aim is to evaluate the relationship
7 between FM% and immune-inflammatory response, after 10th days in ICU.

8 Methods: Prospective observational study in single cohort of 22 adult patients, affected by
9 COVID-19 pneumonia and admitted to the Intensive Care Unit (ICU) and classified (10) lean
10 and (12) obese, according to Percentage of Fat Mass (FM%) and age (De Lorenzo
11 classification). Patients were analyzed at admission in ICU and at 10th day.

Results: Obese have steatosis, impaired hepatic function, compromise immune response and higher inflammation. In addition, they have a reduced prognostic nutritional index (PNI), nutritional survival index for ICU patients.

Conclusion: This is the first study evaluating FM% in COVID-19 patient. We underlined obese characteristic with likely poorly prognosis and an important misclassification of obesity. A not negligible number of patients with normal BMI could actually have an excess of adipose tissue and therefore have an unfavorable outcome such as an obese. Is fundamental personalized patients nutrition basing on disease phases.

Keywords: fat mass, Covid-19, BMI, obesity, ICU, PNI, steatosis

Background

Obesity, measured as body mass index (BMI), is reported to associate with increase the risk of developing severe pneumonia in COVID-19 (1). Indeed, the risk correlated to obesity with COVID-19 severity is greater in metabolic associated fatty liver patients (2). At the basis, we found altered mechanism of inflammation and immune response typic of obesity and correlated with alteration in the levels of circulating cytokines (3). In particular, obese patients have higher concentrations of TNF-alpha, MCP-1 and IL-6 which are produced by visceral and subcutaneous adipose tissue and implicated in innate immunity (4,5).

Furthermore, adipose tissue releases high levels of leptin, which creates an unfavorable inflammatory milieu that leads to dysregulation of the immune response (6).

In 2009, during H1N1 pandemia, in obese patients it was characterized changing of differentiation of B cells (7,8), predisposing to a greater risk of contracting influence, but also of being more contagious towards other people (9). However, they have impaired

memory T cell response and vaccination efficacy (10). Specifically, reduced response of virus-specific CD8 + lymphocytes and suboptimal macrophage functionality have been demonstrated, which could explain the low response to the vaccine stimulus (11).

For COVID-19, in Intensive Care Unit (ICU), it was observed that the highest percentage are patients affected by severe obesity, with BMI > 35 kg/m² (12).

The BMI does not reflect necessarily the fat mass (FM). There are evidence that suggest how FM% rather than BMI, predicts inflammatory (TNF-alpha, MCP-1 and IL-6) and immune (leptin) response. These ultimate variables that relates with immune and inflammatory response in FM% (13).

Despite the relationship between FM% and severity of progression inflammatory response in patients admitted to ICU for underlying infective disease, there are no available data on the relevance of FM% in COVID-19 patients treated in ICU.

Moreover, prognostic nutritional index (PNI) can be used to evaluate the nutritional status and survival for ICU patients (14).

Aim of this prospective observational study, in patients admitted to ICU for COVID-19, is to evaluate the relationship between FM% and immune-inflammatory response, after 10th days in ICU.

Also, we want to investigate the metabolic associated fatty liver and PNI.

Subjects and Methods

After IRB approval (Regional Ethic Committee, Section “Area Sud”, 20th April, 2020) and having obtained the signed informed consent by the next of kin, clinical and anthropometric data of patients with COVID-19 pneumonia admitted to the ICU of the Hospital "Bianchi Melacrino Morelli" Reggio Calabria, Italy between 13 March and 6 April 2020, were recorded and analyzed. Standard therapeutic protocol included, for all patients: low molecular weight heparin (LMWH), azithromycin, hydroxychloroquine, lopinavir/ritonavir.

Prospective observational study in single cohort adult patients, affected by COVID-19 pneumonia and admitted to the ICU. Patients with a history of neutropenia, acquired immunodeficiency, who underwent transplants or who received previous immunosuppressive therapies were excluded.

Rather than considering only the BMI, patients recruited in this study were categorized as being “lean” or “obese” according to FM% and age based on criteria presented by De Lorenzo (15).

Computed tomography (CT) (GE Medical SYSTEMS, Gamma Optima, USA) without intravenous contrast was performed within 24 hours of admission to the emergency department.

Hepatic steatosis was evaluated on CT images in 4 liver segments. Independently by 2 qualified operators (LR and CF) (16). If reported difference between the measures were >5%, a third operator was asked to repeat the measure. Waist circumference was measured at the last rib. Where part of the abdomen was outside the field of the image, waist circumference was estimated with a continuous arc. To determine the FM% we used the Siri equation (17). Body density was obtained by the equation of Durnin, using two subcutaneous fat thicknesses of the chest, suprascapular and suprailiac and the correction factors according to age, sex and folds used (18). The subcutaneous fat thickness was measured at CT, given the agreement

between the CT and plicometry method (19). All CTs were performed with patients in the supine position with arms folded and hands positioned under neck. The subscapular fat thickness was measured in cross section starting from the origin of the scapular spine on the posterior medial edge up to the skin. The suprailiac fat thickness was measured in cross section 2 cm from the last rib on the middle axillary line up to the skin. The subcutaneous fat thickness parameters measured on CT were doubled before insertion into the equation of Durnin (18).

Prognostic nutritional index (PNI) was calculated for each patient as a serum albumin (g/dL) $\times 10 +$ total lymphocyte count (mm^{-3}) $\times 0.005$ (20).

The following blood analysis were performed: C-reactive protein (mg/L), glycemia (mg/dL), creatinine (mg/dL), albumin (g/dL), AST (U/L), ALT (U/L), indirect bilirubin (mg/dl), total bilirubin (mg/dl), direct bilirubin (mg/dl), Platelets ($10^3/\mu\text{L}$), white blood cells (WBC) ($10^3/\mu\text{L}$), neutrophils ($10^3/\mu\text{L}$), Lymphocytes ($10^3/\mu\text{L}$), PNI, fibrinogen (mg/dL), D-dimer (ng/mL).

Statistics

Calculation of the sample size was based on a comparison between matched pairs, a power of 80%, a significance level of 5% (two-tails) and the detection of an effect size of 0.6 between the pairs. According to the study setting, necessary calculated sample size is 20 patients and the G*Power software (Version 3.1.9.6, Germany) was used. (21) Since 20% of the sample may not have all the expected parameters, 24 subjects were enrolled.

All statistical analyzes were conducted with SPSS 23 software (version 23.0, IBM, Armonk, NY, USA). The data collected before statistical evaluations were analyzed for the presence of outliers and for non-normally distribution with the Kolmogorov-Smirnov Test. The

categorical variables have been reported in percentage, while the continuous ones as median and interquartile range. Before, the differences between lean and obese subjects were assessed at admission with the Mann Whitney test for independent samples. Subsequently, the differences in lean and obese subjects were assessed between admission and 10th days with the Wilcox test for matched pairs. Cohen's Kappa was used with binary data to measure the agreement between adiposity classification according to the FM% criterion and the BMI. According to Landis and Koch (22), Cohen's Kappa (κ) values could indicate an agreement: poor ($\kappa < 0.00$), light ($0.00 \leq \kappa \leq 0.20$), discrete ($0.21 \leq \kappa \leq 0.40$), moderate ($0.41 \leq \kappa \leq 0.60$), substantial ($0.61 \leq \kappa \leq 0.80$) or near-perfect ($\kappa > 0.80$). Furthermore, the false-positive rate and false-negative rate were calculated for the different classification methods.

Correlation analysis was conducted with Spearman's rho test. Statistical significance was set to a value of $p < 0.05$. All p values shown are two-tailed.

Results

A total of 27 patients were evaluated for this prospective analytical observational study, 5 subjects were excluded from the study because the following reasons: 1 because COVID-19 negative, 1 died before 10th days; 3 had incomplete data. Finally, 22 patients were included in the study (Figure 1). Mean age of enrolled patients was 58 years (range 49-67), 45% were females and 55% males. Patients were divided into two groups according to FM% and age: 12 patients were "obese" and 10 were "lean" (Table 1).

The BMI-based classification, as compared to FM%, presents a discrete and significant Cohen κ -value ($\kappa = 0.405$ $p < 0.000$). According to the BMI, 36% of the patients were categorized differently, in detail, 9% of the patients that presented a BMI within normal values were detected to have a FM% criteria for being considered "obese" according to FM% and age based on criteria presented by De Lorenzo (15) and 27% of the patients classified

PO were “obese” (Table 2) (Figure 2). The FM%-based categorization coincided to BMI-based categorization for the 10 patients considered “lean” according the FM% and for 4 patients considered “obese”.

Figure 1. Flow Chart Prospective analytical observational study

Figure 2. Wrong classification by BMI of patients detected Lean or Obese according to FM%, gender and age. Values are presented as percentage. The white columns are the lean subjects and the gray columns the obese subjects.

138

139 Baseline characteristics of “lean” vs. “obese” patients. No statistical differences were present
 140 for age and spleen attenuation between groups. Liver and LSR attenuation were lower in
 141 “obese” than in “lean” (Table 1) The ALT and AST were higher “obese” than in “lean”. No
 142 statistical difference was found in blood chemistry parameters between the groups.

143 At day 10th, the C-reactive protein, direct bilirubin, fibrinogen concentrations were lower in
 144 were lower than in “lean” (respectively $p=0.005$; $p=0.033$; $p=0.028$) (Figure 3), the
 145 lymphocytes concentration was increased in “lean” ($p=0.038$). Comparison of baseline to
 146 day 10th, concentrations of albumin and PNI decreased in “obese” while remained unchanged
 147 in “lean” patients (Figures 4 and 5). No statistical difference was present in “lean” and
 148 “obese” groups between baseline and 10 days (Table 3).

149 Finally, an inversely proportional correlation was observed between FM% and liver
 150 attenuation (correlation coefficient = -0.702 and $p = 0.002$) (Figure 6).

Figure 4. Albumin concentration comparison between Baseline and 10th day is shown. The white boxes are the lean subjects and the gray boxes the obese subjects. Values are presented

as median with interquartile range. Statistical significance attributed to results with $*p<0.05$,
 $**p<0.005$ and $***p<0.001$.

Figure 5. PNI comparison between Baseline and 10th day is shown. The white boxes are the lean subjects and the gray boxes the obese subjects. Values are presented as median with

169 interquartile range. Statistical significance attributed to results with $*p<0.05$, $**p<0.005$ and
 170 $***p<0.001$.

171

172

173

174 Figure 6. Correlation Between Liver Attenuation and Fat Mass % using Spearman's rho test.

175 Data are presented as Fitline and 95% confidence interval. Correlation significant was

176 attributed as $p<0.05$. CC: Correlation Coefficient.

177

178

Discussion

The distinctive feature of this prospective observational study, in patients admitted to the ICU for COVID-19, is that more patients evaluated according FM% result to be “obese” than using the BMI criteria. In FM%-based “obese” patients, at the beginning of ICU treatment, the metabolic associated fatty liver, PNI and the immune-inflammatory response are severely compromised. Furthermore, at 10th day follow up these patients present a persistent lymphocyte reaction that suggest a protract inflammatory reaction.

In COVID-19 patients it is crucial to find risk factors associated with worse clinical course to allocate appropriate resources. However, population characteristics are fundamental for prognosis. In Italy, COVID-19 mortality is strongly influenced by different comorbidities (23) and 52% of deaths are above 80 years of age, unlike China, for which only 20% are above the same age threshold (23).

In particular, pre-existing pathologies including obesity, cardiovascular co-morbidity, arterial hypertension and type 2 diabetes mellitus are established risk factors (24-26). Obesity also can be associated to insulin-resistance, that alters immune response (27). Obese patients have greater infectivity correlated with exhalation, since they have higher ventilator volumes., due to a lower expansion capacity of the thoracic cavity, which consequently limits the lung expansion (28). This also results in increased aerosol production (28). In particularly, Maier et al. (29) showed that obese patients have a longer viral interaction.

In the case of COVID-19, it has been observed that the infectious charge has an average duration of 20 days, but it can last up to 37 days after the infection (30). It remains to be shown the pathophysiological characteristics of patients who have a contagious duration of up to 37 days.

Hence, the importance of assessing the inflammatory state, through circulating cytokines, has already been highlighted in patients suffering from acute respiratory distress syndrome. Indeed, it was possible to identify two distinct phenotypes, with two different mortality risks. This is fundamental for indicating the patient future prognosis (31). In obese, lipid metabolism is already altered (13,32) and a COVID-19 infection leads to an overexpression of the genes involved with a further increase in the production of pro-inflammatory cytokines and a reduced capability responding to infection.

In our study, “lean” patients-according FM%, showed a significative reduction in C-protein reactive, direct bilirubin and fibrinogen and an important increase of lymphocytes at day 10th of ICU. Increased FM% is associated with a reduced ICU treatment response. Actually, our data showed that “obese” do not show the same improvement, based on biochemical-clinical parameters, respect to “lean” after the first 10th ICU days.

According to presented data, adipose tissue quantity acts on therapeutic goal achievement. Increased adipose tissue leads to a lipid metabolism modification with increased storage of fat in liver and onset of steatosis in “obese”. Consequently, these patients suffer for high production of pro-inflammatory cytokines (33,34) and conduct to unfavorable condition, requiring defined protocols to counter malnutrition resulting (35).

Additionally, these patients are at higher risk for infection also because of COVID-19 use angiotensin converting enzyme 2 (ACE2) receptors to enter the host cell (36). The ACE2 is expressed in different tissues: kidney, lung, heart and adipose tissue (36). COVID-19 infection leads to an upregulation of the genes associated with lipid metabolism, involved in the regulation of inflammation (37).

Thus, obese have a higher expression of ACE2 and are therefore more susceptible to this infection (38). The complex picture is characterized by increased predisposition to infection

226 and reduced ability to respond to it. In addition, these patients already present organ damage
227 that induces worse response to treatments.

228 According to the results, the main aim is a proper nutritional medical therapy, which takes
229 into account the amount of fat, as a risk factor for complications in COVID-19. Therefore,
230 the therapeutic approach must be customized on the body composition. In addition, the loss
231 of body protein content is a negative prognosis factor and it has been a constant observation
232 in ICU.

233 A further aim, in the not-affected, affected and discharged COVID-19 patients, is the saving
234 and recovery of lean body mass, following an appropriate protein prescription.

235 In pre-COVID-19 patients, a personalized and balanced Italian Mediterranean Reference Diet
236 characterized by anti-inflammatory and antioxidant properties (39), should be adopted as
237 obesity preventive and therapeutic tool. The protein intake required is based on lean mass
238 content (2 g/kg of lean mass/day), a parameter that can be directly measured or calculated
239 with prediction equations, accessible to all users (15).

240 In COVID-19 patients, a macronutrient balance calculated according to the clinical condition,
241 a correct calorie intake based on the metabolic condition and all micronutrients must be
242 guaranteed. In detail, respiratory failure requires hyperlipidic nutritional medical therapy, to
243 counter hypercapnia and promoting metabolic flexibility (35,40). The calorie prescription
244 must be adjusted daily, following the catabolic and anabolic phases of hospitalization.
245 Similarly, the protein prescription must be modulated according to the metabolic phase. In
246 the anabolic phase, the protein administered should not be counted in the daily energy
247 expenditure and the protein intake must be 1.3 g/kg of body weight/day (40).

In post-COVID-19 patients, keeping in mind that the fragility deriving from bedrest and inadequate nutrition, due to the ventilatory support, a specific nutritional and motor rehabilitation must be provided (40). For patients with comorbidity, nutrition support to anabolic and recovery stress represent a complex passage. Diet therapy, personalized based on the body composition (35), must be hyperproteic, 2-2.5 g/kg of lean mass/day, complete with all amino acids and enriched with branched amino acids, to promote anabolism. The meals consistency must be progressively personalized according to the subject ability to feed.

It is hoped that from the COVID-19 lesson, the Public Institutions will promote the prevention and treatment of obesity and sarcopenia, through healthy nutrition and a correct lifestyle.

The comorbidities costs and the obstacle in the clinical treatment of an obese patient, in addition to the known health-care costs (41), has been paid with human lives.

Conclusions

Our data originally demonstrate that FM% and not only BMI correlates with the course of COVID-19 patients admitted to ICU. Of note a not negligible number of patients with normal BMI could actually have an excess of adipose tissue and therefore have an unfavorable outcome such as an obese (42). Since it is the actual representation of adipose tissue that is the driver of pro-inflammatory modulation the FM% might represent a better index than BMI. Our data, even if the sample size required by statistical tests is respected, nevertheless presents a limited number of patients. The fat mass calculated by CT was necessary since other methods such as bioimpedance and anthroplicometry are difficult to apply in ICU. Future studies are required to determine the relevance of FM% in predicting the clinical outcomes of COVID-19 patients and might be a useful approach also for out of hospital surveillance.

Abbreviation List: ACE2, angiotensin converting enzyme 2; BMI, body mass index; CT, computed tomography; FM, fat mass; ICU, Intensive Care Unit; PNI, prognostic nutritional index.

Declarations

Ethics approval and consent to participate: This study was approved by Regional Ethic Committee, Section “Area Sud”, 20th April, 2020. Signed informed consent by the next of kin was obtained for all patients.

Consent for publication: not applicable

Availability of data and material: The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interests.

Funding: This research received no external funding

Author Contributions: Conceptualization, A.D.L. and M.G.T.; Formal analysis, L.R.; Resources, C.F., S.M., A.F., D.L., M.T.; writing—original draft preparation, M.G.T., L.R., F.B. and P.G.; writing—review and editing, A.D.L., M.G.T., C.F., L.D.R., L.R., S.M., A.F., D.L., M.T., F.B., P.G.; supervision and primary responsibility for final content, P.G. All authors have read and agreed to the published version of the manuscript.

Acknowledgments: The authors are in debt with Pierpaolo Correale for the English revision

References

1. Petrilli CM, Jones SA, Yang J, Rajagopalan H, O'Donnell L, Chernyak Y, Tobin KA, Cerfolio RJ, Francois F, Horwitz LI. Factors associated with hospitalization and critical illness among 4,103 patients with COVID-19 disease in New York City. Preprint at medRxiv 2020, <https://doi.org/10.1101/2020.04.08.20057794>.
2. Zheng KI, Gao F, Wang XB, Sun QF, Pan KH, Wang TY, Ma HL, Liu WY, George J, Zheng MH. Obesity as a risk factor for greater severity of COVID-19 in patients with metabolic associated fatty liver disease. *Metabolism* 2020; 154244. doi: 10.1016/j.metabol.2020.154244.
3. De Lorenzo A, Gratteri S, Gualtieri P, Cammarano A, Bertucci P, Di Renzo L. Why primary obesity is a disease? *J Transl Med* 2019;17(1):169. doi: 10.1186/s12967-019-1919-y.
4. Richard C, Wadowski M, Goruk S, Cameron L, Sharma AM, Field CJ. Individuals with obesity and type 2 diabetes have additional immune dysfunction compared with obese individuals who are metabolically healthy. *BMJ Open Diabetes Res Care* 2017;5(1):e000379. doi: 10.1136/bmjdr-2016-000379.
5. Dietz W, Santos-Burgoa C. Obesity and its Implications for COVID-19 Mortality. *Obesity* (Silver Spring) 2020. doi: 10.1002/oby.22818.
6. Ouchi N, Parker JL, Lugus JJ, Walsh K. Adipokines in inflammation and metabolic disease. *Nat Rev Immunol* 2011;11(2):85-97. doi: 10.1038/nri2921.
7. Zhang AJ, To KK, Li C, Lau CC, Poon VK, Chan CC, Zheng BJ, Hung IF, Lam KS, Xu A, et al. Leptin mediates the pathogenesis of severe 2009 pandemic influenza A(H1N1) infection associated with cytokine dysregulation in mice with diet-induced obesity. *J Infect Dis* 2013;207(8):1270-80. doi:10.1093/infdis/jit031.
8. Milner JJ, Beck MA. The impact of obesity on the immune response to infection. *Proc Nutr Soc* 2012;71(2):298-306. doi: 10.1017/S0029665112000158.

9. Luzi L, Radaelli MG. Influenza and obesity: its odd relationship and the lessons for COVID-19 pandemic. *Acta Diabetol* 2020. doi: 10.1007/s00592-020-01522-8.
10. Park S, Jeon JH, Min BK, Ha CM, Thoudam T, Park BY, Lee IK. Role of the Pyruvate Dehydrogenase Complex in Metabolic Remodeling: Differential Pyruvate Dehydrogenase Complex Functions in Metabolism. *Diabetes Metab J* 2018;42(4):270-281. doi: 10.4093/dmj.2018.0101.
11. Honce R, Schultz-Cherry S. Impact of Obesity on Influenza A Virus Pathogenesis, Immune Response, and Evolution. *Front Immunol* 2019;10:1071. doi: 10.3389/fimmu.2019.01071.
12. Simonnet A, Chetboun M, Poissy J, Raverdy V, Noulette J, Duhamel A, Labreuche J, Mathieu D, Pattou F, Jourdain M, et al. High prevalence of obesity in severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) requiring invasive mechanical ventilation. *Obesity (Silver Spring)* 2020. doi: 10.1002/oby.22831.
13. De Lorenzo A, Soldati L, Sarlo F, Calvani M, Di Lorenzo N, Di Renzo L. New obesity classification criteria as a tool for bariatric surgery indication. *World J Gastroenterol* 2016;22(2):681-703. doi: 10.3748/wjg.v22.i2.681.
14. Kos M, Titiz H, Onec B, Soysal T, Kutlucan A, Sahiner Emen S, Kutlucan L. Association of “Controlling Nutritional Status Index” and “Prognostic Nutritional Index” with intensive care unit survival in elderly patients. *European Geriatric Medicine* 2020;7(1):13-17. doi: <https://doi.org/10.1016/j.eurger.2015.07.007>.
15. De Lorenzo A, Siclari M, Gratteri S, Romano L, Gualtieri P, Marchetti M, Merra G, Colica C. Developing and cross-validation of new equations to estimate fat mass in Italian population. *Eur Rev Med Pharmacol Sci* 2019;23(6):2513-2524. doi: 10.26355/eurrev_201903_17399.

16. Wells MM, Li Z, Addeman B, McKenzie CA, Mujoomdar A, Beaton M, Bird J. Computed Tomography Measurement of Hepatic Steatosis: Prevalence of Hepatic Steatosis in a Canadian Population. *Can J Gastroenterol Hepatol* 2016;2016:4930987. doi: 10.1155/2016/4930987.
17. Siri WE. Body composition from fluid spaces and density: analysis of methods. 1961. *Nutrition* 1993;9(5):480-91.
18. Durnin JV, Womersley J. Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *Br J Nutr.* 1974 Jul;32(1):77-97.
19. Orphanidou C, McCargar L, Birmingham CL, Mathieson J, Goldner E. Accuracy of subcutaneous fat measurement: comparison of skinfold calipers, ultrasound, and computed tomography. *J Am Diet Assoc* 1994;94(8):855-8.
20. Lee JY, Kim HI, Kim YN, Hong JH, Alshomimi S, An JY, Cheong JH, Hyung WJ, Noh SH, Kim CB. Clinical Significance of the Prognostic Nutritional Index for Predicting Short- and Long-Term Surgical Outcomes After Gastrectomy: A Retrospective Analysis of 7781 Gastric Cancer Patients. *Medicine (Baltimore)* 2016;95(18):e3539. doi: 10.1097/MD.0000000000003539.
21. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 2007;39(2):175-91.
22. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics* 1977;33(1):159-74.
23. Onder G, Rezza G, Brusaferro S. Case-Fatality Rate and Characteristics of Patients Dying in Relation to COVID-19 in Italy. *JAMA* 2020. doi:10.1001/jama.2020.4683.

24. Peng YD, Meng K, Guan HQ, Leng L, Zhu RR, Wang BY, He MA, Cheng LX, Huang K, Zeng QT. [Clinical characteristics and outcomes of 112 cardiovascular disease patients infected by 2019-nCoV]. *Zhonghua Xin Xue Guan Bing Za Zhi* 2020;48(0):E004. doi: 10.3760/cma.j.cn112148-20200220-00105.
25. Fadini GP, Morieri ML, Longato E, Avogaro A. Prevalence and impact of diabetes among people infected with SARS-CoV-2. *J Endocrinol Invest* 2020. doi: 10.1007/s40618-020-01236-2.
26. Istituto Superiore di Sanità. 2020, <https://www.epicentro.iss.it/coronavirus/sars-cov-2-decessi-italia>. (accessed 13 April 2020).
27. Reidy PT, Yonemura NM, Madsen JH, McKenzie AI, Mahmassani ZS, Rondina MT, Lin YK, Kaput K, Drummond MJ. An accumulation of muscle macrophages is accompanied by altered insulin sensitivity after reduced activity and recovery. *Acta Physiol (Oxf)* 2019;226(2):e13251. doi: 10.1111/apha.13251.
28. Yan J, Grantham M, Pantelic J, Bueno de Mesquita PJ, Albert B, Liu F, Ehrman S, Milton DK; EMIT Consortium. Infectious virus in exhaled breath of symptomatic seasonal influenza cases from a college community. *Proc Natl Acad Sci U S A* 2018;115(5):1081-1086. doi: 10.1073/pnas.1716561115.
29. Maier HE, Lopez R, Sanchez N, Ng S, Gresh L, Ojeda S, Burger-Calderon R, Kuan G, Harris E, Balmaseda A, et al. Obesity Increases the Duration of Influenza A Virus Shedding in Adults. *J Infect Dis.* 2018 Sep 22;218(9):1378-1382. doi: 10.1093/infdis/jiy370.
30. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, Xiang J, Wang Y, Song B, Gu X, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395(10229):1054-1062. doi: 10.1016/S0140-

- 6736(20)30566-3. Erratum in: Lancet. 2020;395(10229):1038. Lancet 2020;395(10229):1038.
31. Bos LD, Schouten LR, van Vught LA, Wiewel MA, Ong DSY, Cremer O, Artigas A, Martin-Loeches I, Hoogendijk AJ, van der Poll T, et al. Identification and validation of distinct biological phenotypes in patients with acute respiratory distress syndrome by cluster analysis. *Thorax* 2017;72(10):876-883. doi: 10.1136/thoraxjnl-2016-209719.
 32. Merra G, Gualtieri P, Cioccoloni G, Falco S, Bigioni G, Tarsitano MG, Capacci A, Piccioni A, Costacurta M, Franceschi F, et al. FTO rs9939609 influence on adipose tissue localization in the Italian population. *Eur Rev Med Pharmacol Sci* 2020;24(6):3223-3235. doi: 10.26355/eurrev_202003_20689.
 33. Marchisello S, Di Pino A, Scicali R, Urbano F, Piro S, Purrello F, Rabuazzo AM. Pathophysiological, Molecular and Therapeutic Issues of Nonalcoholic Fatty Liver Disease: An Overview. *Int J Mol Sci* 2019;20(8). pii: E1948. doi: 10.3390/ijms20081948.
 34. Kwaifa IK, Bahari H, Yong YK, Noor SM. Endothelial Dysfunction in Obesity-Induced Inflammation: Molecular Mechanisms and Clinical Implications. *Biomolecules* 2020;10(2). pii: E291. doi: 10.3390/biom10020291.
 35. Romano L, Bilotta F, Dauri M, Macheda S, Pujia A, De Santis GL, Tarsitano MG, Merra G, Di Renzo L, Esposito E, De Lorenzo A. Short Report - Medical nutrition therapy for critically ill patients with COVID-19. *Eur Rev Med Pharmacol Sci* 2020;24(7):4035-4039. doi: 10.26355/eurrev_202004_20874.
 36. Li G, He X, Zhang L, Ran Q, Wang J, Xiong A, Wu D, Chen F, Sun J, Chang C. Assessing ACE2 expression patterns in lung tissues in the pathogenesis of COVID-19. *J Autoimmun* 2020:102463. doi: 10.1016/j.jaut.2020.102463.

37. Zou X, Chen K, Zou J, Han P, Hao J, Han Z. Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019-nCoV infection. *Front Med* 2020. doi: 10.1007/s11684-020-0754-0.
38. Kassir R. Risk of COVID-19 for patients with obesity. *Obes Rev* 2020;21(6):e13034. doi: 10.1111/obr.13034.
39. Di Renzo L, Gualtieri P, Romano L, Marrone G, Noce A, Pujia A, Perrone MA, Aiello V, Colica C, De Lorenzo A. Role of Personalized Nutrition in Chronic-Degenerative Diseases. *Nutrients* 2019;11(8). pii: E1707. doi: 10.3390/nu11081707.
40. Singer P, Blaser AR, Berger MM, Alhazzani W, Calder PC, Casaer MP, Hiesmayr M, Mayer K, Montejo JC, Pichard C, et al. ESPEN guideline on clinical nutrition in the intensive care unit. *Clin Nutr* 2019;38(1):48-79. doi: 10.1016/j.clnu.2018.08.037.
41. De Lorenzo A, Esposito E. Editorial - Epidemiological transition, crisis of the Italian health system: ethical and logical economic choices. *Eur Rev Med Pharmacol Sci* 2020; 24:4616-4622. doi: 10.26355/eurev_202004_21049.
42. De Lorenzo A, Bianchi A, Maroni P, Iannarelli A, Di Daniele N, Iacopino L, Di Renzo L. Adiposity rather than BMI determines metabolic risk. *Int J Cardiol* 2013;166(1):111-7. doi: 10.1016/j.ijcard.2011.10.006.

Table 1. Descriptive and Compared between group at baseline

	Overall (n 22) Median (IQR)	Lean (n 10) Median (IQR)	Obese (n 12) Median (IQR)	<i>p</i>
Anthropometrics and Body Parameters				
Age (years)	58.50 (49.00; 67.75)	57.00 (47.00; 68.00)	65.00 (52.50; 67.50)	0.49
Subscapular Thickness (mm)	17.51 (11.83; 21.39)	11.30 (9.74; 16.30)	21.34 (17.94; 26.64)	0.001
Suprailiac Thickness (mm)	21.11 (12.00; 25.51)	11.60 (6.70; 16.30)	25.00 (21.23; 28.05)	0.004
Sum Thickness (mm)	38.96 (23.91; 46.32)	23.10 (18.00; 32.80)	46.03 (41.21; 49.21)	0.001
Fat Mass (%)	37.81 (27.71; 46.52)	27.57 (22.49; 33.18)	45.68 (40.26; 47.59)	0.001
Waist Circumference (cm)	99.49 (96.23; 113.34)	97.11 (94.22; 99.50)	101.25 (99.10; 116.90)	0.048
Liver Attenuation (HU)	47.94 (43.04; 51.34)	51.46 (49.94; 54.14)	44.37 (37.13; 47.94)	0.005
Spleen Attenuation (HU)	51.23 (44.50; 54.14)	54.30 (49.45; 57.26)	51.70 (48.10; 55.89)	0.791
LSR Attenuation	0.92 (0.82; 1.00)	0.96 (0.82; 1.06)	0.82 (0.75; 0.88)	0.017
Blood chemistry Parameters				
C-reactive Protein (mg/L)	74.30 (55.20; 113.00)	88.60 (45.18; 112.00)	89.20 (60.5; 107.00)	0.728
Glycemia (mg/dL)	111.00 (97.00; 138.00)	111.00 (96.00; 134.00)	113.00 (96.93; 146.25)	0.855
Creatinine (mg/dL)	0.80 (0.60; 1.09)	0.94 (0.74; 1.33)	0.70 (0.60; 0.90)	0.173
Albumin (g/dL)	3.30 (3.00; 4.05)	3.50 (2.78; 4.03)	3.30 (3.10; 4.30)	0.830
AST (U/L)	39.00 (31.00; 56.50)	33.00 (28.00; 39.25)	55.00 (36.00; 81.00)	0.027
ALT (U/L)	29.00 (16.50; 60.50)	18.00 (12.75; 36.00)	41.00 (21.00; 96.00)	0.034
Indirect bilirubin (mg/dl)	0.71 (0.44; 1.15)	0.80 (0.50; 1.08)	0.58 (0.26; 1.61)	0.431
Total bilirubin (mg/dl)	1.00 (0.77; 1.40)	1.00 (0.90; 1.40)	0.92 (0.52; 2.40)	0.431
Direct bilirubin (mg/dl)	0.36 (0.26; 0.41)	0.40 (0.32; 0.41)	0.31 (0.22; 0.94)	0.352
Platelets (10 ³ /μL)	187.00 (132.75; 301.25)	202.00 (141.00; 363.00)	172.00 (120.00; 285.00)	0.634
WBC (10 ³ /μL)	6.65 (4.67; 11.33)	5.17 (3.79; 8.73)	6.99 (4.83; 13.07)	0.223
Neutrophils (10 ³ /μL)	4.04 (3.16; 7.08)	3.78 (3.04; 7.24)	4.33 (3.47; 8.61)	0.491
Lymphocytes (10 ³ /μL)	0.85 (0.55; 1.21)	0.89 (0.39; 1.53)	0.81 (0.65; 1.00)	0.711
PNI	36.01 (30.00; 44.51)	35.00 (27.75; 40.26)	36.00 (31.00; 41.0)	0.475
Fibrinogen (mg/dL)	527.00 (414.00; 680.01)	592.00 (390.50; 677.50)	521.00 (390.00; 738.50)	0.953
D-Dimer (ng/mL)	387.00 (224.00; 852.50)	336.00 (164.50; 1974.00)	438.00 (255.00; 869.00)	0.549

Differences among groups at baseline. All parameters are presented as median (interquartile range) and were compared by Mann Whitney test. Statistical significance was attributed as $p < 0.05$. IQR: Interquartile Range; LSR: Liver Spleen Ratio; AST: Aspartate Aminotransferase; ALT: Alanine Aminotransferase; WBC: White Blood Cell; PNI: Prognostic Nutritional Index.

Table 2. Contingency table, κ coefficient of BMI for indicating the misclassification of obesity in population sample.

		FM% Classification		
		Lean	Obese	Total
BMI Classification	NW (18.50-24.99)	45.40% (10)	9.10% (2)	54.50% (12)
	PO (25.00-29.99)	0.00% (0)	27.30% (6)	27.30% (6)
	OB (≥ 30.00)	0.00% (0)	18.20% (4)	18.20% (4)
	Total	45.40% (10)	54.60% (12)	100.00% (22)
$\kappa^* = 0.405$				

FM: Fat Mass; BMI: Body Mass Index; NW: Normal Weight; PO: Pre-Obese; OB: Obese. All values are presented as percentage (numbers) and κ have * p -values were < 0.001 .

Table 3. Difference between baseline and 10th days for each group

	Lean (n 10)			Obese (n 12)		
	Baseline	10 th Days	<i>p</i>	Baseline	10 th Days	<i>p</i>
	Median (IQR)	Median (IQR)		Median (IQR)	Median (IQR)	
Blood Chemistry Parameters						
C-reactive Protein (mg/dL)	88.60 (45.18; 112.00)	17.65 (8.29; 30.20)	0.005	89.20 (60.5; 107.00)	54.95 (27.73; 103.35)	0.465
Glycemia (mg/dL)	111.00 (96.00; 134.00)	103.50 (92.50; 127.25)	0.109	113.00 (96.93; 146.25)	108.00 (87.00; 133.00)	0.345
Creatinine (mg/dL)	0.94 (0.74; 1.33)	1.00 (0.80; 1.89)	0.180	0.70 (0.60; 0.90)	0.71 (0.60; 0.96)	0.206
Albumin (g/dL)	3.50 (2.78; 4.03)	3.15 (2.53; 3.48)	0.109	3.30 (3.10; 4.30)	3.00 (2.80; 3.20)	0.018
AST (U/L)	33.00 (28.00; 39.25)	26.00 (18.50; 34.25)	0.109	55.00 (36.00; 81.00)	41.00 (21.00; 78.00)	0.176
ALT (U/L)	18.00 (12.75; 36.00)	37.00 (15.75; 57.50)	0.285	41.00 (21.00; 96.00)	51.00 (29.00; 93.01)	0.866
Indirect bilirubin (mg/dl)	0.80 (0.50; 1.08)	0.46 (0.31; 0.60)	0.176	0.58 (0.26; 1.61)	0.63 (0.36; 1.87)	0.285
Total bilirubin (mg/dl)	1.00 (0.90; 1.40)	0.57 (0.50; 0.80)	0.091	0.92 (0.52; 2.40)	1.15 (0.70; 2.61)	0.285
Direct bilirubin (mg/dl)	0.40 (0.32; 0.41)	0.18 (0.11; 0.23)	0.033	0.31 (0.22; 0.94)	0.55 (0.25; 0.82)	1.000
Platelets (10 ³ /μL)	202.00 (141.00; 363.00)	216.00 (185.00; 378.00)	0.612	172.00 (120.00; 285.00)	263.00 (180.00; 333.50)	0.139
WBC (10 ³ /μL)	5.17 (3.79; 8.73)	6.22 (3.49; 8.89)	0.866	6.99 (4.83; 13.07)	4.55 (3.62; 10.05)	0.173
Neutrophils (10 ³ /μL)	3.78 (3.04; 7.24)	4.24 (2.70; 6.72)	0.866	4.33 (3.47; 8.61)	2.67 (1.72; 5.62)	0.214
Lymphocytes (10 ³ /μL)	0.89 (0.39; 1.53)	1.41 (0.40; 2.20)	0.038	0.81 (0.65; 1.00)	1.01 (0.50; 1.15)	0.176
PNI	35.00 (27.75; 40.26)	32.51 (25.25; 34.76)	0.109	36.00 (31.00; 41.0)	30.01 (28.00; 32.01)	0.022
Fibrinogen (mg/dL)	592.00 (390.50; 677.50)	265.00 (210.00; 358.00)	0.028	521.00 (390.00; 738.50)	499.00 (208.00; 634.00)	0.593
D-Dimer (ng/mL)	336.00 (164.50; 1974.00)	367.01 (250.00; 4380.50)	0.144	438.00 (255.00; 869.00)	435.00 (304.75; 521.50)	0.161

Differences between baseline and 10th days for each group. All parameters are presented as median (interquartile range) and were compared by Wilcoxon test. Statistical significance was attributed as $p < 0.05$. IQR: Interquartile Range; AST: Aspartate Aminotransferase; ALT: Alanine Aminotransferase; WBC: White Blood Cell; PNI: Prognostic Nutritional Index.

Figures

Figure 1

Flow Chart Prospective analytical observational study

Figure 2

Wrong classification by BMI of patients detected Lean or Obese according to FM%, gender and age. Values are presented as percentage. The white columns are the lean subjects and the gray columns the obese subjects.

Figure 3

C-reactive protein concentration comparison between Baseline and 10th day is shown. The white boxes are the lean subjects and the gray boxes the obese subjects. Values are presented as median with interquartile range. Statistical significance attributed to results with * $p < 0.05$, ** $p < 0.005$ and *** $p < 0.001$.

Figure 4

Albumin concentration comparison between Baseline and 10th day is shown. The white boxes are the lean subjects and the gray boxes the obese subjects. Values are presented as median with interquartile range. Statistical significance attributed to results with * $p < 0.05$, ** $p < 0.005$ and *** $p < 0.001$.

Figure 5

PNI comparison between Baseline and 10th day is shown. The white boxes are the lean subjects and the gray boxes the obese subjects. Values are presented as median with interquartile range. Statistical significance attributed to results with * $p < 0.05$, ** $p < 0.005$ and *** $p < 0.001$.

Figure 6

Correlation Between Liver Attenuation and Fat Mass % using Spearman's rho test. Data are presented as Fitline and 95% confidence interval. Correlation significant was attributed as $p < 0.05$. CC: Correlation Coefficient.