

An Outbreak of Gastroenteritis Associated with a Novel GII.8 Sapovirus Variant-Transmitted by Vomit in Shenzhen, China, 2019

Yuxiao Yan

Lanzhou University

Yuan Li

Shenzhen Baoan Center for Disease Control and Prevention

Shi Wen

Yingkou center for Disease Control and Prevention

Xiangyu Kong

Centers for Disease Control and Prevention

Huiying LI

Centers for Disease Control and Prevention

Qing Zhang

Centers for Disease Control and Prevention

Lili Pang

Centers for Disease Control and Prevention

Li Jiang

The First Hospital of Lanzhou University

Junying Liu

Central Hospital of Zhoukou

Miao Jin (✉ jinmiao37@126.com)

Centers for Disease Control and Prevention <https://orcid.org/0000-0002-8442-5051>

Yuning Li

The First Hospital of Lanzhou University

Zhaojun Duan

Centers for Disease Control and Prevention

Research article

Keywords: Sapoviruses, Outbreak, Vomit, Genome, Phylogenetic analysis

Posted Date: September 22nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-33996/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on December 1st, 2020. See the published version at <https://doi.org/10.1186/s12879-020-05643-x>.

Abstract

Human Sapoviruses (SaVs) has been reported as one of the causative agents of acute gastroenteritis (AGE) worldwide. We investigated an outbreak of SaVs that affected 482 students of a primary school during spring activities from 24th February to 11th March 2019 in Shenzhen city, China. The single factor analysis showed that the students which saw others vomiting in classroom or playground, saw vomitus less than 1.5 meters are susceptible ($P < 0.05$). Seven of eleven fecal samples from patients were positive for GII.8 SaV genotype. In this study, we obtained the genome sequence of a SaV GII.8 strain SZ08 comprehensively analyzed the genetic diversity. The phylogenetic analysis showed that the GII.8 strain SZ08 formed an independent branch and became a novel variant of GII.8 genotype.

Strain SZ08 introduced 11 specific amino acid mutations compared with cluster A-D in full-length VP1. Our study confirmed that constant surveillance will permit the identification of the SaVs genetic diversity as well as an understanding of the evolution of SaVs.

1 Introduction

There has been increasing concern about Sapoviruses (SaVs), a member of the family *Caliciviridae*, as a causative agent of gastroenteritis in humans in both sporadic cases and outbreaks worldwide, especially among infants and young children^[1-5]. SaVs are recognized as the second most commonly etiological virus behind Norovirus in children with acute diarrhea after the successful deployment of the Rotavirus vaccine^[6]. SaVs-associated outbreaks have occurred in semi-closed institutions such as kindergartens, schools, hospitals and nursing homes for the elderly, and happen through the fecal-oral route, the primary route of transmission. SaVs transmit via exposure to SaVs-positive aerosols from feces, vomitus or the consumption of SaVs-contaminated food and water^[7,8]. SaVs-associated acute gastroenteritis (AGE) has increased and has been accepted as a major public health problem worldwide, especially in developing countries^[6,9,10].

The genome of SaVs consists of a single-stranded positive-sense RNA genome of approximately 7.1-7.5kb in size and has a 3'-end poly (A) tail^[7]. The SaVs genome contains either 2 or 3 open reading frames (ORFs). ORF1 encodes nonstructural proteins and a major capsid protein (VP1). ORF2 encodes a minor structural protein (VP2). ORF3 predicted for several SaVs strains encodes proteins with unknown function^[2,7,9].

Based on complete capsid gene (VP1) sequences, sapoviruses are classified into at least 15 genogroups (GI-GXV), four of which (GI, GII, GIV and GV) are known to infect humans. The human sapoviruses in genogroups GI, GII, GIV and GV are currently subdivided into 7 GI (GI.1-GI.7), 8 GII (GII.1-GII.8) and one GII.NA, 1 GIV (GIV.1) and 2 GV (GV.1 and GV.2) genotypes^[11], and one additional genotype of GII.NA that has been reported recently^[12].

Recently, a novel genotype of SaVs, GII.8 strains were reported in some countries and one GII.8 SaV strain (GZ2014-I231) was detected in 2014 in Guangzhou city, Guangdong province of China^[13]. In this study, an AGE outbreak caused by a novel GII.8 SaV variant occurred in primary school in the other city, Shenzhen city, Guangdong province of China in 2019. The genome of the GII.8 strain was analyzed and the epidemiology of this outbreak was described. The emerging of novel GII.8 SaV variant made a warning for SaVs outbreaks control and prevention.

2 Materials And Methods

2.1 Ethics statement

This work was conducted in agreement with the research Ethics committee and the Institutional Review Board at the Baoan district Center for Disease Control and Prevention for human subject projection. We informed each potential subject of the details of study and written informed consent was obtained from the parents or legal guardians of all subjects.

2.2 Epidemiological investigation

Cases were defined as at least three bouts of unformed (loose and watery) stool and/or vomiting episodes in a 24-hour period during the outbreaks. A standardized questionnaire was prepared to collect demographic data (gender and age), and illness onset data (symptoms and duration of symptom).

2.3 Nucleic acid extraction

A 0.1-g fecal specimens was diluted with 1.0ml phosphate-buffered saline (pH 7.2) to a 10% suspension. The supernatants were collected and the viral genomes were extracted by using the QIAamp Viral RNA Mini Kit (Qiagen, Hilden, Germany) and the QIAamp DNA Mini Kit (Qiagen), according to the manufacturer's instructions.

2.4 Bacteria isolation

The stool samples were analyzed for major bacterial pathogens (*Escherichia coli*, *Salmonella*, *Shigella*, *Campylobacter* and *Yersinia enterocolitica*) with MacConkey agar (MAC, Oxoid Ltd, Basingstoke, UK), *Salmonella Shigella* agar (Oxoid Ltd), *Campylobacter*-selective agar (Oxoid Ltd) and *Yersinia* Selective agar (Schiemann's CIN agar, Oxoid Ltd), respectively.

2.5 Virus detection

TaqMan real-time reverse-transcription polymerase chain reaction (RT-PCR) have been employed to test all viral nucleic acid for Norovirus and SaVs with primers and probes as described previously^[14,15]. By using enzyme immunoassay Kits and RT-PCR, Rotavirus A was identified^[16]; and Astrovirus was identified using RT-PCR^[17]; PCR was performed to detect Adenovirus^[18]. Conventional RT-PCR was performed using primer sets including P289 and P290 (in polymerase region) and SLV5317 and SLV5749 (in capsid region) to further characterized all positive samples for SaVs as described previously^[19]. The PCR cycling conditions were as follows: 42°C for 30 minutes, 95°C for 15 minutes followed by 40 cycles of 95°C for 1 minute, 50°C for 1 minute, and 72°C for 1 minute, and a final 10-minute elongation at 72°C. The temperature was held at 4°C until use.

2.6 Nearly complete genome amplification

The nearly complete genome of GII.8 strains were amplified by using fourteen primer sets (Table 1) which were designed based on the whole genome of the reference strains MG674584 and MF462287. The primer set of SLV5317 and VN₃T₂₀^[19] were performed to amplify the complete capsid and ORF2 sequences.

2.7 Genotype and sequence analysis

These nucleotide sequences were first verified using Geneious 10.1.3 (www.geneious.com). Phylogenetic analysis of these sequences was performed in the MEGA6.0 software package (<http://www.megasoftware.net/>). Sequence Alignment was performed using Clustal W. Best substitution models for the dataset were chosen based on the

lowest Bayesian Information Criterion score. Information about amino acid substitutions during the viral evolutionary process were analyzed by comparison within and between different clusters. Phylogenetic trees with bootstrap analysis from 1,000 replicates were generated by using the neighbor-joining method.

2.8 Data analysis

Statistical analysis was performed with the IBM SPSS Statistics software (Version 26.0 International Business Machines Corporation, Armonk, New York, United State) was required for statistical analysis of the collected data. Odds ratio (OR) and 95% confidence interval (CI) of categorical variables were calculated using two tailed, and the comparison of categorical variables among different groups are conducted with Chi-square or Fisher's exact test. Quantitative variable was described as mean, median, standard deviation or inter-quartile range (IQR), quantitative variable was compared by rank-sum test, analysis of variance or t test, significant difference was considered as the level of $P < 0.05$ with two-tailed test.

2.9 Nucleotide sequences and accession numbers

The SaV sequences determined in this study were deposited in GenBank under the accession numbers MT561022 for the nearly full genome sequences.

3 Results

3.1 Epidemiological investigation

This outbreak of AGE occurred in a primary school in Shenzhen city in China. It started on Feb 24th and lasted until March 11th 2019. It included 482 cases among the 1850 students and 95 teachers of the school. The school has 34 classes divided into 6 different grades. There were two buildings in which students of grade 3 and 4 were in Building A and students of grade 5-6 were in Building B. The students from grade 4-6 were organized to participate in military training activities during February 26-27 while the students from grade 1-3 were organized to participate in spring outing on February 27 after the school opened on February 25. The age of 482 cases ranged from 9 to 14 years. The first case from class 4 grade five (C4G5) emerged on February 24, 2019 with the occurrence of vomiting at the classroom and the bus on the way to the activity, respectively. The students of grade 5 got sick first and then grade 4, 6 and 3. There was no case from grade 1 and 2. Peak of cases occurred on March 1 on which local district CDC was invited to investigate this outbreak and the second case peak occurred on March 4 on which classes of G4-6 were suspended and were resumed on March 8. Classes of G3 were suspended on March 8 and resumed on March 11 (Fig 1). The characteristic symptoms within the cohort were diarrhea (67%, 321/482), vomiting (45%, 218/482), dizziness (30%, 144/482), stomach pain (22%, 106/482) and nausea (18%, 86/482). The results showed that the attack rate of each grade was 36% in Grade 5, 23% in Grade 6, 21% in grade 4 and 6% in grade 3, respectively ($P < 0.05$). The attack rate of each floor of Building A was 3% for the 4th floor, 20% for the 5th floor ($\chi^2 = 45.09$, $P < 0.01$), and the attack rate of each floor of Building B was 41% for the 3rd floor, 38% for the 4th floor, and 20% for the 2th floor ($\chi^2 = 16.44$, $P < 0.05$). The difference of incidence among different floors was statistically significant (Fig 2A, Fig 2B).

During the epidemic, 218 students vomited. Vomiting occurred in plastic bags (13%), floor (10%) and window (1%) while riding the bus and vomiting occurred in the bathroom (3%) and dormitory (2%) during activity. Vomiting occurred in Classrooms (9%), playgrounds (3%), hallway (3%) after returning to school (Table 2).

In the group with common exposure to vomit, the single factor analysis showed that among the students that saw other peers vomiting in the classroom or playground, being at a distance of 1.5 meters or less of the vomiting event make them more susceptible to the infection ($P < 0.05$) (Table 3).

3.2 Virology investigation

A total of these virologies, including *E. coli* Salmonella, Shigella, Campylobacter, Yersinia enterocolitica, Norovirus, Rotavirus, Adenovirus and Astrovirus were detected to be negative. Sequence analysis of several positive RT-PCR samples revealed identical sequences, which were identified as SaVs. SaVs were detected in 7 of 11 stool specimens collected from the cases of the outbreak.

3.3 Analysis of nearly complete genome sequence

The nearly complete genomic sequence of strain SZ08 was obtained and its length was 7338 nt excluding the sequences of 5'UTRs and the 3'-end poly (A) tail in this study. The sequenced genomes were predicted to contain two major open reading frames (ORF) from 1 to 6837 (ORF1, encoding the nonstructural proteins and major capsid protein VP1) and from 6838 to 7337 (ORF2, encoding the minor structural protein). Its 3'-UTR had 52 nt. The reference strain MF462288/ GII.8/Peru330/PNV010961/2008 shared the highest query cover and nucleotide identities (92.4%) with strain SZ08 based on complete genome. The nucleotide identities of strain SZ08 with strain Peru330 were 93.13% in the ORF1, 94.61% in ORF2 and 93.30% in ORF3, respectively. According to the amino acid identities of the protein alignment, a great degree of similarity (96.95% at the nearly complete genome, 98.46% in the nonstructural proteins and VP1 and 97.54% in the VP2) were shown between them. We observe 48 aa mutations in the nonstructural proteins and VP1 and 25 aa mutations in the minor structural protein.

3.4 Analysis of complete capsid protein

According to the phylogenetic analysis, Strain SZ08 clustered into GII.8 branch and was independent with other clusters (A, B, C and D) named previously^[13] and was named after cluster E (Fig3). The nucleotide identity of Strain SZ08 was ranged from 92.94% to 94.02% with cluster A-D and the amino acid identity was ranged from 96.76% to 97.48%. However, cluster A-D shared the high nucleotide identity (96.87-99.84%) and amino acid identity (98.31-99.88%). Strain SZ08 introduced 11 specific amino acid mutations (S249A, A275S, S299A, A301T, G302T, V335N, L465I, S497N, I503V, A528V and V552I) compared with cluster A-D. However, five original amino acids have re-appeared (S43T, S183A, E341D, T403S, I520V) from the earliest strain Peru330/PNV010961/2008 (cluster A). Cluster B-D had low amino acid mutations (Fig 4).

4 Discussion

The epidemiology of cases of AGE of viral etiology is a relevant public health issue^[20]. Of note, reports on SaVs-associated AGE across all age groups have recently increased worldwide^[6,13,21], which indicated that the cases caused by SaVs may become extensive and global. And Shenzhen is also one of the high-incidence areas of SaVs-associated outbreaks recently in mainland China^[19,22]. Human SaVs cause AGE in all age groups in both sporadic cases and outbreaks worldwide as well as predominantly occurs in children^[23,24], which correspond to this study. On the other hand, the outbreak described by the study occurred in primary school and exposure to vomit was recognized as the cause, which are similar to the previous studies^[25,26].

The investigation showed the difficulty to prove the person-to-person transmission by contact or from aerosols generated by vomit^[27-29]. In this study, we investigated the vomiting places and the single factor analysis showed that the participation in the treatment of vomitus and the distance of vomitus were the main factors causing gastroenteritis infection (logistic regression, $P < 0.05$). It has been proposed that the formation of aerosols that can remain in the air for some time and can then be breathed in and swallowed or through contamination of surface causes the outbreaks^[30-32]. The aerosols transmission caused by vomiting might explain the partial gastroenteritis outbreaks which are not caused by foodborne or waterborne outbreaks as described previously^[32-35]. Vomit was considered only when exposure was at <1 m, however, possible exposures at greater distances have also been reported^[27,28]. Our study showed the distance under 1.5 m could cause the infection. Therefore, the study gave the evidence that the greater distance also could cause the cases. Since vomiting is a common symptom in calicivirus outbreaks, vomiting aerosols transmission may exist in most SaVs outbreaks. Therefore, vomiting aerosols transmission should be valued in the investigation of gastroenteritis outbreaks.

Emerging virus strains often have a risk of causing a pandemic. SaV GII.8 was first identified in two hospitalized children's samples in Peru, in 2008^[36]. In mainland China, GII.8 was first reported in Shenzhen, in 2011^[22]. Additionally, in 2019, Xue et al. acquired the first GII.8 SaV genome from mainland China^[13]. Based on the phylogenetic analysis, the strain SZ08 in this study was classified as a member of GII.8 SaV. In addition, strain SZ08 isolated in 2019 was clustered as independent branch and was different from the strain GZ2014-L231 isolated in 2014 although strain of Shenzhen and strain GZ2014-L231 of Guangzhou were both from the same province, Guangdong. GII.8 strains were also detected in other countries not only in clinical but also in environmental samples^[37,38]. SaVs outbreak caused by GII.8 strains occurred in long term center in previous report^[36] and the GII.8 SaV outbreak occurred in primary school. It suggested the GII.8 strain could cause the children and the elderly to infect. Despite the low detection rate, the wide spread distribution of the virus in different countries and wide age groups, make it as important concern to understand its genetic diversity and evolutionary characteristics.

Recombination and amino acid mutations are the two important way of viral evolution not only in SaVs but also in Noroviruses^[13,39]. According to the genomic analysis, there was no recombination occurred in strain SZ08.

Amino acid point mutations in proteins are the important way of viral evolution. In this study, by comparing the differences between the strain SZ08 as new cluster and GII.8 human SaV representative strains of other clusters, more amino acid mutations occurred in VP1 capsid region. GII.8 strain might obtain new antigenicity and receptor binding activity through the variation of amino acid sites in the capsid region, just as it occurs in norovirus, and gained greater susceptibility in the population. This might explain there was the large scale of cases in this SaVs outbreak in this study.

In conclusion, vomiting could be one of the transmission mechanisms which may explain a large number of cases of SaVs outbreaks. It is noteworthy that SaVs-associated diarrhea is generally mild, while large-scale outbreaks by SaV may also occur. In addition, we obtained the genome of a novel GII.8 SaV variant from China and comprehensively analyzed the genomic characteristics. The results of this study could not only provide reference data for SaVs researches in the future, but also deepen the understanding of evolution mechanisms of the new GII.8 variant. Constant surveillance is required to monitor the emergence of these strains and will permit the identification of changes in major strains and improve our knowledge of the evolution of SaVs among humans.

Declarations

Acknowledgments

We thank The Key Project of Science and Technology (Grant No. 2018ZX10301408-001 and 2018ZX10102-001) for their significant contribution to the outbreak investigation.

Conflicts of interest

The authors declare that they have no conflict of interests.

References

- [1] Cilli A, Luchs A, Leal E, et al. Human sapovirus GI.2 and GI.3 from children with acute gastroenteritis in northern Brazil[J]. *Mem Inst Oswaldo Cruz*, 2019, 114: e180574. doi:10.1590/0074-02760180574.
- [2] Usuku S, Kumazaki M. A gastroenteritis outbreak attributed to sapovirus genogroup v in Yokohama, Japan[J]. *Jpn J Infect Dis*, 2014, 67(5): 411-2. doi:10.7883/yoken.67.411.
- [3] Luo L F, Qiao K, Wang X G, et al. Acute gastroenteritis outbreak caused by a GI.6 norovirus[J]. *World J Gastroenterol*, 2015, 21(17): 5295-302. doi:10.3748/wjg.v21.i17.5295.
- [4] Ike A C, Hartelt K, Oehme R M, et al. Detection and characterization of sapoviruses in outbreaks of gastroenteritis in southwest Germany[J]. *J Clin Virol*, 2008, 43(1): 37-41. doi:10.1016/j.jcv.2008.04.003.
- [5] Kobayashi S, Fujiwara N, Yasui Y, et al. A foodborne outbreak of sapovirus linked to catered box lunches in Japan[J]. *Arch Virol*, 2012, 157(10): 1995-7. doi:10.1007/s00705-012-1394-8.
- [6] Liu X, Jahuiria H, Gilman R H, et al. Etiological Role and Repeated Infections of Sapovirus among Children Aged Less than 2 Years in a Cohort Study in a Peri-urban Community of Peru[J]. *J Clin Microbiol*, 2016, 54(6): 1598-1604. doi:10.1128/jcm.03133-15.
- [7] Oka T, Wang Q, Katayama K, et al. Comprehensive review of human sapoviruses[J]. *Clin Microbiol Rev*, 2015, 28(1): 32-53. doi:10.1128/cmr.00011-14.
- [8] He Y, Jin M, Chen K, et al. Gastroenteritis Outbreaks Associated with the Emergence of the New GI.4 Sydney Norovirus Variant during the Epidemic of 2012/13 in Shenzhen City, China[J]. *PLoS One*, 2016, 11(11): e0165880. doi:10.1371/journal.pone.0165880.
- [9] Iritani N, Yamamoto S P, Abe N, et al. Epidemics of GI.2 sapovirus in gastroenteritis outbreaks during 2012-2013 in Osaka City, Japan[J]. *J Med Virol*, 2016, 88(7): 1187-93. doi:10.1002/jmv.24451.
- [10] Phan T G, Trinh Q D, Yagyu F, et al. Emergence of rare sapovirus genotype among infants and children with acute gastroenteritis in Japan[J]. *Eur J Clin Microbiol Infect Dis*, 2007, 26(1): 21-7. doi:10.1007/s10096-006-0235-7.
- [11] Kumthip K, Khamrin P, Ushijima H, et al. Genetic recombination and diversity of sapovirus in pediatric patients with acute gastroenteritis in Thailand, 2010-2018[J]. *PeerJ*, 2020, 8: e8520. doi:10.7717/peerj.8520. [12] Diez-Valcarce M, Montmayeur A, Tatusov R, et al. Near-Complete Human Sapovirus Genome Sequences from Kenya[J]. *Microbiol Resour Announc*, 2019, 8(7). doi:10.1128/mra.01602-18. [13] Xue L, Cai W, Gao J, et al. Genome

- characteristics and molecular evolution of the human sapovirus variant GII.8[J]. *Infect Genet Evol*, 2019, 73: 362-367. doi:10.1016/j.meegid.2019.05.017.
- [14] Kageyama T, Kojima S, Shinohara M, et al. Broadly reactive and highly sensitive assay for Norwalk-like viruses based on real-time quantitative reverse transcription-PCR[J]. *J Clin Microbiol*, 2003, 41(4): 1548-57. doi:10.1128/jcm.41.4.1548-1557.2003.
- [15] Oka T, Katayama K, Hansman G S, et al. Detection of human sapovirus by real-time reverse transcription-polymerase chain reaction[J]. *J Med Virol*, 2006, 78(10): 1347-53. doi:10.1002/jmv.20699.
- [16] Magzoub M A, Bilal N E, Bilal J A, et al. Detection and sequencing of rotavirus among sudanese children[J]. *Pan Afr Med J*, 2017, 28: 87. doi:10.11604/pamj.2017.28.87.11008.
- [17] Resque H R, Munford V, Castilho J G, et al. Molecular characterization of astrovirus in stool samples from children in Sao Paulo, Brazil[J]. *Mem Inst Oswaldo Cruz*, 2007, 102(8): 969-74. doi:10.1590/s0074-02762007000800012.
- [18] Allard A, Albinsson B, Wadell G. Rapid typing of human adenoviruses by a general PCR combined with restriction endonuclease analysis[J]. *J Clin Microbiol*, 2001, 39(2): 498-505. doi:10.1128/jcm.39.2.498-505.2001.
- [19] Wang Md J, Li Ph D Y, Kong Md X, et al. Two gastroenteritis outbreaks caused by sapovirus in Shenzhen, China[J]. *J Med Virol*, 2018, 90(11): 1695-1702. doi:10.1002/jmv.25236. [20] Oka T, Mori K, Iritani N, et al. Human sapovirus classification based on complete capsid nucleotide sequences[J]. *Arch Virol*, 2012, 157(2): 349-52. doi:10.1007/s00705-011-1161-2.
- [21] Yamashita Y, Ootsuka Y, Kondo R, et al. Molecular characterization of Sapovirus detected in a gastroenteritis outbreak at a wedding hall[J]. *J Med Virol*, 2010, 82(4): 720-6. doi:10.1002/jmv.21646.
- [22] Wu W, Yang H, Zhang H L, et al. Surveillance of pathogens causing gastroenteritis and characterization of norovirus and sapovirus strains in Shenzhen, China, during 2011[J]. *Arch Virol*, 2014, 159(8): 1995-2002. doi:10.1007/s00705-014-1986-6.
- [23] Svraka S, Vennema H, Van Der Veer B, et al. Epidemiology and genotype analysis of emerging sapovirus-associated infections across Europe[J]. *J Clin Microbiol*, 2010, 48(6): 2191-8. doi:10.1128/jcm.02427-09.
- [24] Phan T G, Khamrin P, Quang T D, et al. Emergence of intragenotype recombinant sapovirus in Japan[J]. *Infect Genet Evol*, 2007, 7(4): 542-6. doi:10.1016/j.meegid.2007.02.004.
- [25] Yu Y, Guo X H, Yan H Q, et al. [Systematic review on the characteristics of acute gastroenteritis outbreaks caused by sapovirus][J]. *Zhonghua Liu Xing Bing Xue Za Zhi*, 2019, 40(1): 93-98. doi:10.3760/cma.j.issn.0254-6450.2019.01.019.
- [26] Becker-Dreps S, Bucardo F, Vinje J. Sapovirus: an important cause of acute gastroenteritis in children[J]. *Lancet Child Adolesc Health*, 2019, 3(11): 758-759. doi:10.1016/s2352-4642(19)30270-6.
- [27] Makison Booth C. Vomiting Larry: a simulated vomiting system for assessing environmental contamination from projectile vomiting related to norovirus infection[J]. *J Infect Prev*, 2014, 15(5): 176-180. doi:10.1177/1757177414545390.

- [28] Marks P J, Vipond I B, Carlisle D, et al. Evidence for airborne transmission of Norwalk-like virus (NLV) in a hotel restaurant[J]. *Epidemiol Infect*, 2000, 124(3): 481-7. doi:10.1017/s0950268899003805.
- [29] Widdowson M A, Glass R, Monroe S, et al. Probable transmission of norovirus on an airplane[J]. *Jama*, 2005, 293(15): 1859-60. doi:10.1001/jama.293.15.1859.
- [30] Lopman B. Air sickness: vomiting and environmental transmission of norovirus on aircraft[J]. *Clin Infect Dis*, 2011, 53(6): 521-2. doi:10.1093/cid/cir486.
- [31] Kirking H L, Cortes J, Burrer S, et al. Likely transmission of norovirus on an airplane, October 2008[J]. *Clin Infect Dis*, 2010, 50(9): 1216-21. doi:10.1086/651597.
- [32] Marks P J, Vipond I B, Regan F M, et al. A school outbreak of Norwalk-like virus: evidence for airborne transmission[J]. *Epidemiol Infect*, 2003, 131(1): 727-36. doi:10.1017/s0950268803008689.
- [33] Godoy P, Artigues A, Bartolome R, et al. [Norovirus gastroenteritis outbreak by person-to-person transmission in a nursing home][J]. *Med Clin (Barc)*, 2006, 127(14): 538-41. doi:10.1016/s0025-7753(06)72322-3.
- [34] Harris J P, Lopman B A, Cooper B S, et al. Does spatial proximity drive norovirus transmission during outbreaks in hospitals?[J]. *BMJ Open*, 2013, 3(7). doi:10.1136/bmjopen-2013-003060.
- [35] Wikswo M E, Cortes J, Hall A J, et al. Disease transmission and passenger behaviors during a high morbidity Norovirus outbreak on a cruise ship, January 2009[J]. *Clin Infect Dis*, 2011, 52(9): 1116-22. doi:10.1093/cid/cir144.
- [36] Diez-Valcarce M, Castro C J, Marine R L, et al. Genetic diversity of human sapovirus across the Americas[J]. *J Clin Virol*, 2018, 104: 65-72. doi:10.1016/j.jcv.2018.05.003.
- [37] Kagning Tsinda E, Malasao R, Furuse Y, et al. Complete Coding Genome Sequences of Uncommon GII.8 Sapovirus Strains Identified in Diarrhea Samples Collected from Peruvian Children[J]. *Genome Announc*, 2017, 5(43). doi:10.1128/genomeA.01137-17.
- [38] Kitajima M, Rachmadi A T, Iker B C, et al. Temporal variations in genotype distribution of human sapoviruses and Aichi virus 1 in wastewater in Southern Arizona, United States[J]. *J Appl Microbiol*, 2018, 124(5): 1324-1332. doi:10.1111/jam.13712.
- [39] White P A. Evolution of norovirus[J]. *Clin Microbiol Infect*, 2014, 20(8): 741-5. doi:10.1111/1469-0691.12746.

Tables

Table 1 Primers used in this study for sapovirus detection and amplification of almost whole genome

	primer	Nucleotide sequence (5'-3')	position	source
Amplification of almost complete genome	orf-1	GAGTTGAACACGCAGTCC	42-59 ^a	This study
		ATGATGGCACCAGTAAGG	1009-1026 ^a	
	orf-2	CGCAGTGTCTAGTGGTGTC	880-897 ^a	
		AAGGTATGGTCGAACGAA	1880-1897 ^a	
	orf-3	CGTCGTGTTCTAACTGTTGA	1763-1782 ^a	
		GAATCTTGTGATAACCTCCATA	2740-2761 ^a	
	orf-4	CCAATTAGTAGCTGAAACCCTT	2719-2740 ^a	
		CCCTTCCACGCAAACACG	3634-3651 ^a	
	orf-5	CAGAGGGCACTTATGAGAC	3432-3450 ^a	
		CAGAGGCAGTTATGGGAG	4434-4451 ^a	
orf-6	GGGTCGTGTATTGTTTGG	4335-4352 ^a		
	CCGAGTTTGGCATTCTA	5280-5297 ^a		
orf-7	TAGTGTTTGAAATGGAGGGC	5157-5176 ^a		
	TTGGGAAGTGACTGCTGA	6164-6181 ^a		
orf-8	AGGGCATCATCTTTCCAC	6114-6131 ^a		
	TTTGCGAGACAGTCCATTA	7029-7047 ^a		
orf-9	GGAATCAACTGCGGAAAT	6577-6594 ^a		
	GAATAAGACAGCGATGGTC	7429-7447 ^a		
AP1	CTCGCCACCTATGAAGCA	5081-5098 ^a		
	AATCGGCAGTGATGTCCT	7004-7021 ^a		

Orf1-1	CTTCTAAGCCATTCTACTCAA CGTGCCATGAACTGTTTG	5-25 ^b 1179- 1196 ^b
Orf1-2	GGCAGCATCACATCAGTC TGTTTGCAGACATGAACG	1093- 1110 ^b 2078- 2095 ^b
Orf1- 1,2	CTTCTAAGCCATTCTACTCAA GCTCTGTTTGCAGACATG	5-25 ^b
Orf1-4	ACCGTGGGTGGTATGACT GGCGACAACCGTTGAAAT	

^aPosition in the complete sapovirus GII genogroup sequence (accession no: MG674584).

^bPosition in the complete sapovirus GII genogroup sequence (accession no: MF462287).

Table 2. Distribution of vomiting places of students

Place of vomiting		Vomiting number	%
Bus	Plastic bag	28	13
	ground	10	5
	Outside the window	2	1
Military training	bathroom	6	3
	Dorm	4	2
	Playground	4	2
	Canteen	1	0
School	toilet	61	28
	Classroom	20	9
	Playground	7	3
	doorway	6	3
	Other	13	6

Table 3. Case control study on risk factors of acute gastroenteritis outbreak caused by Sapovirus

No.	case		Non case		OR value	95%CI		χ^2	P value
	Expose	%	Expose	%		lower limit	Upper limit		
See others vomiting	278	58	365	46	1.58	1.26	1.99	15.60	0.00
See others vomiting in the classroom	120	25	138	17	1.56	1.19	2.06	10.14	0.00
See others vomiting in the playground	36	7	35	4	1.74	1.08	2.81	5.22	0.02
Less than 1.5m from vomit	156	32	166	21	1.42	1.04	1.95	4.81	0.03
Handling vomitus	45	9	36	5	2.17	1.37	3.41	11.57	0.00
See others Vomiting on bus	86	18	116	15	1.25	0.92	1.70	2.11	0.15
See others Vomiting during activity	14	3	30	4	0.76	0.40	1.44	0.72	0.40
See others Vomiting in bathroom	56	12	71	9	1.33	0.92	1.93	2.28	0.13
Self protection in handling	12	27	7	19	1.93	0.69	5.40	1.62	0.20
Family members in the same school	116	24	170	22	1.15	0.88	1.51	1.06	0.30

Figures




Figure 1

Epidemic curve of acute gastroenteritis cases in the outbreak occurred in Shenzhen, China, 2019





Figure 2

Schematic diagram of each floor of Building




Figure 3

Phylogenetic analysis of SaVs based on nucleotide of complete VP1.

Cluster	name	43	107	183	249	275	296	299	301	302	335	341	403	445	465	497	503	520	528	552
A	MF462288_GII.8/Peru330/PNV010961/2008	S	Y	A	S	A	F	S	A	G	V	D	S	M	L	S	I	V	A	V
C	KX894314_GII.8/Miaoli/05099515477/2014/TW	T	F	S	S	A	Y	S	S	G	I	E	T	M	L	S	I	I	A	V
D	MK599471_GII.8/Guangzhou/GZ2014L231/CHN/2014	S	F	S	S	A	Y	S	S	G	V	E	T	M	L	S	I	I	A	V
B	MG674583_PE/2016/GII.8/Lima1867	S	F	S	S	A	Y	S	S	G	V	E	T	M	L	S	I	I	A	V
E	SaV_2019008_ShenZhen_2019_CHN	S	F	A	A	S	Y	A	T	T	N	D	S	M	I	N	V	V	V	I

Figure 4

Amino acid variation in complete capsid protein gene compared to reference strains of SaV GII.8