Enabling Culturally Safe Sexual Health Services in Western Sydney: a Protocol to Improve Outcomes for Aboriginal Young People.

Ashley John Ubrihien (✉ aubr4590@uni.sydney.edu.au)

The University of Sydney School of Medicine <https://orcid.org/0000-0002-9883-0891>

Kylie Gwynne

The University of Sydney School of Medicine

David A. Lewis

The University of Sydney School of Medicine

Study Protocol

Keywords: Sexual Health, Aboriginal, Barriers, Equity

Posted Date: June 10th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-34075/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Aboriginal people face challenges on several fronts when it comes to the health and wellbeing of their community, compared to the rest of the Australian population. This is no different in urban areas such as Australia's largest urban Aboriginal community located in Blacktown, NSW, where sexually transmitted infections remain an issue of concern. Across Australia, rates of Infectious Syphilis Human Immunodeficiency Virus (HIV) and Hepatitis C infection have increased by 400%, 260%, and 15% respectively while Gonorrhoea decreased 12% in the 5-year period from 2013 to 2017. This study explores how to address the barriers that prevent young Aboriginal people under 30 years of age accessing public Sexual Health Services.

Methods

Purposeful qualitative sampling will be undertaken to recruit 20 male and 20 female health consumers, 10 Aboriginal Elders and 10 sexual health clinicians. This recruitment will be undertaken with the assistance of local Aboriginal organisations and Western Sydney Local Health District (WSLHD). Participants will be interviewed on a one on one basis using semi structured interviews and participants will be interviewed by someone of the same gender in order to address cultural preferences. Data collected will be analysed using NVivo and by conducting a thematic analysis.

Discussion

This study will seek to add to the literature that explores why young Aboriginal people do not access sexual health services. This study seeks to understand the experience of clinicians, Aboriginal Elders and Aboriginal young people to provide practical policy and clinical redesign evidence that can be used to improve the experience and cultural safety of sexual health services in urban areas of Australia. The results of the qualitative research will be disseminated with the assistance of participating local Aboriginal organisations and the findings will be published through peer-reviewed scientific journals and conference presentations.

Trial registration

The study is approved by Western Sydney Local Health District Human Research Ethics Committee (HREC/16/WMEAD/449) and the New South Wales Aboriginal Health and Medical Research Council's Human Research Ethics Committee (1220/16).

Background

In 2008 *The Close the Gap report* delivered the Australian Parliament a stark picture of the health of Australia's first people (1). The scale of health inequity between Aboriginal people and the rest of the population manifested in a life expectancy gap of about 10 years (1). Over the past 12 years still the release of that report there has been little change in health outcomes for Aboriginal and Torres Strait Islander Australians (hereafter, Aboriginal) despite a whole of government focus on major goals

articulated in the National Closing the Gap strategy (2). In parallel, the rates of sexually transmissible infection (STIs) for Aboriginal people have continued to rise over this period (1). Of particular concern are rates of Infectious Syphilis, Human Immunodeficiency Virus (HIV) and Hepatitis C infection have increased by 400%, 260%, and 15% respectively while Gonorrhoea and decreased 12% in the 5-year period from 2013 to 2017 (3). There is a need to develop novel approaches to identifying how services can better respond to the health care needs of this population.

Rates of STIs amongst the Australian Aboriginal population are a major public health concern. Incidence rates of STIs are increasing in a population that already has higher rates of infection compared to that of the non-Aboriginal people (4). Whilst the Aboriginal population across Australia represents 3% of the total population, incidence rates of STIs can range anywhere from 3 to 30 times higher than that of the non-Aboriginal population (3, 4).

According to the Australian Bureau of Statistics (ABS), the Western Sydney Local Government area (LGA) of Blacktown City Council has the largest urban population of Aboriginal people with approximately 9,526 living in this area at the time of the 2016 Australian Census (5). Although rates of STIs in urban Aboriginal communities are not as high as remote communities, which experience rates of infection that are 5 to 500 times higher than the general population, sexual health should be prioritised within Blacktown LGA due to the relatively high number of Aboriginal people living in this concentrated urban environment (4). Another important factor worthy of consideration when assessing the scale of the public health problem is the likelihood that members of a community would be sexually active and therefore at risk of acquiring and transmitting STIs to other members of that community. The ABS identifies that the median age of Aboriginal people living in this LGA as 19 years of age which means that there is significant number of people who may be at risk of STI acquisition (5). The New South Wales (NSW) Government's *NSW STI Strategy 2016-2020* also identifies Aboriginal people as a Priority Population in the strategy and in particular focuses upon young people (6).

Aboriginal people in urban communities are often not the focus of research because the funding priorities of the Commonwealth Government tend to direct significant funding towards remote communities and leave state and territory governments to address health needs in urban areas (4). Aboriginal people in Western Sydney are highly urbanised yet are still face significant barriers to healthcare. They are younger than the wider population, experience high rates of unemployment and generally have larger families (5).

One of the barriers to accessing sexual health services appears to be shame and stigma (7-9). In a sample of 45 young Aboriginal people, surveyed by Mooney-Somers *et al.*, one third of the participants indicated that shame would be associated with having an STI and having to seek treatment (7). This is also reflected in the mixed method study of Su *et al.* who found that the shame of many Aboriginal men was associated with the possibility of being seen at a sexual health clinics and an inferences of weakness and disease that might be associated with such a visit (9).

There is a significant gap in the published literature in terms of studies that seek to understand the issues that prevent Aboriginal people participating in sexual health care through the collection of directly

obtained evidence. Many of the studies reviewed, including those undertaken by Bailie *et al.*, Nattabi *et al.*, Buhrer-Skinner *et al.* and Richmond *et al.*, rely on the response of clinicians in quality improvement processes to guide the conclusions of their research (10-13). These studies focus on the process aspects of clinical service delivery and how quality improvement principles can be followed to improve quality; however, these tools provide limited insight into the barriers that prevent young Aboriginal people attending the services (10-13). The conclusions of research using this approach may have some merits; however, they are likely to miss the important nuances that could be obtained by engaging health consumers in the research process. It is possible to engage health consumers in quality improvement processes and this would give those undertaking such a study an objective source of evidence.

The status of sexual health amongst young Aboriginal people in Western Sydney is an area that has not been explored in any great depth within the published literature. The study undertaken by Biggs *et al.* provides a cross sectional view of Aboriginal people attending public sexual health services in Western Sydney through an incentive-based program called 'The Deadly Liver Mob' (14). The Deadly Liver Mob program primarily focused on education concerning Hepatitis C infection for people who are currently or have been injecting drug users (14). For these reasons, the group screened for STIs in this study are a distinct sub-population of the Aboriginal Community, making it hard to generalise the results of this study (14). This study also relied on the retrospective review of routine clinical data so there was no ability to explore questions that were not addressed in a limited data set.

This present study will seek to add to the literature by exploring why young Aboriginal people do not access sexual health services. This study seeks to understand the experience of clinicians, Aboriginal Elders and Aboriginal young people to provide practical policy and clinical redesign evidence that can be used to improve the experience and cultural safety of sexual health services in urban areas of Australia.

Methods

This qualitative study will utilise semi-structured one-on-one interviews with 40 health consumers, 10 clinicians and 10 community elders. All cohort groups will be interviewed using a semi structured interviewer assisted questionnaire with the aim of answering the following questions:

1. What prevents Aboriginal people in Western Sydney from accessing mainstream services?
2. What would enable Aboriginal people in Western Sydney to access mainstream services?
3. Would changes in service modality would make services more accessible?
4. What participants see as the most important things that could be implemented to improve sexual health services in Western Sydney?

The study investigators have adequate qualifications and experience to undertake a qualitative study in the area of Sexual Health and have significant expertise in community engagement and co-design. The questionnaire is worded in plain English to ensure that potential participants with low literacy can also

participate. The results of this study will be analysed thematically using the software program, NVivo (QSR International, Melbourne, Australia), so that major themes can be identified and summarised.

This study will provide benefits to Aboriginal people by enabling local health services to better understand and allocate resource to meet the needs of young Aboriginal people through the acquisition of more robust study data concerning how Sexual Health Services may best be delivered. This data will create translational research that will assist service delivery models in publicly funded Sexual Health services in Western Sydney and perhaps more widely.

Study participants will be free to withdraw from the study at any point up to the submission of the questionnaire. Submissions will not be able to be withdrawn after this point because the individual questionnaire will not feature any identifying marks so therefore any individual submission will not be able to be identified for withdrawal after submission.

Purposeful qualitative sampling will be undertaken to recruit an equal balance of 20 male and 20 female healthcare consumers, 10 Aboriginal Elders and 10 sexual health clinicians. This recruitment will be undertaken with the assistance of local Aboriginal Organisations and Western Sydney Local Health District (WSLHD).

This study has been approved by the Western Sydney Local Health District Human Research Ethics Committee (application reference HREC/16/WMEAD/449) and the New South Wales Aboriginal Health and Medical Research Council's Human Research Ethics Committee (application reference 1220/16).

This study will be conducted in a manner that safeguards the anonymity of participants by ensuring that interview responses do not contain any identifying marks apart from an indication of age and sex. This will assist in protecting the privacy of Aboriginal people who choose to participate in the study. The value of Aboriginal interpretation is integral to this project's success and this will be achieved through the engagement of Aboriginal research assistants, consultation about the study design and through the engagement of Aboriginal people in the analysis of data and the final conclusions of the study. Involvement in the governance of the study is also of paramount importance and every effort will be made to ensure participation.

The investigators anticipate that some risks may be posed by participants disclosing sensitive information particularly around the conduct of illegal activities, a history of sexual or physical abuse or need to have sexual health conditions treated. Should a disclosure of illegal activity be made the interview would be immediately terminated in order to ensure that the participant didn't further incriminate themselves. If the disclosure relates to any form of abuse that falls under the definitions of Section 27 of the *NSW Children and Young Persons (Care and Protection) Act 1998* the matter will be immediately reported to the NSW Department of Families and Communities and referrals would be offered to relevant support services (15). In order to maintain the confidentiality of participants all data collected will be anonymous and contain no potentially identifying information apart from age and sex. It is important to

highlight to participants the risk that is posed by them disclosing illegal activities because of the need of study investigators to report such activities to the relevant government agency.

The results of this study will be disseminated through several different media including meetings with Study participants, local Elders and Aboriginal Groups, peer reviewed publications, presentation of results at conferences, and through summaries presented via social media.

The contribution of local Aboriginal people will be acknowledged in all publications to reflect the unique contribution that they provide, and co-authorship of academic papers will be offered, consistent with University of Sydney policy on the matter. The research will be conducted in accordance with the National Health and Medical Research Council (NHMRC) guidelines for ethical research with Aboriginal people (16). Plain English resources and information meetings will be undertaken with communities involved in this study to provide a summary of the results of the findings.

Discussion

The results of this study will be significant in informing urban health services on the needs of young Aboriginal people in terms of how they prefer to access public sexual health services and the barriers that currently exist that may prevent them from using existing services. Much of the research based both in remote areas and urban areas relies on the experiences of clinicians rather than the lived experience of young Aboriginal people themselves and may miss important aspects of interpretation that only occur when Aboriginal people share their insights. These insights will be deepened in this study by involving local Aboriginal people in the interpretation of the de-identified results of the study.

This engagement project will provide a set of documents that will be informed by the local community and provide policy makers with practical measures that can be implemented to improve the cultural appropriateness and clinical care of young Aboriginal people. This study will also address important policy questions and directly inform on the delivery of healthcare to young Aboriginal people in urban settings. The study is approved by Western Sydney Local Health District Human Research Ethics Committee (HREC/16/WMEAD/449) and the New South Wales Aboriginal Health and Medical Research Council's Human Research Ethics Committee (1220/16).

Abbreviations

ABS: Australian Bureau of Statistics

HIV: Human Immunodeficiency Virus

LGA: Local Government Area

NHMRC: National Health and Medical Research Council

NSW: New South Wales

STIs: Sexually Transmitted Infections

WSLHD: Western Sydney Local Health District

Declarations

Availability of data and materials

Not applicable. This manuscript does not contain any data.

Competing interests

The authors declare that they have no competing interests in this section.

Funding

The authors have no funding to declare for this study.

Authors' contributions

AU drafted the manuscript and assistance was provided by KG and DL with editing and redrafting of the manuscript. All authors read and approved the final manuscript."

Acknowledgements

Not applicable.

References

1. Institute TL. Close the Gap report – “Our Choices, Our Voices”. Carlton, Victoria: The Lowitja Institute; 2019.
2. Institute TL. Close the Gap (2020). Carlton, Victoria: The Lowitja Institute; 2020.
3. Institute K. Aboriginal Surveillance Report on HIV, viral hepatitis and STIs 2018. Kensington, New South Wales, Australia: University of New South Wales; 2018.
4. Kirby Institute. HIV, viral hepatitis and sexually transmissible infections in Australia Annual Surveillance Report. Kensington, NSW: University of New South Wales; 2018 2018.
5. Australian Bureau of Statistics. 2016 Census QuickStats - Aboriginal and Torres Strait Islander people - usual residents Canberra, ACT: Commonwealth of Australia; 2016 [Available from: http://www.censusdata.abs.gov.au/census_services/getproduct/census/2016/quickstat/LGA10750?opendocument.
6. NSW Ministry of Health. NSW STI Strategy 2016-2020. North Sydney, NSW: NSW Government; 2016.
7. Mooney-Somers J, Erick W, Scott R, Akee A, Kaldor J, Maher L. Enhancing Aboriginal and Torres Strait Islander young people? s resilience to blood-borne and sexually transmitted infections: Findings from

- a community-based participatory research project. *Health Promotion Journal of Australia*. 2009;20(3):195-201.
8. Forrest B, Plummer D. Factors affecting indigenous Australians' access to sexual health clinical services. *Venereology*. 1999;12(2):47.
 9. Su J-Y, Belton S, Ryder N. Why are men less tested for sexually transmitted infections in remote Australian Indigenous communities? A mixed-methods study. *Culture, Health & Sexuality*. 2016;18(10):1150-64.
 10. Nattabi B, Matthews V, Bailie J, Rumbold A, Scrimgeour D, Schierhout G, et al. Wide variation in sexually transmitted infection testing and counselling at Aboriginal primary health care centres in Australia: analysis of longitudinal continuous quality improvement data. *BMC Infectious Diseases*. 2017;17(1):148.
 11. Bailie C, Matthews V, Bailie J, Burgess P, Copley K, Kennedy C, et al. Determinants and gaps in preventive care delivery for Indigenous Australians: a cross-sectional analysis. *Frontiers in Public Health*. 2016;4:34.
 12. Buhrer-Skinner M, Muller R, Buettner PG, Gordon R, Debattista J. Reducing barriers to testing for Chlamydia trachomatis by mailed self-collected samples. *Sexual Health*. 2013;10(1):32-8.
 13. Richmond JA, Sasadeusz J, Temple-Smith M. The role of primary health care in hepatitis B testing and management: a case study. *Journal of Community Health*. 2018;43(1):38-47.
 14. Biggs K, Walsh J, Ooi C. Deadly Liver Mob: opening the door—improving sexual health pathways for Aboriginal people in Western Sydney. *Sexual Health*. 2016;13(5):457-64.
 15. NSW Children and Young Persons (Care and Protection) Act 1998, Section 27 (1998).
 16. National Health and Medical Research Council. Ethical guidelines for research with Aboriginal and Torres Strait Islander Peoples. Canberra, Australia: Commonwealth of Australia; 2018.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CoverLetter.pdf](#)
- [4901SSA17wmead163AuthorisationLetter.pdf](#)
- [AHMRCMtg616122016projectapproval.pdf](#)
- [FundingLetter.pdf](#)