Covid-19 in the second half of pregnancy: prevalence and clinical relevance

Marta RUGGIERO

Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico and Università degli Studi di Milano, Milan, Italy

Edgardo SOMIGLIANA (✉ dadosomigliana@yahoo.it)

Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico and Università degli Studi di Milano, Milan, Italy <https://orcid.org/0000-0002-0223-0032>

Beatrice TASSIS

Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

Letizia LI PIANI

Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico and Università degli Studi di Milano, Milan, Italy

Sara UCEDA RENTERIA

Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

Giusy BARBARA

Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

Giovanna LUNGI

Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

Enrico FERRAZZI

Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico and Università degli Studi di Milano, Milan, Italy

Research Article

Keywords: Covid-19, Sars-Cov-2, pregnancy, antibodies

Posted Date: June 9th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-34492/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Evidence on the prevalence and outcome of Covid-19 in pregnancy is yet inconclusive. To draw more information on this issue, we prospectively recruited 315 consecutive women delivering in a referral hospital located in Lombardy, Italy and evaluated the proportion of women who had Covid-19 in pregnancy. All recruited subjects underwent a nasopharyngeal swab to assess the presence of Sars-Cov-2 using Real-time PCR. In addition, two different type of antibodies for the virus were evaluated in peripheral blood, those against the spike proteins S1 and S2 of the envelope and those against the nucleoprotein of the nucleocapsid. Women were considered to have had Covid-19 in pregnancy if at least one of the assessment was positive. Overall, 28 women had a diagnosis of Covid-19 in pregnancy (8.9%, 95%CI: 6.2-12.5%), in line with the local general population. Women diagnosed with the infection were more likely to report one or more episodes of symptoms suggestive for Covid-19 (n=11, 39.3%) compared to unaffected women (n=39, 13.6%). The corresponding OR was 4.11 (95%CI: 1.79-9.44). Symptoms significantly associated with Covid-19 in pregnancy included fever, cough, dyspnea and anosmia. Only one woman necessitated intensive care. No maternal deaths occurred. Pregnancy outcome in women with and without Covid-19 did not also differ. In conclusion, women in the second half of pregnancy do not appear to be more susceptible to Covid-19 and its complications. In addition, the study suggests that pregnancy outcome may not be significantly affected.

Introduction

The Severe Acute Respiratory Syndrome Coronavirus type-2 (SARS-Cov-2) was first identified in the area of Wuhan, China, at the end of 2019 but then rapidly spread globally in the beginning of 2020. In Lombardy, a region of Northern Italy that was particularly hit by the epidemic, the first case was identified on February 20th, 2020¹. SARS-Cov-2 causes a potentially fatal infection named Covid-19 (Coronavirus Disease - 2019). Most common symptoms include fever, cough, sore throat, malaise, myalgia, gastrointestinal symptoms, anosmia and ageusia². In more advanced conditions the subject can experience severe dyspnea and respiratory failure requiring admission in intensive care units³. The fatality rate varies widely in the literature (1–22%) and depends on the clinical phase of the disease, gender, age and predisposing conditions such as diabetes, hypertensive disorders, obesity and smoking^{3,4}.

Albeit still limited, there is evidence that SARS-Cov-2 can cause life-threatening situations also in pregnancy, even if fatal cases seem rare^{5–9}. Severe pneumonia occurs in 5–16% of women^{5,8,9} and may require prompt pregnancy termination for improving respiratory assistance. However, this evidence was drawn from case series, a study design that could overestimate the detrimental effects of SARS-Cov-2. Indeed, asymptomatic or poorly symptomatic women are more likely to go undetected, thus boosting the epidemiological relevance of Covid-19 related complications. To assess the clinical relevance of these complications in pregnancy one needs to know the real denominator, i.e. the total number of women being infected with the virus.

The recent availability of antibodies could provide more informative evidence on this issue. Two different types of antibodies can currently be tested, those against the glycoproteins of the virus envelope and those against the nucleoproteins. The former are generally considered neutralizing while the role of the latter in controlling virus replication is not yet established. Both were shown to be valuable markers of infection ¹⁰.

To shed more light on Covid-19 in pregnancy, we prospectively recruited consecutive women delivering in a referral hospital located in Milan, Lombardy. All women underwent a nasopharyngeal swab to investigate the presence of the virus and provided a blood sample to assess the presence of antibodies against SARS-Cov-2. The primary aim was to determine the rate of pregnant women who were exposed to Covid-19 during pregnancy. Secondary aims included the clinical presentation of the infection and the possible impact on pregnancy outcome. To note, the recruitment period lasted from April 7th to May 6th and thus offered us the unique opportunity to study women that could be exposed to SARS-Cov-2 exclusively during the second part of pregnancy.

Materials And Methods

The study took place at the Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, a referral hospital for Obstetrics located in Milan, in the Italian region of Lombardy. The hospital ensures about 5,000 deliveries per year and was committed by the local health and governmental officials to be a referral hospital also for Covid-19 in pregnancy. All women delivering between April 7th to May 06th, 2020 were consecutively considered for study entry. Exclusion criteria were as follows: 1) pregnancy termination before 20 weeks' gestation (miscarriages), 2) referral from other hospitals because of Covid-19 diagnosis. Eligible women were invited to participate the day after delivery. Women agreeing to be recruited signed an informed consent. They were requested to fill a specific questionnaire and provided a blood sample that was centrifuged at 3,500 rpm for 8 minutes at room temperature and stored at -20 °C until assayed for the presence of antibodies. For the nasopharyngeal swab for Sars-Cov-2, we relied on the ascertainment that was systematically performed at hospital admission. The study was approved by the local Institutional Review Board (Comitato Etico Milano Area 2, N. 356_2020). All research was performed in accordance with the guidelines of the International Conference of Harmonization of Good Clinical Practice (ICH/GCP) and related regulations, including the Helsinki declaration of June 1964 and its adaptation done by the World Medical Association General Assembly in Seoul in 2008.

Collected information included general medical and obstetrics history as well as the pregnancy outcome of the index pregnancy. Patients' charts were used to obtain this information. The questionnaire that women had to fill evaluated the presence of symptoms suggestive for Covid-19 over the last three months. More than one episode of symptoms could be reported. The specific items included in the questionnaire are illustrated in details elsewhere ¹¹. Briefly, they include the presence of fever (≥ 37.5 °C), cough, sore throat, rhinitis, headache, diarrhea, vomit, dyspnea (including tachypnea), asthenia, myalgia, anosmia and ageusia. In addition, women were asked if they were exposed to higher risk of infection for job situations (health workers or laboratory technicians) or contacts with persons with a definitive

diagnosis of Covid–19 or cohabitant with persons doing a job at risk or having symptoms suggestive for Covid–19.

Evaluation of Covid–19 in the cohort was based on three determinations: identification of the specific RNA on a nasopharyngeal swab and research of Immunoglobuline-type G (IgG) using two different tests. Women were considered to have had Covid–19 in pregnancy if at least one of the assessment was positive. The presence of the virus in the nasopharyngeal swab was investigated using the multiplex real-time (RT)-PCR for the qualitative detection of three target genes of SARS-CoV–2 (Allplex 2019-nCov Assay, Seegene, Seoul, Korea): SARS-CoV–2 nucleocapsid proteins and RNA polymerase gene fragments (both specific for Sars-Cov–2), and envelope gene fragment for detection of the sarbecovirus subgenus family that includes SARS-CoV–2. The assay has a full-process negative control, positive control and internal control. The first serologic test was a chemiluminescence (CLIA) immunoassay (LIAISON®, Diasorin, Saluggia - Italy) used for the quantitative detection of IgG antibodies against S1 and S2 antigens of SARS-CoV–2. The S1 and S2 proteins are derived from SARS-CoV–2 spike protein that is responsible for entry into the host cell. Antibodies against these two targets are neutralizing. The test has a sensitivity and specificity of 97.4% and 98.5%, respectively. The second test was an electrochemiluminescence (ECLIA) immunoassay (Elecsys®, Roche Diagnostics, Mannheim, Germany) that uses a recombinant nucleoprotein representing the nucleocapsid antigen for the qualitative determination of antibodies against SARS-CoV–2. This nucleoprotein antigen is internal to the envelope. The test has a specificity greater than 99.8% and a sensitivity of 100%.

Data analyses were performed using the Statistical Package for Social Sciences (SPSS 23.0, IL, USA). Women were considered to be affected if at least one of the three assessments resulted positive for Sars-Cov–2. The sample size (at least 250 subjects) was calculated expecting about 20% of women being affected and aiming at a precision in the estimated prevalence of $\pm 5\%$. A binomial distribution model was used to determine the 95% Confidence Interval (CI) of the most relevant proportions. Data was compared using Student *t* Test, Fisher exact test or Chi Square test as appropriate. P values below 0.05 were considered statistically significant. The database is reported as additional material.

Results

Three hundred eighty-five women delivered during the study period. Twelve were excluded because they were referred from other hospitals for Covid–19. Thirty-five were not recruited because of violation of the study protocol (the study was not proposed). Twenty-three women refused to participate. Three hundred fifteen women were ultimately enrolled. Overall, 28 women had a diagnosis of Covid–19 (8.9%, 95%CI: 6.2–12.5%). Specific rates according to the test used are reported in Table 1. Details on the concordance among tests are illustrated in Table 2. To note, among the 24 women who were detected with antibodies against Sars-Cov–2, 17 (71%) were positive to both tests used, three (12%) were positive only for antibodies against the envelop and four (17%) only to those against the nucleoprotein. Baseline characteristics of women who did and did not have Covid–19 are shown in Table 3. The two groups did not significantly differ for any of these characteristics.

Risk factors for Covid-19 are shown in Table 4. Symptoms significantly associated with Covid-19 included fever, cough, dyspnea and anosmia. A trend emerged also for ageusia (Table 4). Women diagnosed with the infection were more likely to report one or more episodes of symptoms suggestive for Covid-19 ($n = 11$, 39.3%) compared to unaffected women ($n = 39$, 13.6%) ($p = 0.001$). The corresponding Odds Ratio (OR) was 4.11 (95%CI: 1.79–9.44). In contrast, no difference emerged for conditions at higher risk of infection including type of job, reported direct contacts or cohabitation with persons at risk (Table 4).

Pregnancy outcome in women with and without Covid-19 diagnosis is shown in Table 5. No statistically significant differences emerged. Only one woman with Covid-19 underwent cesarean delivery because of pneumonia and severe respiratory insufficiency. After delivery, she required admission in the intensive care unit. She has now recovered. No other women needed intensive care. No maternal deaths occurred.

Discussion

Covid-19 in pregnancy was not rare in our area during this first outbreak of Sars-Cov-2. One in 11 women (8.9%) actually entered in contact with the virus. To note, being the pandemic still ongoing, this proportion is expected to rise in the next future, i.e. when the duration of exposure to the outbreak will be longer (we exclusively recruited women who delivered, thus those who were exposed only in the last three months of pregnancy). On the other hand, the clinical course of the disease appeared mostly unremarkable. Sixty-one percent did not report any symptom, prompt preterm delivery because of Covid-19 complications was necessary only in one case and pregnancy outcome was not markedly influenced. The robustness of these latter conclusions is however modest because they emerged from secondary and underpowered analyses.

Interestingly, the rate of infected women observed in our study is very similar to the prevalence observed in a concomitant survey performed in our area and focusing on blood donors. Specifically, Valenti et al. evaluated the presence of antibodies against the nucleocapsid protein and reported for April 2020 a prevalence of 7.1% (95%CI: 4.4–10.8%), thus in line with our findings ¹². In contrast, our reassuring clinical findings are somehow in disagreement with recent evidence from large case series of affected pregnant women. Savasi et al. reported on an Italian multicenter case series of 77 women hospitalized for Covid-19 and highlighted that one in five delivered urgently for respiratory compromise or were admitted to an intensive care unit ⁹. Based on a retrospective case series in the USA, 7 out of 46 pregnant women with Covid-19 necessitated hospitalization ⁷. This rate resulted even higher (69%) in a Spanish cases series of 60 pregnant infected subjects ⁸. In a large case series from China, only 6 out of 119 pregnant women (5%) were asymptomatic and in 9 cases (8%) the course of the disease was severe ⁵. Last but not least, Hantoushzadeh et al. described 9 cases of maternal death due to Covid-19 in Iran ⁶. As already pointed out in the introduction, the most plausible explanation for the discrepancy with our findings is a selection bias. The denominator is radically different. By mainly focusing on the presence of antibodies, we were able to study an unselected population. In contrast, published case series reported on

women who were mainly identified because of Covid–19 related symptoms. Our data could better reflect the real impact of Covid–19 in pregnancy. Nonetheless, we cannot definitely rule out that differences in the characteristics of the population, local environmental conditions and genetic variants of the virus may have also impacted on clinical relevance and could play a role in explaining these inconsistencies.

Some limitations of our study should be acknowledged. Firstly, the reliability of the diagnostic tests is still a source of debate. Even if Covid–19 was investigated using three different modalities, the accuracy of all the tests used is yet uncertain. The nasopharyngeal swab could detect only ongoing infections and the sensitivity in affected cases was reported to be only 63%¹³. Preliminary evidence is comforting for the other two tests employed to detect antibodies against Sars-Cov–2 but available studies for validation are not optimal. In particular, no attempts have yet been made to investigate the accuracy of these tests for asymptomatic or poorly symptomatic cases. Inferring results obtained in patients with moderate or severe forms of the disease to the whole population is arguable. More in general, the clinical and biological significance of the different type of antibodies remain to be ascertained^{10,14,15}. Noteworthy, in our experience, agreement between the two tests was not excellent since only 17 women were found to be positive to both tests while 7 were positive only to one of the two.

Secondly, even if women referred from other hospitals because of Covid–19 were excluded, we cannot rule out some other selection biases. On one hand, some healthy women with unremarkable history may have decided to deliver in other hospitals to avoid an Institution with Covid–19 affected cases while, on the other hand, some women with mild symptoms suggestive for the infection could have been more likely to refer to our hospital. Both biases could lead to overestimate the detected frequency of Covid–19. However, the high proportion of asymptomatic cases tends to rule out a major role of these confounders.

Finally, since Covid–19 related symptoms were retrospectively collected, one cannot exclude a recall bias. In this regard, it has however to be underlined that women were blinded to the results of the antibodies tests when interviewed and that the investigated period of time was limited to only three months. Even if episodes of mild symptoms could be overlooked, it is unlikely that more significant health troubles could be omitted.

In conclusion, women in the second half of pregnancy do not appear to be more susceptible to Covid–19; the observed prevalence overlaps with the non-pregnant population of the same area. In addition, the study suggests that the course of the disease is mostly unremarkable. However, this latter observation warrants confirmation in larger studies.

Declarations

Acknowledgments

We are in debt with Alessia Borgo and Elena Rossi and the whole team of midwives of the involved obstetrics wards for the precious collaboration in running the study

References

1. Grasselli G, Pesenti A, Cecconi M. Critical Care Utilization for the COVID–19 Outbreak in Lombardy, Italy: Early Experience and Forecast During an Emergency Response. *JAMA*. 2020 Mar 13. [Epub ahead of print]
2. Gandhi RT, Lynch JB, Del Rio C. Mild or Moderate Covid–19. *N Engl J Med*. 2020 Apr 24. [Epub ahead of print].
3. Grasselli G, Zangrillo A, Zanella A, Antonelli M, Cabrini L, Castelli A, Cereda D, Coluccello A, Foti G, Fumagalli R, Iotti G, Latronico N, Lorini L, Merler S, Natalini G, Piatti A, Ranieri MV, Scandroglio AM, Storti E, Cecconi M, Pesenti A; COVID–19 Lombardy ICU Network, Baseline Characteristics and Outcomes of 1591 Patients Infected With SARS-CoV–2 Admitted to ICUs of the Lombardy Region, Italy. *JAMA*. 323, 1574–81 (2020).
4. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DSC, Du B, Li LJ, Zeng G, Yuen KY, Chen RC, Tang CL, Wang T, Chen PY, Xiang J, Li SY, Wang JL, Liang ZJ, Peng YX, Wei L, Liu Y, Hu YH, Peng P, Wang JM, Liu JY, Chen Z, Li G, Zheng ZJ, Qiu SQ, Luo J, Ye CJ, Zhu SY, Zhong NS; China Medical Treatment Expert Group for Covid–19. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med*. 382, 1708–20 (2020).
5. Chen L, Li Q, Zheng D, Jiang H, Wei Y, Zou L, Feng L, Xiong G, Sun G, Wang H, Zhao Y, Qiao J. Clinical Characteristics of Pregnant Women with Covid–19 in Wuhan, China. *N Engl J Med*. 2020 Apr 17:NEJMc2009226. [Epub ahead of print].
6. Hantoushzadeh S, Shamshirsaz AA, Aleyasin A, Seferovic MD, Aski SK, Arian SE, Pooransari P, Ghotbizadeh F, Aalipour S, Soleimani Z, Naemi M, Molaei B, Ahangari R, Salehi M, Oskoei AD, Pirozan P, Darkhaneh RF, Laki MG, Farani AK, Atrak S, Miri MM, Koucheh M, Shojaei S, Hadavand F, Keikha F, Hosseini MS, Borna S, Ariana S, Shariat M, Fatemi A, Nouri B, Nekooghadam SM, Aagaard K. Maternal Death Due to COVID–19 Disease. *Am J Obstet Gynecol*. 2020 Apr 28 [Epub ahead of print]
7. Lokken EM, Walker CL, Delaney S, Kachikis A, Kretzer NM, Erickson A, Resnick R, Vanderhoeven J, Hwang JK, Barnhart N, Rah J, McCartney SA, Ma KK, Huebner EM, Thomas C, Sheng JS, Paek BW, Retzlaff K, Kline CR, Munson J, Blain M, Lacourse SM, Deutsch G, Adams Waldorf K. Clinical Characteristics of 46 Pregnant Women with a SARS-CoV–2 Infection in Washington State. *Am J Obstet Gynecol*. May 18:S0002–9378(20)30558–5. 2020
8. Pereira A, Cruz-Melguizo S, Adrien M, Fuentes L, Marin E, Perez-Medina T. Clinical course of Coronavirus Disease–2019 (COVID–19) in pregnancy. *Acta Obstet Gynecol Scand*. 2020 May 22. [Epub ahead of print].
9. Savasi VM, Parisi F, Patanè L, Ferrazzi E, Frigerio L, Pellegrino A, Spinillo A, Tateo S, Ottoboni M, Veronese P, Petraglia F, Vergani P, Facchinetti F, Spazzini D, Cetin I. Clinical Findings and Disease Severity in Hospitalized Pregnant Women With Coronavirus Disease 2019 (COVID–19). *Obstet Gynecol*. 2020 May 19 [Epub ahead of print].
10. French MA, Moodley Y. The role of SARS-CoV–2 antibodies in COVID–19: Healing in most, harm at times. *Respirology*. 2020 May 20. [Epub ahead of print].

11. Tassis B, Lunghi G, Frattaruolo MP, Ruggiero M, Somigliana E, Ferrazzi E. Effectiveness of a COVID–19 screening questionnaire for pregnant women at admission to an obstetric unit in Milan. *Int J Gynaecol Obstet*. 2020 May 5 [Epub ahead of print].
12. Valenti L, Bergna A, Pelusi S, Facciotti F, Lai A, Tarkowski M, Berzuini A, Caprioli F, Santoro L, Baselli G, Della Ventura C, Erba E, Bosari S, Galli M, Zehender G, Prati D. SARS-CoV–2 seroprevalence trends in healthy blood donors during the COVID–19 Milan outbreak. *medRxiv* [submitted]
13. Wang W, Xu Y, Gao R, Lu R, Han K, Wu G, Tan W. Detection of SARS-CoV–2 in Different Types of Clinical Specimens. *JAMA*. 323, 1843–4 (2020)
14. Mallapaty S. Will antibody tests for the coronavirus really change everything? *Nature*. 580, 571–2 (2020).
15. Sethuraman N, Jeremiah SS, Ryo A. Interpreting Diagnostic Tests for SARS-CoV–2. *JAMA*. 2020 May 6. [Epub ahead of print].

Tables

Table 1. Prevalence of women who had Covid-19 in pregnancy in the studied cohort (n=315) according to type of testing.

Diagnostic test	N	Rate	95%CI
Naso-pharyngeal swab RT-PCR	13	4.1%	2.4-6.9%
Antibodies against CoV-2 envelope	20	6.4%	4.2-9.6%
Antibodies against CoV-2 nucleoprotein	21	6.7%	4.4-10.0%
At least one	28	8.9%	6.2-12.5%

Table 2. Distribution of tests results among Covid-19 cases (n=28)

Positive test			N (%)
RNA-PCR	Anti-envelope	Anti-nucleoprotein	
+	-	-	4 (14%)
-	+	-	3 (11%)
-	-	+	3 (11%)
+	+	-	0 (0%)
+	-	+	1 (4%)
-	+	+	9 (32%)
+	+	+	8 (28%)

Table 3. Baseline characteristics of women who did and did not test positive for Covid-19

Characteristics	Covid-19 n=28	Controls n=287	p
Age (years)	31.6 ± 7.0	34.2 ± 5.4	0.07
BMI (Kg/m ²)	27.1 ± 4.5	26.4 ± 4.2	0.41
Ethnicity			0.08
Caucasian	20 (71.4%)	230 (80.1%)	
Black / African	2 (7.2%)	26 (9.1%)	
Asian	0 (0.0%)	9 (3.1%)	
Hispanic	6 (21.4%)	22 (7.7%)	
Smoking in pregnancy	1 (3.6%)	12 (4.2%)	1.00
ART pregnancy	1 (3.6%)	22 (7.7%)	0.71
Previous vaginal deliveries	7 (25.0%)	89 (31.0%)	0.67
Previous cesarean sections	5 (17.9%)	45 (15.7%)	0.79
Previous myomectomy	0 (0.0%)	5 (1.7%)	1.00
Main medical complications			
Thyroid disorders	0 (0.0%)	22 (7.7%)	0.24
Autoimmune disorders	0 (0.0%)	9 (3.1%)	1.00
Cardiovascular problems	2 (7.2%)	6 (2.1%)	0.15
Thrombophilia	0 (0.0%)	3 (1.0%)	1.00
Others	2 (7.2%)	8 (2.8%)	0.22
Multiple pregnancy	1 (3.6%)	10 (3.5%)	1.00
Flu vaccination in pregnancy	9 (32.1%)	96 (33.4%)	1.00

ART: Assisted Reproductive Techniques

Table 4. Symptoms and risk factors in women who did and did not test positive for Covid-19

Symptoms or risk conditions	Covid-19 n=28	Controls n=287	p
Symptoms suggestive for Covid-19			
Fever	5 (17.9%)	13 (4.5%)	0.015
Cough	7 (25.0%)	19 (6.6%)	0.004
Sore throat	3 (10.7%)	17 (5.9%)	0.40
Rhinitis	4 (14.3%)	19 (6.6%)	0.14
Headache	0 (0.0%)	0 (0.0%)	n.a.
Diarrhea	0 (0.0%)	2 (0.7%)	1.00
Vomit	1 (3.6%)	2 (0.7%)	0.24
Dyspnea	3 (10.7%)	0 (0.0%)	0.001
Asthenia	0 (0.0%)	2 (0.7%)	1.00
Myalgias	0 (0.0%)	2 (0.7%)	1.00
Anosmia	4 (14.3%)	1 (0.3%)	<0.001
Ageusia	2 (7.1)	4 (1.4%)	0.09
Episodes of symptoms suggestive for Covid-19			0.001
None	17 (60.7%)	248 (86.5%)	
1	10 (35.7%)	38 (13.2%)	
2	1 (3.6%)	1 (0.3%)	
Risk conditions			
Job (health worker, lab technician)	0 (0.0%)	3 (1.0%)	1.00
Direct contact with Covid-19 affected persons	1 (3.6%)	3 (1.0%)	0.31
Cohabitant with persons at risk	3 (10.7%)	31 (10.8%)	1.00

Table 5. Pregnancy outcome in women who did and did not test positive for Covid-19

Characteristics	Covid-19 n=28	Controls n=287	p
Gestational Diabetes Mellitus	3 (10.7%)	22 (7.7%)	0.48
Hypertensive disorders	2 (7.1%)	6 (2.1%)	0.15
Intrahepatic cholestasis	1 (3.6%)	5 (1.7%)	0.43
Preterm delivery (< 37 weeks' gestation)	2 (7.1%)	22 (7.7%)	1.00
Cesarean section	10 (35.7%)	108 (37.6%)	1.00
SGA newborns ^a	2 (7.4%)	30 (10.8%)	0.75
LGA newborns ^a	1 (3.7%)	23 (8.3%)	0.71
Stillbirth	0 (0.0%)	0 (0.0%)	n.a.
Neonatal death	0 (0.0%)	0 (0.0%)	n.a.
Maternal death	0 (0.0%)	0 (0.0%)	n.a.

^a Multiple pregnancies excluded

SGA: Small for gestational age. LGA: Large for Gestational Age.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [DatabaseRuggieroetal.xls](#)