Daily Precipitation Concentration in Central Coast Vietnam

Aziz Benhamrouche (✉ hydr21@yahoo.fr)

Universite Ferhat Abbas Setif 1 <https://orcid.org/0000-0001-7083-9328>

Javier Martin-Vide

University of Barcelona

Quoc Bao Pham

Ton Duc Thang University

Mostefa E. Kouachi

Ferhat Abbas Setif 1 University

M. Carmen Moreno-Garcia

University of Barcelona

Research Article

Keywords: Concentration Index, Daily precipitation, monsoon climate, Vietnam

Posted Date: August 31st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-346295/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Theoretical and Applied Climatology on October 13th, 2021. See the published version at <https://doi.org/10.1007/s00704-021-03804-9>.

Abstract

Empirical frequency distribution of daily precipitation amounts can be fitted by a negative exponential distribution, because anywhere there are many small daily totals and few large ones. Therefore, the cumulative percentages of days with precipitation, sorted in increasing order according to their amounts, against the cumulative percentage of the rainfall amounts that they contribute are fitted by positive exponential curves $Y = aX^b$, a and b constants. Based on these curves, the Concentration Index (CI) evaluates the contribution of the rainiest days to the total amount. In this study the CI has been calculated for 15 meteorological stations in Da Nang city and Quang Nam province in Central Coast Vietnam, for the 1979–2016 period. The results show high values of CI, ranging from 0.62 to 0.72. Conversely, the linear correlation between altitude and CI is negative ($R = -0.60$, $p < 0.01$). There are no correlations between the latitude nor the annual mean number of precipitation days and the CI. CI change for the sub-periods of 1979–1997 and 1998–2016 is also analyzed.

1. Introduction

The analysis of precipitation in a daily resolution is a subject of great climatic and environmental interest since the irregularity and concentration of rainfall is linked to a lot of phenomena as runoff, floods, drought, soil erosion, water resources, aquifer recharge etc. In all climates, heavy rainfall is concentrated in a few days per year, so the appearance or not of these days can change the character, dry or rainy, of a year. It is not uncommon that after several dry months a few very rainy days totalize more than normal amount. Water management and territorial planning should take into account the daily precipitation concentration for a better territory planning.

The concentration of daily precipitation has been recently studied in different countries and regions, even on a global scale. The main methodological approach for evaluating it is the Concentration Index (CI) proposed by Martin-Vide (2004). The CI has been calculated for Peninsular Spain and Spain (Martin-Vide, 2004; Serrano-Notivol *et al*, 2017), Iran (Alijani *et al*, 2008), Pearl River basin and Xinjiang (China) (Zhang *et al*, 2009; Li *et al*, 2011), central Andes of Peru (Zubieta, 2009), Europe (Cortesi *et al*, 2012), Peninsular Malaysia (Suhaila and Jemain, 2012), Southern Italy (Coscarelli and Caloiero, 2012), New Zealand (Caloiero, 2013), Chile and Central-Southern Chile (Sarricolea and Martin-Vide, 2014; Sarricolea *et al*, 2019), Algeria and Northeast of Algeria (Benhamrouche *et al*, 2015; Bessaklia *et al*, 2018), northern Morocco (Salhi *et al*, 2019), United States (Royé and Martin-Vide, 2017), Argentina (Llano, 2017), Southeastern Brazil (Nery *et al*, 2017), Southern Russia (Vyshkvarkova *et al*, 2018), Puerto Rico (Vélez *et al*, 2018) among others, and finally at a global scale (Monjo and Martin-Vide, 2016).

The analysis of the daily precipitation concentration has special interest in the intertropical zonewhere heavy rains and showers are frequent. They can cause landslides, heavy soil leaching and other damages particularly in Southeastern Asia with rainy tropical climates and a high population density. This article focuses on the central coast of Vietnam, specifically Da Nang city and Quang Nam province.

The article is organized as follows: Sect. 2 establishes the objectives and shows the methodology; the data and the study area are described in Sect. 3; the results are detailed in Sect. 4 and are then discussed in Sect. 5; finally, the conclusions are in Sect. 6.

2. Objectives And Methodology

The main aim of this study is to evaluate the daily concentration of precipitation in Central Coast Vietnam i.e. evaluate the weight of the rainiest days to the annual total. The specific objectives are: 1) calculate the values of the CI for 15 meteorological stations covering the study area; 2) establish the spatial patterns of the CI (correlations with latitude, longitude and altitude); and 3) compare the CI of the two 19-years sub-periods 1979–1997 and 1998–2016.

Apart from linear correlations and comparison between means, the main method used is the CI. The first step for the CI calculation is to sort the daily precipitation values in ascending order. Then the cumulative percentage of rainy days is plotted against the corresponding cumulative percentage of rainfall amounts. The plot has a markedly positive exponential shape. These exponential curves, also called standardized precipitation curves (Jolliffe and Hope, 1996), specifically suggested by Riehl (1949) and Olascoaga (1950), are of this kind:

$$y = a x^b \quad (1)$$

where a and b are constants, that can be calculated by means of the least-squares procedure (Martin-Vide, 2004):

$$\ln a = \frac{\sum x_i^2 \sum \ln y_i + \sum x_i \sum x_i \ln x_i - \sum x_i^2 \sum \ln x_i - \sum x_i \sum x_i \ln y_i}{N \sum x_i^2 - (\sum x_i)^2} \quad (2)$$

$$b = \frac{N \sum x_i \ln y_i + \sum x_i \sum \ln x_i - N \sum x_i \ln x_i - \sum x_i \sum \ln y_i}{N \sum x_i^2 - (\sum x_i)^2} \quad (3)$$

In Table 1 an example is developed.

Table 1: Frequency distribution in classes of 1 mm, where x_i is the accumulated relative frequencies in percentage, and y_i is the corresponding accumulated percentage of total precipitation, of Que Son meteorological station (1979–2016).

Super limit class	Midpoint (m_i)	n_i	$\sum n_i$	$m_i \cdot n_i$	$\sum m_i \cdot n_i$	$\% \sum n_i = x_i$	$\% \sum m_i \cdot n_i = y_i$
0.9	0.5	202	202	101	101	3.604568	0.100388
1.9	1.5	1490	1692	2274.187	2375.187	30.19272	2.360793
2.9	2.5	415	2107	1037.5	3412.687	37.59814	3.392006
3.9	3.5	289	2396	1012.28	4424.967	42.75517	4.398152
4.9	4.5	224	2620	1014.16	5439.127	46.75232	5.406166
5.9	5.5	231	2851	1270.5	6709.627	50.87438	6.668967
6.9	6.5	176	3027	1144.686	7854.314	54.01499	7.806716
7.9	7.5	161	3188	1212.121	9066.434	56.88794	9.011491
8.9	8.5	127	3315	1079.5	10145.93	59.15418	10.08445
9.9	9.5	120	3435	1140	11285.93	61.2955	11.21754
10.9	10.5	164	3599	1726.904	13012.84	64.22198	12.93398
11.9	11.5	88	3687	1012	14024.84	65.79229	13.93985
12.9	12.5	104	3791	1300.655	15325.49	67.64811	15.23262
13.9	13.5	95	3886	1285.455	16610.95	69.34333	16.51028
14.9	14.5	65	3951	942.5	17553.45	70.50321	17.44707
15.9	15.5	98	4049	1519	19072.45	72.25196	18.95687
16.9	16.5	84	4133	1388.713	20461.16	73.75089	20.33717
17.9	17.5	48	4181	840	21301.16	74.60742	21.17207
18.9	18.5	55	4236	1017.5	22318.66	75.58887	22.18341
19.9	19.5	55	4291	1072.5	23391.16	76.57031	23.24941
20.9	20.9	70	4361	1461.957	24853.12	77.81941	24.70251
21.9	21.5	43	4404	924.5	25777.62	78.58672	25.6214
22.9	22.5	45	4449	1014.062	26791.68	79.38972	26.62932
23.9	23.5	40	4489	940	27731.68	80.1035	27.56362
24.9	24.5	42	4531	1029	28760.68	80.85296	28.58639
25.9	25.5	44	4575	1122	29882.68	81.63812	29.70159
26.9	26.5	28	4603	742	30624.68	82.13776	30.43909

Super limit class	Midpoint (m_i)	n_i	$\sum n_i$	$m_i \cdot n_i$	$\sum m_i \cdot n_i$	$\% \sum n_i = x_i$	$\% \sum m_i \cdot n_i = y_i$
27.9	27.5	31	4634	852.5	31477.18	82.69094	31.28643
28.9	28.5	27	4661	769.5	32246.68	83.17273	32.05126
29.9	29.5	23	4684	678.5	32925.18	83.58315	32.72565
30.9	30.5	34	4718	1037	33962.18	84.18986	33.75637
31.9	31.5	31	4749	976.5	34938.68	84.74304	34.72695
32.9	32.5	33	4782	1072.5	36011.18	85.33191	35.79295
33.9	33.5	21	4803	703.5	36714.68	85.70664	36.49219
34.9	34.5	14	4817	483.553	37198.23	85.95646	36.97281
35.9	35.5	23	4840	816.5	38014.73	86.36688	37.78436
36.9	36.5	22	4862	803	38817.73	86.75946	38.58249
37.9	37.5	22	4884	825	39642.73	87.15203	39.4025
38.9	38.5	19	4903	731.5	40374.23	87.49108	40.12956
39.9	39.5	17	4920	671.5	41045.73	87.79443	40.79699
40.9	40.5	23	4943	931.5	41977.23	88.20485	41.72285
41.9	41.5	24	4967	996	42973.23	88.63312	42.71281
42.9	42.5	21	4988	892.5	43865.73	89.00785	43.5999
43.9	43.5	15	5003	652.5	44518.23	89.27552	44.24845
44.9	44.5	12	5015	534	45052.23	89.48965	44.77921
45.9	45.5	21	5036	955.5	46007.73	89.86438	45.72892
46.9	46.5	16	5052	744	46751.73	90.14989	46.46841
47.9	47.5	14	5066	665	47416.73	90.39971	47.12938
48.9	48.5	15	5081	727.5	48144.23	90.66738	47.85248
49.9	49.5	10	5091	495	48639.23	90.84582	48.34448
50.9	50.5	20	5111	1010	49649.23	91.20271	49.34836
51.9	51.5	10	5121	515.219	50164.45	91.38116	49.86045
52.9	52.5	11	5132	577.5	50741.95	91.57744	50.43445
53.9	53.5	11	5143	588.5	51330.45	91.77373	51.01939

Super limit class	Midpoint (m_i)	n_i	$\sum n_i$	$m_i \cdot n_i$	$\sum m_i \cdot n_i$	$\% \sum n_i = x_i$	$\% \sum m_i \cdot n_i = y_i$
54.9	54.5	9	5152	490.7079	51821.16	91.93433	51.50712
55.9	55.5	15	5167	832.5	52653.66	92.202	52.33458
56.9	56.5	10	5177	565	53218.66	92.38044	52.89615
57.9	57.5	13	5190	747.5	53966.16	92.61242	53.63912
58.9	58.5	6	5196	351	54317.16	92.71949	53.98799
59.9	59.5	9	5205	535.5	54852.66	92.88009	54.52025
60.9	60.5	10	5215	605	55457.66	93.05853	55.12158
61.9	61.5	8	5223	492	55949.66	93.20128	55.6106
62.9	62.5	8	5231	500.0152	56449.67	93.34404	56.10759
63.9	63.5	6	5237	381	56830.67	93.45111	56.48628
64.9	64.5	5	5242	322.5	57153.17	93.54033	56.80682
65.9	65.5	10	5252	655	57808.17	93.71877	57.45785
66.9	66.5	9	5261	598.5	58406.67	93.87937	58.05273
67.9	67.5	9	5270	607.5	59014.17	94.03997	58.65655
68.9	68.5	8	5278	548	59562.17	94.18273	59.20122
69.9	69.5	2	5280	139	59701.17	94.21842	59.33938
70.9	70.5	5	5285	352.5	60053.67	94.30764	59.68975
71.9	71.5	3	5288	214.5	60268.17	94.36117	59.90295
72.9	72.5	6	5294	435	60703.17	94.46824	60.33531
73.9	73.5	9	5303	661.5	61364.67	94.62884	60.9928
74.9	74.5	6	5309	447	61811.67	94.7359	61.43709
75.9	75.5	8	5317	604	62415.67	94.87866	62.03743
76.9	76.5	4	5321	306	62721.67	94.95004	62.34158
77.9	77.5	10	5331	775	63496.67	95.12848	63.11188
78.9	78.5	3	5334	235.5	63732.17	95.18201	63.34595
79.9	79.5	4	5338	318	64050.17	95.25339	63.66203
80.9	80.5	7	5345	563.5	64613.67	95.3783	64.22211

Super limit class	Midpoint (m_i)	n_i	$\sum n_i$	$m_i \cdot n_i$	$\sum m_i \cdot n_i$	$\% \sum n_i = x_i$	$\% \sum m_i \cdot n_i = y_i$
81.9	81.5	1	5346	81.5	64695.17	95.39615	64.30312
82.9	82.5	5	5351	412.5	65107.67	95.48537	64.71312
83.9	83.5	3	5354	250.5309	65358.21	95.5389	64.96213
84.9	84.5	5	5359	422.5	65780.71	95.62812	65.38207
85.9	85.5	5	5364	427.5	66208.21	95.71734	65.80698
86.9	86.5	4	5368	346	66554.21	95.78872	66.15088
87.9	87.5	5	5373	437.5	66991.71	95.87794	66.58573
88.9	88.5	2	5375	177	67168.71	95.91363	66.76166
89.9	89.5	2	5377	179	67347.71	95.94932	66.93957
90.9	90.5	3	5380	271.5	67619.21	96.00286	67.20943
91.9	91.5	4	5384	366	67985.21	96.07423	67.57321
92.9	92.5	6	5390	555	68540.21	96.1813	68.12485
93.9	93.5	2	5392	187	68727.21	96.21699	68.31071
94.9	94.5	5	5397	472.5	69199.71	96.30621	68.78035
95.9	95.5	6	5403	573	69772.71	96.41328	69.34988
96.9	96.5	3	5406	289.5	70062.21	96.46681	69.63762
97.9	97.5	3	5409	292.5	70354.71	96.52034	69.92835
98.9	98.5	5	5414	492.5	70847.21	96.60956	70.41787
99.9	99.5	4	5418	398	71245.21	96.68094	70.81346
100.9	100.5	6	5424	603	71848.21	96.78801	71.4128
101.9	101.5	4	5428	406	72254.21	96.85939	71.81634
102.9	102.5	5	5433	512.5	72766.71	96.94861	72.32574
103.9	103.5	6	5439	621	73387.71	97.05567	72.94297
104.9	104.5	4	5443	418.0748	73805.78	97.12705	73.35851
105.9	105.5	2	5445	211	74016.78	97.16274	73.56823
106.9	106.5	3	5448	319.5	74336.28	97.21627	73.8858
107.9	107.5	4	5452	430	74766.28	97.28765	74.31319

Super limit class	Midpoint (m_i)	n_i	$\sum n_i$	$m_i \cdot n_i$	$\sum m_i \cdot n_i$	$\% \sum n_i =$ x_i	$\% \sum m_i \cdot n_i$ $= y_i$
108.9	108.5	4	5456	434	75200.28	97.35903	74.74456
109.9	109.5	3	5459	328.5	75528.78	97.41256	75.07107
111.9	111.5	4	5463	446	75974.78	97.48394	75.51437
112.9	112.5	2	5465	225	76199.78	97.51963	75.73801
113.9	113.5	7	5472	794.5	76994.28	97.64454	76.52769
114.9	114.5	2	5474	229	77223.28	97.68023	76.7553
115.9	115.5	1	5475	115.5	77338.78	97.69807	76.8701
116.9	116.5	3	5478	349.5	77688.28	97.75161	77.21749
117.9	117.5	4	5482	470	78158.28	97.82298	77.68464
118.9	118.5	5	5487	592.5	78750.78	97.91221	78.27355
119.9	119.5	2	5489	239	78989.78	97.94789	78.5111
120.9	120.5	5	5494	602.5	79592.28	98.03712	79.10995
121.9	121.5	4	5498	486	80078.28	98.10849	79.593
123.9	123.5	2	5500	247	80325.28	98.14418	79.8385
124.9	124.5	1	5501	124.5	80449.78	98.16203	79.96225
125.9	125.5	2	5503	251	80700.78	98.19772	80.21173
128.9	128.5	1	5504	128.5	80829.28	98.21556	80.33945
129.9	129.5	2	5506	259	81088.28	98.25125	80.59688
130.9	130.5	1	5507	130.5	81218.78	98.26909	80.72659
131.9	131.5	2	5509	263	81481.78	98.30478	80.988
132.9	132.5	4	5513	530	82011.78	98.37616	81.51478
133.9	133.5	1	5514	133.5	82145.28	98.394	81.64748
135.9	135.5	2	5516	271	82416.28	98.42969	81.91683
136.9	136.5	3	5519	409.5213	82825.8	98.48323	82.32387
137.9	137.5	3	5522	412.5	83238.3	98.53676	82.73387
139.9	139.5	3	5525	418.5	83656.8	98.59029	83.14984
140.9	140.5	2	5527	281	83937.8	98.62598	83.42913

Super limit class	Midpoint (m_i)	n_i	$\sum n_i$	$m_i \cdot n_i$	$\sum m_i \cdot n_i$	$\% \sum n_i =$ x_i	$\% \sum m_i \cdot n_i$ $= y_i$
141.9	141.5	1	5528	141.5	84079.3	98.64383	83.56978
142.9	142.5	2	5530	285	84364.3	98.67951	83.85305
144.9	144.5	1	5531	144.5	84508.8	98.69736	83.99667
145.9	145.5	1	5532	145.5	84654.3	98.7152	84.14129
146.9	146.5	1	5533	146.5	84800.8	98.73305	84.2869
147.9	147.5	2	5535	295	85095.8	98.76874	84.58012
149.9	149.5	3	5538	448.5	85544.3	98.82227	85.0259
152.9	152.5	1	5539	152.5	85696.8	98.84011	85.17747
154.9	154.5	2	5541	309	86005.8	98.8758	85.4846
155.9	155.5	1	5542	155.5	86161.3	98.89365	85.63916
156.9	156.5	1	5543	156.5	86317.8	98.91149	85.79471
157.9	157.5	1	5544	157.5	86475.3	98.92934	85.95126
159.9	159.5	1	5545	159.5	86634.8	98.94718	86.10979
160.9	160.5	1	5546	160.5	86795.3	98.96502	86.26932
161.9	161.5	1	5547	161.5	86956.8	98.98287	86.42984
162.9	162.5	1	5548	162.5	87119.3	99.00071	86.59135
163.9	163.5	1	5549	163.5	87282.8	99.01856	86.75386
164.9	164.5	1	5550	164.5	87447.3	99.0364	86.91737
165.9	165.5	2	5552	331	87778.3	99.07209	87.24636
166.9	166.5	1	5553	166.5	87944.8	99.08994	87.41185
167.9	167.5	1	5554	167.5	88112.3	99.10778	87.57834
168.9	168.5	2	5556	337	88449.3	99.14347	87.91329
171.9	171.5	1	5557	171.5	88620.8	99.16131	88.08375
172.9	172.5	1	5558	172.5	88793.3	99.17916	88.25521
174.9	174.5	1	5559	174.5	88967.8	99.197	88.42865
175.9	175.5	3	5562	526.5	89494.3	99.25054	88.95196
176.9	176.5	1	5563	176.5	89670.8	99.26838	89.12739

Super limit class	Midpoint (m_i)	n_i	$\sum n_i$	$m_i \cdot n_i$	$\sum m_i \cdot n_i$	$\% \sum n_i = x_i$	$\% \sum m_i \cdot n_i = y_i$
181.9	181.5	1	5564	181.5	89852.3	99.28622	89.30779
182.9	182.5	1	5565	182.5	90034.8	99.30407	89.48919
185.9	185.5	1	5566	185.5	90220.3	99.32191	89.67356
192.9	192.5	1	5567	192.5	90412.8	99.33976	89.8649
194.9	194.5	1	5568	194.5	90607.3	99.3576	90.05822
195.9	195.5	1	5569	195.5	90802.8	99.37545	90.25253
196.9	196.5	1	5570	196.5	90999.3	99.39329	90.44784
197.9	197.5	1	5571	197.5	91196.8	99.41113	90.64414
200.9	200.5	1	5572	200.5	91397.3	99.42898	90.84343
205.9	205.5	1	5573	205.5	91602.8	99.44682	91.04768
209.9	209.5	2	5575	419	92021.8	99.48251	91.46414
216.9	216.5	1	5576	216.5	92238.3	99.50036	91.67933
219.9	219.5	1	5577	219.5	92457.8	99.5182	91.8975
220.9	220.5	1	5578	220.5	92678.3	99.53605	92.11667
229.9	229.9	1	5579	229.8711	92908.17	99.55389	92.34514
238.9	238.5	1	5580	238.5	93146.67	99.57173	92.5822
241.9	241.5	1	5581	241.5	93388.17	99.58958	92.82224
249.9	249.5	1	5582	249.5	93637.67	99.60742	93.07022
255.9	255.5	1	5583	255.5	93893.17	99.62527	93.32418
258.9	258.5	1	5584	258.5	94151.67	99.64311	93.58111
260.9	260.5	1	5585	260.5	94412.17	99.66096	93.84003
263.9	263.5	1	5586	263.5	94675.67	99.6788	94.10193
265.9	265.5	1	5587	265.5	94941.17	99.69665	94.36582
270.9	270.5	1	5588	270.5	95211.67	99.71449	94.63468
272.9	272.5	1	5589	272.5	95484.17	99.73233	94.90553
275.9	275.5	1	5590	275.5	95759.67	99.75018	95.17936
280.9	280.5	1	5591	280.5	96040.17	99.76802	95.45816

Super limit class	Midpoint (m_i)	n_i	$\sum n_i$	$m_i \cdot n_i$	$\sum m_i \cdot n_i$	$\% \sum n_i = x_i$	$\% \sum m_i \cdot n_i = y_i$
281.9	281.5	1	5592	281.5	96321.67	99.78587	95.73796
292.9	292.5	1	5593	292.5	96614.17	99.80371	96.02869
294.9	294.5	1	5594	294.5	96908.67	99.82156	96.3214
296.9	296.5	1	5595	296.5	97205.17	99.8394	96.6161
300.9	300.5	1	5596	300.5	97505.67	99.85724	96.91478
329.9	329.5	1	5597	329.5	97835.17	99.87509	97.24229
333.9	333.5	1	5598	333.5	98168.67	99.89293	97.57377
341.9	341.5	1	5599	341.5	98510.17	99.91078	97.9132
352.9	352.5	1	5600	352.5	98862.67	99.92862	98.26356
395.9	395.5	1	5601	395.5	99258.17	99.94647	98.65666
411.9	411.5	1	5602	411.5	99669.67	99.96431	99.06567
467.9	467.5	1	5603	467.5	100137.2	99.98216	99.53034
472.9	472.5	1	5604	472.5	100609.7	100	100

The definite integral of the exponential curve between 0 and 100 is the area, S , under the curve, and 5,000 - S is the area compressed by the curve, $y = x$, and $x = 100$ (5,000 is the area of the triangle in which it is inserted), let's name S' . Then CI is defined as (Martin-Vide, 2004):

$$CI = S' / 5000 \quad (4)$$

The CI measures the relative separation of the exponential curve from the $Y = X$ line. Note that the separation is greater (or CI is higher), when the weight of a few days, the rainiest ones, in the total is higher (Martín-Vide, 2004; Cortesi et al., 2012; Benhamrouche et al., 2015). The value of the CI oscillates between 0 and 1 and it is similar to the well-known Gini index (Monjo and Martin-Vide, 2016). Figure 1 shows the positive exponential curves of the two extreme CI cases.

It is necessary to remember that the CI is very sensitive to the quality of data, as well as the longitude of classes. In general, the CI of meteorological station with no complete daily data should not be analyzed. On the other hand, calculating the CI from classes of, for example, 5 mm or 10 mm, instead 1 mm, produce a diminution of CI values (Benhamrouche and Martin-Vide, 2018; Nery et al, 2017).

Inverse Distance Weighting (IDW) interpolation was performed using 15 measurement points (stations) in ArcMap 10.5 to generate the spatial distribution of daily precipitation Concentration Index over the study

area for the periods 1979–1997, 1998–2016 and 1979–2016.

3. Dataset And Study Area

The data used are the daily precipitation amounts recorded from 1979 to 2016 (38 years) in 15 meteorological stations located in the Da Nang city and Quang Nam province in Central Coast Vietnam (Table 2). These stations belong to the Vietnamese National Center for Hydro-meteorological Forecasting (NCHMF).

Table 2
Meteorological stations and their geographical coordinates.

Name	Longitude	Latitude	Elevation (m a.s.l)
Da Nang	108.207	16.045	4.75
Cam Le	108.206	16.013	11
Hoi An	108.365	15.87	9
Cau Lau	108.279	15.86	2
Giao Thuy	108.127	15.839	9
Ai Nghia	108.104	15.874	16
Hoi Khanh	107.919	15.827	44
Hien	107.646	15.917	490
Thanh My	107.833	15.767	25
Nong Son	108.033	15.7	27
Que Son	108.1	15.7	19
Hiep Duc	108.102	15.578	22
Tien Phuoc	108.3	15.483	55
Kham Duc	107.79	15.427	396
Tra My	108.233	15.35	123.11

Data quality control has been carried out by the National Center for Hydro-meteorological Forecasting (NCHMF), Vietnam.

The quite similar CI values obtained are also a guarantee of its quality. The study area is located on the coast of the central Vietnam, between parallels 15° N and a bit more than 16° N in front of the South China Sea. The climate is predominantly tropical of monsoon, *Am* Köppen's type, although in the highlands farthest from the sea is temperate. The temperatures are high; between 20 and 30 °C all the

monthly averages at sea level. The precipitation is abundant, usually of about or above 2,000 mm. The annual regime shows a maximum in October, with about 600mm (rainy season: from September to December), and a minimum in March, with less than 30 mm (dry season: from February to April). The mean annual relative humidity is over 80%. It is important to highlight the high sea surface temperatures, close to 30°C during the hottest months.

The study area covers 11,700 km² (1,260 km² belongs to Da Nang city and 10,440 km² to Quang Nam province) and has more than 2.1 million people (about 180 inhabitants per square km). The new crops and settlements, and in general the increasing land uses, are sensitive to precipitation concentration.

4. Results

CI values for the 15 stations in the period 1979–2016 are presented in Table 3. They range from 0.62 (in Hien station) to 0.72 (the highest value, in Da Nang, the lowest and closest to the East sea). The mean value of the 15 stations is 0.68, which is considered a high value.

Table 3
CI values for the period 1979–2016 in 15 rain gauges of the study area.

Station	CI value	Station	CI value
Da Nang	0.72	Hien	0.62
Cam Le	0.70	Thanh My	0.67
Hoi An	0.70	Nong Son	0.66
Cau Lau	0.70	Que Son	0.67
Giao Thuy	0.67	Hiep Duc	0.67
Ai Nghia	0.69	Tien Phuoc	0.65
Hoi Khanh	0.67	Kham Duc	0.67
Tra My	0.69		

The spatial pattern of the CI values (Fig. 4) shows increasing values from west to east as from the mountain to the sea. This is evidenced by a statistical positive significant correlation between eastern longitude and CI: $R = +0.65$, $p < 0.01$ (Fig. 3). Conversely, there is no correlation between latitude and CI. The lowland has the high CI values, so there is a statistical negative significant correlation between altitude and CI: $R = -0.60$, $p < 0.01$ (Fig. 4).

The two 19-years sub-periods i.e. 1979–1997 and 1998–2016 have been considered in order to detect any significant trend (Figs. 7, 8 and 9). 11 out of 15 stations increased the CI value from the first sub-period to the second one, and the mean value went from 0.666 to 0.682, that constitutes a non-negligible change.

5. Discussion

The CI proposed by Martin-Vide (2004) has been used in many countries to evaluate the pluviometric weight of the rainiest days in the total amount, i.e. the concentration of a high percentage of precipitation amount in a few very rainy days. It is also an index that can give us a better picture about erosivity or aggressiveness of precipitation and its effects on soil erosion than the usual indices based on monthly precipitation (Bessaklia et al, 2018), such as the monthly precipitation concentration index developed by Oliver (1980) and the modified Fournier Index developed by Arnoldus (1980). This is an important subject in many countries because erosion reduces soil fertility, modifies the conditions for crops, alters agricultural practices and causes the rapid clogging of reservoirs (Scholz et al, 2008; Ibisate et al, 2016). There are two country groups more affected by soil erosion: very arid countries with poor vegetation cover and countries with high precipitation. Vietnam belongs to the second group, so it is interesting to evaluate the daily precipitation concentration.

The values obtained for Central Coast Vietnam (Da Nang and Quang Nam) between 0.62 and 0.72 can be considered high. As a reference, the threshold that separates high CI values from moderate ones is 0.61 (Martin-Vide, 2004). The value of 0.61 means that 25% of the rainiest days account for approximately 70% of the total amount. CI values higher than 0.66 mean that 25% of the rainiest days represent at least approximately 75%, as occurs in most of the stations analyzed. The Vietnam CI values are not so high than the calculated for the Pearl River basin in China (Zhang et al, 2009), over 0.74, but higher in general terms than the obtained in central and north Europe (Cortesi et al, 2012), with oceanic and continental mid latitudes climates. The CI values for Vietnam are quite similar to the Mediterranean regions ones, as those of a large part of Spain (Martin-Vide, 2004; Serrano-Notivoli et al, 2018), central Chile (Sarricolea et al, 2019), as well as some southern USA states (Texas, California) (Royé and Martin-Vide, 2017) and other countries. This is consistent with the fact that subtropical and tropical regions have higher CI than the mid and high latitudes countries (Monjo and Martin-Vide, 2016), although there is any exception such as northwest Puerto Rico with relatively low values (Vélez et al, 2018).

With respect to the most critical areas, in the study area the meteorological stations closer to the sea are those that have the highest CI values, as shown by the statistically significant correlation obtained between eastern longitude and CI, $R = +0.65$, $p < 0.01$. This is also true in the case of peninsular Spain in relation to the Mediterranean sea (Martin-Vide, 2004; Serrano-Notivoli et al, 2018). Heavy rainfall during the humid monsoon period and other intense tropical showers (including the rain produced by typhoons) contributes to the high CI values of Vietnam. The sea humid air flows are particularly effective in coastal sectors, where the CI values are the highest.

Contrary to what happens in Vietnam in many of the studied regions the correlation between the number of rainy days and CI is negative and statistically significant, as for instance occur in United States ($R = -0.25$, $p < 0.001$, using a grid with 13,527 cells) (Royé and Martin-Vide, 2017), Europe ($R = -0.68$, $p < 0.01$, for 530 meteorological stations) (Cortesi et al, 2012) and other areas and countries. In general, higher the number of rainy days, lower the CI value. Nevertheless, for the whole planet, at a global scale, using the

Gini index of daily precipitation, similar to the CI one, there are regions with a few rainy days with relatively low CI values, and the opposite as occurs in Vietnam (Monjo and Martin-Vide, 2016).

Respect to the relationship between altitude and CI, in general the correlation is also negative and significant, in correspondence with what was obtained for Vietnam, $R = -0.60$, $p < 0.01$ ($R = -0.51$, $p < 0.001$, for United States).

Several of the CI studies analyses the recent changes in CI. Most of them show predominant increasing trends, even significant ones, as is the case of Spain (Serrano-Notivol et al, 2018), United States (Royé and Martin-Vide, 2017), Mediterranean basin (Mathbout *et al*, 2019), etc. In Vietnam, the southern part of the country exhibits a dramatic increase in heavy rainfall events in the wet seasons (Ho-Hagemann et al, 2011) that could explain the increase of CI values obtained in the study area. For the years 2001–2050, based on the IPCC SRES A1B and A2 scenarios, heavy rainfall events tend to remain unchanged or to decrease for all Vietnam subregions except R5, where the study area is located (Ho-Hagemann, 2011).

6. Conclusions

The values of Concentration Index (CI) of daily precipitation in Central Coast Vietnam are high, between 0.62 and 0.72. Over the study area, only 25% of the rainiest days account for more than 70% of annual total amount, even in most of the stations analyzed the mentioned 25% represent at least 75% of the total amount –in Da Nang the 80% approximately. Heavy rainfall during the humid monsoon period and other heavy tropical showers (including the rain produced by typhoons) are the most likely causes of the high CI values of Vietnam.

Locations closer to the sea are those that have the highest CI values, as shown by the statistically significant correlation obtained between eastern longitude and CI. The lowest locations are also those that have the highest CI values, as shown by the negative significant correlation obtained between altitude and CI. In general terms, the CI has increased somewhat between the two 19-year sub-periods that make up the study period.

7. Declarations

Acknowledgments

We wish to acknowledge the support received from the Spanish Ministry project CLICES CGL2017-83866-C3-2-R and Climatology Group 2017 SGR 1362. We appreciate the interest in our research shown by the Water Research Institute of the University of Barcelona and the Royal Academy of Science and Arts of Barcelona.

Conflict of Interest: This manuscript has not been published or presented elsewhere in part or in entirety and is not under consideration by another journal. There are no conflicts of interest to declare.

Funding: No external funding

Authors Contributions

Aziz Benhamrouche: Project administration, Conceptualization, Writing- original draft, Software, Formal analysis, Visualization. **Javier Martin-Vide, Quoc Bao Pham:** Formal analysis; Writing- original draft, Visualization. **Mostefa E. Kouachi and M. Carmen Moreno-Garcia:** Data curation, Writing, Review and editing.

Availability of data and materials: The data that support the findings of this study are available from the first author, [Quoc Bao Pham, phambaoquoc@tdtu.edu.vn], upon reasonable request.

Code availability: It is available from the first author, upon reasonable request.

Ethical Approval: Not applicable

Consent to Participate: Not applicable

Consent to Publish: Not applicable

8. References

1. Alijani, B., O'Brien, J., and Yarnal, B. (2008). Spatial analysis of precipitation intensity and concentration in Iran, *Theor. Appl. Climatol.* 94, 107–124.
2. Arnoldus, H.M.J. 1980. An approximation of the rainfall factor in the USLE. In: Assessment of erosion. Eds. M.De Boodt, D. Gabriels. Chichester, England. John Wiley and Sons Ltd p. 127–132.
3. Benhamrouche, A. and Martin-Vide, J. (2012): Avances metodológicos en el análisis de la concentración diaria de la precipitación en la España Peninsular. *Anales de Geografía de la Universidad Complutense*, 32, 1, 11-27. https://doi.org/10.5209/rev_AGUC.2012.v32.n1.39306.
4. Benhamrouche, A., Boucherf, D., Hamadache, R., Bendahmane, L., Martín-Vide, J. and Teixeira Nery, J. (2015). Spatial distribution of the daily precipitation concentration index in Algeria. *Natural Hazards and Earth System Sciences*, 15, 617–625.
5. Bessaklia H., Ghenim A.N., Megnounif A., Martin-Vide J. (2018). Spatial variability of concentration and aggressiveness of precipitation in North-East of Algeria. *Journal of Water and Land Development*, 36, 3–15. doi: 10.2478/jwld-2018-0001.
6. Caloiero, T., 2014. Analysis of daily rainfall concentration in New Zealand. *Nat. Hazards* 72, 389–404. <https://doi.org/10.1007/s11069-013-1015-1>.
7. Cortesi, N., González-Hidalgo, J. C., Brunetti, M., and MartinVide, J. (2012): Daily precipitation concentration across Europe 1971–2010, *Nat. Hazards Earth Syst. Sci.*, 12, 2799–2810, doi:10.5194/nhess-12-2799-2012.

8. Coscarelli, R. and Caloiero, T. (2012). Analysis of daily and monthly rainfall concentration in southern Italy (Calabria region). *Journal of Hydrology*, 416–417, 145–156.
9. Ho-Hagemann, Ha & Phan-Van, Tan & Quan, Le & Nguyen, Q.. (2011). Extreme climatic events over Vietnam from observational data and RegCM3 projections. *Climate Research*, 49, 87-100.
10. Ibisate, A.; Ollero, A.; Ballarín, D.; Horacio, J.; Mora, D.; Mesanza, A.; Ferrer-Boix, C.; Acín, V.; Granado, D.; Martín-Vide, J.P. (2016). Geomorphic monitoring and response to two dam removals: Urumea and Leizaran (Basque Country, Spain). *Earth Surface Processes and Landforms*, 41, 2239-2255.
11. Jolliffe, I. T. and Hope, P. B. (1996). Representation of daily rainfall distributions using normalized rainfall curves. *International Journal of Climatology*. 16, 10 : 1157-1163
12. Li, X.X., Jiang, F.F., Li, L.L., Wang, G.G., 2011. Spatial and temporal variability of precipitation concentration index, concentration degree and concentration period in
13. Xinjian. *Int. J. Climatol.* 31, 1679–1693. <https://doi.org/10.1002/joc.2181>.
14. Llano, M.P., 2018. Spatial distribution of the daily rainfall concentration index in
15. Argentina: comparison with other countries. *Theor. Appl. Climatol.* 133, 997–1007.
16. <https://doi.org/10.1007/s00704-017-2236-0>.
17. Martin-Vide, J. (2004). Spatial distribution of a daily precipitation concentration index in Peninsular Spain. *International Journal of Climatology*, 24, 8, 959-971.
18. Mathbout, S.; Lopez-Bustins, J.A.; Royé, D.; Martin-Vide, J. and Benhamrouche, A. (2020): Spatiotemporal variability of daily precipitation concentration and its relationship to teleconnection patterns over the Mediterranean during 1975-2015. *International Journal of Climatology*, 40, 3, 1435-1455, [10.1002/joc.6278](https://doi.org/10.1002/joc.6278).
19. Monjo, R. and Martin-Vide, J. (2016). Daily precipitation concentration around the world according to several indices. *International Journal of Climatology*, 36(11), 2828–3838.
20. Nery, J.T., Carfan, A.C. and Martin-Vide, J. (2017). Analysis of rain variability using the daily and monthly concentration indexes in southeastern Brazil. *Atmospheric and Climate Sciences*, 7, 176–190.
21. Royé, D. and Martin, V.J. (2017). Concentration of daily precipitation in the contiguous United States. *Atmospheric Research*, 196, 237–247.
22. Olascoaga, M. J. (1950). Some aspects of Argentine rainfall. *Tellus*. 2, 4, 312-318.
23. Oliver, J.E. 1980. Monthly precipitation distribution: A comparative index. *The Professional Geographer*, 32. Iss. 3 p. 300–309.
24. Riehl, H. (1949). Some aspects of Hawaiian rainfall. *Bulletin of the American Meteorological Society*. 30, 5, 176-187.
25. Salhi, A.; Martin-Vide, J.; Benhamrouche, A.; Benabdelouahab, S.; Himi, M.; Benabdelouahab, T.; Casas Ponsati, A. (2019): Rainfall distribution and trends of the daily precipitation concentration index in northern Morocco: a need for an adaptive environmental policy. *SN Applied Sciences*, 1:277, <https://doi.org/10.1007/s42452-019-0290-1>.

26. Sarricolea, P, Martin-Vide, J., 2014. Spatial analysis of rainfall daily trends and concentration in Chile. *Investig. Geogr. Chile*, 47, 53–66. <http://refhub.elsevier.com/>
27. S0169-8095(17)30883-9/rf0115.
28. Sarricolea, P; Meseguer-Ruiz, O.; Serrano-Notivoli, R.; Soto, M.V.; Martin-Vide, J. (2019): Trends of daily precipitation concentration in Central-Southern Chile. *Atmospheric Research*, 215, 85-98, DOI: 10.1016/j.atmosres.2018.09.005
29. Scholz, G.; Quinton, J.N.; Strauss, P. 2008. Soil erosion from sugar beet in Central Europe in response to climate change induced seasonal precipitation variations. *Catena*, 72, 91–105.
30. Serrano-Notivoli, R., Martin-Vide, J., Saz, M.A., Longares, L.A., Beguería, S., Sarricolea, P, Meseguer-Ruiz, O., de Luis, M. (2018). Spatio-temporal variability of daily precipitation concentration in Spain based on a high-resolution gridded data set. *Int. J.Climatol.* 38, e518–e530. <https://doi.org/10.1002/joc.5387>.
31. Suhaila, J. and Jemain, A.A., (2012). Spatial analysis of daily rainfall intensity and concentration index in Peninsular Malaysia. *Theor. Appl. Climatol.* 108, 235–245.
32. doi.org/10.1007/s00704-011-0529-2.
33. Vélez, A.; Martin-Vide, J.; Royé, D.; Santaella, O. (2018): Spatial analysis of daily precipitation concentration in Puerto Rico. *Theoretical and Applied Climatology*, <https://doi.org/10.1007/s00704-018-2550-1>.
34. Vyshkvarkova, E., Voskresenskaya, E., Martin-Vide, J. (2018). Spatial distribution of the daily precipitation concentration index in southern Russia. *Atmospheric Research*, 203, 36–43.
35. Zhang, Q., Xu, C. Y., Gemmer, M., Chen, Y. Q., and Liu, C. L. (2009). Changing properties of precipitation concentration in the Pearl River basin, China, *Stoch. Env. Res. Risk A.*, 23, 377–385.
36. Zubieta, R., Saavedra, M., 2009. Spatial distribution of a daily precipitation concentration index in central andes of Perú, Mantaro River valley. *Tecnia* 19, 13–22.

Figures

Figure 1

Concentration curve of Que Son meteorological station (1979–2016)

Figure 2

Standardized precipitation curves and CI values of Hien Station and Da Nang (1979-2016).

Figure 3

Location of the 15 meteorological stations used in this study.

Figure 4

Spatial distribution of CI values for the 1979-2016 period.

Figure 5

Linear correlation between longitude and CI for the 1979-2016 period.

Figure 6

Linear correlation between altitude and CI for the 1979-2016 period.

Figure 7

Spatial distribution of CI values for the 1979-1997 sub-period.

Figure 8

Spatial distribution of CI values for 1998-2016 the sub-period.

Figure 9

CI differences between the 1979-1997 and 1998-2016 sub-periods.