

In Utero and Peripubertal Metals Exposure in Relation to Reproductive Hormones and Sexual Maturation and Progression among Boys in Mexico City

Pahriya Ashrap

University of Michigan School of Public Health

John D. Meeker

University of Michigan School of Public Health

Brisa N. Sánchez

University of Michigan School of Public Health

Niladri Basu

McGill University

Marcela Tamayo-Ortiz

Instituto Nacional de Salud Publica

Maritsa Solano-González

Instituto Nacional de Salud Publica

Adriana Mercado-García⁴

Instituto Nacional de Salud Publica

Martha M. Téllez-Rojo

Instituto Nacional de Salud Publica

Karen E. Peterson

University of Michigan School of Public Health

Deborah Watkins (✉ debjwat@umich.edu)

University of Michigan School of Public Health <https://orcid.org/0000-0002-6692-2206>

Research

Keywords: Metal, Hormone, In utero exposure, Pregnancy, Puberty

Posted Date: October 23rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-34705/v3>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Environmental Health on November 25th, 2020. See the published version at <https://doi.org/10.1186/s12940-020-00672-0>.

Abstract

Background: Endocrine disrupting chemicals (EDCs) such as metals have been reported to alter circulating reproductive hormone concentrations and pubertal development in animals. However, the relationship has rarely been investigated among humans, with the exception of heavy metals, such as Pb and Cd. Our aim was to investigate measures of *in utero* and peripubertal metal exposure in relation to reproductive hormone levels and sexual maturation and progression among boys from the Early Life Exposure in Mexico to Environmental Toxicants (ELEMENT) cohorts.

Methods: Our analysis included 118 pregnant women and their male children from the ELEMENT study. Essential and non-essential metals were measured in urine collected from the mothers during the third trimester of pregnancy and their male children at 8-14 years. Reproductive hormone levels [serum testosterone, estradiol, dehydroepiandrosterone sulfate (DHEA-S), inhibin B, and sex hormone-binding globulin (SHBG)] were measured in blood samples from the children at 8-14 years. We also assessed Tanner stages for sexual maturation (genital, pubic hair development, and testicular volume), at two time points (8–14, 10-18 years). We used linear regression to independently examine urinary metal levels in relation to each peripubertal reproductive hormones adjusting for child age and BMI. Generalized estimation equations (GEEs) were used to evaluate the association of *in utero* and peripubertal metal exposures with sexual maturation and progression during follow-up based on Tanner staging and testicular volume.

Results: *In utero* and prepubertal concentrations of some urinary metals were associated with increased levels of peripubertal reproductive hormones, especially non-essential metal(loid)s As and Cd (*in utero*), and Ba (peripubertal) as well as essential metal Mo (*in utero*) in association with testosterone. More advanced pubic hair developmental stage and higher testicular volume at the early teen visit was observed for boys with higher non-essential metal concentrations, including *in utero* Al and peripubertal Ba, and essential metal Zn concentration (peripubertal). These metals were also associated with slower pubertal progression between the two visits.

Conclusion: These findings suggest that male reproductive development may be associated with both essential and non-essential metal exposure during *in utero* and peripubertal windows.

Introduction

In recent decades, a trend toward earlier onset of puberty among boys and girls has been described (Anderson et al. 2003; Euling et al. 2008; Eveleth 2017; Herman-Giddens et al. 1997; Herman-Giddens et al. 2001; Karpati et al. 2002; Lee et al. 2016; Paris et al. 2016; Wacharasindhu 2017; Wyshak and Frisch 1982). These trends in the timing of puberty, a period of physical and psychological development, have raised concerns regarding the potential impact of environmental factors, including endocrine disrupting chemicals (EDCs) (Blakemore et al. 2010; Buck Louis et al. 2008; Lee and Styne 2013; Sisk and Foster 2004). Exposure to EDCs prenatally and at the prepubertal stage are thought to play a role in altered

pubertal timing, possibly via their estrogenic or anti-androgenic effects and disruption of normal homeostatic control of the hypothalamic-pituitary-gonadal (HPG) or hypothalamic-pituitary-adrenal (HPA) axis (Bellingham et al. 2009; Buck Louis et al. 2008; Den Hond and Schoeters 2006; Diamanti-Kandarakis et al. 2009; Fudvoye et al. 2019; Jacobson-Dickman and Lee 2009; Jansen et al. 2018; Y. Liu et al. 2019; Massart et al. 2006; Parent et al. 2016; Pescovitz and Walvoord 2007; Rodriguez et al. 2019; Roy et al. 2009; Watkins et al. 2016; Watkins et al. 2017).

Some metals and metalloids, such as cadmium (Cd), lead (Pb), mercury (Hg), and arsenic (As), are non-essential xenobiotics that are known to be harmful to human health (Atsdr 2007a, b; Faroon et al. 2012; Ha et al. 2017). These non-essential metal(loid)s are persistent in the environment and children's exposure to them is nearly ubiquitous (Claus Henn et al. 2014; World Health Organization 2017). Several other metals, such as chromium (Cr), copper (Cu), manganese (Mn), molybdenum (Mo), selenium (Se) and zinc (Zn), are essential for optimal health but may be harmful at insufficient or excessive levels (Gerberding 2005; Greger 1999; Health and Services 2004; Risher 2003; Shenkin 2003). A number of metal(loid)s are reproductive toxicants and have endocrine disrupting properties which interfere with many aspects of endocrine functions through interacting with hormone secretion, transport and binding receptors as well as genomic expression and epigenetic modification (Bloom et al. 2010; De Coster and van Larebeke 2012; Deb and Mandal 2017; Diamanti-Kandarakis et al. 2009; Laretta et al. 2019; Mendiola et al. 2011; Wallace 2019; Yilmaz et al. 2019). Some of these proposed mechanisms of actions are common to different metals, such as binding with estrogen receptor (Cd, As, Pb) and increasing lipid peroxidation (Pb, Hg), while others are specific, for instance, stimulation or inhibition of nuclear transcription activity (As) and inhibition of LH secretion (Pb) (Iavicoli et al. 2009).

Many human and animal studies have focused on elucidating the reproductive effects associated with heavy metal and metalloids, such as Cd, Pb, Hg, and As. Though many studies on Cd were either cross-sectional and/or conducted on adults, there has been some consistency of findings with regard to the positive relationship between male and female Cd exposure and testosterone levels (Bochud et al. 2017; Garcia-Morales et al. 1994; Jurasovic et al. 2004; Meeker et al. 2010; Menke et al. 2008; Nagata et al. 2005; Telisman et al. 2007; Zeng et al. 2002; Zeng et al. 2004). In a group of men with no occupational exposure, positive associations between blood Pb concentrations and testosterone and/or estradiol levels were reported (Jurasovic et al. 2004; Meeker et al. 2010; Telisman et al. 2007). In longitudinal studies of children's reproductive development, childhood Pb exposure was related to later pubertal onset (Hauser et al. 2008; Williams et al. 2010) and delayed sexual maturation (Williams et al. 2019) in Russian boys in Chapaevsk, Russia. In contrast, we have shown early life exposure to Pb was associated with delays in pubertal development in girls but not boys in Mexico City (Jansen 2018, Liu 2019). Hg was found to be associated with increased estradiol levels in both males and females from a small residential population in Cambodia (Agusa et al. 2007), which was in agreement with a previous study among women with repeated miscarriages (Gerhard et al. 1998). A recent study of As exposure through well water consumption in Taiwan suggested that As may impart an increased risk of erectile dysfunction through a reduction of circulating testosterone levels (Hsieh et al. 2008).

Although less attention has been given to the other metals in the past, a growing body of evidence suggests that certain essential or trace metals, including Cu, fluoride (F), Mn, Mo, and Se can also have adverse effects on male reproduction (Jeng et al. 2015; Yun Liu et al. 2019; Meeker et al. 2008; Telisman et al. 2000). In addition, most reports on the male reproductive effects of metals are from experimental animal, epidemiological, and occupational studies usually involving high doses not commonly encountered by children. Due to the widespread exposure of humans and known adverse effects related to essential and non-essential metal exposure, concern is growing that low-level exposure may also adversely affect reproductive developmental outcomes in boys. Moreover, only a few studies have investigated the cross-sectional relationships in boys, with the exception of two recent longitudinal studies in Russia and Mexico (Yun Liu et al. 2019; Williams et al. 2019), and none have examined exposure during *in utero* development and subsequent hormone levels during puberty, a time at which steroid hormones play an essential role in reproductive development (Cahill 2006; Parent et al. 2015; Sisk and Foster 2004; Spear 2000).

Therefore, the present study assessed whether *in utero* and prepubertal exposure to metals at relatively low doses altered reproductive hormone levels or timing and progression of sexual maturation in boys. We extended the limited metals studied on this topic and examined both essential and non-essential metals in relation to reproductive hormone levels and progression of sexual maturation in boys from ages 8-14 years to 10-18 years.

Method

2.1. Study population

Participants in this study are part of the “Early Life Exposure in Mexico to Environmental Toxicants (ELEMENT)” project, a longitudinal cohort study of pregnant women in Mexico City and their children (Perng et al. 2019). ELEMENT recruited 997 pregnant women from maternity hospitals during their first trimester between 1997 and 2004 and their children, as previously described (Ashrap et al. 2019; Lewis et al. 2013; Y. Liu et al. 2019; Perng et al. 2019; Wu et al. 2018). Inclusion criteria included not planning to leave the area within 5 years; no history of infertility, diabetes, or psychosis; not consuming alcoholic beverages daily during pregnancy; no addiction to illegal drugs; no diagnosis of a high-risk pregnancy; being pregnant with singleton. Mothers completed interview-based questionnaires at up to three prenatal visits [mean gestational age at visit 1: 13.5 (range:9–24) weeks, visit 2: 25.1 (range: 19–37) weeks, visit 3: 34.4 (range: 28–43) weeks] and provided spot urine samples at the third trimester visit. Between 2008 and 2011, a subset of their children (n=250), who were then 8–14 years of age, were contacted to participate in a follow-up study (i.e. early-teen visit). The criteria for eligibility included the availability of archived maternal biological specimens for toxicant assay interest (Perng et al. 2019). Between 2013–2017, 223 (89%) of those 250 children participated in a second follow-up study at age 10–18 years (i.e. late-teen visit). Among those 223 children, 109 were boys (92% retention rate from early to late-teen visit). Figure 1 shows the study design and the timing of biological sample collection and physical examination. Participants provided spot urine and fasting blood samples, anthropometry, and reported

socio-demographic information via an interviewer-administered questionnaire. In the current analyses, we included male children who finished the early-teen visit (a majority of these same boys also completed late-teen visit) and had maternal urinary metal concentration measurements and/or their early-teen visit urinary metal measurements available (n=118).

2.2. Metal Concentrations

Urinary metal concentrations of 14 metals and metalloids: aluminum (Al), As, barium (Ba), Cd, cobalt (Co), Cu, iron (Fe), Mn, Mo, nickel (Ni), Pb, antimony (Sb), selenium (Se), and Zn were measured in maternal third trimester urine and urine samples collected during the early-teen visit at age 8–14 years. Prenatal and peripubertal (early-teen) urine samples were collected in sterile cups, aliquoted within one hour after collection, frozen and stored at -80°F, and shipped on dry ice to McGill University (Montreal, Canada) for analysis. Urinary metals were measured using inductively coupled plasma mass spectrometry (ICPMS; Varian 820-MS, Inc., Palo Alto, California) as described previously (Basu et al. 2010; Srigboh et al. 2016). Accuracy and precision were measured using certified reference standards (Institut National de Santé Publique du Québec, or INSPQ) with coefficients of variation (CVs) ranging from 3 to 14%, and each batch run contained procedural blanks and replicate runs (Lewis et al. 2018; Srigboh et al. 2016). More details regarding quality control (QC) were previously described (Lewis et al. 2018). Values below the limit of detection (LOD) were replaced with the LOD/ $\sqrt{2}$. Urinary specific gravity (SG) was measured using a handheld digital refractometer.

Pb exposure in this study was also measured in maternal patella/blood and early peripubertal blood, and results of these biomarkers in relation to pubertal development within this population have already been published (Jansen et al. 2018; Y. Liu et al. 2019). As patella and blood Pb levels are better biomarkers of long term Pb exposure, we excluded urinary Pb from the current analyses.

2.3. Hormones

Children provided fasting blood samples during the early-teen visit at age 8–14 years. Serum aliquots were separated and frozen at -80 °C, and then sent to the Clinical Ligand Assay Service Satellite (CLASS) Laboratory at the University of Michigan (Ann Arbor, MI) for hormone analysis. Estradiol, testosterone, inhibin B, and sex hormone-binding globulin (SHBG) were measured in serum samples as biomarkers of puberty, and dehydroepiandrosterone sulfate (DHEA-S) was measured as a biomarker of secretion of adrenal androgens. Estradiol, total testosterone, SHBG, and DHEA-S were measured using an automated chemiluminescent immunoassay (Bayer Diagnostics ACS:180). Active inhibin B was assayed using Gen II ELISA (Beckman Coulter, Webster, TX). All laboratories that performed hormone analyses employed standard quality control (QC) measures, including the use of blanks and duplicate samples to measure instrument precision and identify potential sources of contamination at different collection and measurement stages. The samples were calibrated with standards to determine the degree of bias and implement actions to prevent calibration drift. The laboratories also followed pre-specified protocols for samples that exceed QC activity control limits. Values below the LOD were replaced with the LOD/ $\sqrt{2}$.

2.4 Sexual Maturation

Two pediatricians evaluated male offspring for Tanner staging and testicular volume using standardized protocols during both follow-up visits at ages 8–14 and 10-18 years. To ensure consistency, pediatricians were trained prior to the start of each follow-up as previously described (Chavarro et al. 2017) to evaluate Tanner staging in male participants using standardized protocols. Genital development stage (GD) was assessed as an indicator of puberty and pubic hair stage (PH) as an indicator of adrenarche, with stage 1 corresponding to no development and stage 5 corresponding to full development (Marshall and Tanner 1969). Right and left testicular volume (TV) were measured with an orchidometer, and the larger of the two measurements was used in analyses. As a testicular volume of 1–3 mL is considered prepubertal (Ankarberg-Lindgren and Norjavaara 2004; Mouritsen et al. 2013) and $TV \geq 20$ mL was used as an indicator of sexual maturity (Ankarberg-Lindgren and Norjavaara 2004; Sergeyev et al. 2017), cutoffs of 3 mL and 20 mL were used to create a 3-level ordinal variable for testicular volume.

2.5 Covariates

Covariates in our analysis included: age at the early-teen visit, BMI z-scores at both visits, and household socioeconomic status at the late-teen visit. BMI z-scores were calculated based on the World Health Organization child reference curves for age and sex (WHO, 2007) for each follow up visit. A 7-level categorical variable for socioeconomic status (SES) was estimated using a validated scale consisting of thirteen questions on housing quality, services, material goods and head of household education (Asociación Mexicana de Agencias de Investigación de Mercados y Opinión Pública, AMAI version 13x6) (AMAI 2000; López 2008).

2.5. Statistical Methods

The geometric mean concentrations of each metal in urine samples from pregnant women and their children were calculated. The percent of urine samples with concentrations below the LOD were reported for each metal, and metals that were detected in less than 50% of samples were excluded from further analysis.

Multiple linear regressions were used to assess associations between urinary metal concentrations and peripubertal serum hormone concentrations, where serum hormones were natural log-transformed prior to analysis to achieve normal distribution. Metal concentrations from prenatal and peripubertal urine samples were entered into regression models separately and each model was adjusted for child age and BMI z-score, and for SG as a measure of urinary dilution. Results were calculated as the percent difference in hormone (95% confidence interval) per interquartile range (IQR) increase in urinary metal concentrations. Because reproductive hormones may differ greatly in boys at pre-pubertal stages (Tanner stage=1) vs pubertal stages (Tanner stage>1), we performed a secondary analysis in which we examined the associations between *in utero* and peripubertal exposures and reproductive hormone levels among subjects who were pre-pubertal (pubic hair Tanner stage= 1, n=94).

Longitudinal analyses were conducted to explore the association between metals and early-teen and late-teen sexual maturation parameters using repeated measures generalized estimating equation (GEE). This allows us to take fuller advantage of the data that have been collected in our longitudinal cohort study and use the additional within-person information to achieve increases in statistical power to detect associations (Sitlani et al. 2015). With separate models for each metal, the GEE approach was used to fit a multinomial (ordinal) regression model for pubertal stages at each visit as a function of metal exposure, age at early-teen visit and change in time, with adjustment for potential confounders:

$$g(E[Y_{ij}]) = \beta_0 + \beta_1 M_{ij} + \beta_2 Age_i + \beta_3 Time_{ij} + \beta_4 M_{ij} * Time_{ij} + \beta_5 Age_i * Time_{ij} \quad (1)$$

$$g(E[Y_{ij}]) = \beta_0 + \beta_1 M_{ij} + \beta_2 Age_i + \beta_3 Time_{ij} + \beta_4 M_{ij} * Time_{ij} + \beta_5 Age_i * Time_{ij}$$

$$+ \beta_6 BMI_{base} + \beta_7 BMI_{var} \quad (2)$$

Model (1) is the crude model where Y is an outcome of interest (Tanner stages/testicular volume), g is a link function (cumulative logit), i denotes subject number (1,...,n), and j denotes visit number (1, 2). M is ln-transformed metal exposure, A is the age at the early-teen visit, $Time$ is the change in time between the early-teen and late-teen visit. Age at early-teen visit and change in time between two visits were included in the model to account for the effect of baseline age on attained Tanner stage or testicular volume, the natural pubertal progression across time, and the effect of baseline age on the natural pubertal progression. Model (2) is the final adjusted model, in which BMI z-score at the early-teen visit (BMI_{base}) and the change in BMI z-score from early-teen to late-teen visit (BMI_{var}) were included. The coefficients of interest are the cross-sectional effect of metal on tanner stage/testicular volume (β_1) and the effect of metal on the progression of tanner stage/testicular volume over time (β_4). Given the possibility that age and BMI z-score at the early-teen visit may be associated with Tanner stage and influence the future progression of Tanner stage, working independence was chosen as the covariance structure to ensure the validity of parameter estimates (Sullivan Pepe and Anderson 1994). We ran the GEE models both with and without BMI z-score (model 1 and 2) as BMI may be on the causal pathway between exposure and puberty. However, the magnitude of estimates from both models were almost identical, therefore we reported results from the models including BMI z-score. Results are presented as odds ratios (OR) and 95% confidence intervals (95% CI) per IQR increase in exposure. As few participants were categorized as having a PH Tanner stage =3 and stage=4 during the early-teen visit, the number of covariates we could reliably enter into models was limited (Peduzzi et al. 1996; Vittinghoff and McCulloch 2007). Thus, to minimize the number of covariates in GEE models, we included SG-corrected metal concentrations (rather than entering SG as a separate covariate which yielded similar results), using the following equation: $P_c = P[(SG_p - 1)/(SG_i - 1)]$ where P_c is the SG corrected metal concentration ($\mu\text{g/L}$), P is the measured metal concentration, SG_p is the median urinary specific gravity, and SG_i is the individual's urinary specific gravity.

Because SES could be a potential confounder, we ran a sensitivity analysis including SES as a covariate in models of hormones and maturation stages. Furthermore, as both low and high levels of essential

metals are of concern, to explore potential non-linear associations we used adjusted generalized additive models (GAM) to graphically depict the cross-sectional relationship between metal concentrations and hormones and sexual maturation measurements. We also considered significance after adjusting for multiple testing using the Benjamini-Hochberg method (Benjamini and Hochberg 1995). Since Tanner stages/testicular volumes were not independent of each other, we calculated q values (adjusted p values) treating each outcome as a family of tests (11 tests per outcome). A cutoff of 0.15 for q value was used to further interpret main results with greater confidence. All analyses were performed using R version 3.5.2 and SAS 9.4.

Results

3.1. Demographics and Exposure Distributions

Characteristics of participants in the original cohort of 997 women were previously described in detail (Perng et al. 2019). Mothers included in our analysis had similar demographic characteristics to the overall ELEMENT population; the mean age of mothers at the time of enrollment was 26.6 (standard deviation=5.3). Mothers had on average 11 years of education, most were married or cohabitating (89%), and all lived within Mexico City. Very few (3%) reported smoking during pregnancy. Average age of the boys at the early-teen and late-teen visits were 10.4 and 13.7, respectively. Mean and standard deviation of the follow up period were 3.5 and 0.5 years. The percent of samples with metal concentrations below the limit of detection, as well as the geometric means, standard deviations, and selected percentiles of sample concentrations from prenatal and peripubertal (early-teen) visits are shown in Table 1. Spearman correlations between the prenatal and peripubertal visit metal concentrations adjusted for SG are also presented in the table. With the exception of Sb and Fe, other metals were detected in > 50% of urine samples, therefore, Sb and Fe were excluded from further analysis. Weak correlations between maternal and peripubertal metal levels and weak to moderate correlations between different urinary metal concentrations within maternal and child samples have been previously reported elsewhere (Lewis et al. 2018).

Table 1. Distribution of urinary metal concentrations ($\mu\text{g/L}$) among ELEMENT mothers and their male children at age 8-14 years^a.

	<i>In utero</i>								Peripubertal				
	LOD	%< LOD	GM	GSD	25%	50%	75%	MAX	%< LOD	GM	GSD	25%	50%
<i>Essential metals</i>													
Co	0.4	0.0	1.2	1.9	0.8	1.2	2.1	5.2	0.0	0.7	1.5	0.6	0.8
Cu	48.2	46.3	86.1	4.4	34.1	50.2	71.9	2742	56.8	43.3	1.3	34.1	34.1
Mn	0.4	7.4	0.8	1.7	0.6	0.7	1.0	8.7	2.5	1.2	1.8	0.8	1.2
Mo	2.9	15.8	19.5	3.3	12.7	25.7	42.9	308	0.0	46.6	1.8	33.7	50.2
Se	10.5	3.2	29.9	1.7	23.0	32.6	43.1	120	1.7	47.2	1.7	36.3	53.0
Zn	0.1	1.2	271	2.2	160	298	451	1253	0.0	366	1.8	257	411
<i>Non-essential Metals</i>													
Al	8.6	11.6	24.0	2.5	12.3	20.3	42.2	304	24.6	14.5	2.1	8.8	14.0
As	0.3	0.0	14.0	2.0	9.3	13.2	20.6	153	0.0	14.3	2.0	10.2	14.4
Ba	1.1	3.2	4.0	2.0	2.6	4.2	5.9	27.6	10.2	2.5	2.0	1.6	2.4
Ni	3.0	0.0	8.7	1.9	5.8	7.4	11.0	107	0.8	8.1	1.6	5.9	8.0
Cd	54.0	2.1	0.2	2.1	0.1	0.2	0.3	2.7	2.5	0.1	1.6	0.1	0.1

^a uncorrected for specific gravity

^b P value from Spearman correlation test between *in utero* and peripubertal metal concentration measurements.

3.2 Hormone and Tanner Stages Distributions

Distributions of reproductive hormones among ELEMENT boys were described previously (Ferguson et al. 2014). Except for 12 total testosterone measurements, all measures were above the LOD. Spearman correlations among hormones were weak to moderate ($R=-0.42$ to 0.63). Distributions of Tanner stages of sexual maturation and testicular volume among the male children at the two follow up visits are reported (Supplementary Table S1). We additionally provided spaghetti plots depicting Tanner stage and testicular volume progression between two visits (Supplementary Figure S1), and the distribution of measures of sexual maturation for different age groups of ELEMENT boys (Supplementary Table S2). At the early-teen visit, the majority of the boys ($n=94$, 81.7%) were at Tanner stage 1 for pubic hair development whereas 57 boys (49.6%) were at Tanner stage 1 for genital development. Most boys who were at Tanner stage 1 moved to more advanced Tanner stages at the late-teen visit- only 28 (26.4%) and 8 (7.5%) were still at Tanner stage 1 for pubic hair development and genital development after 3 years on average since the early-teen visits. Boys who were at Tanner stages 2, 3, and 4 at the early-teen visit all progressed to higher stages at the late-teen visit, with 14 (13.2%) and 18 (13.2%) boys reaching full development (Tanner stage=5) for the two measurements. In terms of testicular volume distribution, the percentage of the boys in the prepubertal stage dropped from 14.8% to 0% from early-teen to late-teen visit.

3.2 *In utero* and peripubertal metal exposure and peripubertal hormone levels

Associations between *in utero* and peripubertal metal concentrations and reproductive hormones are presented in Figure 2 and Supplementary Table S4. Positive associations were observed between some urinary essential metal concentrations and estradiol, testosterone, and SHBG. One IQR increase in *in utero* Zn concentration was associated with 13.7 % higher serum estradiol levels (95% CI: 0.3, 28.8). *In utero* Co

and Mn were positively associated with SHBG levels, with an IQR increase associated with 16% (95% CI: 0.4, 34.2) and 14.2% (95% CI: 1.5, 28.5) higher serum SHBG after adjustment for child age, BMI z-score, and SG, respectively. As shown in Figure 2, effect estimates for the association between both *in utero* essential and non-essential metal concentrations and testosterone were larger compared to other reproductive hormones, 51.3% for Mo, 35% for As, 38.9% for Cd. In models where reproductive hormones were regressed on concurrent peripubertal exposures, the strongest associations were observed also between metal concentrations and testosterone, particularly with non-essential metals Ni (% Δ /IQR: 35.1%, 95% CI: 2.6, 77.8) and Ba (% Δ /IQR: 59.1%, 95% CI: 22.5, 106.8). Peripubertal Ba concentration was also associated with higher estradiol (% Δ /IQR: 10.2%, 95% CI: 0.5, 20.9). However, no significant associations were detected between peripubertal metal exposures and DHEA-S or inhibin B.

After correcting for multiple testing, the associations of *in utero* Mo, As, and Cd with testosterone levels, as well as the association of peripubertal Ba with testosterone level had *q*-values < 0.15 (Figure 2 and Supplementary Table S4), providing greater confidence in these associations.

3.3 *In utero* and peripubertal metal exposure and sexual maturation

We have presented results from multiple ordinal regression models of *in utero* metal concentrations and Tanner stage and testicular volume in Figure 3 and Supplementary Table S5. Tanner stages and testicular volume at the early-teen visit were not associated with *in utero* metal concentrations, with the exception of a non-essential metal, Al; an IQR increase in *in utero* Al concentrations was associated with 3.6 times greater odds (95% CI: 1.67, 7.76) of being at a higher category of testicular volume versus lower categories. In the same figure and table, associations between *in utero* metal concentrations and pubertal development over time in boys are also presented. During the follow-up, an IQR increase in *in utero* concentrations of non-essential metalloid As was associated with 36% (OR/IQR: 0.64, 95% CI: 0.48, 0.85) lower odds of genital development progression per year, adjusting for age, BMI, and Tanner stage at the early-teen visit (GEE longitudinal model). *In utero* concentrations of non-essential metal(loid)s Al (OR/IQR: 0.61, 95% CI: 0.45, 0.83) and As (OR/IQR: 0.64, 95% CI: 0.43, 0.97) were associated with lower odds of progressing to a higher testicular volume category (i.e. 39 and 36% lower odds/IQR). Essential metal Zn was also associated with 38% lower odds of testicular volume progression (OR/IQR: 0.62, 95% CI: 0.44, 0.88).

Similarly, Figure 4 and Supplementary Table S5 show the cross-sectional and longitudinal associations between prepubertal metal concentrations and sexual maturation and progression. No significant associations were found between peripubertal urinary metal concentrations with Tanner stage for genital development or progression over follow-up, although several essential and non-essential metals were associated with pubic hair Tanner stages, testicular volume, and progressions. Peripubertal Zn concentration was associated with higher odds of being at a higher developmental stage for pubic hair (OR/IQR: 6.11, 95% CI: 1.89, 19.69) and testicular volume (OR/IQR: 5.39, 95% CI: 1.88, 15.49) at the early-teen visit, as well as slower progression of pubic hair development (OR/IQR: 0.47, 95% CI: 0.34, 0.67) and testicular volume (OR/IQR: 0.58, 95% CI: 0.40, 0.85) during the follow-up. Higher Mn was associated with

34% (OR/IQR: 0.66, 95% CI: 0.46, 0.96) lower pubic hair development progression only. Regarding non-essential metals, an IQR increase in peripubertal Ba concentration was associated with 2.3 times greater odds (95% CI: 1.15, 4.75) of being at higher Tanner stage for pubic hair development at age 8–14 years, but with 34% (OR/IQR: 0.66, 95% CI: 0.53, 0.83) lower odds of progressing to a higher pubic hair Tanner stage per year of follow-up. Higher peripubertal urinary Al was also associated with 47% (OR/IQR: 0.53, 95% CI: 0.40, 0.70) lower odds of pubic hair development progression per IQR increase.

All the associations described above had q -values < 0.15 (Figure 3, Figure 4, and Supplementary Table S5) after correcting for multiple testing, except for the association between *in utero* As and testicular volume progressions. Output graphics produced from GAM models showed that there was no significant non-linear relationship between essential and non-essential metal concentrations and hormones and sexual maturation outcomes, after adjusting for the same sets of covariates.

3.4 Sensitivity Analysis

Results from the hormone subset analysis when we restricted to boys who were prepubertal (94 out of 118) at the early-teen visit are similar to the main analysis results (Supplementary Table S6). For models regressing *in utero* metal concentrations, some effect estimates for DHEA_S, SHBG, and testosterone were no longer significant, potentially due to the small sample size. For models regressing peripubertal exposure, the notable effect estimates for testosterone were similar to those from the main analysis. The association between DHEA_S and Ni ($\% \Delta / \text{IQR} = 20.2$, 95%CI=1.7, 42.0) and inhibin B and Al ($\% \Delta / \text{IQR} = -19.2$, 95%CI=-31.7, -4.4) became stronger and significant in the peripubertal subset analysis.

The magnitude of estimates from GEE models with and without BMI (Supplementary Table S7) were almost identical. Findings from models adjusting for SES were generally consistent with metal and hormone associations observed in our main analyses; in SES-adjusted models, the association between *in utero* Co concentrations and higher SHBG ($\% \Delta / \text{IQR} = 14.2$, 95%CI=-2.9, 23.3) was slightly attenuated and no longer significant, while the association between peripubertal Al and inhibin B ($\% \Delta / \text{IQR} = -14.1$, 95%CI=-24.6, -2.2) was stronger and significant. In GEE models for Tanner stage or testicular volume status including SES finding remain consistent with the main models.

A high proportion of samples had metal concentrations above the LOD except for Cu, which was below the detection limit in 46% of samples. In a secondary analysis, we categorized urinary Cu concentrations into three groups. The low group consisted of values below the LOD, while the medium and high groups were made up of equalized bins among the detected values. We estimated the model parameters again and found that they were similar to the main parameter estimates.

Discussion

Previous studies of metal exposure and male reproductive development have primarily been cross-sectional and focused on heavy metals, with the exception of “the Russian Children’s Study” and “Early Life Exposure in Mexico to Environmental Toxicants (ELEMENT) Study” that assessed the longitudinal

impact of blood Pb (Yun Liu et al. 2019; Williams et al. 2019). Thus, it is critical to evaluate the associations of essential and non-essential metals with reproductive hormones, sexual maturation, and progression among boys. Based on our longitudinal analysis of boys in a birth cohort in Mexico City who were examined at two peripubertal follow-up visits, we demonstrated a higher pubic hair development stage and testicular volume at the early-teen visit with higher non-essential metal concentrations, including *in utero* Al and peripubertal Ba and essential metal Zn concentration (peripubertal). We also found associations between *in utero* and peripubertal exposure to a number of both essential and non-essential metals and slowed progression of pubertal development across the follow-up period. In relation to hormone levels, higher non-essential metals, *in utero* As and Cd and peripubertal Ba concentrations as well as essential metal Mo measured *in utero* were associated with substantially higher testosterone levels. None of the associations between metals and hormones and measurements of sexual maturations appear to be non-linear.

To our knowledge, this is the first study to investigate the association between various *in utero* and peripubertal metals measured in urine and reproductive hormone levels in teenage boys. We reported a number of positive associations between specific metals measured in the urine and reproductive hormones, particularly testosterone. Some of our findings have not been reported elsewhere, such as relationships of essential minerals, e.g., Zn with estradiol or Co and Mn with SHBG. The positive association between relatively low Cd exposure and testosterone in this study is supported by epidemiological studies of non-occupational exposures among men (Jurasovic et al. 2004; Meeker et al. 2010; Menke et al. 2008; Telisman et al. 2007). However, *in vivo* and *in vitro* observations linked Cd with altered adrenal gland activity and induced oxidative stress, which could result in toxic effects on the testis and subsequently decrease in serum testosterone (Gay et al. 2013; Liu et al. 2009; Patra et al. 2011; Singhal et al. 1976; Siu et al. 2009).

Mo is an essential nutrient that is a cofactor for important enzymes involving in toxin removal (Schwarz 2016; Todd et al. 2017). While no prior studies have investigated the associations between Mo and hormones in boys, contrary to our finding on *in utero* Mo and increased testosterone, decreased testosterone levels and sperm concentrations were observed in males with increasing levels of Mo (Lewis and Meeker 2015; Meeker et al. 2008; Meeker et al. 2010). Several animal studies have also reported the reproductive toxicity of Mo, including declines in sperm concentration, motility and normal morphology, testicular degeneration, and reduced fertility (Jeter and Davis 1954; Pandey and Singh 2002; Thomas and Moss 1951; Vyskočil and Viau 1999). It is worth noting that the prenatal Mo levels were lower in our study population compared to the previous studies of Mo and reproductive outcomes and those reported in the National Health and Nutrition Examination Survey (NHANES) (Centers for Disease Control and Prevention 2018) (Supplementary Table S3). It is not clear whether Mo levels measured *in utero* will have different effects on reproductive hormones measured during different life stages, which warrants further investigation. Although previous studies have not evaluated associations between Ba and reproductive hormones in human, our finding of positive associations between peripubertal Ba concentrations and serum testosterone and estradiol levels is consistent with previous studies among male zebrafish, where

exposure to barium chloride significantly increased estradiol concentrations and transcripts of genes involved in the HPG axis (Kwon et al. 2016).

One of the main observations in this study is that Zn is associated with higher odds of pubic hair and testicular volume stage at the early-teen follow-up, but a slower progression of puberty from the early-teen to late-teen visit. Few cross-sectional studies on boys and girls also reported a significant positive relationship between serum or plasma zinc levels and stages of sexual maturation (Leonard et al. 1998; Onukwuli et al. 2018; Vivoli et al. 1990; Wagner et al. 1985), as well as reproductive hormones (testosterone). Zn is an essential trace metal and is fundamental for the development of the male reproductive system (Kerns et al. 2018; Mendoza et al. 2017). Studies on various animals described the role of Zn in advancing male puberty, through increased **testicular activity**, testosterone production, metabolism, and growth, primarily via activated the hypothalamus and the pituitary functions (Arangasamy et al. 2018; Dance et al. 2016; El-Masry et al. 1994; Fang and Furuhashi 1978; Geary et al. 2016; Underwood and Somers 1969). However, we did not observe a significant association between serum Zn and testosterone levels.

Pubic hair stages at the early-teen visit and progression were consistently associated with elevated concentrations of certain peripubertal metals, including As, Al, Mn, and Zn. The biological mechanisms that underlie the link between these essential and non-essential metals and pubertal development are uncertain but may be related to reproductive hormones. The main hormones responsible for pubic hair development in males are the androgens DHEA and testosterone (Breehl and Caban 2020; Rubinow and Schmidt 1996). In this cohort, the strongest observed associations were between *in utero* and peripubertal metal concentrations and increased testosterone, while no significant associations were seen in relation to DHEA-S. It is possible that these metals impact the appearance of pubic hair through changes in testosterone.

In this study, we observed slower progression of sexual maturation for those boys at higher Tanner stages at the early-teen visit which was also reported in studies explored the association between pubertal onset and progression (German et al. 2018; Llop-Viñolas et al. 2004; Pantiotou et al. 2008). A compensatory mechanism similar to “catch-up growth” was proposed previously as a potential explanation for this observation (Ashrap et al. 2019; Wu et al. 2018). For those boys who had experienced a delayed pubertal development, their body systems may have responded by accelerating the tempo of pubertal progression (the change from lower stages of puberty to higher stages); while others who had experienced an advanced pubertal development may respond by slowing down the pubertal progression. This concept may explain the associations between metal exposure, earlier puberty onset at early-teen, and slower progression we found in this study. Future research is needed to establish this phenomenon and potential mechanisms.

The ELEMENT study, a prospective longitudinal birth cohort study in Mexico City, provided an opportunity to study the relationships of both *in utero* and peripubertal metal exposure on hormone levels and measures of sexual maturation and progression during the peripubertal period. Our results indicate that

the *in utero* and peripubertal periods are vulnerable life stages, during which metal exposures may lead to disruption of male reproductive hormones and pubertal development. The findings also support that essential and non-essential metals have the potential to disrupt the onset and progression of puberty via interrupting the critical hormonal pathways. Several of the associations with hormones and sexual maturation are comparable to previous reports from human and animal research and supported by the current understanding of male pubertal development. However, there are inconsistent findings between the current study and previous studies and this may be due to a number of reasons; 1) the exposure assessment approaches were different [sampling period, exposure matrices (*i.e.*, blood, urine, hair)]. 2) most of the previous studies exploring the relationship between metals and hormones were conducted on an adult population. 3) most studies examined the cross-sectional relationships whereas we examined the longitudinal association between metals and sexual maturation and progression. It is also worth noting that the assessment of sexual maturation markers in this study was conducted by the same observers at both visits in this study to minimize measurement bias. Limitations of our study include a somewhat small sample size and few observations for certain Tanner stages, which may result in imprecise effect estimates. The age range of children at the early-teen and late-teen visits does overlap, however, our results are unlikely biased as the average follow-up period for different age groups is 3.5 years. Some of the metals measured in this study have a relatively short half-life in urine, so urinary levels at the time of measurement may not fully characterize exposure during each specific window of development. Lastly, hormone levels were only measured at one time point and are likely to be subject to non-differential misclassification due to diurnal variation. Further research is warranted to prospectively explore the underlying mechanisms by which metals may affect male sexual maturation and progression in larger study populations.

Declarations

Ethics approval and consent to participate

Research protocols were approved by the Institutional Review Board at University of Michigan, and the Research, Biosafety and Ethics in Research at the Mexico National Institute of Public Health. Prior to enrollment, informed consent from mothers and informed assent from offspring were obtained.

Consent for publication

Not applicable.

Availability of data and materials

All data generated and analyzed during this study are not publicly available due to the Institutional Review Board restrictions.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by U.S. Environmental Protection Agency (US EPA) grants RD834800 and RD83543601 and National Institute for Environmental Health Sciences (NIEHS) grants P20 ES018171, P01 ES02284401, and P30 ES017885. Its contents are solely the responsibility of the grantee and do not necessarily represent the official views of the US EPA. Further, the US EPA does not endorse the purchase of any commercial products or services mentioned in the publication. This work was also supported and partially funded by the National Institute of Public Health, Ministry of Health of Mexico.

Authors' Contributions

PA conducted data analyses and drafted the manuscript. JDM, KEP, MTR and DJW conceived the study and oversaw its coordination. NB helped oversee the study, data acquisition, metals analysis, and advised the interpretation and drafting of the manuscript. BNS helped oversee the study and advised during the data analysis. MSG and AMG participated in the collection and assembly of data. All authors read and approved the final manuscript.

Acknowledgments

The authors acknowledge the research staff at participating hospitals and the American British Cowdray Hospital in Mexico City for providing research facilities. We thank the mothers and children for participating in the study.

References

- Agusa T, Kunito T, Iwata H, Monirith I, Chamnan C, Tana TS, et al. 2007. Mercury in hair and blood from residents of phnom penh (cambodia) and possible effect on serum hormone levels. *Chemosphere* 68:590-596.
- AMAI. 2000. Avances del comité de niveles socioeconómicos. Vol. 2018:Comité de Niveles Socioeconómicos. Asociación Mexicana de Agencias de Investigación de Mercados y Opinión Pública, A.C.
- Anderson SE, Dallal GE, Must A. 2003. Relative weight and race influence average age at menarche: Results from two nationally representative surveys of us girls studied 25 years apart. *Pediatrics* 111:844-850.
- Ankarberg-Lindgren C, Norjavaara E. 2004. Changes of diurnal rhythm and levels of total and free testosterone secretion from pre to late puberty in boys: Testis size of 3 ml is a transition stage to puberty. *Eur J Endocrinol* 151:747-757.
- Arangasamy A, Venkata Krishnaiah M, Manohar N, Selvaraju S, Guvvala PR, Soren NM, et al. 2018. Advancement of puberty and enhancement of seminal characteristics by supplementation of trace

minerals to bucks. *Theriogenology* 110:182-191.

Ashrap P, Sanchez BN, Tellez-Rojo MM, Basu N, Tamayo-Ortiz M, Peterson KE, et al. 2019. In utero and peripubertal metals exposure in relation to reproductive hormones and sexual maturation and progression among girls in Mexico City. *Environ Res* 177:108630.

ATSDR U. 2007a. Toxicological profile for lead (Atlanta, GA: US Department of Health and Human Services, Agency for Toxic Substances and Disease Registry (ATSDR), Public Health Service). US EPA (2006) Air quality criteria for lead.

ATSDR U. 2007b. Toxicological profile for arsenic. Agency for Toxic Substances and Disease Registry, Division of Toxicology, Atlanta, GA.

Basu N, Abare M, Buchanan S, Cryderman D, Nam DH, Sirkin S, et al. 2010. A combined ecological and epidemiologic investigation of metal exposures amongst indigenous peoples near the Marlin mine in western Guatemala. *Sci Total Environ* 409:70-77.

Bellingham M, Fowler PA, Amezcaga MR, Rhind SM, Cotinot C, Mandon-Pepin B, et al. 2009. Exposure to a complex cocktail of environmental endocrine-disrupting compounds disturbs the kisspeptin/gpr54 system in ovine hypothalamus and pituitary gland. *Environ Health Perspect* 117:1556-1562.

Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)* 57:289-300.

Blakemore SJ, Burnett S, Dahl RE. 2010. The role of puberty in the developing adolescent brain. *Hum Brain Mapp* 31:926-933.

Bloom MS, Parsons PJ, Steuerwald AJ, Schisterman EF, Browne RW, Kim K, et al. 2010. Toxic trace metals and human oocytes during in vitro fertilization (IVF). *Reprod Toxicol* 29:298-305.

Bochud M, Jenny-Burri J, Pruijm M, Ponte B, Guessous I, Ehret G, et al. Urinary cadmium excretion is associated with increased synthesis of cortico- and sex steroids in a family-based Swiss population study. In: *Proceedings of the 19th European Congress of Endocrinology, 2017, Vol. 49* BioScientifica.

Breehl L, Caban O. 2020. Physiology, puberty. In: *StatPearls*. Treasure Island (FL).

Buck Louis GM, Gray LE, Jr., Marcus M, Ojeda SR, Pescovitz OH, Witchel SF, et al. 2008. Environmental factors and puberty timing: Expert panel research needs. *Pediatrics* 121 Suppl 3:S192-207.

Cahill L. 2006. Why sex matters for neuroscience. *Nat Rev Neurosci* 7:477-484.

Centers for Disease Control and Prevention. 2018. Fourth report on human exposure to environmental chemicals. Available:

https://www.cdc.gov/exposurereport/pdf/FourthReport_UpdatedTables_Volume1_Mar2018.pdf
[accessed 04/04/2018].

Chavarro JE, Watkins DJ, Afeiche MC, Zhang Z, Sanchez BN, Cantonwine D, et al. 2017. Validity of self-assessed sexual maturation against physician assessments and hormone levels. *J Pediatr* 186:172-178 e173.

Claus Henn B, Coull BA, Wright RO. 2014. Chemical mixtures and children's health. *Curr Opin Pediatr* 26:223-229.

Dance A, Thundathil J, Blondin P, Kastelic J. 2016. Enhanced early-life nutrition of holstein bulls increases sperm production potential without decreasing postpubertal semen quality. *Theriogenology* 86:687-694. e682.

De Coster S, van Larebeke N. 2012. Endocrine-disrupting chemicals: Associated disorders and mechanisms of action. *J Environ Public Health* 2012:713696.

Deb P, Mandal SS. 2017. Endocrine disruptors: Mechanism of action and impacts on health and environment. *Gene Regulation, Epigenetics and Hormone Signaling*:607-638.

Den Hond E, Schoeters G. 2006. Endocrine disrupters and human puberty. *Int J Androl* 29:264-271; discussion 286-290.

Diamanti-Kandarakis E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, Soto AM, et al. 2009. Endocrine-disrupting chemicals: An endocrine society scientific statement. *Endocr Rev* 30:293-342.

El-Masry K, Nasr A, Kamal T. 1994. Influences of season and dietary supplementation with selenium and vitamin e or zinc on some blood constituents and semen quality of new zealand white rabbit males. *World Rabbit Science* 2.

Euling SY, Herman-Giddens ME, Lee PA, Selevan SG, Juul A, Sorensen TI, et al. 2008. Examination of us puberty-timing data from 1940 to 1994 for secular trends: Panel findings. *Pediatrics* 121 Suppl 3:S172-191.

Eveleth PB. 2017. Timing of menarche: Secular trend and population differences. In: *School-age pregnancy and parenthood*:Routledge, 39-52.

Fang VS, Furuhashi N. 1978. Partial alleviation of the antitesticular effect of pipecolinomethylhydroxyindane by zinc in rats. *J Endocrinol* 79:151-152.

Faroon O, Ashizawa A, Wright S, Tucker P, Jenkins K, Ingerman L, et al. 2012. In: *Toxicological profile for cadmium*. Atlanta (GA).

- Ferguson KK, Peterson KE, Lee JM, Mercado-Garcia A, Blank-Goldenberg C, Tellez-Rojo MM, et al. 2014. Prenatal and peripubertal phthalates and bisphenol a in relation to sex hormones and puberty in boys. *Reprod Toxicol* 47:70-76.
- Fudvoye J, Lopez-Rodriguez D, Franssen D, Parent A-S. 2019. Endocrine disrupters and possible contribution to pubertal changes. *Best Practice & Research Clinical Endocrinology & Metabolism*:101300.
- Garcia-Morales P, Saceda M, Kenney N, Kim N, Salomon DS, Gottardis MM, et al. 1994. Effect of cadmium on estrogen receptor levels and estrogen-induced responses in human breast cancer cells. *J Biol Chem* 269:16896-16901.
- Gay F, Laforgia V, Caputo I, Esposito C, Lepretti M, Capaldo A. 2013. Chronic exposure to cadmium disrupts the adrenal gland activity of the newt *triturus carnifex* (amphibia, urodela). *BioMed research international* 2013.
- Geary T, Kelly W, Spickard D, Larson C, Grings E, Ansotegui R. 2016. Effect of supplemental trace mineral level and form on peripubertal bulls. *Animal reproduction science* 168:1-9.
- Gerberding J. 2005. Toxicological profile for zinc. Atlanta: US Department of Health and Human Services, Agency for Toxic Substances and Disease Registry:11-118.
- Gerhard I, Waibel S, Daniel V, Runnebaum B. 1998. Impact of heavy metals on hormonal and immunological factors in women with repeated miscarriages. *Hum Reprod Update* 4:301-309.
- German A, Shmoish M, Belsky J, Hochberg Z. 2018. Outcomes of pubertal development in girls as a function of pubertal onset age. *European journal of endocrinology* 179:279-285.
- Greger JL. 1999. Nutrition versus toxicology of manganese in humans: Evaluation of potential biomarkers. *Neurotoxicology* 20:205-212.
- Ha E, Basu N, Bose-O'Reilly S, Dorea JG, McSorley E, Sakamoto M, et al. 2017. Current progress on understanding the impact of mercury on human health. *Environ Res* 152:419-433.
- Hauser R, Sergeev O, Korrick S, Lee MM, Revich B, Gitin E, et al. 2008. Association of blood lead levels with onset of puberty in russian boys. *Environ Health Perspect* 116:976-980.
- Health UDo, Services H. 2004. Toxicological profile for copper. US Department of Health and Human Services, Atlanta.
- Herman-Giddens ME, Slora EJ, Wasserman RC, Bourdony CJ, Bhapkar MV, Koch GG, et al. 1997. Secondary sexual characteristics and menses in young girls seen in office practice: A study from the pediatric research in office settings network. *Pediatrics* 99:505-512.

- Herman-Giddens ME, Wang L, Koch G. 2001. Secondary sexual characteristics in boys: Estimates from the national health and nutrition examination survey iii, 1988-1994. *Arch Pediatr Adolesc Med* 155:1022-1028.
- Hsieh FI, Hwang TS, Hsieh YC, Lo HC, Su CT, Hsu HS, et al. 2008. Risk of erectile dysfunction induced by arsenic exposure through well water consumption in taiwan. *Environ Health Perspect* 116:532-536.
- Iavicoli I, Fontana L, Bergamaschi A. 2009. The effects of metals as endocrine disruptors. *J Toxicol Environ Health B Crit Rev* 12:206-223.
- Jacobson-Dickman E, Lee MM. 2009. The influence of endocrine disruptors on pubertal timing. *Curr Opin Endocrinol Diabetes Obes* 16:25-30.
- Jansen EC, Zhou L, Song P, Sanchez BN, Mercado A, Hu H, et al. 2018. Prenatal lead exposure in relation to age at menarche: Results from a longitudinal study in mexico city. *J Dev Orig Health Dis* 9:467-472.
- Jeng HA, Huang YL, Pan CH, Diawara N. 2015. Role of low exposure to metals as male reproductive toxicants. *Int J Environ Health Res* 25:405-417.
- Jeter MA, Davis GK. 1954. The effect of dietary molybdenum upon growth, hemoglobin, reproduction and lactation of rats. *The Journal of nutrition* 54:215-220.
- Jurasovic J, Cvitkovic P, Pizent A, Colak B, Telisman S. 2004. Semen quality and reproductive endocrine function with regard to blood cadmium in croatian male subjects. *Biometals* 17:735-743.
- Karpati AM, Rubin CH, Kieszak SM, Marcus M, Troiano RP. 2002. Stature and pubertal stage assessment in american boys: The 1988-1994 third national health and nutrition examination survey. *J Adolesc Health* 30:205-212.
- Kerns K, Zigo M, Sutovsky P. 2018. Zinc: A necessary ion for mammalian sperm fertilization competency. *Int J Mol Sci* 19.
- Kwon B, Ha N, Jung J, Kim PG, Kho Y, Choi K, et al. 2016. Effects of barium chloride exposure on hormones and genes of the hypothalamic-pituitary-gonad axis, and reproduction of zebrafish (*danio rerio*). *Bull Environ Contam Toxicol* 96:341-346.
- Lauretta R, Sansone A, Sansone M, Romanelli F, Appetecchia M. 2019. Endocrine disrupting chemicals: Effects on endocrine glands. *Front Endocrinol (Lausanne)* 10:178.
- Lee EJ, Lee BR, Lee JH, Chang GT. 2016. A study on the recent trend of chief complaint of korean pediatric and adolescent outpatients. *The Journal of Pediatrics of Korean Medicine* 30:45-58.

- Lee Y, Styne D. 2013. Influences on the onset and tempo of puberty in human beings and implications for adolescent psychological development. *Horm Behav* 64:250-261.
- Leonard MB, Zemel BS, Kawchak DA, Ohene-Frempong K, Stallings VA. 1998. Plasma zinc status, growth, and maturation in children with sickle cell disease. *J Pediatr* 132:467-471.
- Lewis RC, Meeker JD, Peterson KE, Lee JM, Pace GG, Cantoral A, et al. 2013. Predictors of urinary bisphenol a and phthalate metabolite concentrations in mexican children. *Chemosphere* 93:2390-2398.
- Lewis RC, Meeker JD. 2015. Biomarkers of exposure to molybdenum and other metals in relation to testosterone among men from the united states national health and nutrition examination survey 2011-2012. *Fertil Steril* 103:172-178.
- Lewis RC, Meeker JD, Basu N, Gauthier AM, Cantoral A, Mercado-Garcia A, et al. 2018. Urinary metal concentrations among mothers and children in a mexico city birth cohort study. *Int J Hyg Environ Health* 221:609-615.
- Liu J, Qu W, Kadiiska MB. 2009. Role of oxidative stress in cadmium toxicity and carcinogenesis. *Toxicology and applied pharmacology* 238:209-214.
- Liu Y, Téllez-Rojo M, Hu H, Sánchez BN, Martínez-Mier EA, Basu N, et al. 2019. Fluoride exposure and pubertal development in children living in mexico city. *Environmental Health* 18:26.
- Liu Y, Tellez-Rojo MM, Sanchez BN, Zhang Z, Afeiche MC, Mercado-Garcia A, et al. 2019. Early lead exposure and pubertal development in a mexico city population. *Environ Int* 125:445-451.
- Llop-Viñolas D, Vizmanos B, Monasterolo RC, Subías JE, Fernández-Ballart J, Martí-Henneberg C. 2004. Onset of puberty at eight years of age in girls determines a specific tempo of puberty but does not affect adult height. *Acta Paediatrica* 93:874-879.
- López H. 2008. Nivel socioeconómico amai. (AMAI, ed):INEGI.
- Marshall WA, Tanner JM. 1969. Variations in pattern of pubertal changes in girls. *Arch Dis Child* 44:291-303.
- Massart F, Parrino R, Seppia P, Federico G, Saggese G. 2006. How do environmental estrogen disruptors induce precocious puberty? *Minerva Pediatr* 58:247-254.
- Meeker JD, Rossano MG, Protas B, Diamond MP, Puscheck E, Daly D, et al. 2008. Cadmium, lead, and other metals in relation to semen quality: Human evidence for molybdenum as a male reproductive toxicant. *Environ Health Perspect* 116:1473-1479.
- Meeker JD, Rossano MG, Protas B, Padmanahban V, Diamond MP, Puscheck E, et al. 2010. Environmental exposure to metals and male reproductive hormones: Circulating testosterone is inversely associated with

blood molybdenum. *Fertil Steril* 93:130-140.

Mendiola J, Moreno JM, Roca M, Vergara-Juarez N, Martinez-Garcia MJ, Garcia-Sanchez A, et al. 2011. Relationships between heavy metal concentrations in three different body fluids and male reproductive parameters: A pilot study. *Environ Health* 10:6.

Mendoza AD, Woodruff TK, Wignall SM, O'Halloran TV. 2017. Zinc availability during germline development impacts embryo viability in *Caenorhabditis elegans*. *Comp Biochem Physiol C Toxicol Pharmacol* 191:194-202.

Menke A, Guallar E, Shiels MS, Rohrmann S, Basaria S, Rifai N, et al. 2008. The association of urinary cadmium with sex steroid hormone concentrations in a general population sample of US adult men. *BMC Public Health* 8:72.

Mouritsen A, Aksglaede L, Soerensen K, Hagen CP, Petersen JH, Main KM, et al. 2013. The pubertal transition in 179 healthy Danish children: Associations between pubarche, adrenarche, gonadarche, and body composition. *Eur J Endocrinol* 168:129-136.

Nagata C, Nagao Y, Shibuya C, Kashiki Y, Shimizu H. 2005. Urinary cadmium and serum levels of estrogens and androgens in postmenopausal Japanese women. *Cancer Epidemiol Biomarkers Prev* 14:705-708.

Onukwuli VO, Chinawa J, Eke CB, Nwokocha AR, Emodi IJ, Ikefuna AN. 2018. Impact of zinc on sexual maturation of female sickle cell anemia (sca) children in Enugu, Southeast Nigeria. *Pediatr Hematol Oncol* 35:145-155.

Pandey R, Singh S. 2002. Effects of molybdenum on fertility of male rats. *Biometals* 15:65-72.

Pantsiotou S, Papadimitriou A, Douros K, Priftis K, Nicolaidou P, Fretzayas A. 2008. Maturation tempo differences in relation to the timing of the onset of puberty in girls. *Acta Paediatrica* 97:217-220.

Parent A-S, Franssen D, Fudvoye J, Pinson A, Bourguignon J-P. 2016. Current changes in pubertal timing: Revised vision in relation with environmental factors including endocrine disruptors. In: *Puberty from bench to clinic*, Vol. 29: Karger Publishers, 174-184.

Parent AS, Franssen D, Fudvoye J, Gerard A, Bourguignon JP. 2015. Developmental variations in environmental influences including endocrine disruptors on pubertal timing and neuroendocrine control: Revision of human observations and mechanistic insight from rodents. *Front Neuroendocrinol* 38:12-36.

Paris F, Gaspari L, Sultan C. 2016. Precocious puberty and environmental endocrine disruptors. In: *Early puberty*: Springer, 9-20.

Patra R, Rautray AK, Swarup D. 2011. Oxidative stress in lead and cadmium toxicity and its amelioration. *Veterinary medicine international* 2011.

- Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. 1996. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol* 49:1373-1379.
- Perng W, Tamayo-Ortiz M, Tang L, Sanchez BN, Cantoral A, Meeker JD, et al. 2019. Early life exposure in Mexico to environmental toxicants (element) project. *BMJ Open* 9:e030427.
- Pescovitz OH, Walvoord EC. 2007. *When puberty is precocious: Scientific and clinical aspects*: Springer Science & Business Media.
- Risher J. 2003. *Toxicological profile for selenium*: Agency for Toxic Substances and Disease Registry.
- Rodriguez DL, Francisco AC, Gerard A, Blacher S, Tirelli E, Bourguignon J-P, et al. Endocrine disruptors transgenerationally alters pubertal timing through epigenetic reprogramming of the hypothalamus. In: *Proceedings of the 21st European Congress of Endocrinology, 2019, Vol. 63* BioScientifica.
- Roy JR, Chakraborty S, Chakraborty TR. 2009. Estrogen-like endocrine disrupting chemicals affecting puberty in humans—a review. *Med Sci Monit* 15:RA137-145.
- Rubinow DR, Schmidt PJ. 1996. Androgens, brain, and behavior. *Am J Psychiatry* 153:974-984.
- Schwarz G. 2016. Molybdenum cofactor and human disease. *Current opinion in chemical biology* 31:179-187.
- Sergeyev O, Burns JS, Williams PL, Korrick SA, Lee MM, Revich B, et al. 2017. The association of peripubertal serum concentrations of organochlorine chemicals and blood lead with growth and pubertal development in a longitudinal cohort of boys: A review of published results from the Russian children's study. *Rev Environ Health* 32:83-92.
- Shenkin A. 2003. Dietary reference values for vitamin A, vitamin K, arsenic, boron, chromium, copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium and zinc. *Journal of Human Nutrition and Dietetics* 16:199-200.
- Singhal R, Merali Z, Hrdina P. Aspects of the biochemical toxicology of cadmium. In: *Proceedings of the Federation Proceedings, 1976, Vol. 35*, 75-80.
- Sisk CL, Foster DL. 2004. The neural basis of puberty and adolescence. *Nat Neurosci* 7:1040-1047.
- Sitlani CM, Rice KM, Lumley T, McKnight B, Cupples LA, Avery CL, et al. 2015. Generalized estimating equations for genome-wide association studies using longitudinal phenotype data. *Stat Med* 34:118-130.
- Siu ER, Mruk DD, Porto CS, Cheng CY. 2009. Cadmium-induced testicular injury. *Toxicology and Applied Pharmacology* 238:240-249.
- Spear LP. 2000. The adolescent brain and age-related behavioral manifestations. *Neurosci Biobehav Rev* 24:417-463.

- Srigboh RK, Basu N, Stephens J, Asampong E, Perkins M, Neitzel RL, et al. 2016. Multiple elemental exposures amongst workers at the agbogbloshie electronic waste (e-waste) site in Ghana. *Chemosphere* 164:68-74.
- Sullivan Pepe M, Anderson GL. 1994. A cautionary note on inference for marginal regression models with longitudinal data and general correlated response data. *Communications in Statistics-Simulation and Computation* 23:939-951.
- Telisman S, Cvitkovic P, Jurasovic J, Pizent A, Gavella M, Rocic B. 2000. Semen quality and reproductive endocrine function in relation to biomarkers of lead, cadmium, zinc, and copper in men. *Environ Health Perspect* 108:45-53.
- Telisman S, Colak B, Pizent A, Jurasovic J, Cvitkovic P. 2007. Reproductive toxicity of low-level lead exposure in men. *Environ Res* 105:256-266.
- Thomas J, Moss S. 1951. The effect of orally administered molybdenum on growth, spermatogenesis and testes histology of young dairy bulls. *Journal of dairy science* 34:929-934.
- Todd GD, Keith S, Faroon O, Buser M, Ingerman L, Hard C, et al. 2017. Toxicological profile for molybdenum: Draft for public comment.
- Underwood E, Somers M. 1969. Studies of zinc nutrition in sheep. I. The relation of zinc to growth, testicular development, and spermatogenesis in young rams. *Australian Journal of Agricultural Research* 20:889-897.
- Vittinghoff E, McCulloch CE. 2007. Relaxing the rule of ten events per variable in logistic and cox regression. *Am J Epidemiol* 165:710-718.
- Vivoli G, Fantuzzi G, Bergomi M, Tonelli E, Gatto MR, Zanetti F, et al. 1990. Relationship between zinc in serum and hair and some hormones during sexual maturation in humans. *Sci Total Environ* 95:29-40.
- Vyskočil A, Viau C. 1999. Assessment of molybdenum toxicity in humans. *Journal of applied toxicology* 19:185-192.
- Wacharasindhu S. 2017. A trend of normal puberty around the world. *Siriraj Medical Journal* 61:1-2.
- Wagner PA, Bailey LB, Christakis GJ, Dinning JS. 1985. Serum zinc concentrations in adolescents as related to sexual maturation. *Hum Nutr Clin Nutr* 39:459-462.
- Wallace DR. 2019. Metals as endocrine disruptors in the environment. *EC Pharmacology and Toxicology* ECO 2:12-14.
- Watkins DJ, Peterson KE, Ferguson KK, Mercado-García A, Tamayo y Ortiz M, Cantoral A, et al. 2016. Relating phthalate and BPA exposure to metabolism in peripubescence: The role of exposure timing, sex,

and puberty. *The Journal of Clinical Endocrinology* 101:79-88.

Watkins DJ, Sánchez BN, Téllez-Rojo MM, Lee JM, Mercado-García A, Blank-Goldenberg C, et al. 2017. Phthalate and bisphenol a exposure during in utero windows of susceptibility in relation to reproductive hormones and pubertal development in girls. *Environmental research* 159:143-151.

Williams PL, Sergeev O, Lee MM, Korrick SA, Burns JS, Humblet O, et al. 2010. Blood lead levels and delayed onset of puberty in a longitudinal study of russian boys. *Pediatrics* 125:e1088-1096.

Williams PL, Bellavia A, Korrick SA, Burns JS, Lee MM, Sergeev O, et al. 2019. Blood lead levels and timing of male sexual maturity: A longitudinal study of russian boys. *Environ Int* 125:470-477.

World Health Organization. 2017. *Inheriting a sustainable world? Atlas on children's health and the environment*:World Health Organization.

Wu Y, Peterson KE, Sanchez BN, Dolinoy DC, Mercado-Garcia A, Tellez-Rojo MM, et al. 2018. Association of blood leukocyte DNA methylation at line-1 and growth-related candidate genes with pubertal onset and progression. *Epigenetics* 13:1222-1233.

Wyshak G, Frisch RE. 1982. Evidence for a secular trend in age of menarche. *N Engl J Med* 306:1033-1035.

Yilmaz B, Terekeci H, Sandal S, Kelestimur F. 2019. Endocrine disrupting chemicals: Exposure, effects on human health, mechanism of action, models for testing and strategies for prevention. *Reviews in Endocrine and Metabolic Disorders*:1-21.

Zeng X, Lin T, Zhou Y, Kong Q. 2002. Alterations of serum hormone levels in male workers occupationally exposed to cadmium. *J Toxicol Environ Health A* 65:513-521.

Zeng X, Jin T, Buchet JP, Jiang X, Kong Q, Ye T, et al. 2004. Impact of cadmium exposure on male sex hormones: A population-based study in china. *Environ Res* 96:338-344.

Figures





Figure 1

Recruitment, sample collection and examination timeline for ELEMENT cohort



^a Results are presented as the percent difference in peripubertal hormone levels associated with an interquartile range (IQR) increase in *in utero* and peripubertal metal


^b Linear regression models were adjusted for child age, BMI z-score and specific gravity

* Significant associations with p value < 0.05 & q value < 0.15

* Significant associations with p value < 0.05 & q value > 0.15

Figure 2

Peripubertal hormone levels associated with in utero and peripubertal metal concentration among ELEMENT boys(ab).




^a GEE models were adjusted for child age and BMI z-score (baseline and change)

* Significant associations with p value<0.05 & q value<0.15

* Significant associations with p value<0.05 & q value>0.15

Figure 3

Odds Ratios (95% CI) for the Generalized Estimating Equations of in utero Metal Exposure and Tanner Stages/testicular volume(a)



^a GEE models are adjusted for child age and BMI z-score (baseline and change)

* Significant associations with p value < 0.05 & q value < 0.15

* Significant associations with p value < 0.05 & q value > 0.15

Figure 4

Odds Ratios (95% CI) for the Generalized Estimating Equations of peripubertal Metal Exposure and Tanner Stages/testicular volume(a)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementalInformation.pdf](#)