Factors influencing early initiation of breastfeeding among mothers in Rajshahi district, Bangladesh

Ummay Ayesha

University of Rajshahi

Abu Sayed Md. Al Mamun

University of Rajshahi

Md. Nurul Islam

University of Rajshahi

Md. Ripter Hossain

University of Rajshahi

Samme Amena Tasmia

University of Rajshahi

Md. Golam Hossain (✉ hossain95@yahoo.com)

University of Rajshahi <https://orcid.org/0000-0002-3822-5489>

Research article

Keywords: Breast feeding, Early initiation, Newborn, Prevalence, Bangladesh

Posted Date: December 16th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-34757/v3>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Early initiation of breastfeeding (EIBF) act as the first immunization for the newborns, fortify their immune defense system and eventually reduce the mortality rate. This study aimed to identify the influential factors of EIBF among mothers in Rajshahi district, Bangladesh.

Methods: A total number of 421 mothers living in Rajshahi district who had at least one child aged 6-24 months were considered as sample. Mothers were selected using multistage random sampling. This study was conducted from January to March, 2019. The EIBF was measured by a principal question, "Did you provide your breast milk to your newborns within one hour after delivery?" Frequency distribution and Chi-square test and multivariable binary logistic regression model were utilized in this study for getting prevalence and influential factors of EIBF respectively.

Results: This study revealed that the prevalence of EIBF among mothers in Rajshahi district was 88.4%. Multivariable logistic model provided eight influential factors of EIBF: (i) husbands' education level, (ii) husbands' occupation, (iii) family monthly income, (iv) mothers' age, (v) mothers' BMI, (vi) place of delivery, (vii) getting pregnancy with planning, and (viii) mothers taking advice regarding the benefit of breastfeeding during their pregnancy.

Conclusions: This study identified several modifiable influential factors of EIBF. The customs, culture and other characteristics are almost the same across the country. These factors could be considered to increase the rate of EIBF among mothers in Bangladesh.

Background

Breastfeeding is considered one of the most important factors for the growth and development of newborns [1]. Breast milk is the best source of nutrition for the newborn which is uniquely tailored to meet all the nutritional needs of human babies for the first six months of life [1]. Early initiation of breastfeeding or timely initiation of breastfeeding is referred to as providing the breast milk to the newborn within one hour of birth which ensures that they receives the colostrum [2]. Colostrum, the "first milk" produced by the mothers during the first postpartum days, endowed with protective antibodies that inevitably work as the first immunization for the newborns. It contains at least ninety known components including amino acids, minerals and vitamins essential for the growth and development of the newborns [3]. Early initiation of breastfeeding ensures skin-to-skin contact between the mothers and the newborns which helps in preventing hypothermia of the new-born baby, establishes the bond between mothers and children, and most importantly boosts the chances of increasing exclusive breastfeeding practice [2].

EIBF also shows a significant protective role to mothers by reducing the risk of postpartum hemorrhage which is a leading cause of maternal mortality in low income countries [4]. Child mortality is comparatively higher in those low and middle income countries where the prevalence of EIBF is low [5]. Recently, South Asia has made more progress in improving the rate of EIBF than any other region of the world. However, still more than 20 million newborns are not being breastfed within the first hour of birth

[6]. Among South Asian countries, Pakistan (18%) was the lowest EIBF [7] followed by Afghanistan (41%), Nepal (41.8%) [8] and India (42%) [9]. According to BDHS-2014, prevalence of EIBF was 51% in Bangladesh. The prevalence of EIBF was 90%, 78% and 64% in Sri Lanka, Bhutan and Maldives respectively [6]. There were some socio-economic, health-related and individual barriers to EIBF in South Asian countries found by Sharma and Byrne. They also found the limited evidence on the health care system in relation to EIBF [10].

In Bangladesh, the trend of practicing early initiation of breastfeeding among the lactating mothers has been increasing during the last one decade [11]. It is important to sort out the factors that influence the early initiation of breastfeeding to implement strategies and interventions that could speed up the government efforts in improving early initiation of breastfeeding trends among mothers in Bangladesh. More recently, Islam et al. [12] investigated the prevalence and factors associated with early initiation of breastfeeding among Bangladeshi mothers using a nationally representative samples collected by BDHS-2014. BDHS collected data in 2014, since then 6 years have already passed, and some indicators related to early initiation of breastfeeding could have been changed in Bangladesh. They found the prevalence of EIBF among mothers in Bangladesh was 51%, and they also reported that 89% of newborns are breastfed within one day after delivery. The household wealth quintile, medical facilities and women education level have been increasing in Bangladesh during the last two decades; these factors are directly or indirectly related to EIBF [11]. Bangladesh Government is providing health care facilities for ensuring the antenatal and postnatal care services of mothers in Bangladesh consequently the rate of EIBF is increasing [11]. Bangladesh Government is trying to achieve 17 goals under SDGs by 2030. One of the important goals of SDGs is to reduce maternal and under-five child mortality, and EIBF is one of the important factors for reducing maternal and under-five child mortality. It is necessary to determine the current prevalence and associated factors of EIBF among Bangladeshi mothers. Rajshahi district is a part of the northern region of Bangladesh. BDHS-2014 did not find a significant variation in the rate of EIBF among geographical locations (divisions) in Bangladesh [11]. To the best of our knowledge, no research has been conducted in Rajshahi district regarding this issue. The present study may help to increase the rate of EIBF among mothers living in this region.

The aim of this study was to determine the prevalence and associated factors of early initiation of breastfeeding among mothers in Rajshahi district, Bangladesh.

Methods

Study area and population: Rajshahi district was the target area of the present study. It is one of the oldest districts of Bangladesh that belongs to the Northern region of the country under Rajshahi division. The area of Rajshahi district is 2407.01 sq km [13]. A total number of population living in Rajshahi district is 22,868,74 including 11,844,48 males and 11,024,26 females [14]. All mothers who had at least one child (age, 6-24 months) were considered as the study population. This was a cross-sectional study.

Sample size determination: The required sample size for this study was estimated using the formula

given by Cochran [15]:
$$n = \frac{z^2 p(1-p)}{d^2},$$
 where n= the required sample size, p is the proportion of prevalence of early initiation of breastfeeding (here, p= 0.514) and z=1.96 at 95% confidence interval, and d is the margin of error, we considered d=0.05. The prevalence of initiation of breastfeeding (p-value) was taken from a previous publication [12]. This formula provided that 384 sample was adequate for our present study. However, 440 samples (15% extra) were considered for the study for allowing some failure cases.

Sampling: Multistage random sampling was utilized for this study. In the first stage, 2 Upazilas were selected randomly from 9 Upazilas in Rajshahi district. In the second stage, 2 unions were selected from each selected Upazila randomly. Similarly, 1 ward was selected randomly from 30 wards of Rajshahi City Corporation. In the third and final stage, 80 mothers were selected from each selected union and 120 mothers were selected from the selected ward randomly. In each stage, a simple random sampling method (by lottery) was used. All necessary information of our respondents was collected from the respective ward councilor's office/union parishad. Before, collecting data, we discussed with selected mothers and their husbands/guardians about our present study. Unfortunately, 19 selected mothers did not agree to provide their information. Finally, 421 mothers provided their information which was analyzed in this study.

Data collection procedures: A self-administered questionnaire was used for collecting information from selected mothers during the period January to March, 2019. The questionnaire was draft and sent to five experts in health sciences for taking their opinions/suggestions to improve it. According to experts' comments/suggestions the questionnaire was modified and finalized for data collection. The original questionnaire was in English, and the revised questionnaire was translated into Bangla (mother tongue of Bangladesh), and the Bangla questionnaire was checked by present authors. A pilot survey had been done for observing whether there was any lacking or drawback in the questionnaire. We did not get lacking or drawbacks.

Measurement of anthropometric data: Digital scales and a portable stadiometer was used to measure weight and height of our samples respectively. Measurement of individuals was taken without shoes and wearing light clothes using the techniques of Martin and Saller [16]. Height and weight were measured to the nearest 1 cm and 0.1 kg, respectively, and body mass index (BMI) was calculated using the formula, $BMI = \text{weight (kg)} / \{\text{height (m)}\}^2$. BMI was classified into three classes: (i) underweight ($18.5 \text{ kg/m}^2 < BMI$), (ii) normal weight ($18.5 \leq BMI < 25 \text{ kg/m}^2$) and (iii) overweight ($BMI \geq 25 \text{ kg/m}^2$) [12]. In addition, mothers' age was calculated by the difference between the date of interview and their date of birth, and we considered the nearest higher integer.

Outcome variable: Early initiation of breastfeeding (EIBF) was considered as the outcome variable in this study. It was measured by a question, "Did you provide your breast milk to your newborn within one hour after delivered? EIBF was expressed as a dichotomous variable with category 1 for initiation of

breastfeeding within one hour (early initiation) and category 0 for initiation of breastfeeding after one hour (late initiation).

Independent variables: Some socio-economic, demographic and anthropometric factors were considered as independent variables for this study on the basis of previous studies [12, 17-18].

Statistical analysis: Frequency distribution (percentage) was used to determine the prevalence of EIBF among mothers living in Rajshahi district, Bangladesh. Chi-square test was conducted to assess the association between independent variables and the EIBF, the significant associated factors were considered as independent variable in multivariable binary logistic regression model. Binary multivariable logistic regression analysis was used to find the effect of socio-economic, demographic, anthropometric and behavioral factors on EIBF among mothers in Rajshahi district.

The underlying multivariable binary logistic regression can be described as follows:

$$\text{Log}[P/1-P] = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \dots + \beta_K X_K$$

where P = Probability of early initiation of breastfeeding (coded as 1)

1-P = Probability of late initiation of breastfeeding (coded as 0)

X_i s are respective independent variables ($i=1,2,\dots,k$), and β_i s are the regression coefficients. We used the magnitude of the standard error (SE) for detecting the multicollinearity problem among the independent variables, if the magnitude of the SE lies between 0.001 and 0.5, it was judged that there was no evidence of multicollinearity [19]. A value of $p < 0.05$ was considered statistically significant in the analysis. All statistical analyses were performed using SPSS (IBM Version 21).

Results

A total number of 421 samples were selected from mothers having at least one child (age, 6-24 months) in Rajshahi district, Bangladesh to determine the prevalence and associated factors of EIBF.

We found that 88.40% of mothers provided their breast milk to their newborns within one hour after delivery (Table 1). Out of the total 421 sample, 47.7% of mothers were primary or uneducated while 11.2% were higher educated, and 47.5% of their husbands were primary or uneducated and 14.7% of got higher education. Most of the mothers (98.57%) were housewife. More than 68% of mothers were living in low-income families (below 15000 Taka) and living in rural areas (71%). Around 50% mothers' age at first birth was below 20 years. More than 67% were normal weight while 25.7% mothers were underweight. Among the respondent, 42% of mothers delivered their child by C-section. Near half of the mothers (47.5%) delivered their child in public hospitals. About 95% of mothers took advice during pregnancy regarding the benefit of breastfeeding from health providers (Table 1). Chi-square test provided that mothers' and their husbands' education level, mothers' occupation, their husbands' occupation, monthly family income, type of residence, mothers' age, mothers' BMI, respondents' age at first birth, mode of delivery, place of

delivery, taking advice regarding the benefit of breastfeeding during pregnancy were significantly associated factors of initial breastfeeding among Rajshahi mothers (Table 1). These factors were considered as independent variables in multivariable binary logistic regression models.

Table 2 shows the effect of socio-economic and demographic factors on EIBF among mothers in Rajshahi district, Bangladesh. Results of multivariable binary logistic model were interpreted using p-value, adjusted odds ratio (AOR) with 95% confidence interval (CI) for AOR. Standard error (SE) showed that there was no evidence of multicollinearity problem among independent variables except the type of residence due to all rural mothers provided EIBF. Following adjustment for other variables, the logistic model demonstrated that delivery in private hospitals was associated with a 91 % decreased in the odds of EIBF (AOR=0.090, 95% CI: 0.010-0.794; $p<0.05$). The odds of EIBF was decreased by 90.3% and 87.0% among mothers who were living in middle-income families (AOR=0.097, 95% CI: 0.030 -0.315; $p<0.01$) and rich families (AOR=0.130, 95% CI: 0.039-0.437; $p<0.01$) compared to mothers living in poor income families respectively. We found that higher educated husbands' wives was likely to decrease the odds of EIBF by 83.3% (AOR=0.167, 95% CI: 0.035 -0.792; $p<0.05$) compared to primary or uneducated husbands' wives. The BMI of mothers was an important predictor of initial breastfeeding, and it was observed that the odds of EIBF was diminished by 89.2% and 92.8% among normal weight (AOR=0.108, 95% CI: 0.013-0.867; $p<0.05$) and overweight mothers (AOR=0.072, 95% CI: 0.008-0.692; $p<0.05$) compared to underweight mothers. It was also found that the odds of EIBF was 7.133, 3.069 and 7.385 fold higher among mothers aged 20-24 years (AOR=7.133, 95% CI: 2.273-22.381; $p<0.01$), 25-29 years [AOR=3.069, 95% CI: 1.047-8.997; $p<0.05$], 30-34 years (AOR=7.385, 95% CI: 1.623-33.607; $p<0.01$) than mothers aged 35 years and above. Farmers' wives provided their initial breast milk to their newborns had 13.568 times (AOR=13.568, 95% CI: 5.237-35.154; $p<0.01$) higher than others professional husbands' wives. Mothers who got pregnancy with proper planning had provided EIBF 5.941 fold higher (AOR=5.941, 95% CI: 2.237-15.780; $p<0.01$) than those who did not do it. Mothers who took advice regarding the benefit of breastfeeding during their pregnancy had 7.502 times (AOR=7.502, 95% CI: 2.877-19.559; $p<0.01$) higher to provide EIBF than mothers who did not get it. Hosmer and Lemeshow test demonstrated that our selected model was good fitted, and the model can able to explain the variation of outcome variable by 56.60% (Nagelkerke R^2 - value =0.566) (Table 2).

Discussion

In the present study, we tried to determine the prevalence of EIBF and investigate the associated factors of EIBF among mothers living in Rajshahi district, Bangladesh. It was noted that the prevalence of EIBF among mothers was 88.40%. The national survey of Bangladesh (BDHS-2014) showed that 51.0% of newborns were breastfed within one hour after birth (EIBF) and 89% within one day after delivery, and 55.0% of newborns under the age of 6 months were exclusively breastfed [11]. However, more recently published key indicators of BDHS 2017-2018 showed that 65% of newborns under the age of 6 months were exclusively breastfed (EB) in 2017, a level markedly higher than that found in BDHS-2014 [20]. The prevalence of EIBF is not available in the report of BDHS 2017-2018 [20]. BDHS-2014 found that 98% of

children born during the two years of their survey were breastfed at some point in their life in Bangladesh where breastfeeding was almost universal [11]. The rate of breastfeeding for every stage has been increasing with increasing medical facilities in Bangladesh. Mothers can easily get advice about the benefit of breastfeeding from health providers and family planning workers. Also, the practice of providing EIBF was increasing with increasing literacy rate among women which enhanced the awareness regarding the benefit of EIBF especially among rural women in Bangladesh [11]. BDHS-2014 conducted their survey about 6 years ago; some indicators such as mother education level, medical facilities, household wealth quintile have continued to increase during the last two decades in Bangladesh which is ultimately helping to increase the rate of EIBF.

The prevalence of EIBF among mothers in Rajshahi district was also higher than nationally representative sample in Bangladesh [11], and some South Asian countries like India (42.0 %) [9], Pakistan (18.0 %) [7], Nepal (41.8) [8], Afghanistan (41.0%) [7], and others developing countries like Nigeria (34.7%) [21], Iran (32.2%) [22] and South Sudan (48%) [23]. The prevalence of EIBF in Ethiopia [24] was close (83.7%) to our finding.

Mothers who delivered at home were more likely to provide EIBF to their newborn than mothers delivered at private hospitals. A Bangladeshi study found that mothers undergoing vaginal delivery were more likely to provide EIBF to their newborns than mothers having cesarean delivery [12]. The present finding coincided with a Chinese study [25]. It was noted that mothers living in poor-income families were more interested to provide initial breast milk to their newborns than those of middle- and high-income families, and the rate of EIBF was decreasing with increasing family income. The same results had been found in another Bangladeshi study that used BDHS-2014 data [12].

We found that primary or uneducated husbands' wives had more chances to give their initial breast milk to their newborns than higher educated husbands' wives. This finding was supported by another study [26]. They also found that the husband's educational level was significantly associated with EIBF.

The BMI of mothers was an important predictor of initial breastfeeding, and it was observed that underweight mothers were more likely to provide their breast milk to their newborns than normal weight and overweight mothers. The same results had been found in our nationally representative samples [12]. A similar observation was also mentioned by several studies [27-30, 23-25]. This study found that mothers aged 20-34 years were more likely to provide EIBF to their children than mothers aged 35 years and above. Our result was also supported by a global survey [31-32]. Again, in this study, it was found that husband's occupation is an important factor for EIBF, i.e. farmers' wives had a higher chance to give EIBF to their children than other professional husbands' wives. Our result was also in line with another study in India [33], but disagreed with a study in Sudan [34].

In this study, we have found that , husbands' education level, family income, mothers' BMI and place of delivery are important factors for providing initial breast milk to newborn among mothers in Rajshahi district. These four factors are very much related to each other in developing countries like Bangladesh. Most of the home delivery mothers are living in poor families. In Bangladesh, the wife is dominated by her

husband, and most of the females are dependent on their husbands' income, and income is dependent on education level. Usually, uneducated or primary educated husbands are farmers or day labors living in rural or slum areas, their income is not sufficient to maintain their family. Therefore, cannot provide sufficient food to their family members, consequently, they become underweight. . Mothers living in poor families cannot go to hospitals/clinics for delivery, and most of them deliver at home in the presence their close relatives without proper nursing. Traditional customs and culture in the society of Bangladesh are that, after delivery, the mother immediately provides her breast milk to the newborn. It is true especially for most of the mothers having vaginal delivery. It is also mentionable that all home deliveries are vaginal. On the other hand, mothers living in middle or rich families usually deliver at hospitals/clinics under proper nursing. But now a days most of the hospital/clinic deliveries are caesarian, and after the delivery, mothers stay at the operation theatre for more than one hour, and cannot provide breast milk to them in time (within one hour of delivery). Mothers living in poor families are more likely to provide breast milk to their newborn than those living in rich families. A similar result was found in another Bangladeshi study [12]. Normal weight and overweight mothers usually live in rich families, and most of them underwent caesarean section. It might create a higher risk of cephalo-pelvic disproportion and relatively poor progress due to maternal fatigue [35]. This is one of the most important reasons for the differences seen in the practice of EIBF between underweight and overweight mothers. Normal weight and overweight mothers should be the focus of education on the potential benefits of EIBF. We found that mothers getting pregnant with proper planning and taking advice regarding the benefit of EIBF were more likely to provide EIBF than their counterparts. Family planning workers closely working with pregnant mothers and healthcare providers can play a good role to increase the rate of EIBF and exclusive breastfeeding in Bangladesh.

This study may recommend health authorities to take necessary steps for making awareness of mothers about benefit of initial breastfeeding. The findings of our study can help Bangladesh Ministry of Health and Family Welfare to enhance their programmes on appropriate breastfeeding practices at the community level.

Strength and limitations of this study: Perhaps, this was the first time we attempted to investigate the early initiation of breastfeeding and its influential factors among mothers in Rajshahi district, Bangladesh. We considered two new factors; (i) getting pregnant with planning and (ii) mothers taking advice regarding the benefit of breastfeeding during their pregnancy, which were not considered in our national survey. However, there are several limitations of our present project. Firstly, in this study, we considered only Rajshahi district as our study area which is a small part of Bangladesh. Secondly, we used a quantitative cross sectional study which can determine only associated factors but cannot do research in-depth. Mixed research both qualitative and quantitative is required for doing in-depth study. Thirdly, we selected some socio-economic, demographic, anthropometric and behavioral factors as independent variables but other important factors such as parity, gestational age at delivery, maternal obstetric complications, and newborn status were not considered in this study. Fourth, we did not consider the pre-lacteal feeding for calculating the prevalence of EIBF, and it would be better to ask mothers about providing EIBF immediately after delivery for getting accurate information; but we asked

after 6-24 months of delivery. On the basis of our limitations, we may proclaim that further researches are required on breastfeeding among Bangladeshi mothers.

Conclusions

In this study, we considered 421 mothers living in Rajshahi district, Bangladesh having at least one child aged 6-24 months as samples to determine of EIBF and its related factors. Our selected statistical models provided that the rate of EIBF among mothers in Rajshahi district was 88.40%; the predictors of EIBF were: husbands' education level and occupation, family monthly income, mothers' age, mothers' BMI, place of delivery, getting pregnant with proper planning and taking advice regarding the benefit of EIBF. These findings would help the health authorities of Bangladesh government and non-government organizations working with mothers and children health and nutrition for improving their strategies to increase the rate of EIBF.

Abbreviations

AOR- Adjusted Odds Ratio; BDHS- Bangladesh Demographic and Health Survey; EB- Exclusively breastfed; EIBF-Early initiation of breastfeeding; IBM- International Business Machines; SE- Standard error; SPSS- Statistical Package for the Social Sciences.

Declarations

Ethics approval and consent to participate: We followed all rules and regulations of Institute of Biological Science (IBSc), University of Rajshahi, Bangladesh, and IBSc approved this study and provided an ethical clearance letter (Memo No: 69/320/IAMEBBC/IBSc). Written consent was taken from each participant. The information was kept in confidential. The consent was taken from the husband or guardian for participants under 16 years old.

Consent for publication: Not applicable to this study.

Availability of data and materials: The study was based on the primary data. The data will be provided when necessary.

Competing interests: The authors declare that they have no competing interests.

Funding: The authors have no support or funding to report.

Authors' contributions: UA and GH created the concept; ASMAM, MNI, MRH and GH created the design of the study; ASMAM and SAT performed the statistical analysis; UA, ASMA, MIN, RH and GH drafted the manuscript. GH, ASMAM, and SAT made revision of the manuscript. All authors read and approved the final manuscript.

Acknowledgement: The authors gratefully acknowledge the authority of the councilor office and union parishad for providing useful information about our subjects. Also thanks to all participants for giving their information.

Authordetails: Department of Statistics, University of Rajshahi, Rajshahi 6205, Bangladesh.

References

1. United Nations Children's Fund. 1990–2005: Celebrating the Innocenti Declaration on the Protection, Promotion and Support of Breastfeeding: Past Achievements, Present Challenges and Priority Actions for Infant and Young Child Feeding, 2006. 2nd ed. UNICEF Innocenti Research Centre, Florence, Italy.
2. World Health Organization e-Library of Evidence for Nutrition Actions (eLENA), Early Initiation of Breastfeeding, WHO; 2014.
3. Tapa BR. Health factors in colostrum. *Indian Journal of Pediatric*. 2005; 72(7):579–581.
4. Berde AS, Yalcin SS. Determinants of early initiation of breastfeeding in Nigeria: a population-based study using the 2013 demographic and health survey data. *BMC Pregnancy Childbirth*. 2016; 16:32.
5. Nkala TE, Msuya SE. Prevalence and predictors of exclusive breastfeeding among women in Kigoma region, Western Tanzania: a community based cross-sectional study. *Int Breastfeed J*. 2011;6:
6. 2016. From the First Hour of Life: Making the Case for Improved Infant and Young Child Feeding Everywhere. UNICEF: New York.
7. 2018. Every Child Alive. The Urgent Need to End Newborn Deaths. UNICEF: New York.
8. Bhandari S, Thorne-Lyman, AL, Shrestha et al. Determinants of infant breastfeeding practices in Nepal: a national study. *Int Breastfeed*. 2019; 14:14.
9. Aguayo VM, Gupta G, Singh G, Kumar R. Early initiation of breast feeding on the rise in India *BMJ Global Health*. 2016; 1:
10. Sharma IK, Byrne A. Early initiation of breastfeeding: a systematic literature review of factors and barriers in South Asia. *Int Breastfeed*. 2016; 11:17.
11. National Institute of Population Research and Training (NIPORT), Mitra and Associates, and ICF International. Bangladesh Demographic and Health Survey 2014: Dhaka, Bangladesh, and Rockville, Maryland, Mitra and Associates, and ICF International.
12. Islam MA, Mamun ASMA, Hossain MM, Bharati P, Saw A, Lestrel PE, et al. Prevalence and factors associated with early initiation of breastfeeding among Bangladeshi mothers: A nationwide cross-sectional study. *PLoS One*. 2019; 14(4):e0215733. doi: 10.1371/journal.pone.0215733.
13. Rahman, MM. Rajshahi District. Islam, S, Ahmed AJ. (eds.). *Banglapedia: National Encyclopedia of Bangladesh*(Second ed.). Asiatic Society of Bangladesh. 2012.
14. Climate Change Cell (CCC). Climate Change and Health Impacts in Bangladesh. Climate Change Cell, DoE, MoEF; Component 4b, CDMP, MoFDM. 2009, Dhaka.

15. Cochran WG. Sampling Techniques. 1963; 2nd Ed., New York: John Wiley and Sons, Inc.
16. Martin, R. & Saller, K. Lehrbuch der anthropologie. 1957. Gustav Fischer Verlag, Stuttgart.
17. Pamarthi K, Palli JK. Factors influencing early initiation of breast feeding among mothers in North Coastal Andhra Pradesh, India. *Int J Community Med Public Health*. 2019; 6:4316-21.
18. Borah M, Baruah J, Baruah R, Baruah M. Prevalence and factors affecting early initiation of breastfeeding in rural areas of Dibrugarh district, Assam. *Int J Community Med Public Health*. 2019; 6:2176-81.
19. Chan YH. Biostatistics 202: logistic regression analysis. *Singapore Med J*. 2004; 45(4):149-53.
20. The DHS Program. Bangladesh: DHS 2017-18 - Key Indicators Report. 2019.
<https://dhsprogram.com/pubs/pdf/PR104/PR104>.
21. Berde AS, Yalcin SS. Determinants of early initiation of breastfeeding in Nigeria: a population-based study using the 2013 demographic and health survey data. *BMC Pregnancy Child birth*. 2016; 16(32). doi: 10.1186/s12884-016-0818-y.
22. Zarshenas M, Zhao Y, Binns CW, Scott JA. Determinants of in-hospital feeding practices in Shiraz, Iran: Results of a prospective cohort study. 2019; 46(1):137-145.
23. Tongun JB, Sebit MB, Mukunya D, Ndeezi G, Nankabirwa V, Tylleskar T, et al. Factors associated with delayed initiation of breastfeeding: a cross-sectional study in South Sudan. *Int Breastfeed J*. 2018; 13:28. <https://doi.org/10.1186/s13006-018-0170-0>.
24. Beyene MG, Geda NR, Habtewold TD, Assen ZM. Early initiation of breastfeeding among mothers of children under the age of 24 months in Southern Ethiopia. *Int Breastfeed J*. 2016; 12 (1). doi: 10.1186/s13006-016-0096-3.
25. Wu Y, Wang Y, Huang J, Zhang Z, Wang J, Zhou L. The association between caesarean delivery and the initiation and duration of breastfeeding: a prospective cohort study in China. *European Journal of Clinical Nutrition*. 2018; 72:1644–1654. <https://doi.org/10.1038/s41430-018-0127-9>.
26. Hossain, M., Islam, A., Kamarul, T. *et al.* Exclusive breastfeeding practice during first six months of an infant's life in Bangladesh: a country based cross-sectional study. *BMC Pediatr* 18,93 (2018).
<https://doi.org/10.1186/s12887-018-1076-0>
27. Kiani SN, Rich KM, Herkert D, Safon C, Pérez-Escamilla R. Delivery mode and breastfeeding outcomes among new mothers in Nicaragua. *Maternal and child nutrition*. 2018; 14(1):e12474.
<https://doi.org/10.1111/mcn.12474>.
28. Krause KM, Lovelady CA, Østbye T. Predictors of Breastfeeding in Overweight and Obese Women: Data From Active Mothers Postpartum (AMP). *Matern Child Health J*. 2011; 15(3):367- doi: 10.1007/s10995-010-0667-7.
29. Liu J, Smith MG, Dobre MA, Ferguson JE. (2010). Maternal obesity and breast-feeding practices among white and black women. *Obesity*. 2010; 18:175–182.
30. Baker JL, Michaelsen KF, Sørensen TI, Rasmussen KM. High prepregnant body mass index is associated with early termination of full and any breastfeeding in Danish women. *Am J Clin Nutr*.

2007; 86: 404–411.

31. Takahashi K, Ganchimeg T, Ota E, Vogel JP, Souza JP, Laopaiboon M, et al. Prevalence of early initiation of breastfeeding and determinants of delayed initiation of breastfeeding: secondary analysis of the WHO Global Survey. *Sci Rep.* 2017; 7:44868. doi:10.1038/srep44868.
32. Radwan H. Patterns and determinants of breastfeeding and complementary feeding practices of Emirati mothers in the United Arab Emirates. *BMC Public Health.* 2013; 13: 171.
33. Sharma A, Thakur PS, Tiwari R, Kasar PK, Sharma R, Kabirpanthi V. Factors associated with early initiation of breastfeeding among mothers of tribal area of Madhya Pradesh, India: a community based cross sectional study. *Int J Community Med Public Heal.* 2016; 3: 194–9.
34. Hassan AA, Taha Z, Ahmed MAA, Ali AAA, Adam I. Assessment of initiation of breastfeeding practice in Kassala, Eastern Sudan: a community-based study. *International Breastfeeding Journal.* 2018; 13: 34. <https://doi.org/10.1186/s13006-018-0177-6>.
35. Yu Z, Han S, Zhu J, Sun X, Ji C, Guo X. Pre-Pregnancy Body Mass Index in Relation to Infant Birth Weight and Offspring Overweight/Obesity: A Systematic Review and Meta-Analysis. *PLoS One.* 2013; 8(4):e61627. doi: 10.1371/journal.pone.0061627.

Tables

Table 1: Association between early initial breastfeeding and socio-economic, demographic and anthropometric factors

Variable	Group, N(%)	Early initiation of breastfeeding status		Chi-square/ Fisher's exact value	p-value
		No (11.60%) N(%)	Yes (88.40%) N(%)		
Mothers' educational level	Uneducated or primary, 201(47.7)	10(5.0)	191(95.0)	25.061	p<0.001
	Secondary, 173(41.1)	25(14.5)	148(85.5)		
	Higher, 47(11.2)	14(29.8)	33(70.2)		
Husbands' educational level	Uneducated or primary, 200(47.5)	8(4.0)	192(96.0)	61.333	p<0.001
	Secondary, 159(37.8)	16(10.1)	143(89.9)		
	Higher, 62(14.7)	25(40.3)	37(59.7)		
Mothers' occupation	House wife, 415(98.57)	46(11.1)	369(88.9)	8.709	0.023
	Others, 6(1.43)	3(50.0)	3(50.0)		
Husbands' occupation	Farmer, 235(55.82)	5(2.1)	230(97.9)	46.789	p<0.001
Monthly family income (Taka)		6(2.1)	280(97.9)	85.918	p<0.001
	<15000, 287(68.2)				
	15000-25000, 58(13.8)	14(24.1)	44(75.9)		
Type of residence	>25000, 76(18.0)	29(38.2)	47(61.8)	135.908	p<0.001
	Urban, 122(28.98)	49(40.2)	73(59.8)		
	Rural, 299(71.02)	0(0)	299(100)		
Mothers' age (year)	15-19, 17(4.04)	9(52.9)	8(47.1)	42.936	p<0.001
	20-24, 181(42.99)	11(6.1)	170(93.9)		
	25-29, 153(36.34)	20(13.1)	133(86.9)		
	30-34, 51(12.11)	3(5.9)	48(94.1)		
	35 and above, 19(4.52)	6(31.6)	13(68.4)		
Mothers' BMI	Underweight, 08(25.7)	1(0.9)	107(99.1)	28.629	p<0.001
	Normal weight, 85(67.7)	38(13.3)	247(86.7)		
	Overweight, 28(6.6)	10(35.7)	18(64.3)		
Age at first birth (year)	<20, 199(47.3)	14(7.0)	185(93.0)	7.777	0.005
	≥20, 222(52.7)	35(15.8)	187(84.2)		
Type of family	Nuclear, 346(82.19)	36(10.4)	310(89.6)	2.877	0.090
	Joint, 75(17.81)	13(17.3)	62(82.7)		
Gender of children	Boy, 212(50.36)	21(9.9)	191(90.1)	1.248	0.264
	Girl, 209(49.64)	28(13.4)	181(86.6)		
	Others, 186(44.18)	44(23.7)	142(76.3)		
Number of children	One, 233 (55.34)	22(9.4)	211(90.6)	2.449	0.118
	More than one, 188(44.66)	27(14.4)	161(85.6)		
Mode of delivery	Cesarean, 177(42)	33(18.6)	144(81.4)	14.572	p<0.001
	Vaginal, 244(58)	16(6.6)	228(93.4)		
Place of delivery	Home, 45(10.7)	1(2.2)	44(97.8)	31.936	p<0.001
	Public, 200(47.5)	9(4.5)	191(95.5)		
	Private, 176(41.8)	39(22.2)	137(77.8)		
Planning of conception	Yes, 352(83.60)	37(10.5)	315(89.5)	2.655	0.103
	No, 69(16.40)	12(17.4)	57(82.6)		
Taking advice during pregnancy	Yes, 399 (94.77)	39(9.8)	360(90.2)	25.810	p<0.0001

Table 2: Effect of socio-economic and demographic factors on initial breastfeeding among mothers in Rajshahi district, Bangladesh

Variable with groups	B	SE	Wald	p-value	AOR	95% CI for AOR	
						Lower	Upper
Mode of delivery (Caesarean Vs Vaginal ^R)	-1.183	0.323	13.450	0.001	0.306	0.163	0.576
Place of delivery			10.031	0.007			
Public Hospital Vs Home ^R	-1.024	1.126	0.827	0.363	0.359	0.040	3.263
Private Hospital Vs Home ^R	-2.405	1.109	4.698	0.030	0.090	0.010	0.794
Monthly family income (Taka)			16.443	0.000			
15000-25000 Vs ≤15000 ^R	-2.330	0.600	15.074	0.000	0.097	0.030	0.315
≥25000 Vs ≤15000 ^R	-2.042	0.620	10.864	0.001	0.130	0.039	0.437
Husbands' education level			6.323	0.042			
Secondary Vs Primary or Uneducated ^R	-0.463	0.669	0.480	0.489	0.629	0.170	2.335
Higher Vs Primary or Uneducated ^R	-1.792	0.796	5.074	0.024	0.167	0.035	0.792
Mothers, education level			4.474	0.107			
Secondary Vs Primary or Uneducated ^R	-0.031	0.625	0.003	0.960	0.969	0.285	3.296
Higher Vs Primary or Uneducated ^R	1.297	0.812	2.548	0.110	3.658	0.744	17.981
Mothers' age at first birth (year) (>20 Vs ≥20^R)	0.262	0.448	0.341	0.559	1.299	0.540	3.125
Mothers' BMI			5.208	0.074			
Normal Vs Under weight ^R	-2.227	1.064	4.383	0.036	0.108	0.013	0.867
Overweight Vs Underweight ^R	-2.630	1.154	5.193	0.023	0.072	0.008	0.692
Mothers' age (year)			31.742	p<0.001			
15-19 Vs 35 and above ^R	-0.891	0.693	1.655	0.198	0.410	0.106	1.594
20-24 Vs 35 and above ^R	1.965	0.583	11.341	p<0.001	7.133	2.273	22.381
25-29 Vs 35 and above ^R	1.121	0.549	4.177	0.041	3.069	1.047	8.997
30-34 Vs 35 and above ^R	1.999	0.773	6.688	0.010	7.385	1.623	33.607
Mothers' occupation, (House wife Vs Others^R)	1.221	.836	2.135	0.144	3.390	0.659	17.438
Husbands' occupation, Farmer Vs Others^R	2.608	0.486	28.823	p<0.001	13.568	5.237	35.154
Planning concept, Yes Vs No^R	1.782	0.498	12.780	p<0.001	5.941	2.237	15.780
Taking advice, Yes Vs No^R	2.015	0.489	16.986	p<0.001	7.502	2.877	19.559
Constant	7.601	1.548	24.110	p<0.001	2000.883		
Nagelkerke R²- value =0.566							
Hosmer and Lemeshow Test							
Chi-sauare value=5.568 p-value=0.696							

N.B: R=Reference Category, B=Co-efficient, SE=Standard Error, AOR=Adjusted Odds ratio, CI=Confidence Interval.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryFile.docx](#)