

Reconstruction of a catalogue of genome-scale metabolic models with enzymatic constraints using GECKO 2.0

Iván Domenzain

Chalmers University of Technology <https://orcid.org/0000-0002-5322-2040>

Benjamín Sánchez

Technical University of Denmark <https://orcid.org/0000-0001-6093-4110>

Mihail Anton

Chalmers University of Technology <https://orcid.org/0000-0002-7753-9042>

Eduard J Kerkhoven

Chalmers University of Technology <https://orcid.org/0000-0002-3593-5792>

Aarón Millán-Oropeza

INRAE, Micalis Institute

Céline Henry

INRA <https://orcid.org/0000-0002-2355-1791>

Verena Siewers

Chalmers University of Technology <https://orcid.org/0000-0002-9502-9804>

John Morrissey

UCC <https://orcid.org/0000-0001-7960-2001>

Nikolaus Sonnenschein

Technical University of Denmark

Jens Nielsen (✉ nielsenj@chalmers.se)

Chalmers University of Technology <https://orcid.org/0000-0002-9955-6003>

Article

Keywords: genome-scale metabolic models, GECKO, enzyme constraints

Posted Date: April 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-351350/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Nature Communications on June 30th, 2022. See the published version at <https://doi.org/10.1038/s41467-022-31421-1>.

Abstract

Genome-scale metabolic models (GEMs) have been widely used for quantitative exploration of the relation between genotype and phenotype. Streamlined integration of enzyme constraints and proteomics data into GEMs was first enabled by the GECKO method, allowing the study of phenotypes constrained by protein limitations. Here, we upgraded the GECKO toolbox in order to enhance models with enzyme and proteomics constraints for any organism with an available GEM reconstruction. With this, enzyme-constrained models (ecModels) for the budding yeasts *Saccharomyces cerevisiae*, *Yarrowia lipolytica* and *Kluyveromyces marxianus* were generated, aiming to study their long-term adaptation to several stress factors by incorporation of proteomics data. Predictions revealed that upregulation and high saturation of enzymes in amino acid metabolism were found to be common across organisms and conditions, suggesting the relevance of metabolic robustness in contrast to optimal protein utilization as a cellular objective for microbial growth under stress and nutrient-limited conditions. The functionality of GECKO was further developed by the implementation of an automated framework for continuous and version-controlled update of ecModels, which was validated by producing additional high-quality ecModels for *Escherichia coli* and *Homo sapiens*. These efforts aim to facilitate the utilization of ecModels in basic science, metabolic engineering and synthetic biology purposes.

Introduction

Genome-scale metabolic models (GEMs) have become an established tool for systematic analyses of metabolism for a wide variety of organisms¹⁻⁶. Their myriads of applications span from model-driven development of efficient cell factories^{3,7-9}, to their utilization for understanding mechanisms underlying complex human diseases¹⁰⁻¹². One of the most common simulation techniques for enabling phenotype predictions with these models is flux balance analysis (FBA), which assumes that there is balancing of fluxes around each metabolite in the metabolic network. This means that fluxes are constrained by stoichiometries of the biochemical reactions in the network, and that cells have evolved in order to operate their metabolism according to optimality principles^{13,14}. Quantitative determination of biologically meaningful flux distribution profiles is a major challenge for constraint-based methods, as optimal phenotypes can be attained by alternate flux distribution profiles¹⁵, caused by the presence of network redundancies that provide organisms with robustness to environmental and genetic perturbations. This limitation is often addressed by incorporation of experimental measurements of exchange fluxes (secretion of byproducts and uptake of substrates) as numerical flux constraints for the FBA problem. However, such measurements are not readily available for a wide variety of conditions and organisms.

In order to overcome these limitations, the concept of enzymatic limitations on metabolic reactions has been explored and incorporated by several constraint-based methods. Some of these have modelled enzyme demands of metabolic reactions by constraining metabolic networks with kinetic parameters and physiological limitations of cells, such as a crowded intracellular volume¹⁶⁻¹⁸, a finite membrane surface

area for expression of transporter proteins¹⁹ and a bounded total protein mass available for metabolic enzymes^{20–25}. All of these modelling frameworks have been successful at expanding the range of predictions of classical FBA, providing explanations for overflow metabolism and cellular growth on diverse environments for *Escherichia coli*^{16–19,21,23,25}, *Saccharomyces cerevisiae*^{22,25,26}, *Lactococcus lactis*²⁷ and even human cells^{20,24}. However, these modelling approaches were applied to metabolic networks of extensively studied model organisms, which are usually well represented in specialized resources for kinetic parameters such as the BRENDA²⁸ and SABIO RK²⁹ databases. Furthermore, collecting the necessary parameters for the aforementioned models was mostly done manually; therefore, no generalized model parameterization procedure was provided as an integral part of these methods.

Enzyme limitations have also been introduced into models of metabolism by other formalisms, for instance, Metabolic and gene Expression models (ME-models), implemented on reconstructions for *E. coli*^{30–33}, *Thermotoga maritima*³⁴ and *Lactococcus lactis*³⁵; and resource balance analysis models (RBA), on reconstructions for *E. coli*³⁶ and *Bacillus subtilis*^{36,37}. These formalisms succeeded at merging genome-scale metabolic networks together with comprehensive representations of macromolecular expression processes, enabling detailed exploration of the constraints that govern cellular growth on diverse environments. Despite the great advances for understanding cell physiology, provided by these modelling formalisms, accuracy on phenotype predictions is compromised by the large number of parameters that are required (rate constants for transcriptional, translational, protein folding and degradation processes), with most of these not being readily available in the literature. Moreover, these models encompass processes that differ radically in their temporal scales (e.g., protein synthesis vs. metabolic rates) and their mathematical representation (presence of non-linear expressions in ME-models), requiring the implementation of more elaborate techniques for numerical simulation.

GECKO, a method for enhancement of GEMs with Enzymatic Constraints using Kinetic and Omics data, was developed in 2017 and applied to the consensus GEM for *S. cerevisiae*, Yeast7³⁸. This method extends the classical FBA approach by incorporating a detailed description of the enzyme demands for the metabolic reactions in a network, accounting for all types of enzyme-reaction relations, including isoenzymes, promiscuous enzymes and enzymatic complexes. Moreover, GECKO enables direct integration of proteomics abundance data, if available, as constraints for individual protein demands, represented as enzyme usage pseudo-reactions, whilst all of the unmeasured enzymes in the network are constrained by a pool of remaining protein mass. Additionally, this method incorporates a hierarchical and automated procedure for retrieval of kinetic parameters from the BRENDA database, which yielded a high coverage of kinetic constraints for the *S. cerevisiae* network. The resulting enzyme-constrained model, ecYeast7, was used for successful prediction of the Crabtree effect in wild-type and mutant strains of *S. cerevisiae* and cellular growth on diverse environments and genetic backgrounds, but also provided a simple framework for prediction of protein allocation profiles and study of proteomics data in a metabolic context. Furthermore, the model formed the basis for modeling yeast growth at different temperatures³⁹.

Since the first implementation of the GECKO method³⁸, its principles of enzyme constraints have been incorporated into GEMs for *B. subtilis*⁴⁰, *E. coli*⁴¹, *B. coagulans*⁴², *Streptomyces coelicolor*⁴³ and even for diverse human cancer cell-lines², showing the applicability of the method even for non-model organisms. Despite the rapid adoption of the method by the constraint-based modelling community, there is still a need for automating the model generation and enabling identification of kinetic parameters for less studied organisms. Here we wanted to build GECKO models for several organisms, and we therefore updated the GECKO toolbox to its 2.0 version. Among other improvements, we generalized its structure to facilitate its applicability to a wide variety of GEMs, and we improved its parameterization procedure to ensure high coverage of kinetic constraints, even for poorly studied organisms. Additionally, we incorporated simulation utility functions, and developed an automated virtual pipeline for update of enzyme-constrained models (ecModels), named ecModels container. This container is directly connected to the original sources of version-controlled GEMs and the GECKO toolbox, offering a continuously updated catalogue of diverse ecModels.

Results

Community development of GECKO

To ensure wide application and enable future development by the research community, we established the GECKO toolbox as open-source software, mostly encoded in MATLAB. It integrates modules for enhancement of GEMs with kinetic and proteomics constraints, automated retrieval of kinetic parameters from the BRENDA database (python module), as well as simulation utilities and export of ecModel files compatible with both the COBRA toolbox⁴⁴ and the COBRapy package⁴⁵. The development of GECKO has been continuously tracked in a public repository (<https://github.com/SysBioChalmers/GECKO>) since 2017, providing a platform for open and collaborative development. The generation of output model files in .txt and SBML L3V1 FBC2⁴⁶ formats enabled the utilization of ecYeastGEM¹ structure as a standard test to track the effects of any modifications in the toolbox algorithm through the use of the Git version control system, contributing to reproducibility of results and backwards compatibility of code.

Interaction with users of the GECKO toolbox and the ecYeastGEM model has also been facilitated through the use of the GECKO repository, allowing users to raise issues related with the programming of the toolbox or even about conceptual assumptions of the method, which has guided cumulative enhancements. Additionally, technical support for installation and utilization of the toolbox and ecYeastGEM is now provided through an open community chat room (available at: <https://gitter.im/SysBioChalmers/GECKO>), reinforcing transparent and continuous communication between users and developers.

New additions to the GECKO toolbox

The first implementation of the GECKO method significantly improved phenotype predictions for *S. cerevisiae*'s metabolism under a wide variety of genetic and environmental perturbations³⁸. However, its development underscored some issues, in particular that quantitative prediction of the critical dilution rate and exchange fluxes at fermentative conditions are highly sensitive to the distribution of incorporated kinetic parameters. Although *S. cerevisiae* is one of the most studied eukaryal organisms, not all reactions included in its model have been kinetically characterized. Therefore, a large number of k_{cat} numbers measured for other organisms (48.35%), or even non-specific to their reaction mechanism (56.03% of k_{cat} values found by introduction of wild cards into E.C. numbers) were needed to be incorporated, in order to fill the gaps in the available data for the reconstruction of the first *S. cerevisiae* ecModel, ecYeast7. Moreover, detailed manual curation of k_{cat} numbers was needed for several key enzymes in order to achieve biologically meaningful predictions.

As the BRENDA database⁴⁷ is the main source of kinetic parameters for GECKO, all of the available k_{cat} and specific activity entries for non-mutant enzymes were retrieved. In total, 38,280 entries for 4,130 unique E.C. numbers were obtained and classified according to biochemical mechanisms, phylogeny of host organisms and metabolic context (**Supp. file 1**), in order to assess significant differences in distributions of kinetic parameters. This analysis showed that not all organisms have been equally studied. Whilst entries for *H. sapiens*, *E. coli*, *R. norvegicus* and *S. cerevisiae* account for 24.02% of the total, very few kinetic parameters are available for most of the thousands of organisms present in the database, showing a median of 2 entries per organism (**Fig. 1A, Supp. file 1**). The analysis also showed that kinetic activity can differ drastically, spanning several orders of magnitude even for families of enzymes with closely related biochemical mechanisms (**Fig. 1B**). Finally, it was also observed that k_{cat} distributions for enzymes in the central carbon and energy metabolism differ significantly from those in other metabolic contexts across phylogenetic groups of host organisms (life kingdoms, according to the KEGG phylogenetic tree⁴⁸), even without filtering the dataset for entries reported exclusively for natural substrates, as previously done by other studies⁴⁹ (**Fig. 1C**).

As k_{cat} numbers depend on biochemical mechanisms, metabolic context and phylogeny of host organisms, a modified set of hierarchical k_{cat} matching criteria was implemented as part of GECKO 2.0. The modified parameterization procedure enables the incorporation of kinetic parameters that have been reported as *specific activities* in BRENDA when no k_{cat} is found for a given query (as the specific activity of an enzyme is defined as its k_{cat} over its molecular weight), adding 8,118 new entries to the catalogue of kinetic parameters in the toolbox. A phylogenetic distance-based criterion, based on the phylogenetic tree available in the KEGG database⁴⁸, was introduced for cases in which no organism-specific entries are available for a

given query in the kinetic parameters dataset. A comparison of the new k_{cat} matching criteria with their predecessor set is shown in **Supp. file 2**.

In order to assess the impact of the modified k_{cat} assignment algorithm on an ecModel, ecYeast7 was reconstructed using both the first and the new version of the GECKO toolbox (GECKO 2.0). A classification of the matched k_{cat} numbers according to the different levels of the new matching algorithm is provided in **Fig. 1D**. The incorporation of specific activity values in the parameter catalogue increased the number of kinetic parameters matched to complete E.C. numbers (no added wild cards) from 1432 to 2696 (**Fig. 1E**). Moreover, the implementation of the phylogenetic distance-based criterion yielded a distribution of kinetic parameters that showed no significant differences when compared to the values reported in BRENDA for all fungi species, in contrast to the kinetic profile matched by the previous algorithm (p -values $<10^{-10}$ and $<10^{-7}$, when compared to the BRENDA fungi and *S. cerevisiae* distributions, respectively, under a Kolmogorov-Smirnov test) (**Fig. 1F**). The quality of phenotype predictions for the ecYeast7 model enhanced by GECKO2.0 was evaluated by simulation of batch growth in 19 different environments, with an average relative error of 23.97% when compared to experimental data (**Fig. 1G**), in contrast, its GECKO1.0 counterpart yielded an average relative error of 32.07%.

The introduction of manually curated k_{cat} numbers in a metabolic network has been proven to increase the quality of phenotype predictions for *S. cerevisiae*^{22,25,38}; nevertheless, this is an intensive and time consuming procedure that is hard to ensure for a large number of models subject to continuous modifications. In order to ensure applicability of the GECKO method to any standard GEM, a unified procedure for curation of kinetic parameters was developed based on parameter sensitivity analysis. For automatically generated ecModels that are not able to reach the provided experimental value for maximum batch growth rate, an automatic module performs a series of steps in which the top enzymatic limitation on growth rate is identified through the quantification of enzyme control coefficients. For such enzymes, the E.C. number is obtained and then its correspondent k_{cat} value is substituted by the highest one available in BRENDA for the given enzyme class. This procedure iterates until the specific growth rate predicted by the model reaches the provided experimental value.

Finally, as the first version of the toolbox relied on the structure and nomenclature of the model Yeast7, its applicability to other reconstructions was not possible in a straightforward way. In order to provide compatibility with any other GEM, based on COBRA⁴⁴ or RAVEN⁵⁰ formats, all of the organism-specific parameters required by the method (experimental growth rate, total protein content, organism name, names and identifiers for

some key reactions, etc.) can be provided in a single MATLAB initialization script, minimizing the modifications needed for the generation of a new ecModel.

ecModels container: an automatically updated repository

Several GEMs that have been published are still subject to continuous development and maintenance^{1-3,5,6}, this renders GEMs to be dynamic structures that can change rapidly. In order to integrate such continuous updates into the enzyme constrained version of a model in an organized way, an automated pipeline named **ecModels container** was developed.

The ecModels container is a continuous integration implementation whose main functionality is to provide a catalogue of ecModels for several relevant organisms that are automatically updated every time a modification is detected either in the original GEM source repository or in the GECKO toolbox, i.e. new releases in their respective repositories. The pipeline generates ecModels in different formats, including the standard SBML and MATLAB files, and stores them in a container repository (<https://github.com/SysBioChalmers/ecModels>) in a version controlled way, requiring minimal human interaction and maintenance. The GECKO toolbox ensures the creation of functional and calibrated ecModels that are compatible with the provided experimental data (maximum batch growth rate, total protein content of cells and exchange fluxes at different dilution rates as an optional input). This whole computational pipeline is illustrated in **Fig. 2**. Further description of the ecModels container pipeline functioning is included in the **Materials and Methods** section.

A catalogue of new ecModels

Following the aforementioned additions to the GECKO toolbox, that have allowed its generalization, we used the toolbox for the reconstruction of four new ecModels from previously existing high-quality metabolic network reconstructions: *iYali4*, for the oleaginous yeast *Yarrowia lipolytica*⁵; *iSM996*, for the thermotolerant yeast *Kluyveromyces marxianus*⁶; *iML1515*, for the widely studied bacterium *E. coli*⁴; and Human1, being the latest and largest network reconstruction available for studying *H. sapiens* metabolism². For the microbial models, all model parameters were calibrated according to the provided experimental data, generated by independent studies^{4,51-53}, yielding functional ecModels ready for simulations. Size metrics for these models can be seen in **Table 1**.

These ecModels, together with ecYeastGEM, are hosted in the ecModels container repository for their continuous and automated update every time that a version change is detected either in the original model source or in the GECKO repository. In the case of microbial species, two different model structures are provided: **ecModel**, which has unbounded individual enzyme usage reactions ready for incorporation of proteomics data; and **ecModel_batch** in which all enzyme usage reactions are connected to a shared protein pool. This pool is then constrained

by experimental values of total protein content, and calibrated for batch simulations using experimental measurements of maximum batch growth rates on minimal glucose media, thus providing a functional ecModel structure ready for simulations.

For ecHumanGEM just the unbounded ecModel files are provided, as this is a general network of human metabolism, containing all reactions from any kind of human tissue or cell type for which evidence is available, and therefore not suitable for numerical simulation. As *H. sapiens* is the most represented organism in the BRENDA database, accounting for 11% of the total number of available k_{cat} values (**Supp. file 1**), kinetic parameters from other organisms were not taken into account for its enhancement with enzyme constraints. ecHuman1 provides the research community with an extensive knowledge base that represents a complete and direct link between genes, proteins, kinetic parameters, reactions and metabolites for human cells in a single model structure, subject to automated continuous update by the *ecModels container* pipeline.

Table 1.- Size metrics summary for the ecModels catalogue.

Original GEMs					
Organism	<i>S. cerevisiae</i>	<i>Y. lipolytica</i>	<i>K. marxianus</i>	<i>E. coli</i>	<i>H. sapiens</i>
Model ID	yeastGEM_8.3.3	iYali4	iSM996	iML1515	Human1
Reactions	3963	1924	1913	2711	13101
Metabolites	2691	1671	1531	1877	8400
Genes	1139	847	996	1516	3628
Enzyme constrained GEMs					
Model ID	ecYeastGEM	eciYali	eciSM996	eciML1515	ecHumanGEM
Reactions	8028	3881	5334	6084	46259
Metabolites	4153	1880	2064	2334	12191
Enzymes	965	647	716	1259	3224
Enzyme coverage	84.72%	76.39%	71.89%	83.05%	88.86%
Reactions w/ k_{cat}	3771	1586	2891	2562	27014
Reactions w/ Isoenzymes	504	205	532	456	3791
Promiscuous Enzymes	572	324	469	673	2184
Enzyme complexes	252	75	27	383	756

Visualization of GECKO simulations in the Caffeine platform

We implemented simulations with ecModels in Caffeine, an open-source software platform for cell factory design. Caffeine, publicly available at <http://caffeine.dd-decaf.eu>, allows user-friendly simulation and visualization of flux predictions made by genome-scale metabolic models. Several standard modelling methods are already included in the platform, such as ^{13}C fluxomics data integration, and simulation of gene deletion and/or overexpression, to interactively explore strain engineering strategies. In order to allow for GECKO simulations, we

added a new feature to the platform for uploading enzyme-constrained models and absolute proteomics data. Additionally, we added a simulation algorithm that recognizes said models, and overlays the selected proteomics data on them, leaving out data that makes the model unable to grow at a pre-specified growth rate. After these inclusions to the platform, enzyme usage can now be computed on the fly and visualized on metabolic maps (**Fig. 2B**), to identify potential metabolic bottlenecks in a given condition. The original proteomics data can be visualized as well, to identify if the specific bottleneck is due to a lack of enzyme availability, or instead due to an inefficient kinetic property. This will suggest different metabolic engineering strategies to the user: if the problem lies in the intracellular enzyme levels, the user can interpret this as a recommendation for overexpressing the corresponding gene, whereas if the problem lies in the enzyme efficiency, the user could assess introducing a heterologous enzyme as an alternative.

GECKO simulation utilities

As ecModels are defined in an irreversible format and incorporate additional elements such as enzymes (as new pseudo-metabolites) and their usages (represented as pseudo-reactions), they might sometimes not be directly compatible with all of the functionalities offered by currently available constraint-based simulation software^{44,45,50,54,55}. We therefore added several new features to the GECKO toolbox that allow the exploration and exploitation of ecModels. These include utilities for: 1) basic simulation and analysis purposes, 2) accessible retrieval of kinetic parameters, 3) automated generation of condition-dependent ecModels with proteomic abundance constraints, 4) comparative flux variability analysis between a GEM and its ecModel counterpart, and 5) prediction of metabolic engineering targets for enhanced production with an implementation of the FSEOF method⁵⁶ for ecModels. Detailed information about the inputs and outputs for each utility can be found on their respective documentation, available at: <https://github.com/SysBioChalmers/GECKO/tree/master/geckomat/utilities>. All of these utilities were developed in MATLAB due to their dependency on some RAVEN toolbox functions⁵⁰.

Predicting microbial proteome allocation in multiple environments

In order to test the quality of the phenotype predictions of an ecModel automatically generated by the *ecModels container* pipeline, batch growth under 11 different carbon sources was simulated with ecML1515 for *E. coli*. **Figure 3A** shows that, for all carbon sources, growth rates were predicted at the same order of magnitude as their corresponding experimental measurements, with the most accurate predictions obtained for growth on D-glucose, mannose and D-glucosamine. Furthermore, batch growth rate and protein allocation predictions, using no exchange flux constraints, were compared between ecML1515 and the J1678 ME-model³², the latter accounting for both metabolism and macromolecular expression processes. The sum squared error (SSE) for batch growth rate predictions across the 11 carbon sources

using *eci*ML1515 was 0.27, a drastic improvement when compared to the 1.21 SSE of *j*JL1678 ME-model predictions³². **Figure 3B** shows the predicted total proteome needed by cells to sustain the provided experimental growth rates for the same 11 environments. Notably *eci*ML1515 predicts values that lie within the range of predictions of the *j*JL1678 ME-model (from the optimal to the generalist case) for 10 out of the 11 carbon sources (see Materials and Methods for simulation details). This shows that the new version of the GECKO toolbox ensures the generation of functional ecModels that can be readily used for simulation of metabolism, due to its systematic parameter flexibilization step which reduces the need of extensive manual curation for new ecModels. Furthermore, *i*ML1515 is a model available as a static file at the BiGG models repository⁵⁷; therefore, its integration to the ecModels container for continuous update demonstrates the flexibility of our pipeline, regarding compatibility with original GEM sources, which can be provided as a link to their *git*-based repositories or even as static URLs.

Proteomics constraints refine phenotype predictions for multiple organisms and conditions

The previously mentioned module for integration of proteomics data generates a condition-dependent ecModel with proteomics constraints for each condition/replicate in a provided dataset of absolute protein abundances [mmol/gDw]. Even though absolute quantification of proteins is becoming more accessible and integrated into systems biology studies⁵⁸⁻⁶², a major caveat of using proteomics data as constraints for quantitative models is their intrinsic high biological and technical variability⁶³, therefore some of the incorporated data constraints need to be loosened in order to obtain functional ecModels. When needed, additional condition-dependent exchange fluxes of byproducts can also be used as constraints in order to limit the feasible solution space. A detailed description of the proteomics integration algorithm implemented in GECKO is given in **Supp. file 2**.

The new proteomics integration module was tested on the three ecModels for budding yeasts available in ecModels container (*ecYeastGEM*, *ecYali*, *eciSM996*). We measured absolute protein abundances for *S. cerevisiae*, *Y. lipolytica* and *K. marxianus*, grown in chemostats at 0.1 h⁻¹ dilution rate and subject to several experimental conditions (high temperature, low pH and osmotic stress with KCl)⁶⁴, and incorporated these data into the ecModels as upper bounds for individual enzyme usage pseudo-reactions. Then, exchange fluxes for CO₂ and oxygen corresponding to the same chemostat experiments were used as a comparison basis to evaluate quality of phenotype predictions. For each organism-condition pair, 3 models were generated and compared in terms of predictions: a pure stoichiometric metabolic model, an enzyme-constrained model with a limited shared protein pool, and an enzyme-constrained model with proteomics constraints. It was found that the addition of the enzyme pool constraint enables major reduction of the relative error in prediction of gaseous exchange fluxes in some of the studied conditions. Additionally, the incorporation of individual protein abundance constraints

improves even further the predictive accuracy of gaseous exchanges, for 10 out of the 11 evaluated cases (**Fig. 4A-C**).

The impact of incorporating enzyme and proteomics constraints on intracellular flux predictions was further assessed by mapping all condition-dependent flux distributions from the tested ecModels to their corresponding reactions in the original GEMs. In general, metabolic flux distributions showed high similarity when comparing ecModel to GEM predictions (**Fig. S1**), as 70-90% of the active fluxes were predicted within the interval of $0.5 <$

fold-change < 2 $\left(FC = \frac{v_i^{ecModel}}{v_i^{GEM}}\right)$ across all conditions (**Fig. S2 A-C, Supp. file 3**). In addition, principal component analysis on absolute enzyme usage profiles predicted by ecModels revealed that, at low dilution rates, predictions of enzyme demands are mostly defined by the selected set of imposed constraints (shared protein pool vs. proteomics constraints) rather than by environmental condition, i.e. exchange fluxes (**Fig. S2 D-F**). However, more straightforward comparison of the models' predictions, by pairwise comparison of predicted absolute enzyme usage profiles, showed that 60 - 80% of the predicted enzyme usages lie within a range of $0.5 < \text{fold-change} < 2$, when comparing ecModels predictions with and without proteomics constraints, across organisms and conditions (**Fig. 4D, Fig. S2 G-I and Supp. file 3**). It was observed that the incorporation of proteomics constraints induces a drastic differential use for a considerable amount of enzymes, as 12-21% of enzyme usages were predicted as either enabled or disabled by these constraints across all the simulated conditions, showing slight enrichment for enabled alternative isoenzymes for already active reactions (**Supp. file 3**). This suggests that upper bounds on enzyme usages induce differentiated utilization of isoenzymes, reflecting well why isoenzymes have been maintained throughout evolution.

The explicit inclusion of enzymes into GEMs by the GECKO method enables prediction of enzyme demands at the protein, reaction and pathway levels. Total protein burden values predicted by ecModels for several relevant metabolic superpathways (central carbon and energy metabolism, amino acid metabolism, lipid and fatty acid metabolism, cofactor and vitamin metabolism and nucleotide metabolism, according to the KEGG metabolic subsystems⁴⁸), showed that central carbon and energy metabolism is the most affected sector in the ecYeastGEM network by integration of proteomics constraints, as protein burden predictions were higher, at least by 20%, for 3 out the 4 simulated conditions when compared with predictions of the ecYeastGEM without proteomics data (**Fig. 4E**).

Relative enzyme usages, estimated as predicted absolute enzyme usage over enzyme abundance for all of the measured enzymes in an ecModel $\left(\frac{e_i}{[E_i]}\right)$, can be understood as the saturation level of enzymes in a given condition. In order to analyze the metabolic mechanisms underlying long-term adaptation to stress in budding yeasts, relative enzyme usage profiles were computed from all the previous simulations of ecModels with proteomics constraints.

Enzymes that display fold-changes higher than 1 for both absolute abundance and their saturation level, when comparing predicted usage profiles between stress and reference conditions, suggest regulatory mechanisms on individual proteins that contribute to cell growth on the analyzed stress condition. **Figure 4F** shows all of the enzymes that were identified as responsive to environmental stress in this study, displaying enrichment for enzymes involved in biosynthesis of diverse amino acids and folate metabolism.

A further mapping of all enzymes in these ecModels to a list of 2,959 single copy protein-coding gene orthologs across the three yeast species⁶⁴ found 310 core proteins across these ecModels. Principal component analysis revealed that variance on absolute enzyme usages and abundance profiles for these core proteins is mostly explained by differences in the metabolic networks of the different species rather than by environmental conditions (**Fig. S3 B-C**), reinforcing previous results suggesting that, despite being phylogenetically related, their long-term stress responses at the molecular level have evolved independently after their divergence in evolutionary history⁶⁴.

Exploring the solution space reduction

A major limitation in the use of GEMs is the high variability of flux distributions for a given cellular objective when implementing flux balance analysis, as this requires solving largely underdetermined linear systems through optimization algorithms^{15,65}. This limitation has usually been overcome with incorporation of measured exchange fluxes as constraints. However, these data are typically sparse in the literature. Previous studies explored the drastic reduction in flux variability ranges of ecModels for *S. cerevisiae* and 11 human cell-lines when compared to their original GEMs due to the addition of enzyme constraints^{1,2,38}. However, the irreversible format of ecModels (forward and backwards reactions are split in order to account for enzyme demands of both directions) hinders their compatibility with the flux variability analysis (FVA) functions already available in COBRA⁴⁴ and RAVEN⁵⁰ toolboxes. As a solution to this, an FVA module was integrated to the utilities repertoire in GECKO, whose applicability has been previously tested on studies with ecModels for *S. cerevisiae*¹ and human cell lines². This module contains the necessary functions to perform FVA on any set of reactions of an ecModel, enabling also a direct comparison of flux variability ranges between an ecModel and its GEM counterpart in a consistent way (see **Supp. file 2**).

The FVA utility was applied on three different ecModels of microbial metabolism and their correspondent GEMs (*iML1515*, *iYali4* and *iSM996*). In all cases the FVA comparisons were carried out for both chemostat and batch growth conditions in order to span different degrees of constraining of the metabolic networks (0.1 h⁻¹ dilution rate and minimal glucose uptake rate fixed for chemostat conditions; biomass production fixed to experimental measurements of and unconstrained uptake of minimal media components, for batch conditions). Cumulative distributions for flux variability ranges for all explored ecModels and GEMs are shown

in **Figure 5**, in which it can be seen that median flux variability ranges are much reduced for all ecModels and conditions, especially at high growth rates where enzyme constraints reduce the variability range 5-6 orders of magnitude when compared to pure GEMs. The cumulative distributions also show a major reduction in the amount of totally variable fluxes (reactions that can carry any flux between -1000 to 1000 mmol/gDw h), which are an indicator of undesirable futile cycles present in the network due to lack of thermodynamic and enzyme cost information⁶⁶⁻⁶⁸. For high growth rates, the amount of totally variable fluxes accounts for 3-12% of the active reactions in the analysed GEMs, in contrast to their corresponding ecModels in which such extreme variability ranges are completely absent.

Further analysis of the FVA results revealed that a reduction of at least 95% of the variability range was achieved for more than 90% of all active fluxes at high growth rates in all ecModel. Interestingly, the aforementioned flux variability metrics were overall improved even for the chemostat conditions, despite a higher degree of constraining (fixed low growth rate and optimal uptake rate), which restrains these models to an energy efficient respiratory mode (**Supp. file 4**).

Discussion

Here we demonstrated how enzyme constrained models for diverse species significantly improve simulation performance compared to traditional GEMs. Furthermore, to enable the community to easily adapt this modelling approach, we upgraded the GECKO toolbox for enhancement of genome-scale models with enzyme and omics constraints to its version 2.0. Major improvements on the k_{cat} matching algorithm were incorporated into the toolbox, based on phylogenetic distance between the modeled organism and the host organisms for data queries, and an automated curation of k_{cat} numbers for over-constrained models were incorporated into the toolbox. Major refactoring of the GECKO toolbox enabled a generalization of the method, allowing the creation of high-quality ecModels for any provided functional GEM with minimal need for case-specific introduction of new code. Additionally, several utility functions were integrated into the toolbox in order to enable basic simulation purposes, accessible retrieval of enzyme parameters, integration of proteomics data as constraints, flux variability analysis and prediction of gene targets for enhanced production of metabolites. Overall, it was shown that these enhancements to the GECKO toolbox improve the incorporation of kinetic parameters into a metabolic model, yielding ecModels with biologically meaningful kinetic profiles without compromising accuracy on phenotype predictions.

Two major limitations of the first version of the GECKO toolbox were its specific customization to the *S. cerevisiae* model, Yeast7, and the need of extensive manual curation for generating an ecModel suited for FBA simulations; thus, its applicability to other GEMs was not a straightforward procedure. To overcome these limitations, we generalized the code with the aim of making GECKO a model-agnostic tool. The development of a procedure for automatic curation of kinetic parameters enabled the generation of

functional ecModels with minimal requirements for experimental data. Recently, ecModels for 11 human cancer cell-lines were generated with this automated procedure, using Human1 as a model input and RNAseq datasets together with the tINIT algorithm¹⁰ to generate cell-line specific networks². These ecModels were used for the prediction of cellular growth and metabolite exchange rates at different levels of added constraints, resulting in remarkable improvements in accuracy when compared with predictions of their original GEMs. This highlights one of the main advantages of ecModels: their capability of yielding biologically meaningful phenotype predictions without an excessive dependency on exchange fluxes as constraints.

In order to further showcase the functionality of the GECKO toolbox 2.0, a family of new high-quality ecModels were generated for *E. coli*, *Y. lipolytica*, *K. marxianus* and *H. sapiens*, based on the original GEMs *i*ML1515, *i*Yali4, *i*SM996 and Human1, respectively. Furthermore, we generated a self-hosted pipeline for continuous and automated generation and update of ecModels, ecModels container, so that each of the currently available ecModels (ecYeastGEM, ec*i*ML1515, ec*i*Yali, ec*i*SM996 and ecHuman1) are integrated to it, providing a version-controlled and continuously updated repository for high-quality ecModels.

Absolute proteomics measurements for the budding yeasts *S. cerevisiae*, *K. marxianus* and *Y. lipolytica* grown under multiple environmental conditions, were incorporated as constraints into their ecModels by using the proteomics integration module added to the GECKO toolbox. Analysis of metabolic flux distributions revealed that net reaction fluxes predicted by GEMs are not significantly affected by the incorporation of kinetic and proteomics constraints, however the explicit integration of enzymes into ecModels extends the range of predictions of classical FBA and enables computation of enzyme demands at the reaction and pathway levels. It was found that incorporation of proteomics constraints does not affect enzyme demand predictions significantly for most of the active enzymes at low dilution rates across the simulated conditions. However, we observed that a diversified utilization of isoenzymes, enforced by proteomics constraints, increases the predicted total protein mass allocated to central carbon and energy metabolism, in comparison to optimal enzyme allocation profiles. This result suggests the relevance of metabolic robustness in contrast to optimal protein utilization for microbial growth under environmental stress and nutrient-limited conditions.

Incorporation of proteomics data allows the use of ecModels as scaffolds for systems-level studies of metabolism, providing a tool for uncovering metabolic readjustments induced by genetic and environmental perturbations, which might be difficult to elucidate by purely data-driven approaches, specially at conditions of relatively low changes at the transcript⁶⁹ and protein levels⁶⁴. For all studied stress conditions in this study, we identified upregulated proteins (increased abundance) that are needed to operate at high saturation levels in stress conditions, while showing low usage at reference conditions, creating lists of potential gene amplification targets for enhancing stress tolerance in three industrially relevant yeast species (**Supp. file 3**). Upregulation and high saturation of enzymes in amino acid and folate metabolism were found to be common across the studied organisms and stress conditions (**Fig. S3 D** and **Supp. file 3**). These results suggests that yeast cells display enzyme expression profiles that

provide them with metabolic robustness for microbial growth under stress and nutrient-limited conditions, in contrast to an optimal protein allocation strategy that prioritizes expression of the most efficient and non-redundant enzymes.

Our results on drastic reduction of median flux variability ranges and the number of totally unbounded fluxes for *ec/Yali*, *ec/SM996* and *ec/ML1515*, together with previous studies^{1,2,38}, suggest that a major reduction of the solution space of metabolic models to a more biologically meaningful subspace is a general property of *ecModels*. However, flux variability is an intrinsic characteristic of metabolism; therefore, metabolic models with highly constrained solution spaces may exclude some biological capabilities of organisms, which are not compatible with the set of constraints used for the analysis (exchange fluxes, growth rates and even profiles of kinetic parameters, considered as condition-independent in *ecModels*).

Here, the predictive capabilities of *ec/ML1515* and *iJL1678* ME-model (both for *E. coli*) for cellular growth and global protein demands on diverse environments were compared. The major improvement in predicted maximum growth rates, together with a comparable performance on quantification of protein demands, shown by *ec/ML1515* suggest that, despite its mathematical and conceptual simplicity, the GECKO formalism is a suitable framework for quantitative probing of metabolic capabilities, compatible with the widely used FBA method and without the need of excessive complexity or computational power. Nevertheless, ME-models provide a much wider range of predictions that explore additional processes in cell physiology with great detail. Direct comparison between the predictions of these modelling formalisms, suggest that ME-models performance can be improved by incorporation of either curated or systematically retrieved kinetic parameters that are suitable for the modelled organisms.

Simpler modelling frameworks that account for protein or enzyme constraints in metabolism, such as flux balance analysis with molecular crowding (FBAwMC)^{16,17}, metabolic modelling with enzyme kinetics (MOMENT)²³ and constrained allocation flux balance analysis (CAFBA)²¹, have also been developed and used to explore microbial cellular growth^{16,17,21} and overflow metabolism^{16,23}. These methods have overcome the lack of reported parameters for some specific reactions either by incorporation of proteomics measurements and prior flux distributions²³, manual curation and sampling procedures^{16,17} or even by lumping protein demands by functionally related proteome groups. In contrast, the new version of the GECKO toolbox provides a systematic and robust parameterization procedure, leveraging the vastly accumulated knowledge of biochemistry research stored in public databases, ensuring the incorporation of biologically meaningful kinetic parameters even for poorly studied reactions and organisms.

The applicability of these other simple modelling formalisms to models for diverse species is limited as none of these methods has been provided as part of a generalized model-agnostic software implementation. Recently, a simplified variant of the MOMENT method (sMOMENT) was developed and embedded into an automated pipeline for generation and calibration of enzyme-constrained models of metabolism (AutoPACMEN)⁷⁰. The pipeline was tested on the generation of an enzyme-constrained version of the *iJO1366* metabolic reconstruction for *E. coli*, which also showed consistency with

experimental data. This work represented a step forward in the field of constrain-based metabolic modelling, as it contributed to standardization of model generation and facilitating their utilization and applicability to other cases. However, due to the intrinsic trade-off between model simplicity and descriptive representation, a limitation of the sMOMENT method is its simplification of redundancies in metabolism, which just accounts for the optimal way of catalysing a given biochemical reaction, discarding the representation of alternative isoforms that might be relevant under certain conditions. In GECKO ecModels, all enzymes for which a gene-E.C. number relationship exists are included in the model structure. As traditional FBA simulations rely on optimality principles one could, in principle, expect the same predicted flux distributions by sMOMENT and GECKO ecModels. Nonetheless, the explicit incorporation of all enzymes in a metabolic network enables explanation of protein expression profiles that deviate from optimality in order to gain robustness to changes in the environment, as it has been recently shown by the integration of a regulatory nutrient-signalling Boolean network together with an ecModel for *S. cerevisiae*'s central carbon metabolism⁷¹.

In conclusion, GECKO2.0 together with the development of the automated pipeline ecModels container facilitates the generation, standardization, utilization, exchange and community development of ecModels through a transparent version-controlled environment. This tool provides a dynamic, and potentially increasing, catalogue of updated ecModels trying to close the gap between model developers and final users and reduce the time-consuming tasks of model maintenance. We are confident that this will enable wide use of ecModels in basic science for obtaining novel insight into the function of metabolism as well as in synthetic biology and metabolic engineering for design of strains with improved functionalities, e.g., for high-level production of valuable chemicals.

Material And Methods

Automation pipeline and version-controlled hosting of the ecModels container

The ecModels repository is used to version-control the pipeline code and the resulting models. The pipeline is restricted to 2 short Python files, whose role is to decide when models need to be updated based on a configuration file **config.ini**, and to consequently invoke the use of GECKO for each model. Updates are deemed necessary when either the underlying dependencies (i.e., GECKO, RAVEN and COBRA toolboxes, the Gurobi solver, and libSMBL) or the source GEMs are independently updated to a new version (release) in their respective repositories.

The pipeline is designed be automatic and to not require supervision. It was developed to work with both version-controlled GEMs and GEMs downloadable from a URL, updating the version in the configuration after a new ecModel is obtained. For easy review, the pipeline log is publicly available under the *Actions* tab of the GitHub repository. The computation is performed through a self-hosted GitHub runner, further leveraging the transparent nature of the GitHub platform and the *git* version-control system. The resulting ecModel and updated configuration are committed to the repository, with the changes being made available for review through a pull request. Additionally, the GECKO output is also replicated in the pull

request body. The *ecModels container* thus continues the transparency and reproducibility of the source models.

Quantification of absolute protein concentrations for *S. cerevisiae*, *Y. lipolytica* and *K. marxianus*

Total protein extraction for the strains *Saccharomyces cerevisiae* CEN.PK113-7D (standard, low pH, high temperature, osmotic stress), *Kluyveromyces marxianus* CBS6556 (standard, low pH, high temperature, osmotic stress) and *Yarrowia lipolytica* W29 (standard, low pH, high temperature) was conducted as described in **Supp. file 2**. Three reference samples (hereafter, 'bulk' samples), one per strain, were constructed by pooling 5 µg of each experimental sample. Aliquots of 15 µg of total protein extract from each sample (3 strains x 4 conditions x 3 replicates) and the three bulks were separated on one-dimensional SDS-PAGE short-migration gels (1x1 cm lanes, Invitrogen, NP321BOX). Yeast proteins digestion was performed on excised bands from gel gradient and digested peptides of UPS2 (Sigma) were used as external standards for absolute protein quantification (more details in **Supp. file 2**). Four µl of the different peptide mixtures (800 ng for yeast peptides and 949 ng for bulks) were analyzed using an Orbitrap Fusion™ Lumos™ Tribrid™ mass spectrometer (Thermo Fisher Scientific).

Protein identification was performed using the open-source search engine X!Tandem pipeline 3.4.4⁷². Data filtering was set to peptide E-value < 0.01 and protein log(E-value) < -3. Relative quantification of protein abundances was carried out using the Normalized Spectral Abundance Factor (NSAF)⁷³ and the NSAF values obtained from UPS2 proteins in bulk samples were used to determine the suitable regression curves that allowed the conversion from relative protein abundance into absolute terms. The regression curves parameters for protein abundance quantification are shown in **Supp. file 2**. MS data is available online on public databases via the PRIDE repository⁷⁴ with the dataset identifier PXD012836.

Simulation of condition-dependent flux distributions

Simulation of cellular phenotypes for conditions of environmental stress at low dilution rates with GEMs were performed by first setting bounds on measured glucose uptake and byproduct secretion rates according to experimental data from previous studies on chemostats⁶⁴. Then the biomass production rate was constrained (both upper and lower bounds) with the experimental dilution rate (0.1 h⁻¹). Maximization of the non-growth associated maintenance pseudo-reaction was set as an objective function for the parsimonious FBA problem as a representation of the additional energy demands for regulation of cellular growth at non-optimal conditions. The same procedure was followed for simulations with ecModels constrained by a total protein pool. For the case of ecModels with proteomics constraints, the same set of constraints was used but the objective function was set as minimization of the total usage of unmeasured proteins, assuming that the regulatory machinery for stress tolerance is represented by the condition-specific protein expression profile.

Prediction of microbial batch growth rates

Batch cellular growth was simulated by allowing unconstrained uptake of all nutrients present in minimal mineral media, enabling a specific carbon source uptake reaction for each case while blocking the rest of the uptake reactions and allowing unconstrained secretion rates for all exchangeable metabolites. Maximization of the biomass production rate was used as an objective function for the resulting FBA problem. For prediction of total protein demands on unlimited nutrient conditions, media constraints were set as expressed above and experimental batch growth rate values were fixed as both lower and upper bounds for the biomass production pseudo-reaction. The total protein pool exchange pseudo-reaction was then unconstrained and set as an objective function to minimize, assuming that when exposed to unlimited availability of nutrients the total mass of protein available for catalyzing metabolic reactions becomes the limiting resource for cells. The **solveLP** function, available in the RAVEN toolbox, was used for solving all FBA problems in this study.

Declarations

Code availability

The source code of the updated **GECKO toolbox** is available at: <https://github.com/SysBioChalmers/GECKO>. The code for **ecModels container** and the whole catalogue of updated ecModel files can be accessed at: <https://github.com/SysBioChalmers/ecModels>. All custom scripts for simulations included in this study can be found at: https://github.com/SysBioChalmers/GECKO2_simulations. All the necessary scripts for reproducing the k_{cat} parameters analysis in Supplementary file 2 are available at: <https://github.com/SysBioChalmers/Enzyme-parameters-analysis>. All of these repositories are public and open to collaborative continuous development.

Acknowledgements

We are grateful to Feiran Li, Raphaël Ferreira, Jonathan Robinson and all the GECKO users that have provided feedback for improving our toolbox and extending its range of applications and to the CHASSY project consortium for having motivated and supported this work. This project has received funding from the European Union's Horizon 2020 Framework Programme for Research and Innovation – Grant Agreements No. 720824 and 686070. This work was also supported by the Knut and Alice Wallenberg Foundation and The Novo Nordisk Foundation - Grant no NNF10CC1016517.

References

1. Lu, H. *et al.* A consensus *S. cerevisiae* metabolic model Yeast8 and its ecosystem for comprehensively probing cellular metabolism. *Nat. Commun.* (2019). doi:10.1038/s41467-019-11581-3
2. Robinson, J. L. *et al.* An atlas of human metabolism. *Sci. Signal.* (2020). doi:10.1126/scisignal.aaz1482
3. Tiukova, I. A., Prigent, S., Nielsen, J., Sandgren, M. & Kerkhoven, E. J. Genome-scale model of *Rhodotorula toruloides* metabolism. *Biotechnol. Bioeng.* (2019). doi:10.1002/bit.27162
4. Monk, J. M. *et al.* iML1515, a knowledgebase that computes *Escherichia coli* traits. *Nature Biotechnology* **35**, 904–908 (2017).
5. Kerkhoven, E. J., Pomraning, K. R., Baker, S. E. & Nielsen, J. Regulation of amino-acid metabolism controls flux to lipid accumulation in *Yarrowia lipolytica*. *npj Syst. Biol. Appl.* **2**, 16005 (2016).
6. Marcišauskas, S., Ji, B. & Nielsen, J. Reconstruction and analysis of a *Kluyveromyces marxianus* genome-scale metabolic model. *BMC Bioinformatics* (2019). doi:10.1186/s12859-019-3134-5
7. Meadows, A. L. *et al.* Rewriting yeast central carbon metabolism for industrial isoprenoid production. *Nature* (2016). doi:10.1038/nature19769
8. Chen, X. *et al.* Fumaric acid production by *Torulopsis glabrata*: Engineering the urea cycle and the purine nucleotide cycle. *Biotechnol. Bioeng.* (2015). doi:10.1002/bit.25334
9. Mishra, P. *et al.* Genome-scale model-driven strain design for dicarboxylic acid production in *Yarrowia lipolytica*. *BMC Syst. Biol.* (2018). doi:10.1186/s12918-018-0542-5
10. Agren, R. *et al.* Identification of anticancer drugs for hepatocellular carcinoma through personalized genome-scale metabolic modeling. *Mol. Syst. Biol.* (2014). doi:10.1002/msb.145122
11. Mardinoglu, A. *et al.* Genome-scale metabolic modelling of hepatocytes reveals serine deficiency in patients with non-alcoholic fatty liver disease. *Nat. Commun.* **5**, 1–11 (2014).
12. Gatto, F., Miess, H., Schulze, A. & Nielsen, J. Flux balance analysis predicts essential genes in clear cell renal cell carcinoma metabolism. *Sci. Rep.* (2015). doi:10.1038/srep10738
13. Orth, J. D., Thiele, I. & Palsson, B. Ø. What is flux balance analysis? *Nat. Biotechnol.* **28**, 245–248 (2010).
14. Ibarra, R. U., Edwards, J. S. & Palsson, B. O. *Escherichia coli* K-12 undergoes adaptive evolution to achieve in silico predicted optimal growth. *Nature* (2002). doi:10.1038/nature01149
15. Mahadevan, R. & Schilling, C. H. The effects of alternate optimal solutions in constraint-based genome-scale metabolic models. *Metab. Eng.* **5**, 264–276 (2003).
16. Beg, Q. K. *et al.* Intracellular crowding defines the mode and sequence of substrate uptake by *Escherichia coli* and constrains its metabolic activity. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 12663–12668 (2007).
17. Vazquez, A. *et al.* Impact of the solvent capacity constraint on *E. coli* metabolism. *BMC Syst. Biol.* **2**, 7 (2008).

18. Molenaar, D., van Berlo, R., de Ridder, D. & Teusink, B. Shifts in growth strategies reflect tradeoffs in cellular economics. *Mol. Syst. Biol.* **5**, 323 (2009).
19. Zhuang, K., Vemuri, G. N. & Mahadevan, R. Economics of membrane occupancy and respiration. *Mol. Syst. Biol.* **7**, (2011).
20. Shlomi, T., Benyamini, T., Gottlieb, E., Sharan, R. & Ruppin, E. Genome-scale metabolic modeling elucidates the role of proliferative adaptation in causing the warburg effect. *PLoS Comput. Biol.* **7**, 1–8 (2011).
21. Mori, M., Hwa, T., Martin, O. C., De Martino, A. & Marinari, E. Constrained Allocation Flux Balance Analysis. *PLoS Comput. Biol.* **12**, (2016).
22. Nilsson, A. & Nielsen, J. Metabolic Trade-offs in Yeast are Caused by F1F0-ATP synthase. *Sci. Rep.* **6**, 1–11 (2016).
23. Adadi, R., Volkmer, B., Milo, R., Heinemann, M. & Shlomi, T. Prediction of microbial growth rate versus biomass yield by a metabolic network with kinetic parameters. *PLoS Comput. Biol.* **8**, (2012).
24. Nilsson, A., Björnson, E., Flockhart, M., Larsen, F. J. & Nielsen, J. Complex I is bypassed during high intensity exercise. *Nat. Commun.* (2019). doi:10.1038/s41467-019-12934-8
25. Chen, Y. & Nielsen, J. Energy metabolism controls phenotypes by protein efficiency and allocation. *Proc. Natl. Acad. Sci. U. S. A.* (2019). doi:10.1073/pnas.1906569116
26. van Hoek, M. J. A. & Merks, R. M. H. Redox balance is key to explaining full vs. partial switching to low-yield metabolism. *BMC Syst. Biol.* (2012). doi:10.1186/1752-0509-6-22
27. Van Hoek, M. J. & Merks, R. M. Redox balance is key to explaining full vs. partial switching to low-yield metabolism. *BMC Syst. Biol.* **6**, 22 (2012).
28. Jeske, L., Placzek, S., Schomburg, I., Chang, A. & Schomburg, D. BRENDA in 2019: A European ELIXIR core data resource. *Nucleic Acids Res.* (2019). doi:10.1093/nar/gky1048
29. Wittig, U., Rey, M., Weidemann, A., Kania, R. & Müller, W. SABIO-RK: An updated resource for manually curated biochemical reaction kinetics. *Nucleic Acids Res.* (2018). doi:10.1093/nar/gkx1065
30. O'Brien, E. J., Lerman, J. A., Chang, R. L., Hyduke, D. R. & Palsson, B. Genome-scale models of metabolism and gene expression extend and refine growth phenotype prediction. *Mol. Syst. Biol.* **9**, (2013).
31. O'Brien, E. J. & Palsson, B. O. Computing the functional proteome: Recent progress and future prospects for genome-scale models. *Curr. Opin. Biotechnol.* **34**, 125–134 (2015).
32. Yang, L. *et al.* Principles of proteome allocation are revealed using proteomic data and genome-scale models. *Sci. Rep.* (2016). doi:10.1038/srep36734
33. King, Z. A., O'Brien, E. J., Feist, A. M. & Palsson, B. O. Literature mining supports a next-generation modeling approach to predict cellular byproduct secretion. *Metab. Eng.* (2017). doi:10.1016/j.ymben.2016.12.004
34. Lerman, J. A. *et al.* In silico method for modelling metabolism and gene product expression at genome scale. *Nat. Commun.* (2012). doi:10.1038/ncomms1928

35. Chen, Y. *et al.* Proteome constraints reveal targets for improving microbial fitness in nutrient-rich 1 environments 2 3 Author list. *bioRxiv* (2020).
36. Bulović, A. *et al.* Automated generation of bacterial resource allocation models. *Metab. Eng.* (2019). doi:10.1016/j.ymben.2019.06.001
37. Goelzer, A. *et al.* Quantitative prediction of genome-wide resource allocation in bacteria. *Metab. Eng.* **32**, (2015).
38. Sánchez, B. J. *et al.* Improving the phenotype predictions of a yeast genome-scale metabolic model by incorporating enzymatic constraints. *Mol. Syst. Biol.* **13**, 935 (2017).
39. Li, G. *et al.* Bayesian genome scale modelling identifies thermal determinants of yeast metabolism. *Nat. Commun.* **12**, 190 (2021).
40. Massaiu, I. *et al.* Integration of enzymatic data in *Bacillus subtilis* genome-scale metabolic model improves phenotype predictions and enables in silico design of poly- γ -glutamic acid production strains. *Microb. Cell Fact.* (2019). doi:10.1186/s12934-018-1052-2
41. Ye, C. *et al.* Improving lysine production through construction of an *Escherichia coli* enzyme-constrained model. *Biotechnol. Bioeng.* **117**, 3533–3544 (2020).
42. Chen, Y. *et al.* Genome-scale modeling for *Bacillus coagulans* to understand the metabolic characteristics. *Biotechnol. Bioeng.* (2020). doi:10.1002/bit.27488
43. Sulheim, S. *et al.* Enzyme-Constrained Models and Omics Analysis of *Streptomyces coelicolor* Reveal Metabolic Changes that Enhance Heterologous Production. *iScience* (2020). doi:10.1016/j.isci.2020.101525
44. Heirendt, L. *et al.* Creation and analysis of biochemical constraint-based models using the COBRA Toolbox v.3.0. *Nat. Protoc.* **14**, 639–702 (2019).
45. Ebrahim, A., Lerman, J. A., Palsson, B. O. & Hyduke, D. R. COBRApy: COntstraints-Based Reconstruction and Analysis for Python. *BMC Syst. Biol.* (2013). doi:10.1186/1752-0509-7-74
46. Olivier, B. G. & Bergmann, F. T. The Systems Biology Markup Language (SBML) Level 3 Package: Flux Balance Constraints. *J. Integr. Bioinform.* **12**, 269 (2015).
47. Placzek, S. *et al.* BRENDA in 2017: New perspectives and new tools in BRENDA. *Nucleic Acids Res.* **45**, D380–D388 (2017).
48. Kanehisa, M., Furumichi, M., Sato, Y., Ishiguro-Watanabe, M. & Tanabe, M. KEGG: integrating viruses and cellular organisms. *Nucleic Acids Res.* (2020). doi:10.1093/nar/gkaa970
49. Bar-Even, A. *et al.* The moderately efficient enzyme: Evolutionary and physicochemical trends shaping enzyme parameters. *Biochemistry* **50**, 4402–4410 (2011).
50. Wang, H. *et al.* RAVEN 2.0: A versatile toolbox for metabolic network reconstruction and a case study on *Streptomyces coelicolor*. *PLOS Comput. Biol.* **14**, e1006541 (2018).
51. Ochoa-Estopier, A. & Guillouet, S. E. D-stat culture for studying the metabolic shifts from oxidative metabolism to lipid accumulation and citric acid production in *Yarrowia lipolytica*. *J. Biotechnol.* (2014). doi:10.1016/j.jbiotec.2013.11.008

52. Aggelis, G. & Komaitis, M. Enhancement of single cell oil production by *Yarrowia lipolytica* growing in the presence of *Teucrium polium* L. aqueous extract. *Biotechnol. Lett.* (1999). doi:10.1023/A:1005591127592
53. Overkamp, K. M. *et al.* In vivo analysis of the mechanisms for oxidation of cytosolic NADH by *Saccharomyces cerevisiae* mitochondria. *J. Bacteriol.* (2000). doi:10.1128/JB.182.10.2823-2830.2000
54. Li, M. & Borodina, I. Application of synthetic biology for production of chemicals in yeast *Saccharomyces cerevisiae*. *FEMS Yeast Research* **15**, (2015).
55. Rocha, I. *et al.* OptFlux: An open-source software platform for in silico metabolic engineering. *BMC Syst. Biol.* (2010). doi:10.1186/1752-0509-4-45
56. Choi, H. S., Lee, S. Y., Kim, T. Y. & Woo, H. M. In silico identification of gene amplification targets for improvement of lycopene production. *Appl. Environ. Microbiol.* **76**, 3097–3105 (2010).
57. Norsigian, C. J. *et al.* BiGG Models 2020: Multi-strain genome-scale models and expansion across the phylogenetic tree. *Nucleic Acids Res.* **48**, (2020).
58. Björkeröth, J. *et al.* Proteome reallocation from amino acid biosynthesis to ribosomes enables yeast to grow faster in rich media. *Proc. Natl. Acad. Sci.* **117**, 21804 LP – 21812 (2020).
59. Yu, R. *et al.* Nitrogen limitation reveals large reserves in metabolic and translational capacities of yeast. *Nat. Commun.* (2020). doi:10.1038/s41467-020-15749-0
60. Campbell, K. *et al.* Building blocks are synthesized on demand during the yeast cell cycle. *Proc. Natl. Acad. Sci. U. S. A.* (2020). doi:10.1073/pnas.1919535117
61. Lahtvee, P. J. *et al.* Absolute Quantification of Protein and mRNA Abundances Demonstrate Variability in Gene-Specific Translation Efficiency in Yeast. *Cell Syst.* **4**, 495–504.e5 (2017).
62. Di Bartolomeo, F. *et al.* Absolute yeast mitochondrial proteome quantification reveals trade-off between biosynthesis and energy generation during diauxic shift. *Proc. Natl. Acad. Sci. U. S. A.* (2020). doi:10.1073/pnas.1918216117
63. Sánchez, B. J. *et al.* Benchmarking accuracy and precision of intensity-based absolute quantification of protein abundances in *Saccharomyces cerevisiae*. *Proteomics* **n/a**, 2000093 (2021).
64. Doughty, T. W. *et al.* Stress-induced expression is enriched for evolutionarily young genes in diverse budding yeasts. *Nat. Commun.* (2020). doi:10.1038/s41467-020-16073-3
65. Kauffman, K. J., Prakash, P. & Edwards, J. S. Advances in flux balance analysis. *Current Opinion in Biotechnology* (2003). doi:10.1016/j.copbio.2003.08.001
66. Teusink, B. *et al.* Analysis of growth of *Lactobacillus plantarum* WCFS1 on a complex medium using a genome-scale metabolic model. *J. Biol. Chem.* (2006). doi:10.1074/jbc.M606263200
67. Beard, D. A., Liang, S. D. & Qian, H. Energy balance for analysis of complex metabolic networks. *Biophys. J.* (2002). doi:10.1016/S0006-3495(02)75150-3
68. Maurice Cheung, C. Y., George Ratcliffe, R. & Sweetlove, L. J. A method of accounting for enzyme costs in flux balance analysis reveals alternative pathways and metabolite stores in an illuminated

arabidopsis leaf. *Plant Physiol.* (2015). doi:10.1104/pp.15.00880

69. Patil, K. R. & Nielsen, J. Uncovering transcriptional regulation of metabolism by using metabolic network topology. *Proc. Natl. Acad. Sci. U. S. A.* (2005). doi:10.1073/pnas.0406811102
70. Bekiaris, P. S. & Klamt, S. Automatic construction of metabolic models with enzyme constraints. *BMC Bioinformatics* (2020). doi:10.1186/s12859-019-3329-9
71. Österberg, L. *et al.* A novel yeast hybrid modeling framework integrating Boolean and enzyme-constrained networks enables exploration of the interplay between signaling and metabolism. *bioRxiv* 2020.09.11.290817 (2020). doi:10.1101/2020.09.11.290817
72. Langella, O. *et al.* X!TandemPipeline: A Tool to Manage Sequence Redundancy for Protein Inference and Phosphosite Identification. *J. Proteome Res.* **16**, (2017).
73. Zybilov, B. *et al.* Statistical analysis of membrane proteome expression changes in *Saccharomyces cerevisiae*. *J. Proteome Res.* (2006). doi:10.1021/pr060161n
74. Vizcaíno, J. A. *et al.* 2016 update of the PRIDE database and its related tools. *Nucleic Acids Res.* **44**, (2016).

Figures




Figure 1

kcat distributions in BRENDA and ecYeast7. A) Number of kcat entries in BRENDA per organism. B) kcat distributions for closely related enzyme families. Sample size and median values (in s-1) are shown after each family identifier. C) kcat distributions for enzymes in BRENDA by metabolic context and life kingdoms. Median values are indicated by red dots in each distribution, statistical significance (under a Kolmogorov-Smirnov test) is indicated by red stars for each pair of distributions for a given kingdom. CEM – central carbon and energy metabolism; ALM – Amino acid and lipid metabolism; ISM – Intermediate and secondary metabolism. D) Number of kcat matches in ecYeast7 per assignment category (GECKO 2.0). E) Comparison of the number of kcat matches for E.C. numbers with 0, 1, 2 and 3 introduced wild-cards by GECKO 2.0 and GECKO kcat matching algorithms. F) Cumulative kcat distributions for: all *S. cerevisiae* entries in BRENDA, all entries for fungi in BRENDA, ecYeast7 enhanced by GECKO and ecYeast7 enhanced by GECKO 2.0. Colored points and vertical dashed lines indicate the value for the median value for each distribution. Statistical significance under a Kolmogorov-Smirnov test of the matched kcat distributions when compared to all entries for *S. cerevisiae* and fungi, is shown with red circles and stars, respectively. p-values below 1×10^{-2} are indicated with red. G) Prediction of batch maximum growth rates on diverse media with ecYeast7 enhanced by GECKO 2.0. Glu – glucose, Fru – fructose, Suc – sucrose, Raf – raffinose, Mal – maltose, Gal – galactose, Tre – trehalose, Gly – glycerol, Ace – acetate, Eth – ethanol.




Figure 2

Extending utilization of ecModels. A) ecModels container: Integrated pipeline for continuous and automated update of ecModels. B) Implementation of GECKO simulations in the Caffeine platform (<https://caffeine.dd-decaf.eu/>) for visualization of enzyme usage. The color of the arrows corresponds to the value of the corresponding fluxes. Genes or reactions connected to enzymes with a usage above 90%

are highlighted with a glow around the corresponding text or arrow, respectively. The chosen usage threshold to highlight can be tuned with the slider on the right.




Figure 3

Comparison of predictive capabilities between ecModel1515 and ME-iJL1678 for *E. coli*. A) Maximum batch growth rate predictions on minimal media with diverse carbon sources. Squared-sum errors when compared to experimental values are shown for both ecModel1515 and ME-iJL1678. B) Prediction of total protein content in the cell by ecModel1515 and ME-iJL1678 using the optimal and generalist approaches.




Figure 4

Evaluation of proteomics-constrained ecModels. Comparison of median relative error in prediction of exchange fluxes for O₂ and CO₂ by GEMs, ecModels and proteomics-constrained ecModels across diverse conditions (chemostat cultures at 0.1 h⁻¹ dilution rate) for A) *S. cerevisiae* B) *K. marxianus* C) *Y. lipolytica*. D) Comparison of absolute enzyme usage profiles [mmol/gDw] predicted by ecYeastGEM (ecM) and ecYeastGEM with proteomics constraints (ecP) for several experimental conditions. The region between the two dashed grey lines indicates enzyme usages predicted in the interval $0.5 \leq E_i^{ecP} / E_i^{ecM} \leq 2$, the region between the two dashed black lines indicates enzyme usages predicted in the interval $0.1 \leq E_i^{ecP} / E_i^{ecM} \leq 10$ when comparing the two ecModels. E) Protein burden for different superpathways predicted by ecYeastGEM (ecM) and ecYeastGEM with proteomics constraints (ecP). F) Highly saturated enzymes at different stress conditions for *S. cerevisiae*, *K. marxianus* and *Y. lipolytica* predicted by their corresponding ecModels constrained with proteomics data. Yellow cells indicate condition-responsive enzymes (relative usage ≥ 0.95). Red asterisks indicate enzymes conserved as single copy orthologs across the three yeast species. Std – Reference condition, HiT – High temperature condition, LpH – Low pH condition, Osm – Osmotic stress condition, AA – amino acid metabolism, NUC – nucleotide metabolism, CEM - central carbon and energy metabolism, CofVit – cofactor and vitamin metabolism, Lip – lipid and fatty acid metabolism.




Figure 5

Cumulative distributions of flux variability ranges for iSM996, iYali4 and iML1515 compared to their respective enzyme-constrained versions at low and high growth rates.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Kcatanalysis.pdf](#)
- [suppfile2.docx](#)
- [suppfile3.xlsx](#)
- [suppfile4.xlsx](#)
- [Sfigures.docx](#)