

# RNA-Seq based transcriptome analysis in oral lichen planus

**Haoyu Wang**

Shanghai 9th Peoples Hospital Affiliated to Shanghai Jiaotong University School of Medicine

**Yiwen Deng**

Shanghai 9th Peoples Hospital Affiliated to Shanghai Jiaotong University School of Medicine

**Siqi Peng**

Shanghai University of Traditional Chinese Medicine

**Li Yan**

Shanghai Jiao Tong University Medical School Affiliated Ruijin Hospital

**Hui Xu**

Shanghai 9th Peoples Hospital Affiliated to Shanghai Jiaotong University School of Medicine

**Qingzhong Wang**

Shanghai University of Traditional Chinese Medicine

**Zhengyu Shen** (✉ [neuronszy@sina.com](mailto:neuronszy@sina.com))

<https://orcid.org/0000-0002-1411-8144>

---

## Research

**Keywords:** oral lichen planus, RNA sequencing, weighted gene co-expression network analysis, pathogenesis

**Posted Date:** March 26th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-351759/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Objectives:** Oral lichen planus (OLP) is a T cell-mediated autoimmune disease recognized as an oral potential malignant disorder (OPMD) with the precise mechanism unknown. This study focused on the transcriptional profiles of OLP to elucidate its potential pathogenesis.

**Methods:** We conducted RNA sequencing on matched 6 OLP tissues (n = 6) and 6 normal oral mucosal tissues (n = 6). Gene Ontology (GO) enrichment analysis, Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway and weighted gene co-expression network analysis (WGCNA) were performed on differentially expressed genes (DEGs). We utilized qRT-PCR to validate the top dysregulated genes and hub genes in another 10 pairs of specimens.

**Results:** A total of 153 DEGs (p-values < 0.05) were detected from RNA-Seq. According to GO and KEGG analysis, the dysregulated genes were mainly related to T cell related pathway and Wnt signaling. Based on the WGCNA analysis, 5 modules with high intramodular connectivity and hub genes in each module were gained.

**Conclusions:** RNA-Seq and bioinformatic methods offered a valuable understanding of the biological pathways and key genes in the regulation of OLP. The identified DEGs and hub genes categorized into 2 groups including T cell regulation and inflammation and Wnt signaling pathway may serve as potential novel molecular targets for therapy.

## Introduction

Oral lichen planus (OLP), a chronic T-cell mediated inflammatory disease, could affect oral sites including lips, buccal mucosa, tongue, palate, and gums with various lesions. With a cancerization rate of 1.65% in the patients newly diagnosed as OLP clinically, this disease was classified as an oral potential malignant disorder (OPMD) by the world health organization (WHO) in 2005 (1). Accompanied by clinical features ranging from interlacing white lines, blisters and erosions to burning tingling sensations in the mouth as well as the absence of curative treatment, OLP has brought great distress to patient's physical and mental health, among which the reticular OLP was the most common type. To date, the lack of solid in vitro cell models to study the pathogenesis of OLP posed major challenges to the treatment of OLP, both in terms of therapeutic strategies and new drug development (2).

Researches have suggested that immune, infection, mental and psychological factors and genetics played roles in the pathogenesis of OLP (3). Recently, increasing studies on OLP have been conducted to identify the potential molecular mechanism, mostly focusing on the transcriptional analysis of OLP and several differentially expressed genes (DEGs) have been reported using microarray experiments (4, 5). RNA-Seq, emerging as a novel approach for gene transcriptional profiling recently, has great advantages over the traditional hybridization-based techniques like microarrays. With no need for transcript-specific probes, the full sequences could be converted into several short reads easily sequenced by RNA-Seq, thus facilitating the detection of novel transcripts. Besides, RNA-Seq is more sensitive to identify the genes

with low expression and the rare transcripts could be easily detected for the increased sequencing coverage depth of this technology.

Currently, bioinformatic approaches such as Gene Ontology (GO) and weighted gene co-expression network analysis (WGCNA) have been applied to detect the roles of each gene in the occurrence and progress of the disease at the transcriptional level and have been used widely in the field of various oral diseases, including oral squamous cell carcinoma (6, 7), periodontitis (8) and oral submucous fibrosis (9). However, to date, studies that focus on the transcriptional profiling of OLP using RNA-Seq, especially utilizing bioinformatics have been scarcely reported. As a result, further researches with expanded samples are urgently needed to find relevant genes and signaling pathways mediated by these genes, which may offer new insights into the molecular mechanism of OLP.

In this study, firstly we constructed mRNA expression profiles in OLP tissues by RNA-Seq. Subsequently, to acquire more information from the RNA-Seq data, the GO analysis and WGCNA based on the DEGs were conducted, and hub genes and enrichment pathways were identified, providing a novel overview of the potential pathogenic genes and the therapeutic target of OLP.

## **Materials And Methods**

### **Patients and samples**

Six patients who were newly diagnosed with reticular OLP from September to December 2018 in the department of Oral Mucosal Diseases, Shanghai Ninth People's Hospital were enrolled in this research, including 3 males and 3 females aged 31–47. All the subjects were diagnosed with reticular OLP by clinical manifestations and histological examinations, and had not been treated for OLP within 3 months. Meanwhile, six volunteers who underwent orthognathic surgery were selected as normal controls, including 3 males and 3 females aged 22–44. No obvious inflammation and other pathological changes were detected by department of Pathology. Subjects with systemic diseases, including but not limited to hypertension, diabetes and tumors were excluded.

Subsequently, we further collected another 10 pairs of OLP tissues and normal oral mucosal tissues for qRT-PCR validation. All the specimens were stored at -80°C before use. This study was approved by Institutional Review Board of Shanghai Ninth People's Hospital (SH9H - [2018]85), and patients' informed consent was obtained prior to sample collection..

### **Total RNA Extraction**

Total mRNA from oral mucosal tissues was extracted with TRIzol reagent (Invitrogen, CA, USA), and then concentration and purity were evaluated by Nanodrop 2000 (Thermo Fisher, CA, USA). A ratio of 260 nm/280 nm between 1.8 and 2.2 was set as the screening standard.

### **RNA-Seq**

Six OLP tissues and six normal oral mucosal tissues were used for RNA sequencing. Approximately 2 µg of total RNA was extracted from each specimen and pretreated with Epicentre Ribo-zero™ rRNA Removal Kit. Then the RNA expression profile library was constructed in line with the manufacturer's protocol of NEBNext R Ultradirectional RNA Library Prep Kit (NEB, USA). The steps are as follows: First, RNA was lysed into small fragments after treated with NEBNext First Strand synthesis reaction buffer at high temperature treat, and the First Strand cDNA was synthesized using random hexmer primers and M-MULV reverse transcriptase. The second strand dsDNA was then obtained and the fragment residues were converted into blunt ends by exonuclease or polymerase. Subsequently, the 3' end of each dsDNA fragment was adenylated and connected to the NEBNext adapter with hairpin structure. After purification by AMPure XP system (Beckman Coulter, Beverly, USA), DNA fragments of 150 to 200 bp in length were obtained, which were then sequenced by HiSeq 2500 (Illumina, CA, USA).

### **RNA-Seq data processing and analysis**

FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc/>) was utilized to check the sequencing quality of all the sample data trimmed with the FASTX-Toolkit. The sequencing reads against the human assembly GRCh37 were mapped by TopHat (v 2.0.9). Then we utilized HTSeq to calculate the read counts for each gene in each sample. The transfer matrix method (TMM) was used to normalize the DEGs. Subsequently, DEGs between the evaluated groups were screened using the software DESeq2. Benjamini-Hochberg method was used for multiple comparison tests and  $p < 0.05$  was set as the threshold.

### **Gene Ontology And Enrichment Analysis**

To better explore the underlying biological mechanism and pathways of the DEGs, Gene Ontology (GO) enrichment analysis and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway were performed using topGO and clusterProfiler packages in R platform respectively.  $P < 0.05$  was set as the threshold.

### **Weighted Gene Co-expression Network Analysis**

Weighted gene co-expression network analysis was conducted to generate the gene co-expression network for the DEGs obtained from the RNA-Seq results. In terms of the WGCNA guidance, firstly, an appropriate soft-threshold  $\beta$  was necessary by testing several candidate powers. Then, we selected 10 as the soft-thresholding power for calculation of the adjacencies, after which the adjacencies were converted into a topological overlap matrix to acquire the corresponding dissimilarity. The function *hclust* was utilized to perform cluster analysis on the expression data in the samples and the discrete samples were deleted. The package *dynamicTreeCut* helped identify a group of genes with high topological overlap index as the same module, conduct hierarchical clustering, and establish a nested hierarchical clustering tree. Subsequently, we applied the function *TopHubInEachModule* to detect the hub gene in each module with high connectivity in the weighted co-expression network. The co-expression network of modules was visualized through *igraph* package. Branches of the cluster tree and different colors were used to represent different gene modules. Finally, the personalized expression perturbation profiles (PEEPs)

algorithm which characterized the heterogeneity of gene expression patterns quantitatively was used to identify expression changes within each individual.

### **Verification of the mRNA expression of the top DEGs and hub genes by qRT-PCR**

To validate the results of RNA-sequence and WGCNA analysis, we selected the top five upregulated and downregulated genes respectively in the mRNA profile, as well as the 5 hub genes of each module. Total mRNA from oral mucosal tissues was extracted with TRIzol reagent (TaKaRa, Japan), and then concentration and purity was evaluated by Nanodrop 2000 (Thermo Fisher, USA). After the RNA was reversely transcribed into cDNA with PrimeScript RT kit (TaKaRa, Japan) according to the instructions, SYBR Premix Ex Taq™ kit (TaKaRa, Japan) was applied for qRT-PCR, with  $\beta$ -actin as the endogenous control gene. The real-time quantitative PCR amplification instrument (ABI StepOne Plus) was used to detect the SYBR Green fluorescence signal level after each amplification cycle. The primer sequences of the up-mentioned genes were synthesized by Sangon, China. The relative expression levels of the candidate genes were calculated using the  $2^{-\Delta\Delta Ct}$  method.

## **Results**

### **Identification of the DEGs**

According to high-throughput sequencing data, we obtained 153 DEGs in total with a p-value < 0.05, of which 72 were upregulated while 81 downregulated in OLP tissues against the normal oral mucosal tissues. The top 10 altered genes were listed in the Table S1. In addition, a heatmap and volcano plot of the 153 DEGs expression were shown in the Fig. 1. Among these genes, NEB was the top-ranked upregulated gene with a 14.9-fold increase.

### **Gene Function Analysis For The Dysregulated Genes**

To better explore the biological functions and pathways of the DEGs, GO and KEGG analysis were performed with topGO and clusterProfiler function. Functional annotation revealed that these DEGs were mainly related to cell regulation process. The GO analysis of the biological functions of the DEGs, including biological process (BP), cellular component (CC) and molecular function (MF) were displayed in Fig. 2a and Table S2. To be noted, these enriched terms were mainly related to autophagy process and regulation of membrane including “autophagy”, “process utilizing autophagic mechanism”, “regulation of protein localization to membrane” and “positive regulation of protein localization to membrane”.

13 KEGG pathways significantly enriched were identified in total, which were involved in T cell regulation and Wnt signaling pathway (Fig. 2b). Taken together, functional ontology analysis revealed that the significant expressed genes mainly mediated into immunoregulation process.

### **Identification of hub genes in the functional modules by WGCNA**

WGCNA analysis was conducted to further explore the co-expression network of DEGs and explain the significant transcriptome profile alterations. Firstly, a soft-thresholding power equaled 10 was set using

the pickSoftThreshold program. Shown in the module assignment dendrogram plot (Fig. 3a), a total of five modules were identified by using the package *dynamicTreeCut* functioned to sort out a set of genes with high topological overlap index. Afterwards, the clinical phenotypes and module relationships were combined and the module trait associations were quantified. The results suggested that the brown and turquoise modules were correlated with the phenotypic comparison negatively, while the green, blue and yellow modules were associated with the phenotypic comparison positively (Fig. 3b). Next, hub gene of each module was identified by means of the package *TopHubInEachModule*. The hub genes of the blue, brown, green, turquoise and yellow modules in OLP tissues were RYK (receptor like tyrosine kinase), SLC8A1 (solute carrier family 8 member A1), WDR7 (WD repeat domain 7), MAP3K5 (mitogen-activated protein kinase kinase kinase 5) and GPBP1 (GC-rich promoter binding protein 1), respectively. To better visualize the intramodular connectivity of the hub genes in each module, we displayed the network applying the adjacency matrix of the eigengenes with the hub genes highlighted in each module (Fig. 4). Finally, the data processed by PEEPs algorithm which presents the inherent heterogeneity of gene transcription patterns in each individual specimen was shown in Table S3.

### **Validation of mRNA expression of the top DEGs and hub genes in OLP**

The mRNA expression of the five hub genes (RYK, SLC8A1, WDR7, MAP3K5, GPBP1) obtained from WGCNA analysis as well as the top DEGs (NEB, TNC, NRIP1, DLG1, PTPN22 and SGMS1, TET2, SMO, PARD3, ATP5C1) from RNA-Seq results was detected by real-time PCR in comparisons between another 10 reticular OLP tissues and 10 normal oral mucosal tissues. Set as the endogenous control gene, the expression of  $\beta$ -action was not significantly altered in the 2 groups ( $p > 0.05$ ). The expression of NEB, TNC, NRIP1, DLG1, PTPN22 was notably increased in OLP compared to normal normal oral mucosa (Fig. 5a) while that of SGMS1, TET2, SMO, PARD3, ATP5C1 was decreased in OLP (Fig. 5b), which was consistent with the RNA-Seq data. As displayed in Fig. 5c the expression of SLC8A1 and MAP3K5 was higher in OLP than in the normal control; and the other 3 hub genes, RYK, WDR7 and GPBP1 were downregulated in OLP, although not all the them were statistically insignificant (SLC8A1: 1.87-fold increase,  $p = 0.0025$ ; MAP3K5: 4.11-fold increase,  $p = 0.0001$ ; RYK: 0.55-fold decrease,  $p = 0.0422$ ; WDR7: 0.68-fold decrease,  $p = 0.1324$ ; GPBP1: 0.73-fold decrease,  $p = 0.3399$ ).

## **Discussion**

Oral lichen planus is a chronic disorder of oral mucosa with a global prevalence of 1.01%(10), and reticular OLP is the type with the highest prevalence. OLP is now generally recognized as a T-cell dysfunction-induced autoimmune disease, while the exact mechanism of this disease still remains poorly understood. To help further explain the underlying molecular mechanism in OLP, we integrated the RNA-Seq and bioinformatics to explore the transcriptional profiling of this disease.

Validated by qRT-PCR, a total of 153 DEGs were detected by RNA-Seq in this current study, of which 72 were increased while 81 were decreased in OLP tissues. The top 5 upregulated and downregulated genes in OLP were NEB, TNC, NRIP1, DLG1, PTPN22 and SGMS1, TET2, SMO, PARD3, ATP5C1, respectively. Five

co-expression modules were constructed by WGCNA method and hub genes of each modules were identified as followed: RYK, SLC8A1, WDR7, MAP3K5, GPBP1. According to the GO and KEGG pathway analysis, genes with aberrant expression in OLP were mainly associated with T cell receptor signaling pathway and the Wnt signaling pathway. These above-mentioned results were in line with the existing OLP pathogenesis, and provided new insights into the potential mechanism of OLP.

In OLP, a key pathogenesis is the activation of CD4<sup>+</sup> T-helper and cytotoxic CD8<sup>+</sup> lymphocytes along with the cytokines, resulting in keratinocytes apoptosis and degeneration of the epithelial basal layer (11). T cell in OLP tend to proliferate rapidly and differentiate into distinct Th cell types, including Th1, Th2, Treg, Th17, etc. Growing evidence showed that Th1/Th2 imbalance influenced by Treg subset as well as the CD4<sup>+</sup> Th subset Th17 took great part in immunopathology of OLP (12). Nuclear receptor interacting protein 1 (NRIP1), also known as RIP140, was significantly increased in OLP. NRIP1 interacts with transcription factors and nuclear receptors, and previous studies focused on its molecular mechanism in regulating the immune response. Zhu Jun Yi et al suggested that the overexpression of NRIP1 in macrophages promoted M1-like polarization and expansion in inflammatory diseases (13). NRIP1 was found to modulate the transcription and secretion of pro-inflammatory cytokines by activating NF-κB (14) and this mechanism could also be found in autoimmune disease like psoriasis(15). Interestingly, the NF-κB p65 signaling pathway was also activated and played a key role in the inflammatory response in OLP(16). DLG1, which was one of the top high expressed DEGs in OLP tissues, is a member of the family of molecular scaffolding proteins and encoded a multi-domain scaffolding protein required for normal development. Altered expression of DLG1 is observed in several immunological disorders. Dlg1 functions as a positive regulator of immunity and facilitated T cell receptor (TCR) signal transduction and T cell function. T cells lacking proper Dlg1 expression were deficient in coupling antigenic stimulation resulting in incapability of producing cytokine (17). TET2, which was the most significantly downregulated gene, is universally considered as a tumor-suppressor gene and could modulate Th1 and Th17 cell differentiation and T-cell cytokine production in vivo in autoimmune diseases (18). Earlier study showed that mutation of TET2 caused aberrant CD4<sup>+</sup> T cell proliferation and disturbance of T cell homeostasis (19), which was closely related to the occurrence of OLP. Besides, Tet2 downregulation was vital for 5-hydroxymethylcytosine (5hmC) regulation in adult T-cell leukemia/lymphoma (ATLL) progression (20) and Tet2-KO mice developed myeloid cancers with an incidence rate of about 30% (21). These results demonstrated that TET2 was strongly associated with immune responses. MAP3K5 (also named ASK1), which was the hub gene in turquoise module acquired from WGCNA, is an important member of mitogen-activated protein kinase (MAPK) family and plays crucial roles in physiological processes associated with autoimmune diseases. Recently, S.J. Mnich et al found that as an essential component for the development of rheumatoid arthritis, MAP3K5 participated in the tumor necrosis factor-alpha (TNF-α)-induced production of inflammatory mediators (22) and MAP3K5 deficiency could reduce neuroinflammation in experimental autoimmune encephalomyelitis (23). In addition, in aberrant ribosomal biogenesis, activated MAP3K5 pathways regulated the stability and activity of p53 protein (24), which acted as a transcription factor mediating the induction of cell apoptosis (25). Taken into

consideration the fact that the expression of p53 has been reported upregulated in OLP tissues (26, 27), the interaction between MAP3K5 and p53 may play roles in the pathogenesis of OLP.

Wnt signaling played key roles during cell proliferation and differentiation and was implicated in multiple autoimmune diseases due to its participation in the development of T cells. In OLP, Wnt3 has been featured with positive expression in cytoplasm and lack of nuclear staining, suggesting a potential involvement of Wnt signaling pathway in OLP progression (28). RYK, the hub gene in blue module, belonged to the atypical receptor tyrosine kinase family and was a receptor of Wnt ligand Wnt5a. The interaction of Wnt ligands and RYK originated the downstream signaling cascades critical for immune regulation (29). Besides, Tenascin C (TNC), ranking as the 2nd top upregulated gene in OLP, encoded an extracellular matrix protein (ECM) and was a promoter of tumor metastasis with multiple functions. TNC expression positively linked to cancer progression and cell proliferation. TNC downregulated Dickkopf-1 (DKK1), a Wnt inhibitor by blocking of actin stress fiber formation, thus activating Wnt signaling and induced Wnt target genes in tumor and endothelial cells (30). Nevertheless, the high expression of TNC induced by Wnt/beta-catenin activation, acting as a downstream target, led to a more tumorigenic cell state in Ewing sarcoma (31). The above researches demonstrated that the interaction between TNC and Wnt signaling participated in the modulation of tumor progression and invasion. Nevertheless, to the best of our knowledge, the rest of the key genes have not been reported highly involved in T cell regulation and inflammation or Wnt signaling pathway.

There were limitations in our study. Firstly, the samples size was small. Only reticular OLP tissues were collected and there was a lack of OLP tissues of erosive and erythematous types. Secondly, we did not validate the functions of the key genes as well as the potential signaling pathways in vitro or in vivo.

All in all, in this study, we strictly selected the matched samples of OLP and normal oral mucosal tissues and a range of genes were observed differentially expressed in OLP tissues. Subsequently, by application of the bioinformatic analysis method, our research also confirmed the key roles of biological activities associated with T cell regulation and inflammation and Wnt signaling pathway that have been reported in earlier studies on the molecular mechanisms of OLP and provided a theoretical basis for the study of novel drug targets.

## Conclusions

Taken together, we constructed a transcriptional profiling of OLP using RNA-Seq and 153 differentially expressed gene were detected. Here, with the combination of the results of GO enrichment annotation analysis, KEGG pathway and WGCNA analysis based on the DEGs, biological process including T cell regulation and inflammation and Wnt signaling pathway were identified highly correlated to the pathogenesis of OLP. Further researches are required to explore the molecular mechanism of the key genes as the candidate biomarkers in this disease.

## Abbreviations

BP biological process

CC cell component

DEGs differentially expressed genes

GO Gene Ontology

MF molecular function

OLP oral lichen planus

OPMD oral potential malignant disorder

PEEPs personalized expression perturbation profiles

WGCNA weighted gene co-expression network analysis

## **Declarations**

### **Acknowledgements**

We are grateful to the patients and volunteers who help with the collection of OLP tissues and normal oral mucosal tissues.

### **Availability of data and materials**

The raw data of RNA sequence is available from the corresponding author on reasonable request.

### **Author contributions**

HW performed the experiments and authored the manuscript. YD and LY performed the experiments. SP analyzed the data. HX revised the paper. QW analyzed the data and prepared figures and tables. ZS conceived the experiments, revised the paper and provided fund support.

### **Funding**

This research was supported by National Natural Science Foundation of China (81200786;31871281), Science and Technology Commission of Shanghai (12ZR1417000), Cross-disciplinary Research Fund of Shanghai Ninth People's Hospital, Shanghai JiaoTong University School of Medicine (JYJC201901) and Scientific Research Foundation for Advanced Talents of Shanghai University of Traditional Chinese Medicine.

### **Ethics approval and consent to participate**

Not applicable.

## Consent for publication

All the authors have consented for the publication.

## Conflict of interest

None to declare.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Haoyu Wang, Yiwen Deng contributed equally to this work and should be considered co-first authors.

## Contributor Information

Haoyu Wang, Email: [imayday\\_why@hotmail.com](mailto:imayday_why@hotmail.com)

Yiwen Deng, Email: [sunny3311@sina.com](mailto:sunny3311@sina.com)

Siqi Peng, Email: [pengsiqi9806@gmail.com](mailto:pengsiqi9806@gmail.com)

Li Yan, Email: [yanlishjd@163.com](mailto:yanlishjd@163.com)

Hui Xu, Email: [2201691904@qq.com](mailto:2201691904@qq.com)

Qingzhong Wang, Phone: +86-13524353016, Email: [wangqingzhong3@gmail.com](mailto:wangqingzhong3@gmail.com)

Zhengyu Shen, Phone: +86-13761361333, Email: [neuronszy@sina.com](mailto:neuronszy@sina.com)


## References

1. van der Waal I. Potentially malignant disorders of the oral and oropharyngeal mucosa; terminology, classification and present concepts of management. *Oral Oncol.* 2009;45(4-5):317-23.
2. Sun HY, Zhou GM, Wang Q, Lin XC, Xu B. In vitro culture system for keratinocytes obtained from oral lichen planus lesions. *Clin Oral Investig.* 2014;18(4):1195-203.
3. Alrashdan MS, Cirillo N, McCullough M. Oral lichen planus: a literature review and update. *Arch Dermatol Res.* 2016;308(8):539-51.
4. Danielsson K, Coates PJ, Ebrahimi M, Nylander E, Wahlin YB, Nylander K. Genes involved in epithelial differentiation and development are differentially expressed in oral and genital lichen planus epithelium compared to normal epithelium. *Acta Derm Venereol.* 2014;94(5):526-30.
5. Tao XA, Li CY, Xia J, Yang X, Chen XH, Jian YT, et al. Differential gene expression profiles of whole lesions from patients with oral lichen planus. *J Oral Pathol Med.* 2009;38(5):427-33.

6. Zhang X, Feng H, Li Z, Li D, Liu S, Huang H, et al. Application of weighted gene co-expression network analysis to identify key modules and hub genes in oral squamous cell carcinoma tumorigenesis. *Onco Targets Ther.* 2018;11:6001-21.
7. Wang J, Wang Y, Kong F, Han R, Song W, Chen D, et al. Identification of a six-gene prognostic signature for oral squamous cell carcinoma. *J Cell Physiol.* 2020;235(3):3056-68.
8. Jin SH, Zhou RH, Guan XY, Zhou JG, Liu JG. Identification of novel key lncRNAs involved in periodontitis by weighted gene co-expression network analysis. *J Periodontal Res.* 2020;55(1):96-106.
9. Chikcooree D, Zhu K, Ram V, Wu HJ, He ZJ, Zhang S. A preliminary microarray assay of the miRNA expression signatures in buccal mucosa of oral submucous fibrosis patients. *J Oral Pathol Med.* 2016;45(9):691-7.
10. Gonzalez-Moles MA, Warnakulasuriya S, Gonzalez-Ruiz I, Gonzalez-Ruiz L, Ayen A, Lenouvel D, et al. Worldwide prevalence of oral lichen planus: A systematic review and meta-analysis. *Oral Dis.* 2020.
11. Nogueira PA, Carneiro S, Ramos-e-Silva M. Oral lichen planus: an update on its pathogenesis. *Int J Dermatol.* 2015;54(9):1005-10.
12. Wang H, Zhang D, Han Q, Zhao X, Zeng X, Xu Y, et al. Role of distinct CD4(+) T helper subset in pathogenesis of oral lichen planus. *J Oral Pathol Med.* 2016;45(6):385-93.
13. Yi ZJ, Gong JP, Zhang W. Transcriptional co-regulator RIP140: An important mediator of the inflammatory response and its associated diseases (Review). *Mol Med Rep.* 2017;16(2):994-1000.
14. Ho PC, Tsui YC, Feng X, Greaves DR, Wei LN. NF-kappaB-mediated degradation of the coactivator RIP140 regulates inflammatory responses and contributes to endotoxin tolerance. *Nat Immunol.* 2012;13(4):379-86.
15. Luan C, Chen X, Hu Y, Hao Z, Osland JM, Chen X, et al. Overexpression and potential roles of NRIP1 in psoriasis. *Oncotarget.* 2016;7(45):74236-46.
16. Deng S, Xu Y, Wang X, Liu M, Li L, Yu X, et al. Study on the Role of Salivary Flora and NF-kappaB Inflammatory Signal Pathway in Oral Lichen Planus. *Inflammation.* 2020;43(3):994-1008.
17. Round JL, Tomassian T, Zhang M, Patel V, Schoenberger SP, Miceli MC. Dlg1 coordinates actin polymerization, synaptic T cell receptor and lipid raft aggregation, and effector function in T cells. *J Exp Med.* 2005;201(3):419-30.
18. Ichiyama K, Chen T, Wang X, Yan X, Kim BS, Tanaka S, et al. The methylcytosine dioxygenase Tet2 promotes DNA demethylation and activation of cytokine gene expression in T cells. *Immunity.* 2015;42(4):613-26.
19. Zang S, Li J, Yang H, Zeng H, Han W, Zhang J, et al. Mutations in 5-methylcytosine oxidase TET2 and RhoA cooperatively disrupt T cell homeostasis. *J Clin Invest.* 2017;127(8):2998-3012.
20. Marçais A, Waast L, Bruneau J, Hanssens K, Asnafi V, Gaulard P, et al. Adult T cell leukemia aggressiveness correlates with loss of both 5-hydroxymethylcytosine and TET2 expression. *Oncotarget.* 2017;8(32):52256-68.


21. Moran-Crusio K, Reavie L, Shih A, Abdel-Wahab O, Ndiaye-Lobry D, Lobry C, et al. Tet2 loss leads to increased hematopoietic stem cell self-renewal and myeloid transformation. *Cancer Cell*. 2011;20(1):11-24.
22. Mnich SJ, Blanner PM, Hu LG, Shaffer AF, Happa FA, O'Neil S, et al. Critical role for apoptosis signal-regulating kinase 1 in the development of inflammatory K/BxN serum-induced arthritis. *Int Immunopharmacol*. 2010;10(10):1170-6.
23. Guo X, Harada C, Namekata K, Matsuzawa A, Camps M, Ji H, et al. Regulation of the severity of neuroinflammation and demyelination by TLR-ASK1-p38 pathway. *EMBO Mol Med*. 2010;2(12):504-15.
24. Kim HD, Kim TS, Kim J. Aberrant ribosome biogenesis activates c-Myc and ASK1 pathways resulting in p53-dependent G1 arrest. *Oncogene*. 2011;30(30):3317-27.
25. Romer L, Klein C, Dehner A, Kessler H, Buchner J. p53—a natural cancer killer: structural insights and therapeutic concepts. *Angew Chem Int Ed Engl*. 2006;45(39):6440-60.
26. Oliveira Alves M, Balducci I, Rodarte Carvalho Y, Cabral L, Nunes F, Almeida J. Evaluation of the expression of p53, MDM2, and SUMO-1 in oral lichen planus. *Oral Dis*. 2013;19(8):775-80.
27. Hadzi-Mihailovic M, Petrovic R, Raybaud H, Stanimirovic D, Ozar Koray M. Expression and role of p53 in oral lichen planus patients. *J BUON*. 2017;22(5):1278-86.
28. Balkees G, Noroz N, Ibrahim G, Mohammad D. CD34 and Wnt3 expression in potentially malignant oral disorders. *J Bagh College Dentistry*. 2017;29(3):59-67.
29. Jati S, Sarraf TR, Naskar D, Sen M. Wnt Signaling: Pathogen Incursion and Immune Defense. *Front Immunol*. 2019;10:2551.
30. Saupe F, Schwenzer A, Jia Y, Gasser I, Spenle C, Langlois B, et al. Tenascin-C downregulates wnt inhibitor dickkopf-1, promoting tumorigenesis in a neuroendocrine tumor model. *Cell Rep*. 2013;5(2):482-92.
31. Pedersen EA, Menon R, Bailey KM, Thomas DG, Van Noord RA, Tran J, et al. Activation of Wnt/beta-Catenin in Ewing Sarcoma Cells Antagonizes EWS/ETS Function and Promotes Phenotypic Transition to More Metastatic Cell States. *Cancer Res*. 2016;76(17):5040-53.

## Figures




**Figure 1**

Cluster heatmap and Volcano plot of the 153 DEGs in OLP and normal oral mucosal tissues. a The cluster heatmap of DEGs in each sample. b The volcano plot of the 153 DEGs.




## Figure 2

Gene Ontology enrichment analysis and KEGG pathway analysis of the DEGs. a The enriched GO terms of MF, CC and BP from GO analysis of the DEGs. b The high-enrichment KEGG pathways of the DEGs




## Figure 3

Clustering gene dendrogram plot and relationship between module and phenotypes from WGCNA. a The clustering gene dendrogram showing 5 co-expression modules represented by different colors in OLP. b Module-trait correlation analysis of the 5 co-expression modules, among which MEblue is highly correlated to OLP.



**Figure 4**

Network plots of the 5 modules with the hub genes highlighted by red nodes (A) Network plot of the blue module with RYK as the hub gene; (B) SLC8A1 as the hub gene of the brown module; (C) WDR7 in the green module; (D) MAP3K5 in the turquoise module; (E) GPBP1 in the yellow module.



**Figure 5**

Comparisons between results of RNA-Seq and qRT-PCR verification for the top dysregulated DEGs and hub genes. A total of 15 genes were chosen for validation in another 10 pairs of OLP and normal oral mucosa. The relative mRNA expression levels were in keeping with the RNA-Seq. a Comparisons between results of RNA-Seq and qRT-PCR verification for the top 5 upregulated DEGs. b Comparisons between

results of RNA-Seq and qRT-PCR verification for the top 5 downregulated DEGs. c Comparisons between results of RNA-Seq and qRT-PCR verification for the 5 hub genes

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TableS13.docx](#)