Education in a new place and social support are needed to improve the health of North Korean defectors: A cross-sectional study

Jihyun Lee

Seoul National University School of Dentistry

Yui Park

Seoul National University School of Dentistry

DONGHUN HAN (✉ dhhan73@snu.ac.kr)

Seoul National University School of Dentistry <https://orcid.org/0000-0001-6427-3137>

Ji Won Yoo

University of Nevada Las Vegas

Wenlian Zhou

University of Nevada Las Vegas

Pearl Kim

University of Nevada Las Vegas

Jay J. Shen (✉ jay.shen@unlv.edu)

University of Nevada Las Vegas

Research

Keywords: Education, North Korean defectors, Self-rated health, Social support

Posted Date: June 18th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-35908/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background The number of North Korean defectors (NKDs) escaping to South Korea has increased. The health status of NKDs is an essential factor for a successful settlement into South Korean society. However, no studies have been conducted on the health status of NKDs in terms of education and social support. The aim of this study was to determine the effects of education and social support on the self-rated health status among NKDs. **Methods** This study utilized data gained from face-to-face interviews with 126 NKDs. A multivariable logistic regression and path analysis were performed to assess the effects of education in South Korea and social support with their self-rated health status and to explore the complex relationships between direct and indirect effects of the variables. **Results** NKDs who did not experience regular education in South Korea responded that they were in poor health compared to their counterpart (OR=5.78). Although a direct association between education in South Korea and self-rated health was not shown, there was an indirect path from education in South Korea to self-rated health through social support. **Conclusions** Participation in regular education is important for the health status of NKDs. Social policies and NKD assistance programs should consider and reflect the combination of education and social support interventions relevant to the health status of NKDs.

Background

North Korea's economic difficulties after the fall and dissolution of communism in Eastern Europe and the Soviet Union and a great flood in 1995 that caused the Great Famine led to the deaths of millions of North Koreans due to starvation and hunger-related illnesses during the late 1990s [1]. Since then, many North Korean defectors (NKDs) have escaped to South Korea. The highest concentration of military power in the world is on the border between North and South Korea, so NKDs often pass through third countries such as China, Thailand, and Laos to enter South Korea. Despite the huge barriers and great danger, the number of NKDs entering South Korea has been steady at 1000 ~ 2000 annually since 2002, and the cumulative total by September 2019 was 33,247 [2].

Despite the financial and medical support by Korean Government, NKDs have experienced various difficulties in settling into South Korean society and managing their health conditions [3]. According to a survey of NKDs in 2018, 62.2% of NKDs suffered from stress in overall daily life, while only 54.4% of South Koreans suffered from stress. Furthermore, 14.6% of NKDs had suicidal thoughts compared to 5.1% of South Koreans. Moreover, a particularly notable part was that the "physical/mental illness and disability" was the second highest cause of suicidal ideation (23.3%) [4].

Education can be considered as one of the most critical factors that potentially affect the adjustment of NKDs to South Korean society and their health. The health effects of education attainment against risk of morbidity and mortality are well established [5–7]. High education is shown to protect against a decline in self-rated health (SRH) [8, 9], chronic disease [10], and mortality [11]. As another essential component of human life, social support has been known to be associated with health and well-being. Social

isolation entails a mortality risk comparable in magnitude to that of smoking, a sedentary lifestyle, obesity, and alcohol abuse [12].

Previous evidence suggests that education and social relationships interact with each other to affect health. Some research in this area has found that the protective effects of social support on health are stronger among individuals with high education levels [13, 14]. These studies have demonstrated that those with high levels of education benefit most from the protective qualities of social support, and the social support-health connection is almost nonexistent among those with lower levels of education. Conversely, one study found that the protective qualities of social support are stronger among individuals who are less educated [15]. Yet, additional empirical evidence has demonstrated that social support is predictive of worse, rather than better health outcomes among individuals with a low level of education [13, 14].

Education is not only a determinant of health but also an opportunity to form social relationships. Especially for NKDs, being a part of a social community in South Korean society, such as a regular education in South Korea, provides them with good opportunities to receive social support and may have an important health-promoting impact. Previous studies [16, 17] showed that facilitating the visit of educational institutions for refugees is an essential factor for successful social inclusion.

However, literature is scarce on the exact nature of the links among education in South Korea, social support, and self-rated health status among NKDs. We hypothesized that education in South Korea, social support, and self-rated health status would be directly associated with each other. We also hypothesized that education in South Korea would be directly and indirectly associated with self-rated health status through social support. Therefore, the aims of this study was to identify the individual effects of education and social support on the self-rated health status, respectively, and then to further assess the path from education in South Korea to self-rated health status through social support among NKDs.

Methods

This was a cross-sectional study to assess the association of education and social support with self-rated health. It was approved by the Institutional Review Board of the Seoul National University School of Dentistry (No.S-D20170041).

Participants

The study participants were recruited with cooperation from the Hana Center, alternative schools for NKDs and the Catholic Bishop's Conference of Korea National Reconciliation Committee in Seoul and Gyeonggi Province. A total of 137 participants were recruited, and the purpose and method of this study were explained to them. All the participants voluntarily agreed to join this study and gave their written consent. The criteria for selection and exclusion were as follows.

- Selection criteria
 - those who completed education at the Hana center
 - those who can read and answer the questionnaire by themselves or who can listen to the explanation of the questionnaire and answers
 - those willing to participate in this study
- Exclusion Criteria
 - those with a severe mental illness condition
 - those who want to quit

Variables

The dependent variable was self-rated health status. It was measured with a 5-point scale (very good, good, moderate, poor, and very poor) and reclassified into two groups (good/moderate and poor).

Independent variables were education in South Korea and social support. Education in South Korea was reclassified into 1) college or more, 2) middle/high school, and 3) no education experience in South Korea. Social support score was measured by the Korean version of the Duke-UNC social support questionnaire which was developed by Suh et al. [18] The tool consists of 13 questions with a 5-point scale. After summing up, the score ranges from 13 to 65 points, and a higher score indicates more social support. The reliability of the Korean version of the Duke-UNC social support questionnaire was a Cronbach's $\alpha=0.89$ at the time of its development. Social support was categorized as a lower/higher group based on the median score (34 point).

Confounding variables were age (continuous), gender (male/female), living with family (yes/no), residence period in South Korea (continuous), smoking (yes/no), and drinking (yes/no). Residence period in South Korea was calculated by months from entrance date into South Korea to October 2017 when the data collection was completed.

Statistical analysis

After excluding missing values, 126 participants were included in the final sample for analysis. The characteristics of the participants were analyzed using frequency, percent, means, and standard deviations. Chi-square tests for categorical variables and independent t-tests for continuous variables were used to assess the associations of education in South Korea and confounders with subjective poor health (Table 1). The association between social support and self-rated health status (Supplementary Table 1 and Fig. 1) and the association between social support and education in South Korea (Supplementary Table 2 and Fig. 2) were evaluated by analysis of variance (ANOVA) and analysis of covariance (ANCOVA) controlling for confounders, respectively. The association between education in South Korea and subjective poor health was assessed using logistic regression models adjusting for the

above mentioned confounders (Model 1 in Table 2). Social support was regarded as a mediator, and logistic regression analysis was done controlling for confounders in model 1 and social support (Model 2 in Table 2). The role of the social support was evaluated by the percent (%) excess odd explained, which was calculated as $[(OR_{adj\ confounders} - OR_{adj\ confounders + mediator}) / (OR_{adj\ confounders} - 1) \times 100 (\%)]$ in this study [19]. The % excess odd explained represents the degree to which a mediator explains the relationship between education in South Korea and subjective poor health. Finally, to assess the direct and indirect effects of the education of NKDs in South Korea on their health, we set up a model that hypothesizes that education in South Korea affects their SRH directly and indirectly by their perceived social support. A path analysis was conducted using AMOS 23.0 [20]. Path coefficients were estimated by maximum likelihood estimation. Maximum Likelihood Bootstrapping was conducted to test the significance of the indirect effect with bootstrap samples of 2,000 and 95% bias-corrected confidence intervals. Model fit was evaluated with proper indices.

Table 1
Related factors of self-rated health status among North Korean defectors (n = 126)

Self-rated health status				
Independent Variables		good/moderate	poor	
		(n = 98)	(n = 28)	<i>p</i> -Value
Age (years)		29.88 [14.71]	48.89 [16.98]	< 0.001
Gender	Male	29 [29.6]	2 [7.1]	0.029
	Female	69 [70.4]	26 [92.9]	
Living with family	No	25 [25.5]	10 [35.7]	0.288
	Yes	73 [74.5]	18 [64.3]	
Residence period in South Korea (months)		67.35 [53.50]	97.93 [52.20]	0.009
Smoking	Yes	19 [19.4]	3 [10.7]	0.286
	No	79 [80.6]	25 [89.3]	
Drinking	Yes	66 [67.3]	14 [50.0]	0.093
	No	32 [32.7]	14 [50.0]	
Education in South Korea	No	16 [16.3]	16 [57.1]	< 0.001
	Middle/high school	46 [46.9]	7 [25.0]	
	College or more	36 [36.7]	5 [17.9]	
<i>Notes.</i> Means for continuous variables are reported with standard deviation in brackets. Frequencies for categorical variables are reported with column percent in brackets. <i>p</i> -values were obtained by independent t-test in continuous variables and chi-square test in categorical variables.				

Table 2
Unadjusted and adjusted social support score according to self-rated health status

Social support score	Self-rated health status					<i>p</i> -Value
	Very good	Good	Moderate	Poor	Very poor	
	(n = 10)	(n = 39)	(n = 49)	(n = 18)	(n = 10)	
Unadjusted						
Mean	27.60 ^a	32.00 ^{ab}	34.27 ^{abc}	36.67 ^{bc}	41.30 ^c	0.001
[SD]	[9.66]	[8.15]	[6.76]	[8.27]	[7.51]	
Adjusted						
Mean	28.47 ^a	32.06 ^{ab}	34.13 ^{bc}	37.20 ^{cd}	39.86 ^d	0.007
[SE]	[2.37]	[1.18]	[1.07]	[1.84]	[2.47]	
Unadjusted means are reported with standard deviation in brackets. Adjusted means are reported with standard error in brackets. <i>p</i> -values were obtained by ANOVA and ANCOVA controlling for age, gender, living with family, residence period in South Korea, smoking, and drinking. SD = standard deviation; SE = standard error.						

Results

Descriptive results are listed in Table 1. Age ranges from 13 to 74 years old (mean age: 34.1 years old) and 41.3% were in their twenties. The NKDs were dominantly females (75.4%) and most of them lived with family (72.2%). The range of the residence period in South Korea was 1 to 213 months (mean length of residence in South Korea: 74.14 months), and two thirds of the participants lived in South Korea for 1 ~ 5 years (37.3% for 12 ~ 59 months and 31.7% for 60 ~ 119 months). Smokers and drinkers were 17.5% and 63.5%, respectively.

Additionally, 78.6% of the participants completed a regular educational curriculum in South Korea. The average score of social support was 33.9 with a minimum score of 13 and a maximum score of 51 points. Subjective health conditions showed that "good" and "moderate" were 38.9%, respectively, which surpassed "poor" (22.2%).

ANOVA and ANCOVA results are presented in Tables 2 and 3. The association between social support and self-rated health status is shown in Table 2. Social support scores increased as self-rated health status deteriorated. Individuals with at least a college degree reported a lower social support score (that is, a high social support status) than those who had less than a college degree (Table 3).

Table 3
Unadjusted and adjusted social support score according to education in South Korea

Social support score	Education in South Korea			<i>p</i> -Value
	College or more	Middle/high school	No	
	(n = 41)	(n = 53)	(n = 32)	
Unadjusted				
Mean	31.29 ^a	33.77 ^{ab}	37.59 ^b	0.004
[SD]	[8.45]	[7.77]	[7.50]	
Adjusted				
Mean	31.90	33.05	38.01	0.020
[SE]	[1.27]	[1.14]	[1.63]	
Unadjusted means are reported with standard deviation in brackets. Adjusted means are reported with standard error in brackets. <i>p</i> -values were obtained by ANOVA and ANCOVA controlling for age, gender, living with family, residence period in South Korea, smoking, and drinking. SD = standard deviation; SE = standard error.				

In an adjusted logistic regression analysis controlling for age, gender, living with family, residence period in South Korea, and smoking and drinking, those who did not experience a regular educational curriculum in South Korea showed 5.78 odds of self-reported poor health (Model 1 in Table 4, *p*-value = 0.024). This association was attenuated to 3.95 in model 2 (*p*-value = 0.096). The % excess odd explained for no educational experience in South Korea was 38.23% when social support was included in model 2.

Table 4
Association between education in SK and subjective poor health

Independent Variables	Model 1		Model 2	
	Odds Ratio	95% CI	Odds Ratio	95% CI
Education in SK				
College or more	1.00	[Reference]	1.00	[Reference]
Middle/high school	2.05	[0.47–8.95]	2.11	[0.46–9.67]
No	5.78	[1.26–26.57]	3.95	[0.78–19.93]
Age				
(years, continuous)	1.05	[1.00-1.09]	1.04	[1.00-1.09]
Gender				
(reference = male)	5.53	[0.56–54.47]	8.20	[0.70-96.59]
Living with family				
(reference = yes)	2.48	[0.84–8.51]	1.58	[0.46–5.39]
Residence period in SK				
(months, continuous)	1.00	[0.99–1.02]	1.01	[0.99–1.02]
Smoking				
(reference = no)	3.23	[0.40-26.03]	5.85	[0.57–60.34]
Drinking				
(reference = no)	1.25	[0.36–4.35]	1.05	[0.28–3.90]
Social support				
(score, continuous)			1.10	[1.02–1.18]
Model1 was adjusted for age, gender, living with family, residence period in SK, smoking, and drinking.				
Model2 was adjusted for age, gender, living with family, residence period in SK, smoking, drinking, and social support. CI = confidence interval; SK = South Korea.				

We evaluated the path model of direct and indirect effects between the three variables of interest, namely education in South Korea, social support, and self-rated health status along with a control variable age. As shown in Fig. 1 and Table 5, the path from Education in South Korea to SRH was not significant, in other words, the direct effect of education was not supported ($p = 0.064$). Paths from education to social support ($\beta = 0.30$, $p = 0.001$) and from social support to health ($\beta = 0.30$, $p = 0.001$) were all statistically significant, and the strength was big as the effect of age on health ($\beta = -0.31$, $p = 0.001$). Thus, education

in South Korea did not directly affect SRH but did indirectly affect it through social Support. The fit of the path model was assessed using the indices: $\chi^2/df = 2.64$; Normed Fit Index (NFI) = 0.945; Comparative Fit Index (CFI) = 0.961; and Standardized Root Mean Square Residual (SRMR) = 0.0107. All the indices were above the recommended criteria for good or fit [21].

Table 5
Direct, indirect and total effects of Education in South Korea on Self-rated Health

Dependent Variable	Independent Variable	Direct Effect	Indirect Effect	Total Effect	R ²
Self-rated Health	Education in South Korea	0.16	0.089**	0.25**	0.29
	Social Support	0.30***		0.30**	
	Age	-0.31***		-0.31**	
Social Support	Education in South Korea	0.30***		0.30**	0.09
* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$					

Discussion

This study analyzed the association of receiving education in South Korea and social support with self-reported health status among NKDs and the link among those three variables. This is the first study that observed the independent main effect of receiving education in South Korea on health among NKDs. Our findings indicate that receiving education in South Korea is positively associated with self-rated health status, even when potential confounding factors are taken into consideration. Like other populations, educational attainment is a significant predictor of morbidity, mortality, and other indices of physical health among NKDs [22–24]. Receiving education in South Korea among NKDs is also significantly associated with social support. Those with college education in South Korea, relative to those without, had a higher level of social support. It is evident that education in South Korea is associated with psychological aspects of health that may be considered as an important part of social relationships.

Our findings also indicate that social support is positively associated with better health. The buffering hypothesis suggests that social support affects health by attenuating the physiological effects of psychosocial stress [25]. Social support has been shown to have both buffering effects and direct beneficial effects on general morbidity [26, 27]. Evidence suggests that socioeconomic status (SES) and social relationships can interact with each other to affect health. Some studies have found that the protective qualities of social support are stronger among individuals with a low SES [15, 28]. Social support is particularly important for individuals with a lower SES. However, some research reports that the protective effects of social support on health are stronger among individuals with a high SES [13, 14, 29].

This study's finding further supports that not only receiving education in South Korea is associated with the health status of NKDs—independence of socio-demographic factors and health behaviors—but that also social support has an important role in the association between education in South Korea and self-rated health among NKDs. However, we are not clear on the exact pathway from education to health which remains to be elucidated.

The hypothesis that is the most interesting to us is the potential effects of receiving education in South Korea on self-rated health either through social support or not. Our finding showed that education in South Korea did not directly affect their health, but education leads to social support, which affects their health. In other words, education affects health indirectly through social support. This result is not consistent with other studies [5–9] that argued a direct education effect on health. However, the significant path from education to social support and the one from social support to health were in agreement with previous studies. Previous evidence has supported that greater social support is associated with more education [30, 31]. Formal education may promote interpersonal skills and friendships and provide more resources (e.g., income and leisure time) to develop and maintain relationships [32]. Social support may also promote healthier lifestyles by enhancing individuals' motivation, normative environment, and access to information [33–35]. The significant indirect effect of education on health through social support is in the same line with studies showing that education and social relationships interact with each other to affect health [13, 14]. It is noteworthy that the strength of this indirect effect of education (0.25) and direct effect of social support (0.30) on health is comparable to the strength of the age effect (-0.31) on health. There could be many other mediating factors between education and health. In this study, we proposed social support as one of the factors. Further study is recommended to discover other factors that link education and health. Therefore, based on our findings, there is a need for a policy enforcing education and social integration activities in the NKD group for their health.

This study has limitations. First, the selection of study participants was restricted. There are 33247 NKDs throughout South Korea. However, the participants in this study were selected by a snowball sampling method mainly from the residents in the Seoul and Gyeonggi metropolitan areas. Therefore, the study participants may not be considered as a representative sample of the NKD population. This led to a wide range for the confidence interval in the results of the logistic regression analysis in this study. Therefore, the interpretation of the study results should be performed carefully. However, considering the limited opportunities to contact NKDs and restrictions on information collection from NKDs, these limitations are inevitable. Second, because this study was a cross-sectional survey research, it is difficult to infer causality among the variables. A well-designed longitudinal follow-up study will be needed to find underlying factors affecting the health status of NKDs on which proper health promotion and intervention strategies and programs can be developed. Third, although the questionnaires used in this study had proven validity and reliability, it was still challenging for some NKDs with different social and cultural backgrounds to fully understand and answer the questionnaires. It is necessary to develop tools to better measure the health status of this vulnerable group.

Conclusions

Despite the aforementioned limitations, this study has some vital findings. The NKDs encounter challenges in adapting to a new political and social system in South Korea. To assist NKDs to overcome these challenges and improve their health condition, social integration policies and interventions focusing on enrollment in regular education curriculums/programs and participating in social interaction activities are essential. Further research should develop related intervention programs for NKDs and assess their outcomes.

Abbreviations

ANCOVA

analysis of covariance

ANOVA

analysis of variance

CFI

Comparative Fit Index

NFI

Normed Fit Index

NKDs

North Korean defectors

SES

socioeconomic status

SRH

self-rated health

SRMR

standardized root mean square residual

Declarations

Ethics approval and consent to participate

This study was approved by the Institutional Review Board of the Seoul National University School of Dentistry (No.S-D20170041). Informed consent was obtained from all participants.

Consent for publication

Informed consent was obtained from all participants.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This was a self-funded study. Dental Research Institute supported the cost of English editing.

Authors' contributions

All authors have made substantial contributions to conception and design of the study. Han DH and Shen JJ contributed equally to conception, design, and interpretation of study. Lee J drafted this manuscript. Park Y carried out the survey and made a dataset. Yoo JW, Kim P, and Zhou W revised it critically for important intellectual content. All authors have given final approval of the version to be published.

Acknowledgements

Not applicable.

References

1. Noland M, Robinson S, Wang T. Famine in North Korea: Causes and Cures. *Econ Dev Cult Change*. 2001;49:741–67.
2. Ministry of Unification. Settlement Support for North Korean Defectors. 2020. https://www.unikorea.go.kr/eng_unikorea/whatwedo/support/ Accessed 06 May 2020.
3. Park SM. Primary health care for vulnerable population. *Korean J Fam Med*. 2009;30:91–7.
4. Korea Hana Foundation. 2018 Social Integration Survey of North Korean Refugees in South Korea. Seoul: Korea Hana Foundation; 2019. p. 85–87.
5. Baker DP, Leon J, Smith Greenaway EG, Collins J, Movit M. The education effect on population health: A reassessment. *Popul Dev Rev*. 2011;37:307–32.
6. Herd P, Goesling B, House JS. Socioeconomic position and health: The differential effects of education versus income on the onset versus progression of health problems. *J Health Soc Behav*. 2007;48:223–38.
7. Marmot MG, Shipley MJ. Do socioeconomic differences in mortality persist after retirement? 25 year follow up of civil servants from the first Whitehall study. *BMJ*. 1996;313:1170–80.
8. Meyer OL, Castro-Schilo L, Aguilar-Gaxiola S. Determinants of mental health and self-rated health: A model of socioeconomic status, neighborhood safety, and physical activity. *Am J Public Health*. 2014;104:1734–41.

9. Assari S, Lankarani MM. Demographic and Socioeconomic Determinants of Physical and Mental Self-rated Health across 10 Ethnic Groups in the United States. *Int J Epidemiol Res.* 2017;3:185–93.
10. Choi AI, Weekley CC, Chen SC, et al. Association of educational attainment with chronic disease and mortality: The Kidney Early Evaluation Program (KEEP). *Am J Kidney Dis.* 2011;58:228–34.
11. Gakidou E, Cowling K, Lozano R, Murray CJ. Increased educational attainment and its effect on child mortality in 175 countries between 1970 and 2009: A systematic analysis. *Lancet.* 2010;376:959–74.
12. Holt-Lunstad J, Smith TB, Layton JB. Social relationships and mortality risk: A meta-analytic review. *PLoS Med.* 2010;7:e1000316.
13. Gorman BK, Sivaganesan A. The role of social support and integration for understanding socioeconomic disparities in self-rated health and hypertension. *Soc Sci Med.* 2007;65:958–75.
14. Fagundes CP, Bennett JM, Alfano CM, et al. Social support and socioeconomic status interact to predict Epstein-Barr virus latency in women awaiting diagnosis or newly diagnosed with breast cancer. *Health Psychol.* 2012;31:11–9.
15. Brummett BH, Barefoot JC, Vitaliano PP, Siegler IC. Associations among social support, income, and symptoms of depression in an educated sample: the UNC Alumni Heart Study. *Int J Behav Med.* 2003;10:239–50.
16. Thomas R. The Right to Quality Education for Refugee Children Through Social Inclusion. *J Hum Rights Soc Work.* 2016;1:193–201.
17. Gušić S, Malešević A, Cardeña E, Bengtsson H, Søndergaard H. “I Feel Like I do Not Exist:” A Study of Dissociative Experiences Among War-Traumatized Refugee Youth. *Psychol Trauma.* 2018;10:542–50.
18. Suh SY, Im YS, Lee SH, Park MS, Yoo TW. A study for the development of Korean version of the Duke-UNC functional social support questionnaire. *J Korean Acad Fam Med.* 1997;18:250–60.
19. Szklo M, Nieto J. *Epidemiology. Beyond the basics.* 3rd ed. London: Jones & Bartlett Publishers; 2014. pp. 161–2.
20. Amos [computer program]. Version 23.0. Chicago: IBM SPSS Inc; 2014.
21. Hu L, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Struct Equ Model.* 1999;6:1–55.
22. Strand BH, Grøholt EK, Steingrimsdóttir ÓA, Blakely T, Graff-Iversen S, Naess Ø. Educational inequalities in mortality over four decades in Norway: Prospective study of middle aged men and women followed for cause specific mortality, 1960–2000. *BMJ.* 2010;340:c654.
23. Case A, Deaton A. Rising morbidity and mortality in midlife among white non-Hispanic Americans in the 21st century. *Proc Natl Acad Sci USA.* 2015;112:15078–83.
24. Viner RM, Hargreaves DS, Ward J, Bonell C, Mokdad AH, Patton G. The health benefits of secondary education in adolescents and young adults: An international analysis in 186 low-, middle-and high-income countries from 1990 to 2013. *SSM Popul Health.* 2017;3:162–71.

25. Cohen S, Wills TA. Stress, social support, and the buffering hypothesis. *Psychol Bull.* 1985;98:310–57.
26. McKay DA, Blake RL Jr, Colwill JM, et al. Social supports and stress as predictors of illness. *J Fam Pract.* 1985;20:575–81.
27. Blake RL Jr, McKay DA. A single-item measure of social supports as a predictor of morbidity. *J Fam Pract.* 1986;22:82–4.
28. North RJ, Holahan CJ, Moos RH, Cronkite RC. Family support, family income, and happiness: a 10-year perspective. *J Fam Psychol.* 2008;22:475–83.
29. Krause N. Anticipated support, received support, and economic stress among older adults. *J Gerontol B Psychol Sci Soc Sci.* 1997;52:P284–93.
30. Mirowsky J, Ross CE. Education, social status, and health. Hawthorne: Aldine de Gruyter; 2003.
31. Cutler DM, Lleras-Muney A. Understanding differences in health behaviors by education. *J Health Econ.* 2010;29:1–28.
32. Cutler DM, Lleras-Muney A. Education and health: Evaluating theories and evidence. In: Schoeni RF, House JS, Kaplan GA, Pollack H, editors. *Making Americans healthier: Social and economic policy as health policy.* New York: Russell Sage Foundation; 2008. pp. 29–60.
33. Uchino BN. Social support and health: a review of physiological processes potentially underlying links to disease outcomes. *J Behav Med.* 2006;29:377–87.
34. Chatzisarantis NLD, Hagger MS, Brickell T. Using the construct of perceived autonomy support to understand social influence within the theory of planned behavior. *Psychol Sport Exerc.* 2008;9:27–44.
35. Gillison FB, Standage M, Cumming SP, Zakrzewski-Fruer J, Rouse PC, Katzmarzyk PT. Does parental support moderate the effect of children's motivation and self-efficacy on physical activity and sedentary behaviour? *Psychol Sport Exerc.* 2017;32:153–61.

Figures

Figure 1

Path diagram of direct and indirect effects of Education in South Korea on Self-rated Health. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$