The Value of Urinary RBP4 in The Diagnosis of FSGS and other Renal Diseases.

Karolina Marek-Bukowiec (✉ marek-bukowiec@wssk.wroc.pl)

Regional Specialized Hospital in Wrocław <https://orcid.org/0000-0003-0606-9466>

Andrzej Konieczny

Wrocław Medical University

Krzysztof Ratajczyk

Regional Specialized Hospital in Wrocław

Katarzyna Macur

Univeristy of Gdańsk and Medical Univeristy of Gdańsk

Paulina Czaplewska

University of Gdańsk and Medical Univeristy of Gdańsk

Agnieszka Czyżewska-Buczyńska

Regional Specialized Hospital in Wrocław

Paweł Kowal

Regional Specialized Hospital in Wrocław

Wojciech Witkiewicz

Regional Specialized Hospital in Wrocław

Research article

Keywords: FSGS, renal diseases, urinary RBP4

Posted Date: June 24th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-35997/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Focal segmental glomerulosclerosis (FSGS) is the leading cause of chronic renal failure worldwide. Unequivocal diagnosis of FSGS can only be made with a renal biopsy, which is an invasive, risk-associated medical procedure. The discovery of non-invasive molecular biomarkers for the diagnosis of FSGS still remains an important scientific goal. This study examines the urinary proteome profile of FSGS patients and reference groups, in order to identify urinary protein expression alterations indicative of FSGS.

Methods

Urine samples were collected from subjects representing FSGS, IgA nephropathy (IgAN), clear cell renal cell carcinoma (ccRCC), chromophobe renal cell carcinoma (chRCC), and healthy control group, respectively. The samples were pooled and subjected to SWATH-MS proteomics analysis. ELISA was utilized to validate the expression level of Retinol-binding protein 4 (uRBP4) in 130 urine supernatant samples (21 FSGS, 20 IgAN, 31 ccRCC, 21 chRCC, 7 prostate cancer patients and 30 healthy volunteers).

Results

The MS study identified 194 (FSGS), 179 (IgAN), 271 (ccRCC), 255 (chRCC), and 275 (healthy controls) urinary proteins. The comparative proteomic analysis revealed that Retinol-binding protein 4 clearly discriminates FSGS from the rest of the groups. Increased levels of uRBP4 in FSGS urine specimens were also detected by ELISA (FSGS vs IgAN $p=0.0244$, FSGS vs ccRCC $p=0.004$, FSGS vs chRCC $p=0.013$, FSGS vs prostate cancer $p=0.02$, FSGS vs healthy controls $p<0.00001$). Significantly elevated levels of uRBP4 were also observed for IgAN, ccRCC and chRCC versus healthy individuals.

Conclusions

Urinary RBP4 protein is significantly upregulated in FSGS in comparison to IgAN, ccRCC, chRCC, prostate cancer patients and healthy subjects. Determining FSGS diagnosis based on uRBP4 expression alone is not possible. Specific uRBP4 concentration cut-off can be applied to accurately distinguish individuals with renal disorder (in general) from healthy subjects. Thus, urinary RBP4 could serve as a screening biomarker identifying people at risk of renal disorders, who should undergo more detailed diagnostics.

1 Background

Focal segmental glomerulosclerosis is the predominant subtype of primary glomerular diseases (primary glomerulopathies, PGs) in adults and the commonest cause of chronic renal failure (chronic kidney disease, CKD) [1]. The worldwide incidence of this rare condition is 0.8 cases per 100,000 subjects per year [2]. The precise diagnosis of FSGS is based entirely on kidney biopsy that is an aggressive procedure associated with risk of complications [1]. Presently, there are no molecular diagnostic strategies available

for FSGS that could serve as a non-invasive alternative to renal biopsy. Undoubtedly, urinary proteome comprises a valuable source of biomarkers of kidney diseases, as much of the proteins come directly from the kidneys and reflect their actual physiological status. Urinary proteins are relatively stable comparing to the proteins of sera, and are more desired as biomarkers due to a non-invasive way of urine collection [3]. Interestingly, urinary proteins proposed hitherto as possible candidates of chronic kidney diseases seemed to outperform the 'biomarkers' found in the blood [4].

There is a very small number of studies performed to date exploring the urinary proteome of FSGS. All of them are based on a small spectrum of reference groups i.e. healthy controls and/or other glomerulopathies [5, 6, 7, 8]. In the past studies the urinary proteome of FSGS has been never confronted simultaneously with the proteomic profile of the normal status, IgA nephropathy (the second most common glomerulopathy worldwide) and the commonest subtypes of renal cancers i.e. ccRCC and chRCC. Such strategy i.e. to mine and compare the urinary proteomes of distinct renal disease states (glomerulopathies, kidney cancers, normal status) may enhance the chances of finding quantitative/qualitative proteomic mark truly specific for particular renal disorder. Within the framework of this study we aimed to characterize the urinary proteome of FSGS subjects, and compare it to the protein expression profiles of the following reference groups: healthy controls, IgAN subjects, ccRCC, chRCC, and prostate cancer subjects in order to extract proteins specifically indicative of the FSGS.

2 Methods

2.1 Ethics Statement

This study was approved by the Bioethics Committee of Regional Specialized Hospital in Wrocław (approval ID: KB/nr 9/rok 2016). All samples were collected at Regional Specialized Hospital after obtaining written informed consent from study participants.

2.2 Description of Participants

Participants in this *study* included healthy volunteers (age range; 23 – 60, n=30, 16 men and 14 women), patients with biopsy-proven primary FSGS (age range 24-74, n=20, 16 men and 4 women), and primary IgAN (age range 21-66, n=19, 11 men and 8 women), patients with histopathologically confirmed primary ccRCC (age range 48-89, n=31, 18 men and 13 women), primary chRCC (age range 19-86, n=21, 12 men and 9 women), and prostate cancer (age range 57-79, n=7 men).

2.3 Urine Sample Collection and Handling

All participants provided first-morning urine (20 ml - 50 ml) in a sterile urine containers. Urine samples were obtained in the morning before scheduled renal biopsy (IgAN, FSGS patients) or surgery (renal cancer and prostate cancer patients). The specimens were processed at room temperature within 1-3 hours post-collection. All samples were tested for the following parameters: leukocyte esterase, nitrite, urobilinogen, protein, pH, blood, specific gravity, ketone, bilirubin, glucose with a Multistix 10 SG Reagent

Strips (Siemens Healthcare Diagnostics Inc, NY, USA). Urine specimens of healthy controls and prostate cancer patients had normal urine parameters. The summary of urine dipstick test results is included in Supplementary table 1. Urine samples were centrifuged at 4300g for 30 minutes at room temperature and the supernatants were aliquoted into 2 ml sterile LoBind tubes, and stored at -80°C until further processing.

2.4 Sample Preparation for SWATH-MS/MS (quantitative analysis)

Urine supernatants were thawed in a waterbath (37°C) and vigorously vortexed. Equal volumes of urine supernatant from four subjects from the same group were pooled. The pooled samples were concentrated and desalted using 10-kDa molecular weight cutoff membranes (Merck KGaA, Darmstadt, Germany). Concentrated proteins were washed twice and eluted with equal volumes of molecular biology grade water (4000g/room temperature/20 minutes). The protein quantity in each sample was assessed in replicates by Qubit 2.0 fluorometer using Qubit Protein Assay Kit (Thermo Fisher Scientific, MA, USA). Adequate volume of each pooled sample (corresponding to 15µg of proteins) was vacuum dried using speedvac (miVac, Genevac, UK). The protein pellets were suspended in 15 µl 50 mM NH₄HCO₃ and sonicated in a water bath (15 minutes/no heating). Samples were reduced by 10 mM dithiotreitol (DTT) at 56°C for 30 min, and subsequently alkylated by iodoacetamide in the final concentration of 20 mM (darkness/room temperature/30 minutes). After adding 20 µl of 50 mM NH₄HCO₃ to each sample, the samples were incubated overnight with MS-grade trypsin (1:50 trypsin to substrate ratio) with constant shaking (750 rpm) at 37°C. Digestion reaction was quenched by 5% formic acid/50% acetonitrile solution. The samples were desalted using a ZipTip C18 (Millipore) according to manufacturer's instruction. Peptides were eluted into 50% acetonitrile/0,1% formic acid solution and subjected to LC-MS/MS.

2.5 Sample preparation for spectral library generation

The spectral library was generated by IDA-MS from the unfractionated peptide samples (prepared as in section 'Sample Preparation for SWATH-MS/MS') and fractionated peptide specimens.

Fractionated peptide samples were prepared from 120 µg and 100 µg protein pellets using Multiple Affinity Removal System (MARS-14, Agilent Technologies), and following the manufacturers' protocol. The low-abundant (LAP) and the high-abundant protein fraction (HAP) were subjected to proteolytic digestion and desalting as described in the section 'Sample Preparation for SWATH-MS/MS'.

2.6 LC-MS/MS

The samples were separated with the Ekspt MicroLC 200 system (Eksigent, CA, USA) using ChromXP C18CL column (3 µm, 120 Å, 150 x 0,5 mm). The injection volume was 5µl. The mobile phases consisted of LC-MS grade formic acid (0,1%) in water [A] and acetonitrile [B]. The separation was performed by a 30 minute gradient (0-2 min 10% B; 2-23 min 10-90% B; 23-28 min 90% B, 28,1-30 min 10% B). Data acquisition was performed with a TripleTOF 5600+ mass spectrometer with a DuoSpray Ion Source

in positive ion mode (AB SCIEX, CA, USA). The microLC-MS/MS system was controlled by AB SCIEX Analyst TF 1.6 software.

- LC-MS/MS in IDA (information-dependent acquisition) mode

LC-MS/MS analyses in IDA mode were performed for the unfractionated and fractionated (LAP and HAP) samples. The TOF MS survey scan was conducted in the m/z range: 100-2000 with the accumulation time of 50 ms. Top 10 precursor ions, with the charge states from +2 to +5 were chosen for the collision-induced dissociation (CID) fragmentation. Product ion spectra were collected in the m/z range of 100-2000 with the accumulation time of 40 ms. The collision energy was automatically adjusted to the particular ion using rolling collision energy function. Precursor ions were excluded from reselection for 5 s after two occurrences. The duty cycle time was 1.11 seconds. The qualitative analyses in IDA mode were performed in three biological replicates, and three technical replicates per single biological replicate.

- LC-MS/MS in SWATH (Sequential Windowed of All Theoretical Fragment Ion Spectra) mode

The SWATH-MS analyses were performed for unfractionated protein samples. The parameters of SWATH-MS1 survey scan were as follows: high sensitivity mode, the m/z range: 100-2000, accumulation time of 50 ms. The parameters in the fragmentation mode were: m/z range 400-1000, divided into 25 windows, each 25 Da wide, accumulation time of 40 ms. The collision energy for each window was determined for a +2 to +5- ions centered upon the window with a spread of 2. The duty cycle time was 1.11 seconds. The quantitative SWATH analyses were performed in three biological replicates (each protein pool was subjected to three independent digestion and desalting procedures), and three technical replicates per each biological sample.

2.7 Data analysis

- Database search: IDA-MS

IDA MS/MS spectra were searched against SwissProt database (species: Homo sapiens, version 31.07.2017) by ProteinPilot (V4.5, SCIEX) using the Paragon algorithm. The search parameters were: TripleTOF 5600+ platform, Cys alkylation - iodoacetamide, digestion – trypsin, ID Focus: allow biological modifications, Search effort-thorough ID, Confidence>10%, automatic FDR (false discovery rate). Protein was considered a true positive if the following criteria were met: at least two peptides identified per protein with confidence >95% , and the FDR value < 0,1%.

- Data analysis: SWATH-MS

SWATH-MS data were processed against the MS/MS spectra libraries extracted from SwissProt database. The ProteinPilotgroup file was loaded into MS/MS All with SWATH Acquisition MicroApp 2.01 in PeakView 2.2 (SCIEX) to automatically generate a spectral library, according to the following parameters: maximum number of proteins-10000, modified peptides-allowed, peptides shared across proteins-not allowed, a maximum of 6 peptides per protein and 6 transitions per peptide. The MS/MS

spectra and chromatograms of the chosen ions (XIC) from SWATH-MS analyses were compared with the spectral libraries. Starting parameters for the peptides were: [Conf] $\geq$ 99, FDR < 1%, XIC width: 75 ppm, offset XIC extraction window: 10 min. Retention time calibration was performed based on 3-7 peptides, uniformly distributed according to their elution time. The peptides and transitions were manually selected for further quantitative analyses.

- Statistical analysis

The t-test analysis were performed by MarkerView software version 1.2. (SCIEX).

2.8 ELISA

The expression level of RBP4 in urine supernatant of healthy subjects, FSGS, IgAN, renal cancer and prostate cancer patients was measured using Human RBP4 ELISA kit (Fine Test, Wuhan, China) following manufacturer's instruction. The samples were diluted 3 times with Sample Dilution Buffer included in the kit. The urine creatinine levels were assessed in parallel with Creatinine (Cr) Colorimetric Assay Kit (Sarcosine oxidase method) (Elabscience Biotechnology, TX, USA) and QuantiChrom Creatinine Assay kit, DICT-500 (BioAssay Systems, CA, USA), both provided consistent results. The samples were diluted 30 and 10 times, respectively using double-distilled water. The level of uRBP4 (ng/ml) was normalized against the averaged amount of creatinine (ng/ml).

2.9 Statistical analysis of ELISA results

The box-whisker plots were done with Microsoft Excel 2013 (version 15.0.5207.1000, Microsoft Corp.). Statistical analysis was performed using the statistical package for the social scientists (SPSS version 16, IL, USA) (Pearson correlation test, nonparametric Mann-Whitney U Test with $p < 0.05$).

3 Results

3.1 Urinary protein identification with SWATH-MS

We performed SWATH-MS analysis of 5 urine sample pools, each composed of four specimens from FSGS, IgAN, ccRCC, chRCC and healthy subject group, respectively. The pooled urine samples of FSGS and IgAN were characterized by similar degree of proteinuria (moderate-level proteinuria), as assessed by dipstick test. ccRCC, chRCC and healthy subjects had normal urine parameters. The SWATH-MS performed on sample pools provided quantitation for 194 (FSGS), 179 (IgAN), 271 (ccRCC), 255 (chRCC) and 275 (healthy controls) urinary proteins (Confidence Interval (CI) > 95%, FDR < 1%) (Supplementary table 2). A total of 447 different proteins were identified across all groups of interest, from which 87 proteins were common for each group (Supplementary table 3). 38 proteins were found to be unique for FSGS, including 4 components of PPAR signaling pathway which is implicated in this condition (Supplementary table 3). 23 molecules were found to be exclusively secreted into the urine of IgAN (13% of the total protein content), 37 proteins were specific for chRCC (14.5% of the total protein content), 33

for ccRCC (11% of the total protein content) and 33 for healthy controls (12% of the total protein content) (Supplementary table 3). FSGS shared more proteins with IgAN (66%), and healthy controls (66%), than with chRCC (48%), and ccRCC (44%). The comparative, quantitative analysis of the overlapping proteomes between FSGS and the reference groups revealed that Retinol-binding protein 4 is the only molecule that clearly distinguishes FSGS from the rest of the groups ($p < 0.05$). uRBP4 protein, the carrier of vitamin A showed the highest, significant fold-change between FSGS vs healthy controls (FCH = 9.85, $p = 0.004$), and between FSGS vs IgAN (FCH = 4.8, $p = 0.012$).

3.2 Urbp4-elisa

For validation, urinary levels of RBP4 were assessed with ELISA for FSGS, IgAN, ccRCC, chRCC, prostate cancer and healthy control group. The amount of uRBP4 was found to be significantly increased in FSGS in relation to IgAN ($p = 0.0244$), ccRCC ($p = 0.0042$), chRCC ($p = 0.0128$), prostate cancer ($p = 0.02$) and healthy subjects ($p < 0.00001$) (Fig. 1). The levels of uRBP4 also differed significantly between IgAN ($p = 0.0076$), ccRCC ($p = 0.015$), and chRCC ($p = 0.007$) versus healthy controls (Fig. 1). Specific concentration cut-off value of uRBP4 i.e. ≥ 200 ng/ml allowed unequivocal discrimination of renal pathologies (41% of FSGS cases, 26% of IgAN cases, 13% ccRCC cases and 14% of chRCC cases) from prostate cancer and the normal status.

Number of the asterisks indicates the level of significance (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$)

There was no significant relationship between the level of Retinol-binding protein 4 and the degree of proteinuria and hematuria in chRCC, ccRCC, FSGS and IgAN patients (Pearson Correlation Coefficient $p < 0.05$). After stratification by gender it appeared that concentration ranges of uRBP4 in glomerulopathic, renal cancer and healthy individuals are lower in women than in men (Fig. 2). When only men were considered, significant differences in uRBP4 level were noted between FSGS versus healthy controls, ccRCC, chRCC, and prostate cancer patients (but not vs IgAN). In case of women, uRBP4 distribution was significantly different between healthy controls vs IgAN, ccRCC and chRCC. Because this study included only 4 FSGS females, this subgroup was omitted in the statistical analysis.

*SD-standard deviation of the mean.

4 Discussion

This study was initially aimed to identify urinary proteins which expression level could help distinguish FSGS patients from healthy individuals, IgAN (the second most common glomerulopathy), and renal cancer patients. Comparative analysis of the SWATH-MS protein expression profiles revealed that urinary Retinol-binding protein 4 (uRBP4), a carrier of retinol shows the highest fold change between FSGS and the rest of the groups. RBP4-ELISA performed on urine supernatant specimens from FSGS, IgAN, renal cancer, prostate cancer patients, and healthy controls (validation set) confirmed upregulation of uRBP4 in FSGS. However, it also revealed that FSGS diagnosis cannot be clearly stated based on uRBP4

concentration alone. Interestingly, it appeared that an ongoing renal disease, both glomerular disease and renal cancer can be 'confirmed' using specific cut-off value of uRBP4 concentration i.e. ≥ 200 ng/ml.

RBP4 is a small 21 kDa protein belonging to lipocalins, a family of proteins that facilitate the transport of small hydrophobic compounds. RBP4 is predominantly synthesized in the liver, and in lesser amount in adipose tissue (20–40%) and immune cells [9, 10]. This abundant, plasma protein is mainly responsible for the redistribution of retinol (vitamin A alcohol) from the liver to peripheral tissues [11]. Possibly it also binds and transports fatty acids [12]. In circulation $\approx 86\%$ of RBP4 remains complexed with transthyretin (TTR) as a 76 kDa unit (retinol transport complex), what prevents its leakage through the glomerular filtration barrier, and assures maintaining stable levels of retinol in the plasma [13]. The remaining 14% of the uncomplexed RBP4 is being freely filtered through the renal glomerulus, and contributes approximately 8.5% of the proteins in the glomerular ultrafiltrate (on a molar basis) [14]. A number of studies performed in recent two decades provided evidence that uRBP4 is probably the most sensitive functional marker of the proximal renal tubule, which can detect a minor tubular damage even earlier than KIM-1 'biomarker' [15,16,17]. Interestingly, the increase in amount of uRBP4 precedes the development of albuminuria, what indicates that uRBP4 could be potentially implemented into diagnostics and serve as a biomarker of the earliest, asymptomatic stages of renal damage. Urinary levels of RBP4 increases (even up to 10^4 -fold) in many conditions that affect kidneys e.g. renal glomerular diseases, allograft rejection, acute kidney injury (AKI), lupus nephritis, Fanconi syndrome and diabetes [16,18,19,20]. Elevated levels of urinary RBP4 are also observed in tuberculosis, head and neck cancers and bladder cancer [21,22,23]. It has not been yet established, whether particular concentration ranges of urinary RBP4 allow identification of specific renal disorders, or help discriminate individuals with a renal/non-renal pathologies among healthy subjects.

Our study is the first report of the expression distribution of Retinol-binding protein 4 in urine samples of FSGS, ccRCC and chRCC patients, providing novel insights into the diagnostic potential of uRBP4 in renal conditions. Here we provide evidence that specific cut-off point of uRBP4 concentration (≥ 200 ng/ml) could be applied to detect individuals at high risk of kidney disease diagnosis (i.e. glomerulopathy, renal cancer), who should be provided with a detailed examination. We cannot exclude that the cut-off value could be lower, as the healthy controls included in our study have not undergone diagnostics for urologic diseases. The level of uRBP exceeded 100 ng/ml in only 4 from 30 healthy subjects, and in 30% individuals the protein was not or almost not detectable.

Interestingly, in our parallel project, in which we had profiled the proteomic content of urinary exosomes from FSGS, IgAN, renal cancer patients and healthy controls, no significant differences in RBP4 concentration have been observed between the groups, in fact there was a large variation in RBP4 concentration between samples from the same cohort (data not shown).

Hopefully, assessing the changes in the level of urinary RBP4 will become a detection mode of the earliest FSGS and IgAN cases (normal creatinine, nonproteinuric subjects), and the asymptomatic renal cancers (no physical signs, normal urine parameters), at the highly curable stage.

This study has several limitations i.e. no inclusion of other glomerular diseases (i.e. membranous nephropathy, minimal change disease), large variation in the degree of proteinuria and hematuria among FSGS and IgAN patients (patients included in ELISA validation study), small sample size, lack of validation of proteins found specifically in the urine of FSGS.

5 Conclusions

This study demonstrated that the amount of urinary Retinol-binding protein 4 is significantly raised in FSGS in comparison to IgAN, ccRCC, chRCC, prostate cancer and healthy status. Although, it shown that the expression profile of uRBP4 couldn't make clear distinction between FSGS and the rest of diseases and the normal status, it revealed that a particular concentration cut-off value of uRBP4 could be useful in 'recognizing' subjects with renal pathologies (in general) among healthy individuals and prostate cancer patients. Two questions need to be addressed in prospective, validation studies: 1. Does urinary uRBP4 can be used to predict the ongoing renal disorder (in general)? 2. Which group of renal disorders can be accurately recognized by assessing the expression status of uRBP4? Further, multicenter validation studies, involving broad spectrum of patients (all types of glomerulopathies, urological cancer patients, non-urological cancer patients etc.) need to be performed to ultimately prove, whether uRBP4 possesses a diagnostic value.

Abbreviations

uRBP4

urinary retinol binding protein 4

FSGS

focal segmental glomerulosclerosis

IgAN

IgA nephropathy

ccRCC

clear cell renal cell carcinoma

chRCC

chromophobe renal cell carcinoma

SWATH-MS

sequential window acquisition of all theoretical fragment ion spectra mass spectrometry

PG

primary glomerulopathy

CKD

chronic kidney disease

DTT

dithiotreitol

IDA

information dependent acquisition

LAP

low-abundant fraction

HAP

high-abundant fraction

CID

collision-induced dissociation

FDR

false discovery rate

CI

confidence interval

FCH

fold change

TTR

transthyretin

AKI

acute kidney injury

Declarations

Ethics approval and consent to participate: This study was approved by the Bioethics Committee of Regional Specialized Hospital in Wrocław (approval ID: KB/nr 9/rok 2016). All samples were collected at Regional Specialized Hospital after obtaining written informed consent from study participants.

Consent for publication: Not applicable

Availability of data and materials: The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests: The authors declare that they have no competing interest

Funding: This work is a result of research supported in part by the Leading National Research Centre (KNOW) for years 2014-2018 (Project number U17/1/4/2016), and by National Science Center, Sonata 12 (Project number UMO-2016/23/D/NZ5/01448). The funders did not influence the study design, data generation and analysis, or manuscript preparation.

Authors' contributions: KMB prepared samples for MS, performed ELISA experiments, analyzed and interpreted the experimental data, and was a major contributor in writing the manuscript; AK and KR collected the biological material and took part in manuscript preparation, KM and PC conducted SWATH-MS analysis and performed proteomic data analysis, ACB, PK and WW collected the biological material. All authors have read and approved the manuscript.

References

1. Stratta P, Canavese C, Marengo M, Mesiano P, Besso L, Quaglia M, et al. Risk management of renal biopsy: 1387 cases over 30 years in a single centre. *Eur J Clin Invest*. 2007;37:954–63.
2. McGrogan A, Franssen CF, de Vries CS. The incidence of primary glomerulonephritis worldwide: a systematic review of the literature. *Nephrol Dial Transplant*. 2011;26(2):414–30.
3. Good DM, Zürgbilg P, Argilés A, Bauer HW, Behrens G, Coon JJ, et al. Naturally occurring human urinary peptides for use in diagnosis of chronic kidney disease. *Mol Cell Proteomics*. 2010;9(11):2424–37.
4. Bidin MZ, Shah AM, Stanslas J, Seong CLT. Blood and urine biomarkers in chronic kidney disease: An update. *Clin Chim Acta*. 2019;495:239–50.
5. Goligorsky MS, Addabbo F, O'Riordan E. Diagnostic potential of urine proteome: a broken mirror of renal diseases. *J Am Soc Nephrol*. 2007;18(8):2233–9.
6. Nafar M, Kalantari S, Samavat S, Rezaei-Tavirani M, Rutishuser D, Zubarev RA. The novel diagnostic biomarkers for focal segmental glomerulosclerosis. *Int J Nephrol*. 2014;2014:574261.
7. Wang Y, Zheng C, Wang X, Zuo K, Liu Z. Proteomic profilebased screening of potential protein biomarkers in the urine of patients with nephrotic syndrome. *Mol Med Rep*. 2017;16(5):6276–84.
8. Choi YW, Kim YG, Song MY, Moon JY, Jeong KH, Lee TW, et al. Potential urine proteomics biomarkers for primary nephrotic syndrome. *Clin Proteomics*. 2017;16(14):18.
9. Flower DR. The lipocalin protein family: structure and function. *Biochem J*. 1996;318:1–14.
10. Tsutsumi C, Okuno M, Tannous L, Piantadosi R, Allan M, Goodman DS, et al. Retinoids and retinoid-binding protein expression in rat adipocytes. *J Biol Chem*. 1992;267(3):1805–10.
11. Cho YM, Youn BS, Lee H, Lee N, Min SS, Kwak SH, et al. Plasma retinol-binding protein-4 concentrations are elevated in human subjects with impaired glucose tolerance and type 2 diabetes. *Diab Care*. 2006;29(11):2457–61.
12. Perduca M, Nicolis S, Mannucci B, Galliano M, Monaco HL. High resolution crystal structure data of human plasma retinol-binding protein (RBP4) bound to retinol and fatty acids. *Data Brief*. 2018;18:1073–81.
13. Monaco HL, Rizzi M, Coda A. Structure of a complex of two plasma proteins: transthyretin and retinol-binding protein. *Science*. 1995;268(5213):1039–41.
14. Norden AGW, Burling KA, Zeni L, Unwin RJ. A New Estimate of the Glomerular Sieving Coefficient for Retinol-Binding Protein 4 Suggests It Is Not Freely Filtered. *Kidney Int Rep*. 2019;4(7):1017–18.
15. Domingos MA, Moreira SR, Gomez L, Goulart A, Lotufo PA, Benseñor I, et al. Urinary Retinol-Binding Protein: Relationship to Renal Function and Cardiovascular Risk Factors in Chronic Kidney Disease. *PLoS One*. 2016;11(9):e0162782.
16. Gonzalez-Calero L, Martin-Lorenzo M, Ramos-Barron A, Ruiz-Criado J, Maroto AS, Ortiz A, et al. Urinary Kininogen-1 and Retinol binding protein-4 respond to Acute Kidney Injury: predictors of

- patient prognosis? Sci Rep. 2016;21(6):19667.
17. Musial K, Zwolińska D. Is urinary kim-1 the best marker of tubular damage in children with chronic kidney disease ? Nephrol Dial Transplant. 2018;33(1):i302.
 18. Turnier JL, Brunner HI, Bennett M, Aleed A, Gulati G, Haffey WD, et al. Discovery of SERPINA3 as a candidate urinary biomarker of lupus nephritis activity. Rheumatol. 2019;58(2):321–30.
 19. Park SE, Lee NS, Park JW, Rhee EJ, Lee WY, Oh KW, et al. Association of urinary RBP4 with insulin resistance, inflammation, and microalbuminuria. Eur J Endocrinol. 2014;171(4):443–49.
 20. Norden AG, Lapsley M, Unwin RJ. Urine retinol-binding protein 4: a functional biomarker of the proximal renal tubule. Adv Clin Chem. 2014;63:85–122.
 21. Wong SY, Cabodi M, Rolland J, Klapperich CM. Evaporative concentration on a paper-based device to concentrate analytes in a biological fluid. Anal Chem. 2014;86:11981–85.
 22. Ferrari E, Wittig A, Basilico F, Rossi R, De Palma A, Di Silvestre D, et al. Urinary Proteomics Profiles Are Useful for Detection of Cancer Biomarkers and Changes Induced by Therapeutic Procedures. Molecules. 2019;24(4):794.
 23. Li F, Yu Z, Chen P, Lin G, Li T, Hou L, et al. The increased excretion of urinary orosomucoid 1 as a useful biomarker for bladder cancer. Am J Cancer Res. 2016;6(2):331–40.
 24. **Additional files.**
 25. Supplementary Table 1. 'The summary of urine dipstick test results'; Supplementary Table 2: 'The list of proteins identified in each sample pool'; Supplementary Table 3: 'Proteins common for FSGS, IgAN, ccRCC, chRCC and healthy controls' and 'Proteins found to be exclusively secreted into the urine of FSGS, IgAN, ccRCC, chRCC, and healthy controls'.

Figures

Figure 1 Concentration distribution of RBP4 in urine supernatant samples (Men+Women).

Number of the asterisks indicates the level of significance(*p<0.05, **p<0.01, ***p<0.001)

Figure 1
Concentration distribution of RBP4 in urine supernatant samples (men+women).

Figure 2 Concentration distribution of urinary RBP4 analysed separately for men (A) and women (B).

*SD - standard deviation of the mean

Figure 2
Concentration distribution of urinary RBP4 analysed separately for men and women.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryTable1.xls](#)
- [SupplementaryTable2.xls](#)
- [SupplementaryTable3.xls](#)