Muscarinic Acetylcholine Receptor M5 Involved in Spermatogenesis through the Modification of Cell-Cell Junctions

Yong Zhao (✉ yzhao818@hotmail.com)

Qingdao Agricultural University

Xiao Han

QAU

Pengfei Zhang

QAU

Wei Shen

QAU

Hongfu Zhang

CAAS

Cong Zhang

QAU

Bohui Xiong

QAU

Shen Yin

QAU

Zhaojia Ge

QAU

Research

Keywords: Muscarinic acetylcholine receptor M5, Spermatogenesis, BTB, ES, Phldb2

Posted Date: June 23rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-36073/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Muscarinic acetylcholine receptor (mAChR) antagonists have been reported to decrease male fertility, however the roles of mAChRs in spermatogenesis and the underlying mechanisms are not understood. During spermatogenesis, extensive remodeling between Sertoli cells and/or germ cells interfaces takes place to accommodate the transport of developing germ cells across the blood-testis barrier (BTB) and adluminal compartment. The cell-cell junctions play vital role in the spermatogenesis process.

Methods

This study used ICR male mice and spermatogonial cells (C18-4) as well as Sertoli cells (TM-4). shRNA of control or M5 gene was injected into 5-week-age ICR mice testes. Ten days post viral grafting, mice were deeply anesthetized with pentobarbital and the testes were collected. One testicle was fresh frozen in liquid nitrogen then total RNA was extracted for RNA-seq analysis or total protein was isolated for Western blotting (WB). The second testicle was fixed in 4% paraformaldehyde (PFA) for immunofluorescence staining (IHF). C18-4 or TM-4 cells were treated with shRNA of control or M5 gene. Then the cells were collected for RNA-seq analysis, WB, or IHF.

Results

Knockdown of mAChR M5 disrupted mouse spermatogenesis and damaged the actin-based cytoskeleton and many types of junction proteins in both Sertoli cells and germ cells. M5 knockdown decreased Phldb2 expression in both germ cells and Sertoli cells which suggested that Phldb2 may regulate the cytoskeleton and other junctional proteins to control blood-testis barrier (BTB) and ectoplasmic specializations (ES) formation to regulate spermatogenesis.

Conclusion

Recent epidemiological studies suggest 20–35% of couples are infertile, and many of them have idiopathic failed gametogenesis. Moreover, the human spermatozoa quality has been declined significantly. Environmental contaminants have been found to play a major role in these phenomena because they disrupt the cell-cell junctions in testes to diminish spermatogenesis. Our investigation has elucidated a novel role for mAChR M5 in the regulation of spermatogenesis through Phldb2 regulation of the BTB and ES. M5 may be an attractive future therapeutic target in the treatment of male reproductive disorders.

Background

Both the superior (from the coeliac and aortic plexuses) and inferior (from the inferior mesenteric and hypogastric plexuses) spermatic nerves innervate the testes and are involved in fine-tuning the regulation of testicular functions (Gerendai 2004; Gerendai and Halász 1997). Other studies have shown that chronic testicular denervation in mature rats leads to the disruption of male germ cell development (Chow et al., 2000), thus emphasizing the vital role of the nervous system in spermatogenesis. Moreover, cholinergic fibers are known to regulate steroidogenesis in the testis (Zhu et al., 2002).

Cellular actions of the ancient signaling molecule acetylcholine (ACh) are mediated by two types of membrane receptors: nicotinic receptors and muscarinic ACh receptors (mAChRs; Eglen 2006; Wessler et al., 1998). These receptors are widely present in the central and peripheral nervous systems (Eglen 2006; Wessler et al., 1998). mAChRs are members of the G protein-coupled receptor (GPCR) family and include five subtypes (M1–M5) that are encoded by five different genes (Eglen 2006).

M1, M3, and M5 mAChRs mainly couple with $G_{q/11}$ to activate the phosphoinositide-specific phospholipase C β (PLC β) to produce inositol 1,4,5-triphosphate (IP3) and 1,2-diacylglycerol (DAG), followed by the elevation of intracellular Ca^{2+} and the activity of protein kinase C (PKC). M2 and M4 mainly couple with G_i/o to produce protein-dependent signaling. M1–M4 have been studied for a long time, however, M5 is the least investigated mAChR subtype and was the last one to be cloned (Bender et al., 2019; Vuckovic et al., 2019). However, mAChR M5 appears to have many important roles which makes it an attractive therapeutic target (Vuckovic et al., 2019; Fujii et al., 2017; Berizzi et al., 2016). mAChRs are present in the male reproductive system; in particular, M1–M4 are found in efferent ductules, epididymides, vas deferens, seminal vesicles, and the prostate (Lucas et al., 2008). The mRNAs of M1–M5 mAChRs have been identified in rat testis Sertoli cells (Borges et al., 2001). mAChR antagonists are reported to disrupt male fertility (defects in efferent ductules, epididymides, vas deferens, and seminal vesicles) which indicates that mAChRs play vital roles in male reproduction (Lucas et al., 2008). Meanwhile the roles of mAChRs in spermatogenesis and the underlying mechanisms are not understood.

Spermatogenesis is a complex process that involves meiosis and differentiation of germ cells from spermatogonia to spermatocytes, then on to spermatids (Sohni et al., 2019; Wang et al., 2018; Zhou et al., 2016). During spermatogenesis, the developing germ cells migrate progressively from the basal to the adluminal compartment with extensive junction restructuring between germ cell-Sertoli cell or Sertoli cell-Sertoli cell of the seminiferous epithelium. There are three major types of junctions found in the testis between germ cells and Sertoli cells, and among Sertoli cells: anchoring junctions (adherens junctions; AJs), tight junctions (TJs), and gap junctions (GJs) (Jegou 1993; Skinner 1991). It has been reported that Sertoli and germ cells develop an intimate and elaborate bidirectional trafficking system to communicate together through paracrine factors and signaling molecules that have been reported to play important roles in the regulation of cell-cell junction restructuring. Many investigations found that the following play vital roles in the cell-cell junctions: proteins like occludin, zonula occludens-1, claudin-11, and filamin A (Su et al., 2012; Chung et al., 1999); proteases and protease inhibitors such as cathepsin L, tryptase, cystatin C, and α 2-macroglobulin (Wong et al., 2000; Mruk et al., 1997); cytokines such as transforming growth factor (TGF) β 2 and TGF β 3; kinases and phosphatases such as myotubularin (Li et al., 2001);

small GTPases, such as Cdc42, N-Ras, Rac2, and RhoB (Lui et al., 2003a,b); transcriptional regulation (Lie et al., 2013; Lui et al., 2012); signaling pathways such as the nitric oxide synthase (NOS)-cGMP-protein kinase G (PRKG)-b-catenin (CATNB) pathway (Lee et al., 2005), FAK pathway (Lie et al., 2013), and many other pathways (Li et al., 2018a,b; Wen et al., 2018a; Lie et al., 2013).

Although it is well known that during the development of spermatogonial cells into spermatids the destructuring and restructuring of cell-cell junctions takes place, the intriguing cross-talk mechanisms of regulation of this restructuring are not yet fully understood. In the current study, we report that gene knockdown of M5 disrupted mouse spermatogenesis and damaged the actin-based cytoskeleton and many types of junction proteins in both Sertoli cells (TM4) and germ cells (C18-4); however, knockdown of M1 or M3 had little effect on spermatogenesis even though there are many similarities between mAChRs M1, M3, and M5. Phldb2 (pleckstrin homology-like domain, family B, member 2, alternatively called LL5 β) not only plays important roles in acetylcholine receptor (AChR) aggregation in the postsynaptic membrane, it is also involved in cell adhesion formation and extracellular matrix formation (Xie et al., 2019; Lim et al., 2016; Stehbens et al., 2014). Knockdown of M5 decreased PHLDB2 expression in both germ cells and Sertoli cells. Phldb2 may regulate cytoskeleton (actin) and other junctional proteins to control both BTB and ES formation to regulate spermatogenesis. Our data show that M5 is important for spermatogenesis, and the interactions of M5 and PHLDB2 in the regulation of the BTB and ES may contribute to this process.

Materials And Methods

Mice. All procedures involving live mice were performed in accordance with the NIH Guide for the Care and Use of Laboratory Animals and the protocols approved by the Qingdao Agricultural University Animal Care and Use Committee. ICR mice were used in this investigation.

Production of lentivirus. Lentivirus production was performed as described previously (Shen et al., 2019). Lenti-*shM5*, *shM1*, *shM3*, and *shNC* were cloned using the lentivirus-*shNC* vector as a backbone (Supplementary Fig. 1a). There were three knockdown shRNAs at three different positions for each gene. The sequences for each gene (M1, M3, and M5) and NC are listed below:

shNC (5'-3'): TTCTCCGAACGTGTCACGT

shM1 #1 (5'-3'): GCATTCATCGGGATCACCACA

shM1 #2 (5'-3'): GGCCTACAGCTGGAAAGAAGA

shM1 #3 (5'-3'): GGACACCATATAACATCATGG

shM3 #1 (5'-3'): GCAACATCCTTGTCATTGTGG

shM 3 #2 (5'-3'): GCAGTGACAGTTGGAATAACA

shM 3 #3 (5'-3'): GGCCCAGAAGAGTATGGATGA

shM 5 #1 (5'-3'): GGAGTCTTATCACAATGAAAC

shM 5 #2 (5'-3'): GGACTCCTTATAACATCATGG

shM 5 #3 (5'-3'): GGACCCAGGAGACAAACAATG

The efficiency and specificity of shRNA knockdown were determined by transfecting into 293T cells using Lipofectamine 2000 (Invitrogen, Waltham, MA, USA; #11668-027), followed by analysis at 60 h post-transfection by qPCR. Lentivirus production was then performed as shown in supplementary Fig. 1b. Briefly, the lentiviral DNA was cotransfected with packaging plasmid pG-P1-VSVG, pG-P2-REV, and pG-P3-RRE HEK293T cells using RNAi-mate (GenePharma, Shanghai, China). The medium containing lentivirus was collected at 72 h post-transfection, pooled, filtered through a 0.2- μ m filter, and concentrated using an ultracentrifuge at 19 k rpm for 2 h (4 °C). The virus was washed once and then resuspended in PBS. Approximately 10^9 infectious viral particles/ml were obtained.

In vivo virus grafting and sample collection. In vivo virus grafting was performed as previously described (Shen et al., 2019). Briefly, 5-week-old ICR male mice were anesthetized with isoflurane. Microinjections were performed using 26-gauge needles connected to a 100 μ L syringe. Virus (3 μ L with titer greater than 3×10^8 /ml) for each position and for each shRNA (in total 9 μ L with a titer $> 9 \times 10^8$ /ml for shNC, or shM1, or shM3, or shM5 individually) were mixed and then injected into the testes. Ten days post viral grafting, mice were deeply anesthetized with pentobarbital and the testes were collected. One testicle was fresh frozen in liquid nitrogen then total RNA was extracted for RNA-seq analysis or total protein was isolated for Western blotting (WB). The second testicle was fixed in 4% paraformaldehyde (PFA). Subsequently, the tissues were processed to be embedded in paraffin wax for immunochemical analyses as reported in our previous articles (Zhao et al., 2020a).

Cell culture, transfection, and growth on cover slips. The TM4 cell line (mouse Sertoli cells; purchased from ATCC) was cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (FBS; Gibco; Thermo Fisher Scientific, Inc.) at 37 °C in 5% CO₂ (Hofmann et al., 2005). The C18-4 cell line (mouse spermatogonia stem cells; Donated by Dr. Wenxian Zeng, Northwest A&F University) was held in DMEM/F12 (Gibco) supplemented with 10% (FBS), 2 mM L-glutamine (Invitrogen), and 100 U/ml penicillin and streptomycin (Invitrogen) (He et al., 2009; Hofmann et al., 2005; Li et al., 2018c). The cells were transfected with shRNA in 6-well plates. Similarly, three respective shRNAs for each gene were mixed together (titer $> 3 \times 10^8$ /ml) with RNAi-mate for the transfection for both C18-4 and TM4 cells. The transfection medium was changed after 12 h. Stable transfected cells were cultured in a similar manner to the non-transfected cells in their respective media. The transfected cells were plated on the cover slips in a 6-well plate for 2 d, after which the cover slips with the cells were collected and fixed in 4% PFA for immunofluorescence staining.

RNA Isolation and RNA-seq analyses as reported in our earlier article (Zhang et al., 2016). Briefly, total RNA was isolated using TRIzol Reagent (Invitrogen) and purified using a Pure-Link1 RNA Mini Kit (Cat: 12183018A; Life Technologies) following the manufacturers' protocol. Total RNA samples were first treated with DNase I to degrade any possible DNA contamination. Then the mRNA was enriched using oligo(dT) magnetic beads. Mixed with the fragmentation buffer, the mRNA was broken into short fragments (about 200 bp), after which, the first strand of cDNA was synthesized using a random hexamer-primer. Buffer, dNTPs, RNase H, and DNA polymerase I were added to synthesize the second strand. The double strand cDNA was purified with magnetic beads. Subsequently, 3'-end single nucleotide A (adenine) addition was performed. Finally, sequencing adaptors were ligated to the fragments. The fragments were enriched by PCR amplification. During the QC step, an Agilent 2100 Bioanalyzer and ABI StepOnePlus Real-Time PCR System were used to qualify and quantify the sample library. The library products were prepared for sequencing in an Illumina HiSeqTM 2500. The reads were mapped to reference genes using SOAPaligner (v. 2.20) with a maximum of two nucleotide mismatches allowed at the parameters of "-m 0 -x 1000 -s 40 -l 35 -v 3 -r 2". The read number of each gene was transformed into RPKM (reads per kilo bases per million reads), and then differentially expressed genes were identified using the DEGseq package and the MARS (MA-plot-based method with random sampling model) method. The threshold was set as $FDR \leq 0.001$ and an absolute value of $\log_2$ ratio ≥ 1 to judge the significance of the difference in gene expression. Then on the data were analyzed by GO enrichment, KEGG enrichment or Metascape (<http://metascape.org/gp/index.html#/main/step1>).

Western blotting. Western blotting analysis of proteins was carried out as previously reported (Zhang et al., 2016; Zhao et al., 2020b). Briefly, testicular tissue samples were lysed in RIPA buffer containing the protease inhibitor cocktail from Sangong Biotech, Ltd. (Shanghai, China). Protein concentration was determined using a BCA kit (Beyotime Institute of Biotechnology, Shanghai, China). Goat anti-glyceraldehyde 3-phosphate dehydrogenase (GAPDH; Cat #: sc-48166, Santa Cruz Biotechnology, Inc., Dallas, Texas, USA) was used as a loading control. The remaining primary antibodies (Abs) were purchased from Abcam or Beijing Biosynthesis Biotechnology CO., LTD, (Beijing, China; Supplementary Table 1). Secondary donkey anti-goat Ab (Cat no.: A0181) was purchased from Beyotime Institute of Biotechnology, and goat anti-rabbit (Cat no.: A24531) Abs were bought from Novex[®] by Life Technologies (USA). Fifty micrograms of total protein per sample were loaded onto 10% SDS polyacrylamide electrophoresis gels. The gels were transferred to a polyvinylidene fluoride (PVDF) membrane at 300 mA for 2.5 h at 4 °C. The membranes were then blocked with 5% bovine serum albumin (BSA) for 1 h at RT, followed by three washes with 0.1% Tween-20 in TBS (TBST). The membranes were incubated with primary Abs diluted with 1:500 in TBST with 1% BSA overnight at 4 °C. After three washes with TBST, the blots were incubated with the HRP-labelled secondary goat anti-rabbit or donkey anti-goat Ab respectively for 1 h at RT. After three washes, the blots were imaged. The bands were quantified using Image-J software. The intensity of the specific protein band was normalized to actin first, then the data were normalized to the control. The experiment was repeated > 6 times.

Detection of protein levels and location in testis using immunofluorescence staining. The methodology for immunofluorescence staining of testicular samples is reported in our recent publications (Zhang et al., 2016; Zhao et al., 2020b). Sections of testicular tissue (5 µm) were prepared and subjected to antigen retrieval and immunostaining as previously described (Wang et al., 2016). Briefly, sections were first blocked with normal goat serum in PBS, followed by incubation with primary Abs (Supplementary Table 1; 1:100 in PBS-0.5% Triton X-100; Bioss Co. Ltd. Beijing, PR China) at 4 °C overnight. After a brief wash, sections were incubated with an Alexa 546-labeled goat anti-rabbit secondary Ab (1:100 in PBS; Molecular Probes, Eugene, OR, USA) at RT for 30 min and then counterstained with 4',6-diamidino-2-phenylindole (DAPI). The stained sections were examined using a Leica Laser Scanning Confocal Microscope (LEICA TCS SP5 II, Germany). Ten animal samples from each treatment group were analysed. Positively stained cells were counted. A minimum of 1000 cells were counted for each sample of each experiment. The data were then normalized to the control.

Immunofluorescence staining for cells on cover slips. The cells on cover slips were fixed with 4% PFA overnight (Zhao et al., 2020b). Subsequently, the cells were treated with 2% Triton X-100 in PBS for 30 min. After three washes with PBS, the cells were blocked with normal goat serum in PBS, followed by incubation with primary Abs (Supplementary Table 1) at 4 °C overnight. After a brief wash, sections were incubated with an Alexa 546-labeled goat anti-rabbit secondary Ab (1:100 in PBS) at RT for 30 min and then counterstained with DAPI. For actin formation, the cells were incubated with Phalloidin-TRITC for 1 h at RT, then counterstained with DAPI. The stained sections were examined using a Leica Laser Scanning Confocal Microscope (LEICA TCS SP5 II, Germany). The experiments were repeated at least six times.

Statistical analysis. Data were statistically analysed using the ANOVA function in SPSS statistical software (IBM Co., NY). Comparisons between groups were tested by One-Way ANOVA analysis and the LSD test. All groups were compared with each other for every parameter (mean ± SEM). Differences were considered significant at $p < 0.05$.

Data Availability

Source data associated with Fig. 3, 5, and 6 can be accessed through GEO: GSE137628 and GSE142618. All data are reported in the main text and supplementary materials and are available from the corresponding author upon request.

Results

Time dependent expression of M5, M1, and M3 mAChRs in mouse testes. The expression of M5, M1, and M3 mAChRs have been detected in mouse testes using specific antibodies (Fig. 1). M5 and M1 receptor expression levels were increased in murine testes between 1 and 6 wk of age. However, M3 receptor expression was low in 1 wk old mouse testes while it was elevated at a consistent level between 3 and 6 wk of age. These three receptors are expressed in Sertoli cells and germ cells. M5 and M1 receptors are

mainly expressed in the perinuclei region, while M3 is mainly expressed in the nuclei, which is constant with the findings of Lucas et al (2008). Moreover, M1 is mainly present in spermatids of 6 wk old mice, while M5 and M3 are present mainly in the early stages of germ cells (spermatogonia or spermatocytes; Fig. 1).

M5 knockdown disrupted mouse spermatogenesis, while M1 and M3 knockdown did not. In order to differentiate the function of M5 from M1 or M3, as they all couple to $G_{q/11}$, the expression of these receptors was modified in mouse testes during the current investigation. After M5 knockdown using short hairpin RNA (shRNA), the protein level of M5 receptor was decreased (Fig. 2a, b; Supplementary Fig. 1); the gene expression profile of mouse testes was then determined by RNA-seq analysis. Compared to the control (shNC), 661 genes were increased while 622 genes were decreased in shM5 mouse testis samples (Fig. 2c). Principal Component Analysis (PCA) showed that the shM5 and shNC groups were clearly separated (Fig. 2d). The genes decreased by shM5 were mainly enriched in the gamete generation, spermatogenesis, and male reproduction functional pathways in gene ontology (GO) enrichment analysis (Fig. 2e), while those genes increased by shM5 were enriched in others not related to spermatogenesis signaling pathways (Fig. 2f). In order to search more deeply for the mechanisms of shM5 knockdown disruption of spermatogenesis, Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis was performed. There were two signaling pathways “cell adhesion molecules (CAMs)” and “ECM-receptor interaction” that were significantly enriched in the decreased genes (Fig. 2g), while they were not found in the increased genes (Fig. 2h). As we know, cell-cell junctions (especially in the BTB) play vital roles in spermatogenesis and male fertility (Li et al., 2018b; Wen et al., 2018a; Lui et al., 2012; Lee et al., 2005). The data suggested that shM5 may damage cell-cell junctions in the testis to disrupt spermatogenesis.

However, shM1 or shM3 had little effect on spermatogenesis (Supplementary Fig. 2 and Supplementary Fig. 3). There were 263 genes increased and 550 genes decreased by shM1 (Supplementary Fig. 2a). Furthermore, shM1 and shNC were clearly separated by PCA analysis (Supplementary Fig. 2b). A total of 404 genes were increased and 724 genes were decreased by shM3 and PCA analysis showed that shM3 and shNC were also clearly separated (Supplementary Fig. 3a, b). GO enrichment analysis showed that there was one signaling pathway “gamete generation” enriched in shM1 decreased genes (Supplementary Fig. 2c). In addition, no pathways related to spermatogenesis or male fertility were enriched by shM3 knockdown genes (Supplementary Fig. 3c). The data here suggested that M1 or M3 may have a different role from M5 in mouse spermatogenesis.

Using GO enrichment analysis, the signaling pathways related to spermatogenesis and male reproduction for the shM5 group were separated and analyzed for their protein-protein network. The proteins in these pathways were connected together with the above mentioned signaling pathways (Fig. 3a). These proteins could be divided into three groups as shown in Fig. 3b. Furthermore, we found that the levels of proteins important to spermatogenesis (Li et al., 2018b; Wen et al., 2018a; Lee et al., 2005) (TP1, PGK2, CREM, and p-FSCN1) were decreased in shM5 mouse testis samples (Fig. 3c, d; Supplementary Fig. 4). Meanwhile, Sertoli cell marker SOX9 remained unchanged by shM5 (Fig. 3c). The data further suggested that shM5 upset spermatogenesis.

The genes (decreased by shM5) in “cell adhesion molecules” and “ECM-receptor interaction” pathways from KEGG enrichment analysis were further analyzed to determine their interactive network. These genes were connected together with the signaling pathways (Fig. 4a). Furthermore, the protein levels of many cell-cell junction proteins such as claudin, occludin, Cx43, Cx37, JAM1, ZO-1, E-cadherin, and catenin were determined in mouse testis samples. Results showed that the protein levels of occludin, E-cadherin, Cx37, and Cx43 were decreased by shM5 (Fig. 4b, c; Supplementary Fig. 4). The data here further indicated that shM5 may damage cell-cell junctions to disrupt spermatogenesis.

Knockdown M5 decreased cell-cell junction proteins in spermatogonia stem cells. To further search for the M5 mechanism during spermatogenesis, M5 expression was modified in spermatogonia stem cells (C18-4 cell line) (He et al., 2009; Hofmann et al., 2005) and Sertoli cells (TM4 cell line) (Ge et al., 2018). The protein level of M5 in C18-4 cells was decreased by shM5 (Fig. 5a) and the gene expression profile in C18-4 cells after shM5 treatment was significantly altered. An increase was seen in 239 genes while 261 genes were decreased by shM5 compared to shNC (Fig. 5b). The PCA data showed that shM5 was clearly separated from shNC (Fig. 5c). Reactome and GO enrichment analyses showed that many signaling pathways were related to the extracellular matrix and cell adhesion for the genes decreased by shM5 (Fig. 5d, e), but not for those genes that were increased (Supplementary Fig. 5a, b). Furthermore, cell cytoskeleton (actin) was detected in C18-4 cells by shM5, and it was noteworthy that actin formation was dramatically damaged by shM5 (Fig. 5f). Subsequently, the cell-cell junction proteins were determined to confirm the disruption to the cytoskeleton. All eight cell-cell junction proteins occludin, claudin, JAM1, Cx37, Cx43, E-cadherin, ZO-1, and catenin were decreased by shM5 compared to shNC (Fig. 5g; Supplementary Fig. 6). The data here suggested that M5 plays a vital role in cell-cell junction formation in germ cells.

M5 knockdown decreased cell-cell junction proteins in Sertoli cells. M5 protein levels in Sertoli cells (TM4 cell line) were decreased by shM5 (Fig. 6a). shM5 significantly changed the gene expression profile in TM4 cells. In total, 90 genes were increased and 566 genes were decreased by shM5 in TM4 cells (Fig. 6b). PCA analysis found that shNC and shM5 could be clearly separated (Fig. 6c). GO enrichment analysis found that 174 out of 566 genes (decreased genes) were enriched into “protein binding” functional signaling pathways which suggested that Sertoli cell junctions may be affected by shM5 (Fig. 6d), but not for those genes that were increased (Supplementary Fig. 5c, d). The next step was to determine the cytoskeleton (actin formation) in TM4 cells. It was found that actin formation was disrupted by shM5 in TM4 cells (Fig. 6f). At the same time, the protein levels of the eight cell junctional proteins occludin, claudin, JAM1, Cx37, Cx43, E-cadherin, ZO-1, and catenin were significantly reduced by shM5 in Sertoli cells (Fig. 6g; Supplementary Fig. 6). Furthermore, there were a few functional pathways related to Sertoli cell functions such as the MAPK signaling pathway, PI3K-AKT pathway, and mTOR pathway that have been enriched in KEGG analysis for the genes decreased by shM5 in TM4 cells (Fig. 6e) (Ni et al., 2019)³⁷. The data here suggested that shM5 upset the function of Sertoli cells, in particular the cell junctions.

Deep analysis discovered the overlap of functional pathways in both C18-4 and TM4 cells caused by shM5. Twenty-seven genes overlapped between C18-4 and TM4 cells by shM5 (Fig. 7a). Enrichment

analysis of those 27 genes showed that they play important roles in intracellular signaling transduction and the extracellular matrix (Fig. 7b). One of the genes, Phldb2, not only plays an important role in AChR aggregation in the postsynaptic membrane (Xie et al., 2019), but it is also involved in cell adhesion formation and extracellular matrix formation (Lim et al., 2016; Stehbens et al., 2014). Phldb2 interacts with CLASP and filamins during its involvement in focal adhesion and extracellular matrix turnover (Fig. 7c) (Stehbens et al., 2014; Su et al., 2012). The expression of Phldb2 was decreased in both C18-4 and TM4 cells. Furthermore, the protein levels of PHLDB2 were significantly decreased by shM5 in C18-4 cells (Fig. 7d; Supplementary Fig. 6) and TM4 cells (Fig. 7e; Supplementary Fig. 6).

Discussion

mAChRs include five subtypes (M1–M5) which are known to play various roles in the physiology and pathophysiology of different autocrine and neuronal systems. Activation of mAChRs is known to be involved in cell proliferation, differentiation, growth, and other functions in male reproductive systems. This activation could release epidermal growth factor receptor (EGFR) ligands to bind to EGFR and could activate the extracellular signal-regulated kinase (ERK)1/2 pathway (Lucas et al., 2008) to stimulate Sertoli cell proliferation. However, there are currently no reports regarding the activation of mAChR on spermatogenesis. Although there are many similarities between M1, M3, and M5, their functions in mouse testes are very different. In the current investigation we found that knockdown of M1 or M3 had little effect on spermatogenesis, whereas M5 knockdown disrupted spermatogenesis and damaged the expression of proteins (claudin, occludin, ZO-1, JAM1 Cx43, Cx37, E-cadherin, and catenin) in the BTB and ES.

Spermatogenesis is a choreographed process of diploid spermatogonia undergoing differentiation to produce haploid germ cells. During spermatogenesis, extensive remodeling at Sertoli cell-cell and Sertoli cell-germ cell interfaces takes place to accommodate the transport of developing germ cells across the BTB and adluminal compartment (Wen et al., 2018b). The BTB is made up of actin-based tight junctions (TJs) and gap junctions (GJs). The extracellular matrix (ECM) is involved in spermatogenesis through regulating the BTB since Sertoli cells are in physical contact with the basement membrane. Although the BTB plays vital roles in the control of Sertoli cell-cell interactions, ES is a unique actin-rich AJ in the testes that regulates Sertoli cell-germ cell interactions. ES includes basal ES at the Sertoli cell–cell interface of the BTB, and the apical ES at the Sertoli–spermatid interface (Li et al., 2015). Actin-based cytoskeletons in Sertoli and germ cells play crucial roles in the regulation of homeostasis of the BTB and the cytoskeletal elements at the basal ES (Wen et al., 2018c). Even though there have been many recent reports that have highlighted the factors involved in the regulation of the BTB and ES (Wen et al., 2018; Li et al., 2015), the intriguing cross-talk mechanism(s) between basal and apical ES are still not fully understood, because the restructuring of the BTB close to the basement membrane and the disruption of the apical tubulobulbar complex (TBC) at the luminal edge of the epithelium happen almost concurrently (Li et al., 2015). In the current investigation, we aimed to explore the underlying mechanisms regulating the basal and apical ES. We found that shM5 disrupted mouse spermatogenesis *in vivo* and damaged the actin-based cytoskeleton and many types of junction proteins in both Sertoli cells (TM4) and germ

cells (C18-4). Claudin and occludin are major players in Sertoli cell TJs (McCabe et al., 2016). Cadherins, AJ transmembrane proteins, are reported to be present at the basal ES (Lee et al., 2003). Catenins and ZO-1, the peripheral adaptors of basal ES and TJs, are involved in the engagement/disengagement mechanism between basal ES and TJs to make the BTB one of the “tightest” barriers in the mammalian body (Yan and Cheng 2005). Connexin 43, connexin 37, and junctional adhesion molecules (JAMs) have also been found to play important roles in cell-cell junctions (Li et al., 2016; Zhang and Lui 2015). In the current investigation, claudin, occludin, ZO-1, JAM1 Cx43, Cx37, E-cadherin, and catenin were found to be decreased by shM5 in spermatogonial cells (C18-4) and Sertoli cells (TM4). Our data suggested that M5 may be involved in ES and BTB restructuring to regulate spermatogenesis.

Phldb2 is a PH domain-containing protein that is highly sensitive to phosphatidylinositol 3,4,5-triphosphate (PIP3) as well as PIP2 (Xie et al., 2019). Phldb2 plays important roles in AChR aggregation in the postsynaptic membrane (Xie et al., 2019). Moreover, Phldb2 can associate with CLIP-associating proteins (CLASPs), Prickle 1, and Liprin α 1 to be involved in focal adhesion disassembly and cell polarization and migration (Lim et al., 2016; Astro et al., 2014; Lansbergen et al., 2006). Podosomes are actin-rich, dynamic structures capable of remodeling the extracellular matrix (ECM), and have been found in many types of cells such as osteoclasts, macrophages, and epithelial cells. Podosomes present around AChR aggregates are called “synaptic podosomes” (Proszynski and Sanes 2013). The cytoskeleton- and membrane-associated protein Phldb2 is known to be one of the key components of synaptic podosomes, and its upset perturbs AChR clustering in cultured myotubes (Kishi et al., 2005). In the current study, we found that shM5 decreased Phldb2 expression in both germ cells and Sertoli cells. Therefore, Phldb2 may regulate the cytoskeleton (actin) and other junctional proteins (claudin, occludin, Cx43, ZO-1, cadherin, and catenin) to control BTB and ES formation that in turn affect spermatogenesis. Since Phldb2 interacts with AChR, the mAChR M5 knockdown may upset Phldb2 and disrupt the BTB and ES to damage spermatogenesis.

Conclusion

In summary, our investigation has elucidated a novel role for mAChR M5 in the regulation of spermatogenesis through Phldb2 regulation of the BTB and ES. Further studies on the cross-talk between M5 and Phldb2, and among Phldb2, BTB, and ES will shed light on our understanding of the mechanisms of M5 in the regulation of the BTB/ES and spermatogenesis.

Abbreviations

Abs
antibodies
Ach
acetylcholine
AJs
adherens junctions

ATCC
American Type Culture Collection
BSA
bovine serum albumin
BTB
blood-testis barrier
CATNB
catenin
Cx37
connexin 37
Cx43
connexin 43
DAG
1,2-diacylglycerol
DAPI
4',6-diamidino-2-phenylindole
DMEM
Dulbecco's Modified Eagle Medium
ECM
extracellular matrix
EGFR
epidermal growth factor receptor
ERK
extracellular signal-regulated kinase
ES
ectoplasmic specializations
FBS
Fetal bovine serum
GAPDH
glyceraldehyde 3-phosphate dehydrogenase
GJs
gap junctions
GO
gene ontology
GPCR
G protein-coupled receptor
HRP
horse radish peroxidase
IHF
immunofluorescence staining

IP3
inositol 1,4,5-triphosphate
JAMs
junctional adhesion molecules
KEGG
Kyoto Encyclopedia of Genes and Genomes
mAChR
muscarinic acetylcholine receptor
NC
negative control
NOS
nitric oxide synthase
PCA
Principal Component Analysis
PFA
paraformaldehyde
Phldb2
pleckstrin homology-like domain, family B, member 2, also called LL5 β
PKC
protein kinase C
PLC β
phosphoinositide-specific phospholipase C β
PRKG
protein kinase G
PVDF
polyvinylidene fluoride
RT
room temperature
shRNA
short hairpin RNA
TBC
tubulobulbar complex
TBS
Trice buffer saline
TBST
TBS with Tween-20
TGF
transforming growth factor
TJs
tight junctions

WB
Western blotting
ZO-1
Zonula occludens-1

Declarations

Ethics approval and consent to participate

All procedures involving live mice were performed in accordance with the NIH Guide for the Care and Use of Laboratory Animals and the protocols approved by the Qingdao Agricultural University Animal Care and Use Committee. ICR mice were used in this investigation.

Consent for publication

None.

Availability of data and materials

Source data associated with Fig. 3, 5, and 6 can be accessed through GEO: GSE137628 and GSE142618. All data are reported in the main text and supplementary materials and are available from the corresponding author upon request.

Competing interest

The authors declare no competing interests.

Funding

This study was supported by the National Natural Science Foundation of China (31772408 to YZ; 31672428 to HZ).

Author's contribution

YZ conceived and designed the study. XH and PZ performed animal experiments and RNA-seq data analysis. CZ, and BX performed histological analyses and IHF and WB. SY and ZG performed the cell experiments. WS and HZ did the data analysis and help on the manuscript preparation. YZ prepared the manuscript with input from co-authors. All authors read and approved the final manuscript.

Acknowledgements

We thank the investigators and staff of The Beijing Genomics Institute (BGI) for technical support. We thank Dr. Wenxian Zeng from Northwest A&F University for kindly donation of C18-4 cells.

References

1. Astro V, Chiaretti S, Magistrati E, Fivaz M, de Curtis I. Liprin-alpha1, ERC1 and LL5 define polarized and dynamic structures that are implicated in cell migration. *J Cell Sci.* 2014;127:3862–76.
2. Bender AM, Garrison AT, Lindsley CW. The Muscarinic acetylcholine receptor M5: therapeutic implications and allosteric modulation. *ACS Chem Neurosci.* 2019;10:1025–34.
3. Berizzi AE, Gentry PR, Rueda P, Den Hoedt S, Sexton PM, Langmead CJ, Christopoulos A. Molecular mechanisms of action of M5 muscarinic acetylcholine receptor allosteric modulators. *Mol Pharmacol.* 2016;90:427–36.
4. Borges MOR, Abreu MLC, Avellar MCW, Porto CS. Characterization of muscarinic acetylcholine receptor in rat Sertoli cells. *Endocrinology.* 2001;142:4701–10.
5. Chow SH, Giglio W, Anesetti R, Ottenweller JE, Pogach LM, Huang HF. The effects of testicular denervation on spermatogenesis in the Sprague-Dawley rat. *Neuroendocrinology.* 2000;72:37–45.
6. Chung SSW, Lee WM, Cheng CY. A study on the formation of specialized inter-Sertoli cell junctions in vitro. *J Cell Physiol.* 1999;181:258–72.
7. Eglen RM. Muscarinic receptor subtypes in neuronal and non-neuronal cholinergic function. *Auton Autacoid Pharmacol.* 2006;26:219–33.
8. Fujii T, Mashimo M, Moriwaki Y, Misawa H, Ono S, Horiguchi K, Kawashima K. Physiological functions of the cholinergic system in immune cells. *J Pharmacol Sci.* 2017;134:1–21.
9. Ge X, Pan P, Jing J, Hu X, Chen L, Qiu X, Ma R, Jueraitetibaike K, Huang X, Yao B. Rosiglitazone ameliorates palmitic acid induced cytotoxicity in TM4 Sertoli cells. *Reprod Biol Endocrinol.* 2018;16:98.
10. Gerendai I. Supraspinal connections of the reproductive organs: structural and functional aspects. *Acta Physiol Hung.* 2004;91:1–21.
11. Gerendai I, Halász B. Neuroendocrine asymmetry. *Front Neuroendocrinol.* 1997;18:354.
12. He Z, Jiang J, Kokkinaki M, Dym M. Nodal signaling via an autocrine pathway promotes proliferation of mouse spermatogonial stem/progenitor cells through Smad2/3 and *Oct-4* activation. *Stem Cells.* 2009;27:2580–90.
13. Hofmann MC, Braydich-Stolle L, Dettin L, Johnson E, Dym M. Immortalization of mouse germ line stem cells. *Stem Cells.* 2005;23:200–10.
14. Jegou B. The Sertoli-germ cell communication network in mammals. *Int Rev Cytol.* 1993;147:25–96.
15. Kishi M, Kummer TT, Eglen SJ, Sanes JR. LL5beta: a regulator of postsynaptic differentiation identified in a screen for synaptically enriched transcripts at the neuromuscular junction. *J Cell Biol.* 2005;169:355–66.

16. Lansbergen G, Grigoriev I, Mimori-Kiyosue Y, Ohtsuka T, Higa S, Kitajima I, Demmers J, Galjart N, Houtsmuller AB, Grosveld F, et al. CLASPs attach microtubule plus ends to the cell cortex through a complex with LL5beta. *Dev Cell*. 2006;11:21–32.
17. Lee NP, Mruk DD, Wong CH, Cheng CY. Regulation of Sertoli-Germ Cell Adherens Junction Dynamics in the Testis Via the Nitric Oxide Synthase (NOS)/cGMP/Protein Kinase G (PRKG)/b-Catenin (CATNB) Signaling Pathway: An In Vitro and In Vivo Study. *Biol Reprod*. 2005;73:458–71.
18. Lee NP, Mruk D, Lee WM, Cheng CY. Is the cadherin/catenin complex a functional unit of cell-cell actin-based adherens junctions in the rat testis? *Biol Reprod*. 2003;68:489–508.
19. Li SYT, Yan M, Chen H, Jesus T, Lee WM, Xiao X, Cheng CY. mTORC1/rpS6 regulates blood-testis barrier dynamics and spermatogenetic function in the testis in vivo. *Am J Physiol Endocrinol Metab*. 2018a;314:E174–90.
20. Li XY, Zhang Y, Wang XX, Jin C, Wang YQ, Sun TC, Li J, Tang JX, Batool A, Deng SL, et al. Regulation of blood-testis barrier assembly in vivo by germ cells. *FASEB J*. 2018b;32:1653–64.
21. Li B, He X, Zhuang M, Niu B, Wu C, Mu H, Tang F, Cui Y, Liu W, Zhao B, et al. Melatonin Ameliorates Busulfan-Induced Spermatogonial Stem Cell Oxidative Apoptosis in Mouse Testes. *Antioxid Redox Signal*. 2018c;28:385–400.
22. Li N, Mruk DD, Mok KW, Li MW, Wong CK, Lee WM, Han D, Silvestrini B, Cheng CY. Connexin 43 reboots meiosis and reseals blood-testis barrier following toxicant-mediated aspermatogenesis and barrier disruption. *FASEB J*. 2016;30:1436–52.
23. Li N, Mruk DD, Wong CK, Lee WM, Han D, Cheng CY. Actin-bundling protein plastin 3 is a regulator of ectoplasmic specialization dynamics during spermatogenesis in the rat testis. *FASEB J*. 2015;29:3788–805.
24. Li JC, Mruk D, Cheng CY. The inter-Sertoli tight junction permeability barrier is regulated by the interplay of protein phosphatases and kinases: an in vitro study. *J Androl*. 2001;22:847–56.
25. Lie PP, Cheng CY, Mruk DD. Signalling pathways regulating the blood-testis barrier. *Int J Biochem Cell Biol*. 2013;45:621–5.
26. Lim BC, Matsumoto S, Yamamoto H, Mizuno H, Kikuta J, Ishii M, Kikuchi A. Prickle1 promotes focal adhesion disassembly in cooperation with the CLASP–LL5β complex in migrating cells. *J Cell Sci*. 2016;129:3115–29.
27. Lucas TFG, Siu ER, Royer C, Trindade ES, Nader HB, Lazari MFM, Porto CS. Muscarinic acetylcholine receptors: relevance to infertility and male contraception. *Immunology Endocrine Metabolic Agents in Medicinal Chemistry*. 2008;8:42–50.
28. Lui WY, Lee WM, Cheng CY. Transforming growth factor beta3 regulates the dynamics of Sertoli cell tight junctions via the p38 mitogen-activated protein kinase pathway. *Biol Reprod*. 2003a;68:1597–612.
29. Lui WY, Lee WM, Cheng CY. Minireview: Rho GTPases and spermatogenesis. *Biochim Biophys Acta*. 2003b;1593:121–9.

30. Lui WY, Cheng CY. Transcriptional regulation of cell adhesion at the blood-testis barrier and spermatogenesis in the testis. *Adv Exp Med Biol.* 2012;763:281–94.
31. McCabe MJ, Foo CF, Dinger ME, Smooker PM, Stanton PG. Claudin–11 and occludin are major contributors to Sertoli cell tight junction function, in vitro. *Asian J Androl.* 2016;18:620–6.
32. Mruk D, Zhu LJ, Silvestrini B, Lee WM, Cheng CY. Interactions of proteases and protease inhibitors in Sertoli-germ cell cocultures preceding the formation of specialized Sertoli-germ cell junctions in vitro. *J Androl.* 1997;18:612–22.
33. Ni F, Hao S, Yang W. Multiple signaling pathways in Sertoli cells: recent findings in spermatogenesis. *Cell Death Dis.* 2019;10:541.
34. Proszynski TJ, Sanes JR. Amotl2 interacts with LL5 β , localizes to podosomes and regulates postsynaptic differentiation in muscle. *J Cell Sci.* 2013;126:2225–35.
35. Shen M, Wang F, Li M, Sah N, Stockton ME, Tidei JJ, Gao Y, Korabelnikov T, Kannan S, Vevea JD, et al. Reduced mitochondrial fusion and Huntingtin levels contribute to impaired dendritic maturation and behavioral deficits in Fmr1-mutant mice. *Nat Neurosci.* 2019;22:386–400.
36. Skinner MK. Cell-cell interactions in the testis. *Endocr Rev.* 1991;12:45–77.
37. Sohni A, Tan K, Song HW, Burow D, de Rooij DG, Laurent L, Hsieh TC, Rabah R, Hammoud SS, Vicini E, et al. The neonatal and adult human testis defined at the single-cell level. *Cell Rep.* 2019;26(6):1501–17.e4.
38. Stehbens SJ, Paszek M, Pemble H, Ettinger A, Gierke S, Wittmann T. CLASPs link focal adhesion-associated microtubule capture to localized exocytosis and adhesion site turnover. *NatCell Biol.* 2014;16:561–73.
39. Su W, Mruk DD, Lie PP, Lui WY, Cheng CY. Filamin A Is a Regulator of Blood-Testis Barrier Assembly during Postnatal Development in the Rat Testis. *Endocrinology.* 2012;153:5023–35.
40. Vuckovic Z, Gentry PR, Berizzi AE, Hirata K, Varghese S, Thompson G, van der Westhuizen ET, Burger WAC, Rahmani R, Valant C, et al. Crystal structure of the M5 muscarinic acetylcholine receptor. *Proc Natl Acad Sci USA.* 2019;116:26001–7.
41. Wang M, Liu X, Chang G, Chen Y, An G, Yan L, Gao S, Xu Y, Cui Y, Dong J, et al. Single-cell RNA sequencing analysis reveals sequential cell fate transition during human spermatogenesis. *Cell Stem Cell.* 2018;23:599–614.
42. Wen Q, Tang EI, Gao Y, Jesus TT, Chu DS, Lee WM, Wong CKC, Liu YX, Xiao X, Silvestrini B, et al. Signaling pathways regulating blood–tissue barriers-Lesson from the testis. *Biochim Biophys Acta Biomembr.* 2018a;1860:141–53.
43. Wen Q, Li N, Xiao X, Lui WY, Chu DS, Wong CKC, Lian Q, Ge R, Lee WM, Silvestrini B, et al. Actin nucleator Spire 1 is a regulator of ectoplasmic specialization in the testis. *Cell Death Dis.* 2018b;9:208.
44. Wen Q, Tang EI, Li N, Mruk DD, Lee WM, Silvestrini B, Cheng CY. Regulation of Blood-Testis Barrier (BTB) Dynamics, role of actin-, and microtubule-based cytoskeletons. *Methods Mol Biol.* 2018c;1748:229–43.

45. Wessler I, Kirkpatrick CJ, Racke K. Non-neuronal acetylcholine, a locally acting molecule, widely distributed in biological systems: expression and function in humans. *Pharmacol Ther.* 1998;77:59–79.
46. Wong CC, Chung SS, Grima J, Zhu LJ, Mruk D, Lee WM, Cheng CY. Changes in the expression of junctional and nonjunctional complex component genes when inter-Sertoli tight junctions are formed in vitro. *J Androl.* 2000;m21:227–37.
47. Xie MJ, Ishikawa Y, Yagi H, Iguchi T, Oka Y, Kuroda K, Iwata K, Kiyonari H, Matsuda S, Matsuzaki H, et al. PIP3-Phldb2 is crucial for LTP regulating synaptic NMDA and AMPA receptor density and PSD95 turnover. *Sci Rep.* 2019;9:4305.
48. Yan HH, Cheng CY. Blood-testis barrier dynamics are regulated by an engagement/disengagement mechanism between tight and adherens junctions via peripheral adaptors. *Proc Natl Acad Sci USA.* 2005;102:11722–7.
49. Zhang WD, Zhao Y, Zhang HF, Wang SK, Hao ZH, Liu J, Yuan YQ, Zhang PF, Yang HD, Shen W, et al. Alteration of Gene Expression by Zinc Oxide Nanoparticles or Zinc Sulfate *In vivo*, and Comparison with *In vitro* Data—a Harmonious Case. *Theriogenology.* 2016;86:850–61.
50. Zhang X, Lui WY. Transforming growth factor- β 3 regulates cell junction restructuring via MAPK-mediated mRNA destabilization and Smad-dependent protein degradation of junctional adhesion molecule B (JAM-B). *Biochim Biophys Acta.* 2015;1849:601–11.
51. Zhao Y, Feng Y, Liu M, Chen L, Meng Q, Tang X, Wang S, Liu L, Li L, Shen W, et al. Single-cell RNA sequencing analysis reveals alginate oligosaccharides preventing chemotherapy-induced mucositis. *Mucosal Immunol.* 2020a. doi:10.1038/s41385-019-0248-z.
52. Zhao Y, Zhang P, Ge W, Feng Y, Li L, Sun Z, Zhang H, Shen W. Alginate oligosaccharides improve germ cell development and testicular microenvironment to rescue busulfan disrupted spermatogenesis. *Theranostics* 2020b; 10(7), 3308–3324.
53. Zhou Q, Wang M, Yuan Y, Wang X, Fu R, Wan H, Xie M, Liu M, Guo X, Zheng Y, et al. Complete meiosis from embryonic stem cell-derived germ cells in vitro. *Cell Stem Cell.* 2016;18(3):330–40.
54. Zhu C, Palmada MN, Aguado LI, Cavicchia JC. Administration of acetylcholine to the spermatic nerve plexus inhibits testosterone secretion in an in vitro isolated rat testis-nerve plexus system. *Int J Androl.* 2002;25:134–8.

Supplementary Figure Legends

Supplementary Fig. 1 Structure of the lentivirus vector used in this study and its production protocol for subsequent infection of cells and animals.

Supplementary Fig. 2 Quantitative data for IHF and WB of mouse testis samples. **a**, Quantitative data for IHF of mouse testis samples (for M5 detection). **b**, Quantitative data for WB of mouse testis samples (for M5 detection). (n = 6/group). Data are presented as mean $\pm$ SEM. a, b indicate a significant difference among different treatments (p < 0.05).

Supplementary Fig. 3 Gene expression in mouse testes after M1 knockdown by short hairpin RNA (shRNA) in vivo. **a**, Gene expression heatmap of mouse testis samples after M1 shRNA treatment for 10 days. **b**, PCA analysis for the gene expression of mouse testes. **c**, GO enrichment analysis of the genes decreased by shM1 treatment in mouse testes. **d**, GO enrichment analysis of the genes increased by shM1 treatment in mouse testes. **e**, KEGG enrichment analysis of the genes decreased by shM1 treatment in mouse testes. **f**, KEGG enrichment analysis of the genes increased by shM1 treatment in mouse testes.

Supplementary Fig. 4 Gene expression in mouse testes after M3 knockdown by short hairpin RNA (shRNA) in vivo. **a**, Gene expression heatmap of mouse testis samples after M3 shRNA treatment for 10 days. **b**, PCA analysis for the gene expression of mouse testes. **c**, GO enrichment analysis of the genes decreased by shM3 treatment in mouse testes. **d**, GO enrichment analysis of the genes increased by shM3 treatment in mouse testes. **e**, KEGG enrichment analysis of the genes decreased by shM3 treatment in mouse testes. **f**, KEGG enrichment analysis of the genes increased by shM3 treatment in mouse testes.

Supplementary Fig. 5 Quantitative data for WB of mouse testis samples. **a**, Quantitative data for WB of mouse testis samples for the detection of PGK2. **b**, Quantitative data for WB of mouse testis samples for the detection of TP1. **c**, Quantitative data for WB of mouse testis samples for the detection of CREM. **d**, Quantitative data for WB of mouse testis samples for the detection of p-FSCN1. **e**, Quantitative data for WB of mouse testis samples for the detection of occludin. (n = 6/group). Data are presented as mean ± SEM. **a**, **b** indicate a significant difference among different treatments (p < 0.05).

Supplementary Fig. 6 The data summary of increased gene in C18-4 and TM4 cells after shM5 treatment in vitro. **a**, Reactome enrichment analysis of the genes increased by shM5 treatment in C18-4 cells. **b**, GO enrichment analysis of the genes increased by shM5 treatment in C18-4 cells. **c**, Reactome enrichment analysis of the genes increased by shM5 treatment in TM4 cells. **d**, GO enrichment analysis of the genes increased by shM5 treatment in TM4 cells.

Supplementary Fig. 7 Quantitative data for IHF of C18-4 and TM4 cell samples. **a**, Quantitative data for IHF of C18-4 and TM4 cell samples for the detection of occludin. **b**, Quantitative data for IHF of C18-4 and TM4 cell samples for the detection of claudin. **c**, Quantitative data for IHF of C18-4 and TM4 cell samples for the detection of E-cad (E-cadherin). **d**, Quantitative data for IHF of C18-4 and TM4 cell samples for the detection of ZO-1. **e**, Quantitative data for IHF of C18-4 and TM4 cell samples for the detection of catenin. **f**, Quantitative data for IHF of C18-4 and TM4 cell samples for the detection of Cx43. **g**, Quantitative data for IHF of C18-4 and TM4 cell samples for the detection of Cx37. **h**, Quantitative data for IHF of C18-4 and TM4 cell samples for the detection of JAM1. **i**, Quantitative data for IHF of C18-4 and TM4 cell samples for the detection of Phldb2. (n = 6 times). Data are presented as mean ± SEM. **a**, **b**, **c**, **d** indicate a significant difference among different treatments (p < 0.05).

Supplementary Table 1. Primary antibodies information.

Figures

Figure 1

Protein expression of muscarinic acetylcholine receptors (mAChRs) M1, M3, and M5 in mouse testes at the age of 1, 2, 3, and 6 weeks. Scale bar: 25 μm.

Figure 2

M5 knockdown by short hairpin RNA (shRNA) in mouse testes disrupted spermatogenesis in vivo. a, The protein level of M5 detected by immunofluorescence staining (IHF) in mouse testes after treatment with M5 shRNA for 10 d. Scale bar: 50 µm. b, The protein level of M5 detected by Western blotting (WB) in mouse testes after treatment with shRNA for 10 days. c, Gene expression heatmap of mouse testicular samples after shRNA treatment for 10 days. d, Principal Component Analysis (PCA) for gene expression

in mouse testes. e, Gene ontology (GO) enrichment analysis of the genes decreased by shM5 treatment in mouse testes. f, GO enrichment analysis of the genes increased by shM5 treatment in mouse testes. g, Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis of the genes decreased by shM5 treatment in mouse testes. h, KEGG enrichment analysis of the genes increased by shM5 treatment in mouse testes.

Figure 3

Network analysis of the genes related to spermatogenesis and male fertility in mouse testes. a, The network of genes related to spermatogenesis and male fertility, and the signaling pathways enriched by GO analysis. b, The network of genes related to spermatogenesis and male fertility enriched by GO

analysis. c, The protein level of TP1, PGK2, and SOX9 detected by IHF in mouse testes after shM5 treatment. Scale bar: 25 μ m. d, The protein level of TP1, PGK2, CREM, and p-FSCN1 detected by WB in mouse testes after shM5 treatment.

Figure 4

Network analysis of the genes related to cell-cell junctions in mouse testes. a, The network of genes related to cell-cell junctions, and the signaling pathways enriched by GO analysis. b, Protein levels of

occludin, claudin, Cx43, and Cx37 detected by IHF in mouse testes after shM5 treatment. Scale bar: 25 μ m. d, Protein level of occludin detected by WB in mouse testes after shM5 treatment.

Figure 5

shM5 disrupted the blood-testis barrier and ectoplasmic specialization (ES) in mouse spermatogonia cells (C18-4) in vitro. a, Protein level of M5 detected by IHF in C18-4 cells after shM5 treatment. Scale bar: 25 μ m. b, Gene expression heatmap of C18-4 cells after shM5 treatment. c, PCA analysis for gene

expression of C18-4 cells. d, Reactome enrichment analysis of the genes decreased by shM5 treatment in C18-4 cells. e, GO enrichment analysis of the genes decreased by shM5 treatment in C18-4 cells. f, Status of the cell cytoskeleton (actin) in C18-4 cells detected by IHF after shM5 treatment. g, Protein levels of the blood-testis barrier (BTB) and ectoplasmic specializations (ES) specific proteins occludin, claudin, E-cadherin, ZO-1, catenin, Cx43, Cx37, and JAM1 detected by IHF in C18-4 cells after shM5 treatment. Scale bar: 25 μ m.

Figure 6

shM5 disrupted the blood-testis barrier (BTB) and ectoplasmic specializations (ES) in Sertoli cells (TM4) in vitro. a, Protein level of M5 detected by IHF in TM4 cells after shRNA treatment. Scale bar: 25 μ m. b, Gene expression heatmap of TM4 cells after shRNA treatment. c, PCA analysis for the gene expression of TM4 cells. d, Reactome enrichment analysis of the genes decreased by shM5 treatment in TM4 cells. e, GO enrichment analysis of the genes decreased by shM5 treatment in TM4 cells. f, Status of the cell cytoskeleton (actin) in TM4 cells detected by IHF after shM5 treatment. g, Protein levels of the BTB and ES specific proteins occludin, claudin, E-cadherin, ZO-1, catenin, Cx43, Cx37, and JAM1 detected by IHF in TM4 cells after shM5 treatment. Scale bar: 25 μ m.

Figure 7

Phldb2 pathway involved in the disruption of the blood-testis barrier (BTB) and ectoplasmic specialization by knockdown M5 in C18-4 and TM4 cells in vitro. a, Summary of gene expression after shM5 treatment in C18-4 and TM4 cells. b, The enrichment data of the 27 genes (overlay in both C18-4 and TM4 cells) by Metascape online. c, Gene network of PHLDB2 signaling pathway. d, Protein levels of

PHLDB2 in C18-4 cells after shM5 treatment. Scale bar: 25 μ m. e, Protein levels of PHLDB2 in TM4 cells after shM5 treatment. Scale bar: 25 μ m.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryFig1.tif](#)
- [SupplementaryFig2.tif](#)
- [SupplementaryFig3.tif](#)
- [SupplementaryFig4.tif](#)
- [SupplementaryFig5.tif](#)
- [SupplementaryFig6.tif](#)
- [SupplementaryFig7.tif](#)