

Differential expression of IDA (INFLORESCENCE DEFICIENT IN ABSCISSION)-like genes in *Nicotiana benthamiana* during corolla abscission, stem growth and water stress

Daniel Ventimilla

Instituto Valenciano de Investigaciones Agrarias

Concha Domingo

Instituto Valenciano de Investigaciones Agrarias

Daniel Gonzalez-Ibeas

Instituto Valenciano de Investigaciones Agrarias

Manuel Talon

Instituto Valenciano de Investigaciones Agrarias

Francisco R. Tadeo (✉ tadeo_fra@gva.es)

<https://orcid.org/0000-0002-5839-4255>

Research article

Keywords: abiotic stress, cell separation, cis-acting regulatory elements, gene expression, LRR-RLKs, phylogeny, signaling peptide, Solanaceae, vegetative growth

Posted Date: January 3rd, 2020

DOI: <https://doi.org/10.21203/rs.2.12900/v3>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 20th, 2020. See the published version at <https://doi.org/10.1186/s12870-020-2250-8>.

Abstract

Background IDA (INFLORESCENCE DEFICIENT IN ABSCISSION)-like signaling peptides and the associated HAE (HAESA)-like family of receptor kinases were originally reported in the model plant *Arabidopsis thaliana* (*Arabidopsis*) to be deeply involved in the regulation of abscission. Actually, IDA peptides, as cell-to-cell communication elements, appear to be implicated in many developmental processes that rely on cell separation events, and even in the responses to abiotic stresses. However, the knowledge related to the molecular machinery regulating abscission in economically important crops is scarce. In this work, we determined the conservation and phylogeny of the IDA -like and HAE -like gene families in relevant species of the Solanaceae family and analyzed the expression of these genes in the allopolyploid *Nicotiana benthamiana*, in order to identify members involved in abscission, stem growth and in the response to drought conditions. Results The phylogenetic relationships among the IDA -like members of the Solanaceae studied, grouped the two pairs of NbenIDA1 and NbenIDA2 protein homeologs with the *Arabidopsis* prepropeptides related to abscission. Analysis of promoter regions searching for regulatory elements showed that these two pairs of homeologs contained both hormonal and drought response elements, although NbenIDA2A lacked the hormonal regulatory elements. Expression analyses showed that the pair of NbenIDA1 homeologs were upregulated during corolla abscission. NbenIDA1 and NbenIDA2 pairs showed tissue differential expression under water stress conditions, since NbenIDA1 homeologs were highly expressed in stressed leaves while NbenIDA2 homeologs, especially NbenIDA2B , were highly expressed in stressed roots. In non-stressed active growing plants, nodes and internodes were the tissues with the highest expression levels of all members of the IDA -like family and their putative HAE -like receptors. Conclusion Our results suggest that the pair of NbenIDA1 homeologs are involved in the natural process of corolla abscission while both pairs of NbenIDA1 and NbenIDA2 homeologs are implicated in the response to water stress. The data also suggest that IDA peptides may be important during stem growth and development. These results provide additional evidence that the functional module formed by IDA peptides and its receptor kinases, as defined in *Arabidopsis*, may also be conserved in Solanaceae.

Background

The significance of the *INFLORESCENCE DEFICIENT IN ABSCISSION* (IDA)-like gene family is primary associated with the observation that *AtIDA* was deeply involved in the regulation of the abscission of floral organs and cauline leaves in *Arabidopsis thaliana* (*Arabidopsis*) [1–3]. Abscission is an active, organized and highly coordinated cell separation process allowing the detachment of entire vegetative and reproductive organs through the modification of cell-to-cell adhesion and breakdown of cell walls at specific sites on the plant body known as abscission zones (AZs), a discrete group of functionally specialized cells (for a review, see [4]). From an evolutionary point of view, abscission is a very favorable process that has several advantages such as seed dispersal as well as the shedding of no longer needed, damaged or infected organs. In addition, the abscission process is related to other processes such as senescence, pathogen defense and drought stress tolerance [5]. Abscission of aerial organs, on the other

hand, may become a major limiting factor of yield in an agricultural context. It is widely accepted that the control of abscission in Arabidopsis requires physical interaction of the hormonal peptide AtIDA, a pair of redundant receptor-like protein kinases, HAESA (HAE) and HAESA-LIKE2 (HSL2), and SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASE (SERK) co-receptors (for a review, see [6]). This interaction activates a signal transduction through a downstream MAP kinase cascade that leads to the expression of multiple abscission-related hydrolytic enzymes such as pectin-methylesterases, polygalacturonases, cellulases, xyloglucan-endotransglycosylase/hydrolases and expansins [7, 8]. The release of this set of enzymes cause the disassembly of the cell wall and the dissolution of the middle lamella, resulting in the detachment of floral organs in Arabidopsis. Furthermore, in Arabidopsis *ida* mutants, petals remain indefinitely attached to the flower [1]. It has been also shown that synthetic IDA peptides were able to induce early floral abscission in Arabidopsis flowers [9].

The function of IDA peptides, as cell-to-cell communication elements, does not appear to be solely restricted to their roles in abscission. The activity of these peptides has also recently been involved in other developmental processes in Arabidopsis all of them settled on cell separation events such as the emergence of lateral roots and the root cap sloughing [10, 11]. In addition, IDA peptides have recently been involved in the response to abiotic stresses. Expression analysis of Arabidopsis *AtIDL6* and *AtIDL7*, for instance, revealed that these two genes are rapidly induced during various stress treatments [12]. It was subsequently determined experimentally that these peptides were involved in the stress response as modulators of reactive oxygen species (ROS) signaling [13]. Furthermore, treatments with *AtIDL6* and *AtIDL7* peptides caused downregulation of important stress-response key regulators like ZINC FINGER PROTEIN and WRKY transcription factors.

In addition to Arabidopsis, *IDA*-like genes have been identified in some crop species associated with organ abscission and the emergence of lateral roots [14] as well. Regarding organ abscission, particular members of the *IDA*-like gene family of tomato (*SllDA1*), soybean (*GmIDA2a*), citrus (*CitIDA3*), litchi (*LcIDL1*), oil palm (*EgIDA5*) and yellow lupine (*LlIDA*) were highly expressed in leaf, flower or fruit abscission zones during abscission [15–19]. In addition, the *AtIDA* homologues of citrus (*CitIDA3*) and litchi (*LcIDA1*) were able to induce earlier floral organ abscission and to rescue the *ida2* abscission deficiency when ectopically expressed in Arabidopsis [16, 20]. All these data, together with the positive effect of treatments with synthetic IDA peptides on triggering organ abscission [9, 19, 21] strongly suggest conserved functions of *IDA*-like genes in regulating cell separation events during organ abscission.

Solanaceae is a large plant family with approximately 90 genera comprising more than 3000 species found on almost all continents. *Solanaceae* is also one of the most economically important families worldwide. Some species of this family such as tomato (*Solanum lycopersicum*), potato (*S. tuberosum*), eggplant (aubergine; *S. melongena*) and pepper (*Capsicum annuum*) are of great relevance as a human food source. Overall, more than 29 million hectares of these Solanaceae food species were cultivated globally in 2016, producing 644 million metric tons with a net production value of more than 146 billion US dollars (<http://www.fao.org/faostat>). Thus, in addition to being important in human nutrition, they are

also relevant in economic and social terms. Other Solanaceae such as tobacco (*Nicotiana* spp.) have medical importance as a source of plant drugs while *Nicotiana benthamiana* is considered a relevant model organism for the study of plant-microbe interactions and also in plant molecular research and biotechnology [22, 23]. In this work, firstly, we determined the conservation and phylogeny of the *IDA*-like and *HAE*-like gene families by taking advantage of the free availability of the diploid genome sequences of tomato, eggplant, pepper, *N. Sylvestris*, *N. tomentosiformis*, the allopolyploids *N. tabacum* and *N. benthamiana*, and the double haploid genome sequence of potato in the Solanaceae Genomic Network (SGN; <https://solgenomics.net/>). Allopolyploidy is a type of whole genome duplication derived from hybridization of two or more diverged taxa, that primarily occurs through the fusion of unreduced (2n) gametes. The result of this kind of genome merging is the occurrence of pairs of homologous genes from each of the diploid parents in the allopolyploid genome, called homeologs [24]. Therefore, in the allopolyploid genomes of *N. tabacum* and *N. benthamiana* we will likely find pairs of homeologs for many of the members of the gene families.

In order to identify and discriminate members involved in organ abscission, stem growth and in the response to drought conditions, we examined the expression of the homeolog genes of the *N. benthamiana* *IDA*-like and *HAE*-like families. The abscission of the corolla, the only organ that undergoes abscission in *N. benthamiana*, should be highly similar to that reported in *N. tabacum* [25]. The detachment of the corolla is due to the dissolution of the middle lamella and apparently to the disintegration of the parenchymal cells in its basal zone, a process that results in the detachment of the senescent corolla. Furthermore, the effect of water stress on the species of this contrasted genus has been the subject of major research and the physiological responses of these plants are also well known [26–29].

Results

The *IDA*-like gene family in the Solanaceae

Table 1 summarizes all *IDA*-like genes identified in our search in representative species of the genus *Nicotiana* such as *N. sylvestris*, *N. tomentosiformis*, *N. tabacum* and *N. benthamiana* in addition to other Solanaceae of agronomic interest such as tomato, potato, eggplant and pepper. All prepropeptides identified share two relevant characteristics, a signal peptide targeting the protein to the apoplast through the secretory pathway and a highly conserved C-terminal signature termed PIP motif typical of this gene family [1].

IDA-like families of the *Nicotiana* species *N. sylvestris* and *N. tomentosiformis* consisted of 5 members, while in the allopolyploids *N. benthamiana* and *N. tabacum* these families are formed by 5 pairs of homeologs, with one exception corresponding to *NbenIDA4* whose homeolog pair was not found in the genomic screening. All *IDA*-like genes found in *Nicotiana* are new identifications, as the six members found in *S. melongena* and *C. annuum* and the seven members of the *S. tuberosum* family. In *S.*

Lycopersicum, five out of the eight *IDA*-like genes detected, members 1 to 5, were already described in [15] and named *SIIDA1-5*, while the other three peptides, *SlycIDA6-8*, are incorporated in the current work.

Phylogenetic relationship among IDA-like prepropeptides in Solanaceae

The phylogenetic relationships among the IDA-like members of the species of Solanaceae studied, in addition to those of Arabidopsis, are grouped in three major clades (Figure 1). Clade I (shadowed in green colors) was divided in two subclades. The subclade shadowed in green contained the two Arabidopsis prepropeptides involved in floral organ abscission, *AtIDA* and *AtIDL1* [1, 9]. The largest subclade grouped members of all eight Solanaceae species studied, as well as *AtIDL8*, the most divergent IDA-like peptide from Arabidopsis. In this subclade, Solanaceae members are further divided in two major groups. The group shadowed in lime green contained *SIIDA1*, the IDA-like member of tomato that has been associated with leaf abscission [15], other prepropeptides of potato (*StubIDA4*), eggplant (*SmelIDA5*) and pepper (*CalIDA4*) as well as the *IDA1* members of the *Nicotiana* species under study. These were *NsylIDA1* and *NtomIDA1* of the diploid species *N. sylvestris* and *N. tomentosiformis*, respectively, and the two pairs of *NtabIDA1* and *NbenIDA1* homeolog prepropeptides corresponding to *N. tabacum* and *N. benthamiana*. The 5'-UTR regions and the predicted CDSs of all these *IDA1* genes from the genus *Nicotiana* showed high degree of conservation (see Additional files 1 and 3). The other group shadowed in light green included other prepropeptides from the *Nicotiana*, *Solanum* and *Capsicum* genera, with a small subdivision composed of *AtIDL8* together with *SlycIDA6*, *SlycIDA7* and *StubIDA1* (Figure 1). A second clade, clade II (shadowed in light orange), appeared to be limited to the Solanaceae family. This clade included prepropeptides from the *Nicotiana*, *Solanum* and *Capsicum* genera, but none from Arabidopsis, an observation suggesting that it might have diverged before the irruption of the Brassicaceae family 40 million years ago [30]. The third clade, clade III (shadowed in light gold), included *AtIDL6* and *AtIDL7*, two IDA-like members of Arabidopsis that have been associated with processes different than cell separation, such as stress response [13]. The topology of the clade showed that there was a great diversification in Arabidopsis that generated at least six members, *AtIDL2-7*. It also included prepropeptides from the *Nicotiana* and *Solanum* genera, but none from *Capsicum*.

Cis-acting regulatory elements in the promoter regions of the *N. benthamiana* *IDA*-like family

Figure 2 shows a schematic representation of the *cis*-acting regulatory elements along 1000 bp of the 5'-UTR region of the *IDA*-like family members of *N. benthamiana* and *AtIDA* and *AtIDL1* of Arabidopsis. Searches for response elements to hormones related to ABA, methyl jasmonate (MeJa), AUXs or GAs, as well as response elements to biotic and abiotic stresses were performed. Interestingly, the pair of *NbenIDA1* homeologs contained similar promoter regions carrying response elements to ABA, MeJa and AUX as *AtIDA* in similar locations; these phytohormones have been involved in the abscission process in different ways [1, 31, 32]. The pairs of *NbenIDA1* and *NbenIDA2* homeologs also carry drought response elements in their promoter regions. On the other hand, *NbenIDA2B*, *NbenIDA3A*, *NbenIDA4*, and the pair of *NbenIDA5* homeologs are characterized by the occurrence of GA response elements (Figure 2).

Expression patterns of *IDA*-like and *HAE*-like genes in *Nicotiana benthamiana* during growth and abscission

Expression analysis of the family of *IDA*-like ligand peptides and their putative *HAE*-like receptors in different plant tissues of *N. benthamiana* are presented in Figure 3. *HAE*-like receptors were identified through the analyses of the phylogenetic relationships between the *HAE*-like receptor-like kinases (RLKs) of Arabidopsis and *Nicotiana* (see Additional files 2 and 3). The plant material selected for gene expression analysis is shown in the panel A of Figure 3. This plant material included different vegetative tissues of a plant in active growth (apical buds, young and mature leaves, nodes and internodes, and roots), as well as reproductive tissues (anthers, styles, stigmas, and fruits) at different developmental stages, including samples of the base of the flower corollas, a tissue that in tobacco (*N. tabacum*) has been shown to respond to the abscission process [25]. Panel B in Figure 3 shows the expression of the *N. benthamiana* *IDA*-like and *HAE*-like homeologs in apical buds, nodes, internodes, the whole corolla, the ensemble formed by the stigma and the style and also in roots relative to the lowest expression level of each gene. Panel C in Figure 3 shows the expression pattern of each *IDA*-like and *HAE*-like homeolog in leaves, anthers and fruits relative to that at the earliest developmental stage in every organ and panel D shows the expression patterns of each homeolog gene at the corolla base in developmental stages 2, 4 and 5 relative to the corolla developmental stage 1. Virtually all members of the *IDA*-like family of *N. benthamiana* were mainly expressed in nodes and internodes, although *NbenIDA1A* expression levels were not especially high in internodes (Figure 3B). No changes in the expression patterns of *IDA*-like homeolog genes were observed in leaves and fruits (Figure 3C) but the expression level for all of them except *NbenIDA3B* and *NbenIDA4* showed a tendency to increase between closed and dehiscent anthers (Figure 3C). Interestingly, expression of both *NbenIDA1* homeologs at the base of the flower corolla increased with the stage of development of the tissue, in parallel to the progress of the abscission process (Figure 3D). The expression pattern of *NbenIDA1B* was similar to that detected in *NbenIDA1A* although at a much more modest level. The expression levels of the *NbenIDA2* homeologs were transiently high in stage 2 when the corolla tube is fully elongated and the limb lobes are still closed to return later in stages 4 and 5 to almost the basal level of expression (Figure 3D).

The highest expressions levels of the putative receptors of the *IDA*-like peptides, *NbenHAE.1*, *NbenHAE.2*, *NbenHSL2.1* and *NbenHSL2.2*, were also registered in nodes and internodes (Figure 3B). Additionally, their expression levels also showed a tendency to increase between closed and dehiscent anthers (Figure 3C) and a slight increase was observed at the corolla base associated with corolla development (Figure 3D).

Expression patterns of *IDA*-like genes in *Nicotiana benthamiana* during water stress

The presence of drought response elements in the promoter regions of some particular *IDA*-like members, e.g. *NbenIDA1A*, *NbenIDA1B*, *NbenIDA2A* and *NbenIDA2B* (Figure 2), suggested that their expression might be regulated by the water status of the plant. Therefore, we exposed actively growing plants of *N. benthamiana* to 6 (mild stress) and 8 (severe stress) days of water stress and the expression levels of all members of the *IDA*-like family in axillary buds, roots and leaves were determined (Figure 4). While no

differences in gene expression were found in axillary buds, those of the pair of *Nben/DA1* homeologs dramatically increased in leaf blades of plants subjected to severe water stress. In contrast, this condition resulted in higher increases in transcripts belonging of both *Nben/DA2* homeologs in roots, indicating differential roles of this gene family in response to water stress. Changes in the expression of the rest of genes were of minor relevance although it is worth to mention that these members tended to repress their expression levels in roots of plants subjected to water stress, although *Nben/DA5A* expression was also reduced in stressed leaves.

Discussion

IDA-like genes were searched in relevant genera of the Solanaceae family including several species of *Nicotiana* (*N. sylvestris*, *N. tomentosiformis*, *N. tabacum* and *N. benthamiana*), and other crops of agronomic interest such as tomato (*Solanum lycopersicum*), potato (*S. tuberosum*), eggplant (*S. melongena*) and pepper (*Capsicum annuum*) (Table 1). This gene family was identified in a large number of Angiosperms [17] and their members contained a signal peptide targeting the protein to the apoplast through the secretory pathway and a conserved C-terminal part, the PIP motif (Additional file 4). The presence of a signal peptide in the sequence of all identified genes suggests a mechanism of posttranslational maturation in the apoplast similar to that described in Arabidopsis, where the prepropeptide is proteolytically processed by subtilisin-like serine proteinases to yield a bioactive peptide [33].

The reduced size of the mature *IDA*-like peptides (an alignment of the complete coding sequences of these genes can be seen in Additional file 4), precluded the study of the phylogenetic relationships based on these premises and, therefore, a circular phylogenetic tree was generated using complete sequences encoding prepropeptides including a signal peptide and a variable region (Figure 1). This tree shows that the Arabidopsis *IDA*-like gene family exhibits a higher degree of diversification than the Solanaceae genera studied, with the exception of *Nicotiana*. The two Arabidopsis prepropeptides related to abscission, AtIDA and AtIDL1 [1, 31], nested in a small clade grouped with a larger clade including many *IDA*-like members from the different species of Solanaceae. This large clade was divided in two subgroups in one of which was nested the tomato *SlycIDA1*, associated with abscission [15], and its homolog genes in *S. melongena*, *C. annuum*, *S. tuberosum* and in the four *Nicotiana* species studied. The genus *Nicotiana* originates in South America and its members have spread over four continents. This genus consists of diploid species and several allopolyploid species of different ages and paternity. Thus, cultivated tobacco (*N. tabacum*) is an allotetraploid ($2n = 4x = 48$) representing a hybridization event involving the diploid species *N. sylvestris* ($2n = 2x = 24$) and *N. tomentosiformis* ($2n = 2x = 24$) as their female and male parents, respectively. This hybridization event is believed to have occurred recently, less than 200,000 years ago [34, 35]. The fact that all homeologs of the *IDA*-like family of *N. tabacum* have corresponding counterparts in *N. sylvestris* and *N. tomentosiformis* (Table 1 and Figure 1) is in line with this observation (Figure 1).

In contrast to tobacco, *N. benthamiana* has been described as an ancient allotetraploid whose polyploidy level ($2n = 4x = 38$) likely evolved through genome re-arrangements and fractionation giving rise to a remarkable descending dysploidy [35]. Parentals of *N. benthamiana* are unknown, although it is believed that it comes from a hybridization event that occurred >10 Myr ago between species belonging to the *Sylvestres* and *Noctiflorae* sections of *Nicotiana* [35]. Two homeologs were identified for most of the analysed genes in our work, except for *NbenIDA4*. Genomic responses to polyploidy are complex in *Nicotiana* species, ranging from small to large genome re-sizing depending on the polyploid age and the similarity of parental genome donors [35]. Reduction in the number of chromosomes in *N. benthamiana* strongly suggests a considerable genome downsizing, probably as a consequence of being an older polyploid. The size reduction involves 1 Gb in length relative to tobacco (4.5 Gb genome size of *N. tabacum*, 3.5 Gb *N. benthamiana*). Therefore, we suggest that gene loss might explain the absence of a second copy of *NbenIDA4* in our data, rather than a misrepresentation in the draft assembly of the genome used for the analysis. Interestingly, several genetic studies estimated that the genome of *N. tabacum* had lost DNA from its progenitors since polyploidization and that this genomic loss was greater and biased towards the genome of the male parental *N. tomentosiformis* [35–38]. Therefore, we also think that similar biased gene loss may have happened in *N. benthamiana* involving the copy of *NbenIDA4* belonging to the parent of the *Noctiflorae* section of *Nicotiana*.

The analyses of the *cis*-acting regulatory elements in the 5'-UTR regions of the *N. benthamiana* *IDA*-like family and *Arabidopsis AtIDA* and *AtIDL1* (Figure 3) failed to identify ethylene response elements, in agreement with the idea that *IDA*-like genes regulating the abscission process are not directly dependent upon ethylene [39, 40]. In contrast, the presence of response elements to AUXs, ABA, MeJa and GAs in the promoter regions of the *IDA*-like family members of *N. benthamiana* and *AtIDA* and *AtIDL1* suggests that these phytohormones might play a role in regulating the expression of these genes. The occurrence of functional indole-3-acetic acid (IAA) signaling in the abscission zone during organ separation, for instance, has been demonstrated by Basu and co-workers [41]. It has also been determined that ABA and MeJa have abscission-promoting effects, while the role of GAs is not entirely clear [6, 42]. However, it has been shown in citrus that flower pollination increased bioactive gibberellin A1 (GA₁) levels and reduced ovary abscission and that the treatment of unpollinated ovaries with GA₃ also suppressed ovary abscission [43, 44].

Our bioinformatic analyses also indicated that the coding and promoter sequences of the pair of *NbenIDA1* homeologs are highly similar and that promoters share the same hormonal response elements in similar positions, in addition to the same drought response element (Figure 2). High conservation of the coding and promoter sequences of *IDA1* duplicated genes in *N. sylvestris*, *N. tomentosiformis* and *N. tabacum* (see Additional file 3) suggests that they may be very important in the regulation of cell separation processes and response to stressful conditions. Furthermore, the pair of *NbenIDA2* homeologs also contains drought stress response elements in their promoter regions (Figure 2). Likewise, the coding and promoter sequences of the *IDA1* genes in *N. attenuata*, *N. sylvestris*, *N. tomentosiformis*, *N.*

benthamiana and *N. tabacum* are very similar and have the same response elements in the same positions in their promoters, except *N. attenuata* (Additional file 1).

The results described above (Figure 1) suggested that in *N. benthamiana*, NbenIDA1A and NbenIDA1B peptides may be involved in the abscission process. This suggestion is also supported by the gene expression patterns found at the corolla base of the flowers during the process of natural abscission (Figure 3D). Similarly, there seems to be a correlation between the expression of the *IDA*-like genes and that of their putative receptors of the *HAE*-like family, *NbenHAE.1*, *NbenHAE.2* and *NbenHSL.2.2* (Figure 3D), that also increased during the last phases of the corolla abscission.

As described for *IDA*-like families in other species [12], the different members of the *N. benthamiana* family are also expressed in multiple plant tissues (Figure 3). This is not a surprise since the *IDA*-like signaling peptides, as cell-to-cell communication elements, function in several cell separation events, including lateral root emergence and root cap sloughing [10, 11]. Interestingly, in plants of *N. benthamiana* actively growing, the highest expression level of most members of the *IDA*-like family was found in nodes and internodes. It is worth mentioning that the promoter regions of *NbenIDA2B*, *NbenIDA3A*, *NbenIDA4*, *NbenIDA5A* and *NbenIDA5B* genes contain GAs response elements, and that these hormones are pivotal regulators of stem growth [45]. Moreover, all *HAE*-like genes analyzed also show higher expression levels in nodes and internodes, in parallel with the pattern observed for the *IDA*-like genes. These expression patterns might be linked to the formation of vascular bundles and to the cell elongation and division associated with the process of stem elongation implying cell wall remodeling.

The occurrence of cis-acting elements related to the drought response in the two pairs of *NbenIDA1* and *NbenIDA2* homeologs (Figure 2) also suggested to test the response of the *IDA*-like genes to water stress conditions. In the experiment reported in Figure 4 it is clearly observed that the first pair of homeologs was highly expressed in leaves from *N. benthamiana* plants severely stressed while in roots, the genes that responded to water deficit were the members of the second pair. Furthermore, these two pairs of homeologs are phylogenetically close to *AtIDA* and *AtIDL1*, two *Arabidopsis* genes that are induced under abiotic stress conditions [12].

Our gene expression data showed that while most pairs of homeologs showed similar expression patterns, some of them exhibited divergence in certain organs and tissues studied (see Figure 3). This might well be linked to the frequent observation that some duplicated genes, after a whole genome duplication event, evolve to undertake different functions or partition the function of the ancestral gene in a process of subfunctionalization. This process can include epigenetic, coding sequence or promoter modifications that alters regulatory mechanisms (e.g. silencing) and give rise for example to differential level of expression or tissue specificity. Subfunctionalization becomes more relevant when gene dosage is not an adaptive advantage for the polyploid [46]. Therefore, our gene expression data might be revealing a putative subfunctionalization for the homeolog pairs *NbenIDA1* at the corolla base and *NbenIDA3* in leaves and anthers. We took special care in primer specificity during qPCR assays in order to

distinguish between both homeologs, since gene expression artifacts may be recurrent among genes derived from genome duplicated areas due to high sequence similarity.

It has been recently observed that IDA signaling peptides can certainly regulate important developmental processes as well as fundamental plant responses to environmental conditions [13]. Our data indicate that in the allopolyploid *N. benthamiana*, the two pairs of *NbenIDA1* and *NbenIDA2* homeologs are differentially involved in the responses to drought stress while only *NbenIDA1* homeologs are apparently implicated in the natural process of corolla abscission. These data suggest that IDA-like signaling peptides can play different biological roles in various tissues and under distinct abiotic conditions.

Conclusions

We have investigated the *IDA*-like and *HAE*-like gene families of different Solanaceae species, *S. lycopersicum*, *S. melongena*, *C. annuum*, *S. tuberosum*, and four species of the genus *Nicotiana*, *N. sylvestris*, *N. tomentosiformis*, *N. benthamiana*, and *N. tabacum* and determined their phylogenetic relationships. In the allopolyploid *N. benthamiana*, specific analyses of the the *cis*-acting regulatory sequences and the examinations of the gene expression patterns of the *IDA*-like family have identified putative candidate *IDA*-like genes implicated in corolla abscission and in the response to water stress. The results suggest that the pair of *NbenIDA1* homeologs are both involved in the natural process of corolla abscission. Interestingly, they also show specific differential expression under water stress conditions. *NbenIDA1* homeologs are highly expressed in stressed leaves while *NbenIDA2* homeologs, especially *NbenIDA2B*, are highly expressed in stressed roots. In addition, nodes and internodes are the tissues with the highest expression of the *IDA*-like and *HAE*-like genes in normal active growing plants, suggesting that these peptides are also essential during stem growth and development. These results add new evidence that the functional module formed by IDA-like peptides and its receptor kinases as defined in *Arabidopsis*, may be conserved in Solanaceae.

Methods

Retrieval and sequence analysis

The EPIP motif of AtIDA (FGYLPKGVPPIPPSAPSKRHNSFVNSLPH) was used to identify the *IDA*-like members of the selected Solanaceae species (*N. sylvestris*, *N. tomentosiformis*, *N. tabacum*, *N. benthamiana*, *Solanum lycopersicum*, *S. tuberosum*, *S. melongena* and *Capsicum annuum*) by tBLASTn and BLASTp inquiries in the Sol Genomics [47] web platform (<https://solgenomics.net/tools/blast/>), depending on the databases status. “*N.sylvestris Genome*”, “*N.tomentosiformis Genome*”, “*N.tabacum BX Genome*”, “*N.benthamiana v1.0.1*”, “*Tomato ITAG release 3.20*”, “*Potato PGSC DM v3 scaffolds*”, “*Eggplant draft genome (release 2.5.1)*” and “*Capsicum annuum UCD10X genome chromosomes (v1.0)*” databases [48–53] were used, respectively. *Arabidopsis* AtHAE, AtHSL1 and AtHSL2 protein sequences were retrieved from Phytozome v12.1, TAIR10 database and were used to identify the *HAE*-like members of the selected Solanaceae species in the same way as described above. Newly identified genes were

named numerically, adding an “A”, “B”, “.1” or “.2” termination to the *IDA*-like or *HAE*-like gene pairs for the allopolyploids *N. tabacum* and *N. benthamiana*.

Sequence alignments were performed through MEGA7 software [54] using the ClustalW algorithm with default parameters (DNA Data Bank of Japan, DDBJ; <http://clustalw.ddbj.nig.ac.jp/>). Phylogenetic trees were created using the Neighbor-Joining method [55] using 1000 bootstrap replicates. The trees are drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic trees. The evolutionary distances were computed using the Poisson correction method [56] and are in the units of the number of amino acid substitutions per site. All ambiguous positions were removed for each sequence pair (pairwise deletion option).

Peptide localization prediction and presence of signal peptides in the IDL amino acid sequences were analyzed using the TargetP [57] and SignalP-5.0 [58] services. Up to 1000 base pairs of promoter regions upstream of the start codon of the IDL genes of the *Nicotiana* species available in Sol Genomics databases were retrieved and submitted for cis-acting regulatory element analysis in PlantCARE [59]. Schematic representations of regulatory elements of the promoter sequences were created using IBS1.0.3 software [60].

Plant materials and growth conditions

N. benthamiana seeds were obtained from Dr. José Guerri and Dr. Karelia Velázquez of the Centro de Protección Vegetal y Biotecnología (IVIA, Moncada, Spain). The seeds were germinated on nutrient soil and transplanted individually in small pots with an artificial potting mix (50% vermiculite and 50% peat moss) in a plant growth chamber at 20/24 °C (night/day), 60% relative humidity and a 16/8-h light/dark regime. Water stress was induced by not watering the plants for 6 and 8 days for mild and severe stress conditions respectively.

RNA extraction

Basal portion of the corollas at different flower development stages as well as the rest of tissues used in gene expression analysis were manually collected from the plants and frozen with liquid nitrogen. Tissues were grinded using Thomas Scientific’s Liquid Nitrogen Cooled Mortar. Total RNA was extracted using Macherey-Nagel’s NucleoSpin® RNA Plant, following the manufacturer’s instructions. cDNA was synthesized from the RNA extraction using Thermo Fisher Scientific’s SuperScript™ II Reverse Transcriptase, following the manufacturer’s instructions.

qPCR analysis

Quantitative PCR analysis were performed using LightCycler® FastStart DNA MasterPLUS SYBR Green I reaction mix and a LightCycler 2.0 instrument (Roche, Basel, Switzerland) using gene-specific primers designed based on transcriptome sequences using the Primer3Plus software [61]. Primer pairs are listed in Additional file 5. Specificity of all the primer pairs used in this work was assessed by primer BLAST and melting curve analysis [62, 63]. The fluorescence intensity data was obtained through LightCycler

Software version 4.1. The *N. benthamiana* housekeeping gene *NbenPP2A* [64] was used for normalization in all qPCR reactions carried out in this work. Three biological replicates were run for assessing the expression values of each gene. The averaged expression values were obtained in the form of Ct (cycle threshold) and all the analyzes were performed through $2^{-\Delta\Delta CT}$ method.

For gene expression relative quantification of *IDA*-like and *HAE*-like genes in different plant tissues and organs in wild type plants, every gene expression was normalized respect to *NbenPP2A* expression. In apical buds, nodes, internodes, corolla, style and stigma, and root tissues, gene expression values were relative to the lowest expression value of each gene in the relevant tissue or organ, within primer sets. In leaf, anthers, fruits and corolla base tissues, gene expression values were relative to the expression value of the earliest developmental stage relevant for that tissue (young leaf, anthers and fruits in stage 2, and corolla base of a stage 1 flower, respectively) within primer sets. Units were represented as the log2 of the fold change. In the case of leaf, anthers, fruits and base of the corolla tissues, red color indicates that the gene is upregulated (values over 0); white, that remain unchanged (values close to 0); and blue, that the gene is downregulated (values under 0); all respect to *NbenPP2A* expression in the corresponding tissue in its earliest developmental stage.

In the water stress experiment, *NbenPP2A* was also used as a housekeeping gene for normalization, and watered tissue expression values as a relative reference, thus constituting our control conditions. Conditions were appropriate for each measurement, using the corresponding control tissue (leaf or root of watered plants) as a relative reference. Units were represented as fold change. Red color indicates that the gene is upregulated (values over 1); white, that remain unchanged (values close to 1); and blue, that the gene is downregulated (values under 1); all respect to *NbenPP2A* expression in the corresponding watered (control) tissue.

Abbreviations

ABA: abscisic acid, AUX: auxins, GA: gibberellin, HAE: HAESA, IDA: INFLORESCENCE DEFICIENT IN ABSCISSION

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Not applicable

Competing interests

The authors declare that they have no competing interests

Funding

Financial support was provided by the Instituto Nacional de Investigación y Tecnología Agraria y Alimentaria, INIA, and Agencia Estatal de Innovación (Ministerio de Ciencia, Innovación y Universidades, Spain) through grants # RTA-00071-C06-01 and # RTI2018-097790-R-100, respectively, and the Instituto Valenciano de Investigaciones Agrarias (IVIA), through grant # 51915. Funds were used for the design of the study and collection, analysis, and interpretation of data and in writing the manuscript, as well as in the open access payment.

Authors' contributions

DV, CD, MT and FRT designed the research, DV, CD, and FRT performed the research and analyzed the data, DV, DGI, MT and FRT drafted and substantively revised the article.

Acknowledgements

We want to acknowledge Dr. Karelia Velázquez and Àngel Boix for preparing and taking care of the *N. benthamiana* plants.

Author details

Instituto Valenciano de Investigaciones Agrarias (IVIA), Centro de Genómica; CV-315, Km 10,7; E-46113 Moncada (Valencia), Spain

References

1. Butenko MA, Patterson SE, Grini PE, Stenvik G, Amundsen SS, Mandal A, et al. INFLORESCENCE DEFICIENT IN ABSCISSION Controls Floral Organ Abscission in Arabidopsis and Identifies a Novel Family of Putative Ligands in Plants. *Plant Cell*. 2003;15:2296–307.
2. Patharkar OR, Walker JC. Core Mechanisms Regulating Developmentally Timed and Environmentally Triggered Abscission. *Plant Physiol*. 2016;172:510–20.
3. Shi C, Alling RM, Hammerstad M, Aalen RB. Control of Organ Abscission and Other Cell Separation Processes by Evolutionary Conserved Peptide Signaling. *Plants*. 2019;8:225.
4. Estornell LH, Agustí J, Merelo P, Talón M, Tadeo FR. Elucidating mechanisms underlying organ abscission. *Plant Sci*. 2013;199–200 February:48–60.
5. Patharkar OR, Walker JC. Connections between abscission, dehiscence, pathogen defense, drought tolerance, and senescence. *Plant Sci*. 2019;284 July:25–9.
6. Patharkar OR, Walker JC. Advances in abscission signaling. *J Exp Bot*. 2018;69:733–40.

7. Liu B, Butenko M a., Shi C-L, Bolivar JL, Winge P, Stenvik G-E, et al. NEVERSHED and INFLORESCENCE DEFICIENT IN ABSCISSION are differentially required for cell expansion and cell separation during floral organ abscission in *Arabidopsis thaliana*. *J Exp Bot.* 2013;64:5345–57.
8. Niederhuth CE, Patharkar OR, Walker JC. Transcriptional profiling of the *Arabidopsis* abscission mutant hae hsl2 by RNA-Seq. *BMC Genomics.* 2013;14:37.
9. Stenvik G-E, Tandstad NM, Guo Y, Shi C-L, Kristiansen W, Holmgren A, et al. The EPIP Peptide of INFLORESCENCE DEFICIENT IN ABSCISSION Is Sufficient to Induce Abscission in *Arabidopsis* through the Receptor-Like Kinases HAESA and HAESA-LIKE2. *Plant Cell.* 2008;20:1805–17.
10. Kumpf RP, Shi C-L, Larrieu A, Stø IM, Butenko MA, Péret B, et al. Floral organ abscission peptide IDA and its HAE/HSL2 receptors control cell separation during lateral root emergence. *Proc Natl Acad Sci.* 2013;110:5235–40.
11. Shi C-L, von Wangenheim D, Herrmann U, Wildhagen M, Kulik I, Kopf A, et al. The dynamics of root cap sloughing in *Arabidopsis* is regulated by peptide signalling. *Nat Plants.* 2018;4:596–604.
12. Vie AK, Najafi J, Liu B, Winge P, Butenko MA, Hornslien KS, et al. The IDA/IDA-LIKE and PIP/PIP-LIKE gene families in *Arabidopsis*: Phylogenetic relationship, expression patterns, and transcriptional effect of the PIPL3 peptide. *J Exp Bot.* 2015;66:5351–65.
13. Vie AK, Najafi J, Winge P, Cattan E, Wrzaczek M, Kangasjärvi J, et al. The IDA-LIKE peptides IDL6 and IDL7 are negative modulators of stress responses in *Arabidopsis thaliana*. *J Exp Bot.* 2017;68:3557–71.
14. Liu C, Zhang C, Fan M, Ma W, Chen M, Cai F, et al. GmIDL2a and GmIDL4a, Encoding the Inflorescence Deficient in Abscission-Like Protein, Are Involved in Soybean Cell Wall Degradation during Lateral Root Emergence. *Int J Mol Sci.* 2018;19:2262.
15. Tucker ML, Yang R. IDA-like gene expression in soybean and tomato leaf abscission and requirement for a diffusible stelar abscission signal. *AoB Plants.* 2012;2012:pls035.
16. Estornell LH, Wildhagen M, Pérez-Amador MA, Talón M, Tadeo FR, Butenko MA. The IDA Peptide Controls Abscission in *Arabidopsis* and Citrus. *Front Plant Sci.* 2015;6 November:1–7.
17. Stø IM, Orr RJS, Fooyontphanich K, Jin X, Knutsen JMB, Fischer U, et al. Conservation of the abscission signaling peptide IDA during Angiosperm evolution: withstanding genome duplications and gain and loss of the receptors HAE/HSL2. *Front Plant Sci.* 2015;6 October:931.
18. Kim J, Yang J, Yang R, Sicher RC, Chang C, Tucker ML. Transcriptome Analysis of Soybean Leaf Abscission Identifies Transcriptional Regulators of Organ Polarity and Cell Fate. *Front Plant Sci.* 2016;7 February:1–16.
19. Wilmowicz E, Kućko A, Ostrowski M, Panek K. INFLORESCENCE DEFICIENT IN ABSCISSION-like is an abscission-associated and phytohormone-regulated gene in flower separation of *Lupinus luteus*. *Plant Growth Regul.* 2018;85:91–100.
20. Ying P, Li C, Liu X, Xia R, Zhao M, Li J. Identification and molecular characterization of an IDA-like gene from litchi, LcIDL1, whose ectopic expression promotes floral organ abscission in *Arabidopsis*. *Sci Rep.* 2016;6:37135.

21. Tranbarger TJ, Domonh  do H, Cazemajor M, Dubreuil C, Fischer U, Morcillo F. The PIP Peptide of INFLORESCENCE DEFICIENT IN ABSCISSION Enhances *Populus* Leaf and *Elaeis guineensis* Fruit Abscission. *Plants*. 2019;8:143.
22. Goodin MM, Zaitlin D, Naidu RA, Lommel SA. *Nicotiana benthamiana*: Its History and Future as a Model for Plant–Pathogen Interactions. *Mol Plant-Microbe Interact*. 2008;21:1015–26.
23. Bally J, Jung H, Mortimer C, Naim F, Philips JG, Hellens R, et al. The Rise and Rise of *Nicotiana benthamiana*: A Plant for All Reasons. *Annu Rev Phytopathol*. 2018;56:405–26.
24. Glover NM, Redestig H, Dessimoz C. Homoeologs: What Are They and How Do We Infer Them? *Trends Plant Sci*. 2016;21:609–21.
25. Wu X-M, Yu Y, Han L-B, Li C-L, Wang H-Y, Zhong N-Q, et al. The Tobacco BLADE-ON-PETIOLE2 Gene Mediates Differentiation of the Corolla Abscission Zone by Controlling Longitudinal Cell Expansion. *Plant Physiol*. 2012;159:835–50.
26. Giorio P, Guida G, Mistretta C, Sellami MH, Oliva M, Punzo P, et al. Physiological, biochemical and molecular responses to water stress and rehydration in Mediterranean adapted tomato landraces. *Plant Biol*. 2018;20:995–1004.
27. Tani E, Kizis D, Markellou E, Papadakis I, Tsamadia D, Leventis G, et al. Cultivar-Dependent Responses of Eggplant (*Solanum melongena* L.) to Simultaneous *Verticillium dahliae* Infection and Drought. *Front Plant Sci*. 2018;9 August:1181.
28. Wang Z, Xu W, Kang J, Li M, Huang J, Ke Q, et al. Overexpression of alfalfa Orange gene in tobacco enhances carotenoid accumulation and tolerance to multiple abiotic stresses. *Plant Physiol Biochem*. 2018;130 September:613–22.
29. L  pez-Serrano L, Canet-Sanchis G, Vuletin Selak G, Penella C, San Bautista A, L  pez-Galarza S, et al. Pepper Rootstock and Scion Physiological Responses Under Drought Stress. *Front Plant Sci*. 2019;10 January:38.
30. Bailey CD, Koch MA, Mayer M, Mummenhoff K, O’Kane Jr SL, Warwick SI, et al. Toward a Global Phylogeny of the Brassicaceae. *Mol Biol Evol*. 2006;23:2142–60.
31. Stenvik G-E, Butenko MA, Urbanowicz BR, Rose JKC, Aalen RB. Overexpression of INFLORESCENCE DEFICIENT IN ABSCISSION activates cell separation in vestigial abscission zones in *Arabidopsis*. *Plant Cell*. 2006;18:1467–76.
32. Aalen RB, Wildhagen M, St   IM, Butenko MA. IDA: a peptide ligand regulating cell separation processes in *Arabidopsis*. *J Exp Bot*. 2013;64:5253–61.
33. Schardon K, Hohl M, Graff L, Pfannstiel J, Schulze W, Stintzi A, et al. Precursor processing for plant peptide hormone maturation by subtilisin-like serine proteinases. *Science (80-)*. 2016;354:1594–7.
34. Clarkson JJ, Knapp S, Garcia VF, Olmstead RG, Leitch AR, Chase MW. Phylogenetic relationships in *Nicotiana* (Solanaceae) inferred from multiple plastid DNA regions. *Mol Phylogenet Evol*. 2004;33:75–90.
35. Leitch IJ, Hanson L, Lim KY, Kovarik A, Chase MW, Clarkson JJ, et al. The Ups and Downs of Genome Size Evolution in Polyploid Species of *Nicotiana* (Solanaceae). *Ann Bot*. 2008;101:805–14.

36. Skalická K, Lim KY, Matyasek R, Matzke M, Leitch AR, Kovarik A. Preferential elimination of repeated DNA sequences from the paternal, *Nicotiana tomentosiformis* genome donor of a synthetic, allotetraploid tobacco. *New Phytol.* 2005;166:291–303.
37. Renny-Byfield S, Chester M, Kovařík A, Le Comber SC, Grandbastien M-A, Deloger M, et al. Next Generation Sequencing Reveals Genome Downsizing in Allotetraploid *Nicotiana tabacum*, Predominantly through the Elimination of Paternally Derived Repetitive DNAs. *Mol Biol Evol.* 2011;28:2843–54.
38. Renny-Byfield S, Kovařík A, Chester M, Nichols RA, Macas J, Novák P, et al. Independent, Rapid and Targeted Loss of Highly Repetitive DNA in Natural and Synthetic Allopolyploids of *Nicotiana tabacum*. *PLoS One.* 2012;7:e36963.
39. Butenko MA, Stenvik G-E, Alm V, Saether B, Patterson SE, Aalen RB. Ethylene-dependent and -independent pathways controlling floral abscission are revealed to converge using promoter::reporter gene constructs in the ida abscission mutant. *J Exp Bot.* 2006;57:3627–37.
40. Wang G-Q, Wei P-C, Tan F, Yu M, Zhang X-Y, Chen Q-J, et al. The Transcription Factor AtDOF4.7 Is Involved in Ethylene- and IDA-Mediated Organ Abscission in Arabidopsis. *Front Plant Sci.* 2016;7:863.
41. Basu MM, González-Carranza ZH, Azam-Ali S, Tang S, Shahid AA, Roberts JA. The Manipulation of Auxin in the Abscission Zone Cells of Arabidopsis Flowers Reveals That Indoleacetic Acid Signaling Is a Prerequisite for Organ Shedding. *Plant Physiol.* 2013;162:96–106.
42. Marciak K, Kućko A, Wilmowicz E, Świdziński M, Przedniczek K, Kopcewicz J. Gibberellic acid affects the functioning of the flower abscission zone in *Lupinus luteus* via cooperation with the ethylene precursor independently of abscisic acid. *J Plant Physiol.* 2018;229:170–4.
43. Ben-Cheikh W, Perez-Botella J, Tadeo FR, Talon M, Primo-Millo E. Pollination Increases Gibberellin Levels in Developing Ovaries of Seeded Varieties of Citrus. *Plant Physiol.* 1997;114:557–64.
44. Mehouachi J, Iglesias DJ, Tadeo FR, Agustí M, Primo-Millo E, Talon M. The role of leaves in citrus fruitlet abscission: Effects on endogenous gibberellin levels and carbohydrate content. *J Hortic Sci Biotechnol.* 2000;75:79–85.
45. Rizza A, Jones AM. The makings of a gradient: spatiotemporal distribution of gibberellins in plant development. *Curr Opin Plant Biol.* 2019;47:9–15.
46. Force A, Lynch M, Pickett FB, Amores A, Yan Y, Postlethwait J. Preservation of Duplicate Genes by Complementary, Degenerative Mutations. *Genetics.* 1999;151:1531–45.
47. Mueller LA, Solow TH, Taylor N, Skwarecki B, Buels R, Binns J, et al. The SOL Genomics Network. A Comparative Resource for Solanaceae Biology and Beyond. *Plant Physiol.* 2005;138:1310–7.
48. Bombarely A, Rosli HG, Vrebalov J, Moffett P, Mueller LA, Martin GB. A Draft Genome Sequence of *Nicotiana benthamiana* to Enhance Molecular Plant-Microbe Biology Research. *Mol Plant-Microbe Interact.* 2012;25:1523–30.
49. Edwards KD, Fernandez-Pozo N, Drake-Stowe K, Humphry M, Evans AD, Bombarely A, et al. A reference genome for *Nicotiana tabacum* enables map-based cloning of homeologous loci implicated in nitrogen utilization efficiency. *BMC Genomics.* 2017;18:448.

50. Tomato Genome Consortium T. The tomato genome sequence provides insights into fleshy fruit evolution. *Nature*. 2012;485:635–41.
51. Sierro N, Battey JN, Ouadi S, Bovet L, Goepfert S, Bakaher N, et al. Reference genomes and transcriptomes of *Nicotiana sylvestris* and *Nicotiana tomentosiformis*. *Genome Biol*. 2013;14:R60.
52. Sierro N, Battey JND, Ouadi S, Bakaher N, Bovet L, Willig A, et al. The tobacco genome sequence and its comparison with those of tomato and potato. *Nat Commun*. 2014;5:3833.
53. Hulse-Kemp AM, Maheshwari S, Stoffel K, Hill TA, Jaffe D, Williams SR, et al. Reference quality assembly of the 3.5-Gb genome of *Capsicum annuum* from a single linked-read library. *Hortic Res*. 2018;5:4.
54. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol Biol Evol*. 2016;33:1870–4.
55. Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol*. 1987;4:406–25.
56. Troadec JP, Gervois A, Oger L. Statistics of Voronoi cells of slightly perturbed face-centered cubic and hexagonal close-packed lattices. *Europhys Lett*. 1998;42:167–72.
57. Emanuelsson O, Nielsen H, Brunak S, von Heijne G. Predicting Subcellular Localization of Proteins Based on their N-terminal Amino Acid Sequence. *J Mol Biol*. 2000;300:1005–16.
58. Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, et al. SignalP 5.0 improves signal peptide predictions using deep neural networks. *Nat Biotechnol*. 2019;37:420–3.
59. Lescot M. PlantCARE, a database of plant cis-acting regulatory elements and a portal to tools for in silico analysis of promoter sequences. *Nucleic Acids Res*. 2002;30:325–7.
60. Liu W, Xie Y, Ma J, Luo X, Nie P, Zuo Z, et al. IBS: an illustrator for the presentation and visualization of biological sequences: Fig. 1. *Bioinformatics*. 2015;31:3359–61.
61. Untergasser A, Cutcutache I, Koressaar T, Ye J, Faircloth BC, Remm M, et al. Primer3—new capabilities and interfaces. *Nucleic Acids Res*. 2012;40:e115.
62. Bustin S, Huggett J. qPCR primer design revisited. *Biomol Detect Quantif*. 2017;14:19–28.
63. Terol J, Nueda MJ, Ventimilla D, Tadeo F, Talon M. Transcriptomic analysis of *Citrus clementina* mandarin fruits maturation reveals a MADS-box transcription factor that might be involved in the regulation of earliness. *BMC Plant Biol*. 2019;19:47.
64. Liu D, Shi L, Han C, Yu J, Li D, Zhang Y. Validation of Reference Genes for Gene Expression Studies in Virus-Infected *Nicotiana benthamiana* Using Quantitative Real-Time PCR. *PLoS One*. 2012;7:e46451.

Table

Table 1. *lDA-like* gene families in species of the Solanaceae family (genome localization from different Sol Genomics Network databases [47]). All prepropeptides are predicted to be localized in the secretory pathway according to TargetP [56] and SignalP-5.0 [57].

Gene name	Genome localization	Prepropeptide length (aa)	Predicted signal peptide length (aa)	PIP domain
NsylIDA1	Nsyl_KD945166.1:74265..74582 forward	105	39	PIPPSAPSKRHN
NsylIDA2	Nsyl_KD978144.1:88678..88980 reverse	100	32	PIPPSAPSKRHN
NsylIDA3	Nsyl_KD951180.1:40337..40579 forward	80	32	PIPPSAPSKRHN
NsylIDA4	Nsyl_KD977536.1:13349..13576 forward	75	22	PIPPSAPSQRHN
NsylIDA5	Nsyl_KD962079.1:38313..38564 forward	83	30	PIPASGPSRKHN
NtomIDA1	Ntom_KB972926.1:26032..26325 forward	97	37	PIPPSAPSKRHN
NtomIDA2	Ntom_KB954314.1:53025..53614 forward	96	32	PIPPSAPSKRHN
NtomIDA3	Ntom_KB969023.1:33965..34204 forward	79	31	PIPPSAPSKRHN
NtomIDA4	Ntom_KB956501.1:19193..19405 reverse	70	22	PIPPSAPSQRHN
NtomIDA5	Ntom_KB958630.1:30910..31161 reverse	83	30	PIPASGPSRKHN
NtabIDA1A	Ntab-BX_AWOK-SS18147:707412..707735 reverse	107	48	PIPPSAPSKRHN
NtabIDA1B	Ntab-BX_AWOK-SS9960:271769..272062 forward	97	37	PIPPSAPSKRHN
NtabIDA2A	Ntab-BX_AWOK-SS12153:24919..25221 reverse	100	32	PIPPSAPSKRHN
NtabIDA2B	Ntab-BX_AWOK-SS20685:67080..68370 reverse	96	32	PIPPSAPSKRHN
NtabIDA3A	Ntab-BX_AWOK-SS473:166199..166441 reverse	80	32	PIPPSAPSKRHN
NtabIDA3B	Ntab-BX_AWOK-SS2799:946688..946927 forward	79	31	PIPPSAPSKRHN
NtabIDA4A	Ntab-BX_AWOK-SS18001:26098..26325 forward	75	22	PIPPSAPSQRHN
NtabIDA4B	Ntab-BX_AWOK-SS12176:491822..492033 reverse	70	22	PIPPSAPSQRHN
NtabIDA5A	Ntab-BX_AWOK-SS18104:315608..315859 reverse	83	30	PIPASGPSRKHN
NtabIDA5B	Ntab-BX_AWOK-SS9524:125323..125574 forward	83	30	PIPASGPSRKHN
NbenIDA1A	Niben101Scf00570:62104..62373 reverse	90	36	PIPPSAPSK----
NbenIDA1B	Niben101Scf01338:640730..641035 forward	101	35	PIPPSAPSKRHN
NbenIDA2A	Niben101Scf23219:7370..7663 reverse	97	32	PIPPSAPSKRHN
NbenIDA2B	Niben101Scf03368:114599..114892 reverse	97	32	PIPPSAPSKRHN
NbenIDA3A	Niben101Scf18667:206436..206678 forward	80	32	PIPPSAPSKRHN
NbenIDA3B	Niben101Scf01180:267334..267576 reverse	80	32	PIPPSAPSKRHN
NbenIDA4	Niben101Scf19133:87532..87771 forward	79	25	PIPPSAPSQRHN
NbenIDA5A	Niben101Scf03848:699324..699575 forward	83	30	PIPASGPSRKHN
NbenIDA5B	Niben101Scf02135:404883..405122 reverse	79	26	PIPASGPSRKHN
SlycIDA1	SL3.0ch05:4200134..4200439 forward	101	36	PIPPSAPSKRHN
SlycIDA2	SL3.0ch06:38623220..38623453 forward	77	30	PIPPSAPSKRHN
SlycIDA3	SL3.0ch04:5799910..5800149 forward	79	27	PIPPSSPSKRHN
SlycIDA4	SL3.0ch07:58068277..58068558 reverse	93	34	PIPPSAPSKRCN
SlycIDA5	SL3.0ch05:1629558..1629893 forward	111	29	LIPPSGPSRRHN
SlycIDA6	SL3.0ch09:540104..540379 reverse	91	26	PIPPSAPCSRSS
SlycIDA7	SL3.0ch09:546577..546855 reverse	92	27	PLPPSAPCSRSS
SlycIDA8	SL3.0ch11:533813..534061 reverse	82	28	PIPASGPSRKHN
StubIDA1	PGSC0003DMB000000071:252349..252627 reverse	92	27	PIPPSAPCSRSS
StubIDA2	PGSC0003DMB000000131:879373..879621 forward	82	28	PIPASGPSRKHN
StubIDA3	PGSC0003DMB000000243:905236..905523 forward	95	29	PVPPSGPSRRHN
StubIDA4	PGSC0003DMB000000410:16621..16935 reverse	104	36	PIPPSAPSKRHN
StubIDA5	PGSC0003DMB000000420:159814..160050 forward	78	26	PIPPSSPSKRHN
StubIDA6	PGSC0003DMB000000461:377302..377535 forward	77	30	PIPPSAPSKRHN
StubIDA7	PGSC0003DMB000000592:149451..149714 forward	87	34	PIPPSAPSERCN
SmelIDA1	Sme2.5_00993.1:18248..18481 forward	77	30	PIPPSAPSKRHN

SmelIDA2	Sme2.5_04429.1:34294..34539 forward	81	28	PIPPSAPSLRHN
SmelIDA3	Sme2.5_04724.1:40347..40592 forward	81	27	PIPASGPSRKHN
SmelIDA4	Sme2.5_06686.1:19811..20078 forward	85	25	PIPPSAPSDRCN
SmelIDA5	Sme2.5_08129.1:7336..7444 forward	102	34	PIPPSGPSKRHN
SmelIDA6	Sme2.5_09763.1:10983..11228 reverse	81	26	PVPPSAPSDRCN
CalDA1	PepperUCD10Xch04:178438292..178438525 forward	77	27	PIPPSAPSKRHN
CalDA2	PepperUCD10Xch06:176812434..176812673 reverse	79	29	PIPPSAPSKRHN
CalDA3	PepperUCD10Xch11:6480406..6480714 forward	102	33	PIPPSGPSKRHN
CalDA4	PepperUCD10Xch11:4624209..4624499 forward	96	35	PIPPSAPSKRHN
CalDA5	PepperUCD10Xch11:6480457..6480714 forward	85	18	PIPPSGPSKRHN
CalDA6	PepperUCD10Xch11:27920042..27920356 reverse	104	24	PIPPSEPRHN

Additional Files

Additional file 1. Nicotiana *IDA1* promoters and coding sequences alignment.pdf. Alignment of the 5'-UTR sequences (500 bp) and the CDS of *Nben/IDA1A*, *Nben/IDA1B*, *Ntab/IDA1A*, *Ntab/IDA1B*, *Nsyl/IDA1* and *Ntom/IDA1* genes. Start codon is highlighted in green, and cis-acting regulatory elements are highlighted as follows: brown line, abscisic acid; blue line, methyl jasmonate; red line, auxins; grey line, drought.

Additional file 2. Solanaceae HSL family.pdf. *HAE*-like gene families in species of the Solanaceae family, (genome localization from different Sol Genomics Network databases [47]).

Additional file 3. HAE-like peptides phylogenetic tree.pdf. Circular phylogenetic tree of HAE-like peptides of *Arabidopsis thaliana* and several species of the Solanaceae family (*N. benthamiana*, *N. tabacum*, *N. sylvestris*, *N. tomentosiformis*, *S. lycopersicum*, *S. tuberosum*, *S. melongena* and *C. annuum*).

Additional file 4. Alignment IDA-like prepropeptides.pdf. Sequence alignment of IDA-like peptides from several species of the Solanaceae family (*N. benthamiana*, *N. tabacum*, *N. sylvestris*, *N. tomentosiformis*, *S. lycopersicum*, *S. tuberosum*, *S. melongena* and *C. annuum*) and from *Arabidopsis*. The CINEMA color scheme is used to inform about the chemical nature of the amino acid residues in the EPIP domain (blue, polar positive; red, negative; green, neutral; white, non-polar aliphatic; purple, ocher and yellow, aromatic residues).

Additional file 5. Gene expression raw and final processed data.xlsx.

Additional file 6. qPCR primers.pdf. Primers used for quantitative PCR analysis.

Figures




Figure 1

Unrooted circular phylogenetic tree of IDA-like prepropeptides of *Arabidopsis thaliana* and relevant species of the Solanaceae family such as *N. Sylvesteris*, *N. tomentosiformis*, *N. benthamiana*, *N. tabacum*, *S. lycopersicum*, *S. melongena*, *C. annuum* and *S. tuberosum*. Clade I shadowed in green color is divided in two subclades containing abscission-related genes (shadowed in lime green) and another one that grouped members of all eight Solanaceae species studied, as well as AtIDL8. The last subclade is subdivided in a group containing genes from Nicotiana, eggplant and pepper closed to SIIDA1, a tomato gene previously associated to abscission (shadowed in lime green) and other group (shadowed in light green) including from the Nicotiana, Solanum and Capsicum genera, with a small subdivision composed of AtIDL8 together with peptides from *S. lycopersicum* and *S. melongena*. Clade II (shadowed in light orange) includes exclusively Solanaceae genes. Clade III (shadowed in light gold) includes most of the *Arabidopsis* IDA-like family as well as Nicotiana and Solanum IDA-like genes. Bootstrap values are shown in each node.




Figure 2

Schematic representation of cis-acting regulatory elements of the promoter regions of the *N. benthamiana* IDA-like gene family and *Arabidopsis* AtIDA and AtIDL1. Light grey boxes represent 1000 bp long promoter regions while dark grey boxes represent the 5' part of the gene. In-silico searches of response elements to hormones related to abscission such as abscisic acid, methyl jasmonate, auxins or gibberellins, as well as response elements to biotic and abiotic stresses were performed. Response elements color scheme: red, auxins; brown, abscisic acid; blue, methyl jasmonate; green, gibberelins; grey, drought stress; white, defense response. The promoter region of NbenIDA3B is 493 bp long since the rest of the sequence is not available yet.




Figure 3

Expression patterns of IDA-like and HAE-like genes based on quantitative real-time PCR in several organs and tissues of *N. benthamiana* at different stages of corolla development. (A) Floral organs (sepals were removed), fruits and vegetative tissues utilized for gene expression analysis. The developmental stages of the corolla that were selected as a source of floral organs and fruits showed the corolla tube bulb above the calyx and the corolla limb closed (stage 1); fully elongated corolla tube and closed corolla limb

and anthers (stage 2); open flower, corolla limb fully expanded and anthers dehiscent (stage 4); lobes of the corolla limb curl inwards, significant loss of turgor of the corolla tube which easily detached from the receptacle (stage 5). Dash lines mark the tissue collected from the base of the corolla. (B) Gene expression levels in apical buds (AB), nodes (N), internodes (In), whole corolla (Co), the ensemble formed by the style and the stigma (S+S) and roots (R) relative to the lowest expression level of each gene. (C) Gene expression patterns in leaves, anthers and fruits. Gene expression levels were relative to that at the earliest developmental stage in every organ (YL, young leaf; An2, anthers at stage 2; Fr2, fruit at stage 2, respectively). (D) Gene expression patterns in the corolla base during corolla developmental stages 1, 2, 4 and 5: Gene expression levels were relative to that at stage 1. Gene expression levels were normalized with respect to those of NbenPP2A gene, applying the 2- $\Delta\Delta Ct$ method. Relative gene expression levels (log2 fold change) are given next to the color scale column. Upregulation and downregulation of gene expression is shown by red and blue color, respectively. Gene expression raw and processed final data are shown in Additional File 5.





Figure 4

Expression patterns of IDA-like genes based on quantitative real-time PCR in control and water stressed plants of *N. benthamiana*. General appearance of a well-watered plant (A) and plants subjected to water stress during 6 days (B) and during 8 days (C). Expression patterns in roots (R) and mature leaves (L) of water stressed plants during 6 or 8 days (D). Expression levels were calculated through the $2-\Delta\Delta Ct$ method, normalized to that of *NbenPP2A* gene and relative to gene expression in control (watered) plants.

For each stressed organ the appropriate reference was used (well-watered leaf or root). Expression levels relative to watered plants (fold change) are given next to the color scale. Red, white and blue colors indicate, respectively, gene induction (values over 1), unchanged (values close to 1) and repression (values under 1) all of them regarding to that of NbenPP2A gene in the relevant watered (control) organ. Gene expression raw and processed final data are shown in Additional File 5.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1NbenandAtIDApromotersandCDSs.pdf](#)
- [Additionalfile3HAElikepeptidesphylogenetictree.pdf](#)
- [AdditionalFile5Geneexpressiondata.xlsx](#)
- [Additionalfile6qPCRprimers.pdf](#)
- [Additionalfile2SolanaceaeHSLfamily.pdf](#)
- [Additionalfile4AlignmentIDAlikeprepropeptides.pdf](#)