

Surgical Management and Outcome of Large and Giant Intracranial Aneurysm

Mohammed M. Al Barbarawi (✉ mmalbarbarawi@just.edu.jo)

Jordan University of Science & Technology

Khaled Z. Alawneh

Jordan University of Science and Technology

Abdelhalim Z. Bqour

Ministry of Health

Walla Abdelaziz

Jordan University of Science and Technology

Suhair M. Qudsieh

Yarmouk University

Ala M. Barbarawi

Jordan University of Science & Technology

Sultan M. Jarrar

Jordan University of Science & Technology

Amer A. Jaradat

Jordan University of Science & Technology

Anwar Megdadi

Jordan University of Science & Technology

Owais Ghammaz

Jordan University of Science & Technology

Research Article

Keywords: large aneurysm, Giant intracranial aneurysm (GIA), Modified Rankin Scale (MRS), clipping, coiling

Posted Date: November 28th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3654403/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Large and giant intracranial aneurysms, exceeding 10 mm and 25 mm in diameter, respectively, are rare in neurosurgery. These lesions carry substantial morbidity and mortality risks, often causing mass effects, intracranial hemorrhage, stroke, or seizures. We present our experience with 54 patients diagnosed with these aneurysms at King Abdullah University Hospital. We conducted neurological assessments and preoperative evaluations using the modified Rankin Scale. Individualized treatment strategies, employing surgical techniques or endovascular embolization, were tailored to each patient. Surgical interventions, including direct clipping, aneurysmal wall resection, aneurysmorrhaphy with clipping, and cross-clamping (average duration: 20 minutes), were performed in 32 cases (59.2%). Endovascular coiling was utilized in 22 patients (40.8%), with 10 requiring flow diverter stents. Of the cases, 32 were giant aneurysms, and 22 were large aneurysms. The cohort included 28 females and 26 males, averaging 46.9 years. Aneurysms were predominantly in the anterior circulation (80.6%), with initial symptoms including mass effect (78.7%), cerebral ischemia (8.3%), intracranial hemorrhage (3%), and seizures (2.8%). Outcomes ranged from favorable to excellent in 70.4% of patients, while 18.5% experienced poor outcomes, primarily in posterior circulation aneurysms or due to poor preoperative conditions. The overall mortality rate was 11.1%, with eight fatalities. Management of giant intracranial aneurysms remains challenging, with higher morbidity and mortality rates compared to other neurosurgical conditions. No universally effective technique exists, emphasizing the importance of meticulous perioperative planning and surgical expertise. Further research and treatment advancements are needed to enhance the management of these complex aneurysms.

Introduction

Despite multiple advances in microsurgical and endovascular techniques, the treatment of large and giant intracranial aneurysms poses a significant challenge for vascular neurosurgeons. These rare lesions, accounting for approximately 5% of all intracranial aneurysms¹⁻³, predominantly occur in the anterior circulation and affect females more frequently than males. Patients with large and giant aneurysms often present with subtle symptoms, including slow cranial nerve dysfunction, particularly involving the optic and oculomotor nerves, progressive neurological dysfunction, chronic headache, or slowly progressive dementia. Intracranial hemorrhage, encompassing subarachnoid or intraventricular hemorrhage (SAH or IVH), is the second most common presenting symptom. Some patients may experience a sentinel headache before a major bleeding event, while others may deteriorate due to acute hydrocephalus. Seizures or ischemic strokes secondary to thrombus formation within the giant aneurysm, subsequent embolus, or vascular compromise due to aneurysm bulkiness are also observed in a few cases. Incidental discovery of these lesions is rare³⁻¹³. The pathogenesis of giant intracranial saccular aneurysms remains poorly understood. It is believed that these aneurysms evolve from smaller saccular aneurysms, which may or may not rupture. The continual hemodynamic changes and blood flow turbulence within the sac contribute to the enlargement of these small aneurysms, leading to wall vibration, degeneration, and further expansion. Repeated minor hemorrhages also contribute to the

thickening of the aneurysmal wall through fibrin or calcium deposition. Moreover, giant aneurysms often contain multi-laminated thrombus^{3,7,14-20}. In contrast, fusiform giant aneurysms affect the entire arterial wall, extending over a substantial length of the artery. This pathology is likely a consequence of atherosclerotic ectasia or dissection^{4,10,14,21-23}.

The management strategy for large or giant intracranial aneurysms varies^{1,3-7,9-12,20,22-25}. Ideally, the treatment aims to isolate the neck of the aneurysm from blood flow and its hemodynamic forces without sacrificing the main artery or the perforators, decompress the neural tissue, and ultimately restore adequate blood flow³. Treatment methods can be categorized as direct or indirect. Direct clipping of the aneurysm is the preferred technique when feasible, offering favorable short and long-term outcomes^{5,11}. However, direct clipping alone may not be possible in cases where the neck is large or heavily calcified. In such instances, alternative techniques are employed. Although these alternatives are considered substandard to direct clipping, they become necessary to avoid sacrificing the parent vessel, which could lead to devastating infarction or incomplete debulking of the aneurysmal mass. These options include microsurgical resection of the aneurysmal mass, aneurysmorrhaphy with arterial reconstruction through suturing the sac, or using a wrapping reinforcement technique. Indirect approaches involve trapping the aneurysm using clips or ligature above and below the aneurysm or obliterating the internal carotid artery (ICA) in the neck using the Hunterian technique if the patient can tolerate parent artery occlusion^{4,10,21}. Alternatively, extracranial-intracranial revascularization surgery utilizing a long saphenous vein graft bypass (JSVG)^{2,26}, or superficial temporal artery to middle cerebral artery anastomosis (STA-MCA) may be necessary to restore blood flow^{16,27}.

In recent years, the endovascular procedure has gained popularity in the treatment of large or giant aneurysms. Balloon trapping techniques, previously used by placing balloons above and below the aneurysm endovascularly^{13,28}, have been replaced by GDC embolization or flow diverter stent and coiling techniques due to the high morbidity associated with balloon trapping^{5,29-33}. The long-term efficacy of these endovascular techniques is still under evaluation. In certain cases, a combined surgical and endovascular approach may be employed as an alternative, with the endovascular technique serving as part of a multimodal treatment plan^{11,18,25}.

In this study, we focused on this variety of management strategies used by the senior author when treating giant aneurysms in different locations to improve the outcome and to reduce the morbidity and mortality rate related to the complexity of these lesions that can be attained by selecting the best suitable technique for a particular location.

Materials and methods

Patient Demographics and Clinical Materials:

This retrospective study was conducted at King Abdullah University Hospital and was approved by the Institutional Research Board (IRB) committee at Jordan University of Science and Technology (JUST). The study group comprised 54 cases who were treated for large and giant aneurysms between December 2005 and December 2020.

Inclusion Criteria:

Patients with large aneurysms measuring between 10 and 25 mm or giant aneurysms measuring more than 25 mm were included in the study. Only cases that received surgical or radiological interventions were included.

Exclusion Criteria:

Patients with small aneurysms were excluded from the study. Additionally, cases with large or giant aneurysms that remained untreated were also excluded.

Patient Assessment:

Each patient was individually assessed, considering factors such as age, presenting symptoms, exact size and location of each aneurysm, and the patient's pre-operative neurological condition. The Modified Rankin Scale (MRS) provided in Table 1 was utilized to determine the most suitable surgical approach for each patient. Table 2 provides information on the preoperative Modified Rankin Scale (MRS) grades of patients.

Table 1
Modified Rankin Scale (MRS)

Modified Rankin Scale (MRS)	
0	Asymptomatic
1	No significant disability despite symptoms, able to perform all routine tasks
2	Slight disability, cannot perform all previous tasks but can look after own affairs without help.
3	Moderate disability needs some help but can walk unaided
4	Moderate-severe disability, cannot walk without help, cannot attend to body needs without help
5	Severe disability, bedridden, incontinent, needs a constant nursing care
6	Dead

Table 2
Preoperative Modified Rankin (MRS) Scale
Grades of Patients

MRS	Pre-operative (n = 54)	%
Grade 0	13	23.4
Grade 1	19	34.6
Grade 2	14	26.2
Grade 3	3	3.7
Grade 4	2	1.9
Grade 5	3	5.6
Grade 6	0	0

Preoperative Investigations:

Preoperative investigations were conducted for all patients, except in cases requiring emergency interventions, following the established protocol in our institution. These investigations included various imaging modalities and tests to gather essential information:

Computed Tomography (CT): CT scans were performed without and with contrast to obtain a detailed understanding of the aneurysms. These scans helped determine their size, shape, and location, as well as identify any calcifications or recent hemorrhage.

Digital Subtraction Angiogram (DSA): A four-vessel DSA with a 3D study was conducted to provide comprehensive information about the aneurysm's site and morphology. DSA also facilitated the evaluation of adjacent arteries, the anatomy of the area, the condition of distal branches, the presence of collaterals, and the identification of laminated thrombus in giant aneurysms.

Magnetic Resonance Imaging (MRI): MRI scans from different angles were utilized to assess the relationship between the aneurysms and surrounding brain structures. This imaging technique also aided in identifying the layers of thrombus with varying ages of hemorrhage.

Patients with non-thrombosed large or giant aneurysms underwent additional elective tests, including internal carotid artery (ICA) balloon occlusion tests with single-photon emission computerized tomography (SPECT) scans, electroencephalography (EEG), and neuro-psychiatric consultations. These tests were performed to assess the cerebral blood flow (CBF) and cerebral perfusion state preoperatively, particularly when temporary or permanent occlusion of the parent artery was anticipated.

Operative Approaches:

For anterior circulation and posterior cerebral artery large and giant aneurysms, the cranial pterional or orbito-zygomatico-pterional transsylvian approach was selected. This approach provided a wide

exposure, minimized brain retraction, and offered a direct route to the objective. Vertebral artery and PICA aneurysms were managed using the far lateral suboccipital craniectomy approach.

Surgical Techniques:

Out of the 32 cases that underwent surgical intervention for large or giant aneurysms, 22 cases (64%) were directly clipped using SKS (Sundt-Kees) clips. In 11 patients (36%), resection of the aneurysmal wall and suturing of the aneurysm with clipping technique were performed.

Endovascular Embolization Techniques:

Endovascular embolization was performed in 22 cases. The procedure involved the patient lying supine on the X-ray table. The anesthesia administered varied based on the patient's medical condition, ability to follow instructions and the complexity of the case. Heparin, an anti-clotting medication, was administered to prevent blood clots during the procedure. To ensure stability, the patient's head is positioned securely to minimize movement during the endovascular embolization procedure.

Femoral artery access is obtained to perform endovascular embolization. Once the catheter is optimally placed, the neurointerventionalist administers contrast dye while capturing X-ray images in multiple dimensions. Dye injections are repeated until all necessary arteries are visualized, allowing for accurate measurements of the aneurysm, particularly the neck and aneurysm dome, as well as their relationship with the parent artery.

A microcatheter is then advanced through the standard catheter and into the aneurysm lumen. With continuous contrast injection, the microcatheter enables precise coil delivery and better assessment of the coils. Coiling was performed in four cases using this technique.

In cases where the aneurysm exhibits a wide neck or irregular shape, a stent may be utilized to assist in holding the coils in place. The stent is advanced through the catheter and positioned in the normal artery adjacent to the aneurysm.

In recent developments, a flow-diverter stent has been employed to isolate arterial blood flow from the intraluminal flow within the aneurysm. This leads to clot formation and subsequent shrinkage of the aneurysm. The flow-diverter stent was utilized in five patients with positive results.

Results

A total of 54 patients were included in this study. Table 3 provides information on the patients' demographics and location of aneurysms. The mean age of the patients was 46.9 years, ranging from 33 to 68 years. Out of the 54 patients, 26 (48%) were male, and 28 (52%) were female. The location of aneurysms was categorized into anterior circulation and posterior circulation. A total of 43 patients (79.6%) had aneurysms in the anterior circulation. Among these, the majority were located in the internal cerebral artery (ICA), accounting for 30 cases (55.6%). Within the ICA, the most common subtypes were ICA cavernous, with 17 cases (56.7%), followed by ICA paraclinoid with 8 cases (26.7%). Other subtypes

included ICA p-com (3 cases, 10%), and ICA terminal (2 cases, 6.7%). ACA had 3 cases (5.6%), and MCA had 10 cases (18.5%). Within the ACA subtype, 1 case (33.3%) involved the A1 segment, and 2 cases (66.7%) were related to the anterior communicating artery (A-com). A total of 11 patients (20.4%) had aneurysms in the posterior circulation. Among these, the most common location was the basilar artery (BASA), accounting for 6 cases (11.1%). Within the BASA subtype, 4 cases (66.7%) were located in the trunk, and 2 cases (33.3%) were located in the caput. Other locations in the posterior circulation included the posterior inferior cerebellar artery (PICA) with 1 case (1.9%), vertebral artery (VA) with 2 cases (3.7%), and posterior cerebral artery (PCA) with 2 cases (3.7%).

Table 3
Patients' demographics and location of aneurysms

	All Patients (n = 54)
Age, mean	46.9
Gender, n (%)	
Male	26 (48)
Female	28 (52)
Location, n (%)	
<i>Anterior Circulation</i>	43 (79.6)
ICA	30 (55.6)
ICA cavernous	17 (56.7)
ICA paraclinoid	8 (26.7)
ICA p-com	3 (10)
ICA terminal	2 (6.7)
ACA	3 (5.6)
A1 segment	1 (33.3)
A-com	2 (66.7)
MCA	10 (18.5)
<i>Posterior Circulation</i>	11(20.4)
PICA	1 (1.9)
BASA	6 (11.1)
BASA trunk	4 (66.7)
BASA caput	2 (33.3)
VA	2 (3.7)
PCA	2 (3.7)
<i>Abbreviations: ICA internal cerebral artery, MCA middle cerebral artery, ACA anterior cerebral artery, PCA posterior cerebral artery, BASA basilar artery, VA vertebral artery, PICA posterior inferior cerebellar artery, A-com anterior communicating artery.</i>	

Table 4 presents the relationship between aneurysm location and initial presenting symptoms. Among patients with aneurysms in the ICA 26 (86.7%) presented with symptoms related to mass effect, 2 cases (6.7%) presented with ICH, 1 case (3.3%) presented with stroke, and 1 case (3.3%) was incidentally

discovered. No cases of seizures were reported for ICA aneurysms. In patients with aneurysms in MCA, 5 (50%) presented with symptoms related to mass effect, 2 cases (20%) presented with ICH, 2 cases (20%) presented with stroke, and 1 case (10%) presented with seizures. No incidental findings were reported for MCA aneurysms. Among patients with ACA aneurysms, 1 case (33.3%) presented with symptoms related to mass effect, 1 case (33.3%) presented with ICH, and 1 case (33.3%) presented with seizures. No seizures or incidental findings were reported for ACA aneurysms. Among PCA aneurysms 1 case (50%) presented with symptoms related to mass effect, and 1 case (50%) presented with stroke. No cases of ICH, seizures, or incidental findings were reported for PCA aneurysms. Among cases with BASA aneurysms, 5 cases (83.3%) presented with symptoms related to mass effect, and 1 case (16.7%) presented with stroke. No cases of ICH, seizures, or incidental findings were reported for BASA aneurysms. Among VA aneurysms, 1 case (50%) presented with symptoms related to mass effect, 1 case (50%) presented with ICH, and no cases of stroke, seizures, or incidental findings were reported. There was only 1 case of PICA aneurysm, and the case presented with symptoms related to mass effect, and no other symptoms were reported. The distribution of initial presenting symptoms varied depending on the aneurysm location. Mass effect was the most common symptom observed across multiple locations, followed by ICH, stroke, seizures, and incidental findings.

Table 4
Relationship of aneurysm location with initial presenting symptoms

Initial presenting symptom, n (%)	Location of aneurysm						
	ICA (n = 30)	MCA (n = 10)	ACA (n = 3)	PCA (n = 2)	BASA (n = 6)	VA (n = 2)	PICA (n = 1)
Mass Effect	26 (86.7)	5 (50)	1 (33.3)	1 (50)	5 (83.3)	1 (50)	0 (0)
ICH	2 (6.7)	2 (20)	1 (33.3)	0 (0)	0 (0)	1 (50)	0 (0)
Stroke	1 (3.3)	2 (20)	0 (0)	1 (50)	1 (16.7)	0 (0)	1 (100)
Seizure	0 (0)	1 (10)	1 (33.3)	0 (0)	0 (0)	0 (0)	0 (0)
Incidental	1 (3.3)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)

Abbreviations: ICA internal cerebral artery, MCA middle cerebral artery, ACA anterior cerebral artery, PCA posterior cerebral artery, BASA basilar artery, VA vertebral artery, PICA posterior inferior cerebellar artery, ICH intracranial hemorrhage

Surgical interventions were performed in 59.2% (32 cases) of patients, either through direct clipping or resection of the aneurysmal wall and aneurysmorrhaphy with clipping. Endovascular coiling was conducted in 40.8% (22 cases) of patients, with 10 of them requiring a flow diverter stent.

In this study, all patients underwent post-operative angiogram on the first or second day to ensure complete obliteration of the aneurysm. Additionally, cerebral CT scans were performed on all patients on

the first day after the operation. Two patients required reoperation due to incomplete occlusion of the aneurysm. One patient underwent a redo procedure with direct clipping, resulting in successful aneurysmal stability. Another patient initially underwent endovascular coiling but had a large residual neck, leading to subsequent direct clipping after neurological improvement.

Although multi-modality techniques were employed to address large and giant aneurysms in different locations, the results still demonstrated a higher mortality and morbidity rate compared to smaller lesions. Among the patients, 70.4% achieved excellent to good results, with 33.3% being asymptomatic and classified as grade 0, 14.8% classified as grade I, and the rest falling into grade II. Poor outcomes were observed in 17.6% of patients, ranging from grade III to V. The overall mortality rate was 11.1%, with most deaths occurring in patients with basilar or posterior communicating giant aneurysms.

Ten patients experienced a poor neurological status, either due to devastating initial presentation or deterioration following treatment. Among these, eight patients developed ischemic changes, with four of them showing improvement and minor deficits. One patient who underwent excision of an MCA giant aneurysm experienced postoperative severe vasospasm, which responded well to angiographic intra-arterial nicardipine injection, leading to improvement without deficits. The remaining four patients ended up with a poor neurological condition. For instance, one patient with a left giant MCA aneurysm had right hemiplegia and aphasia that did not improve despite aggressive treatment due to MCA artery occlusion. Another patient with a right ICA cavernous aneurysm developed cavernous sinus syndrome and continued to decline without improvement. Four cases did not exhibit any neurological improvement after endovascular intervention, likely due to their poor preoperative status.

There was a case of a patient who experienced a fall in the hospital, resulting in a small intracerebral hemorrhage. However, this patient recovered well from the hemorrhage. Another patient presented with SAH grade V from a large ICA posterior communicating aneurysm. Initially, coiling of the aneurysm was performed, but one of the coils herniated through the dome, requiring surgical intervention. During the operation, the ICA was trapped for arteriotomy, and the herniated coil was successfully removed. The patient underwent craniotomy for clipping and eventually recovered with minimal hemiparesis, although the course was complicated.

Six patients in the study, unfortunately, passed away. Among them, two had BASA trunk or tip aneurysms treated with endovascular embolization. These patients had a stormy post-operative course and poor recovery, and complications such as massive pulmonary embolus and basilar thrombosis leading to pontine infarction were observed on angiogram and CT scan. Another patient with a left ICA paraclinoid aneurysm developed extensive infarction after endovascular embolization. Additionally, one patient experienced ICA dissection and diffuse bilateral cerebral infarction during the operation for clipping of a right ICA posterior communicating aneurysm, resulting in a worsened condition and eventual death.

Among the patients in the study, two individuals experienced an intracerebral hemorrhage. The first patient initially presented with headache and double vision, which was attributed to right abducent palsy. Unfortunately, the patient's condition worsened, and they bled in the ICU, resulting in a Glasgow Coma

Scale (GCS) score of 3. An urgent endovascular embolization procedure was performed to address the BASA trunk aneurysm, but regrettably, the patient did not recover and passed away shortly thereafter. The second patient, who had a large ICA posterior communicating aneurysm, also had a poor recovery following coiling and subsequently succumbed to the condition at a later stage.

In terms of surgical outcomes, our results indicate a tendency for better outcomes in patients with anterior circulation giant aneurysms. Among these patients, 75.6% achieved good outcomes, while 14% had poor outcomes. The mortality rate in this group was 5.9%. In contrast, for patients with posterior circulation giant aneurysms, 33.3% achieved excellent to good results, 33.3% had poor outcomes, and the mortality rate was notably higher at 28.6%.

When analyzing the outcomes based on aneurysm location, the best results were observed in patients with PCA, MCA, and ICA aneurysms, with success rates of 90%, 80%, and 72.9% respectively. Respectable outcomes were also achieved in patients who underwent clip and suture with debulking as well as direct clipping and endovascular embolization, with success rates of 80.5%, 66.7%, and 62.1% respectively. Patients with basilar aneurysms suffered from a higher morbidity and mortality rate, with 36.7% experiencing severe disability and a matching death rate. Interestingly, patients with PCA giant aneurysms displayed favorable outcomes in this study.

Discussion

Large and giant intracranial aneurysms are relatively rare, comprising only 5% of all intracranial aneurysms¹⁻³. They are more commonly observed in females in most series^{4,9,10,14,21,24,34}. These aneurysms typically present with symptoms related to compression of local neural structures due to their size^{13,28}. Unlike smaller aneurysms, they have a lower tendency to rupture. However, in some series, subarachnoid hemorrhage (SAH) has been reported as the initial presenting symptom in 25% of cases^{8,13,23,34}. The annual risk of bleeding in large and giant aneurysms is estimated to be around 3%, which is higher than that of smaller aneurysms, and this risk increases with the size of the aneurysm^{24,27}. Recent reports have shown that rebleeding from these aneurysms is common, with an incidence of 18.4% within 2 weeks and a high mortality rate of 33%, which is comparable to smaller aneurysms^{32,35}. Therefore, early identification and intervention are recommended for better outcomes^{12,20,25,36}.

Two types of giant intracranial aneurysms are recognized: saccular and fusiform. Saccular aneurysms are more common and predominantly affect the anterior circulation, often located at arterial bifurcations. These aneurysms may arise from the growth of smaller aneurysms that continue to expand due to blood pressure changes. Alternatively, they can develop from repeated minor bleeds that cause thickening of the arterial walls through fibrin deposits^{3,7,14-20}. Fusiform aneurysms, on the other hand, are less common and typically arise from the posterior circulation. They involve pathological changes affecting all layers of the arterial wall and can extend along the length of the artery. The etiology of fusiform aneurysms remains uncertain, although some authors propose atherosclerosis or arterial dissection as possible

causes^{14,21-23}. Pathogenesis studies by Drake et al demonstrated this type in only nine out of 120 patients with fusiform giant aneurysms^{14,21}.

The prognosis of untreated giant aneurysms is poor¹⁻³. If left untreated, these aneurysms continue to enlarge due to turbulence and wall vibration, leading to hemorrhage or progressive compression of blood vessels and neural structures. The mortality rate can be high within a few years after discovery of the aneurysm, reaching approximately 85% in some reports^{7,12}. Thromboembolic stroke or cerebral compression can cause death, but subarachnoid hemorrhage is the most common contributing factor to the high mortality rate^{1-3,6,7}, although rare cases of spontaneous remission have been reported³.

The treatment of giant aneurysms has significantly improved with advancements in microvascular neurosurgical techniques and approaches. These advancements have led to substantial results, with the majority of patients achieving excellent or good outcomes and a much lower mortality rate compared to the natural course of unoperated giant aneurysms in most series^{1,4,6,9,10,14,21,23,24,34}. Nevertheless, there are still a few patients who experience poor outcomes.

When dealing with these complex lesions, the treatment strategy aims to achieve several goals. Firstly, it involves isolating the aneurysm from the circulation. Secondly, it focuses on preserving the parent arteries and their perforators. Lastly, it aims to decompress the adjacent structures affected by the aneurysm. These goals can be accomplished through the utilization of appropriate surgical techniques by skilled and experienced surgeons, along with thorough preoperative planning^{7,8,12,13}.

In the past, Hunterian ligation was a commonly used method to treat giant aneurysms. This technique involved suturing the parent vessel to reduce blood flow to the aneurysm, relying on collateral circulation for nourishment^{4,8,10,13,21}. While effective in some cases, there were risks associated with this approach, including the possibility of delayed hemodynamic stroke or continued expansion and bleeding of the aneurysm, potentially caused by retrograde blood flow from opposite or posterior circulation or the external carotid artery^{10,14,21,26,37,38}.

Yasargil et al. introduced the extracranial-intracranial bypass technique by anastomosing the superficial temporal artery to the middle cerebral artery (STA-MCA)^{16,27}. Surgeons began combining STA-MCA anastomosis with Hunterian occlusion of the internal carotid artery (ICA) to treat giant aneurysms. Another technique involved using a long saphenous vein graft bypass (LSVGB) to restore blood flow between the common carotid artery and the intracranial segment of the ICA in cases of complete ICA occlusion^{2,26}. These techniques and their variants have proven effective in treating occlusive vascular diseases and giant aneurysms, particularly when parent vessel trapping is not tolerated by the patient^{6,26,33,36,37,39-42}. Revascularization procedures should be considered when reserve capacity is compromised, and the decision can be based on clinical aspects and preoperative physiological studies^{26,43,44}.

Direct surgical neck occlusion through clipping the aneurysm while preserving the original artery is considered the ideal treatment method, but its feasibility varies in different series due to factors such as a broad, bulbous, firm, or calcified neck^{1,4-6,8,10,11,13}. Additional clips may be required to reinforce the closing tension, with longer clips being less effective at the tip compared to shorter clips^{4,10,22,23}.

The basic principles of aneurysm surgery apply to giant aneurysms as well. Proximal and distal control of the parent artery, sharp dissection, and temporary clipping when necessary are crucial. Decompression of the aneurysm through thrombus removal or blood aspiration is important before applying the clip. Total excision of the aneurysm wall is unnecessary since these lesions are not neoplastic and do not recur once the sac is isolated from the bloodstream. The focus is on decompressing the aneurysm and isolating the neck from circulation^{1,3,6}.

Wrapping reinforcement has been successfully used by encasing the aneurysm with surgical gauze or plastic and applying purse string sutures around the base and dome to restrain it. However, concerns exist regarding the need for extensive dissection around the base of the brain and the possibility of continued aneurysm growth^{4,5,8,11,13}.

Trapping the aneurysm can be an effective technique in selected cases where sufficient collateral circulation is present and direct clipping is not feasible. Care must be taken to consider the branches arising from the trapped segment, and the sacrificed portion should be minimized by applying clips close to the aneurysm^{1,4,6,8,10,13}. Excision of the aneurysm with primary reconstruction could be an alternative for MCA aneurysms whereby this vessel can be repaired easily. It might be impossible to repair if the artery is friable.

Endovascular techniques have shown promising results in the treatment of inoperable giant aneurysms. Serbinenko introduced the detachable balloon occlusion method for carotid cavernous fistula and aneurysms, preserving the internal carotid artery^{13,28}. However, this approach is suitable only for selected cases. In 1989, Hilal reported successful endovascular occlusion of inoperable intracranial aneurysms using platinum coils with Dacron fibers, inducing intra-aneurysmal thrombotic occlusion^{5,31}. More recently, Guglielmi introduced intra-aneurysmal electro-thrombosis by injecting thrombogenic detachable platinum coils via an endovascular approach^{29,30}. While effective for smaller aneurysms, this technique is unsuitable for giant aneurysms due to their size, irregular shape, and presence of thrombus^{25,29}. However, there are varying reports on its efficacy^{9,18,32,33}.

According to a review by Campos et al., flow diversion is a newly developed endovascular technique that has shown success in treating giant aneurysms. Among 22 patients who underwent endovascular embolization, 10 required flow diverters. Previous studies have demonstrated that endovascular embolization with flow diversion is a safe and effective solution for large or giant, wide-neck internal carotid artery (ICA) aneurysms, with a primary effectiveness rate of 62.8%. The major complications or neurological death rate was found to be 8.3%, consistent with our own series⁴.

In certain cases, a combined surgical and endovascular approach may be beneficial^{11,35}. Some reports suggest endovascular ICA occlusion following extracranial-intracranial bypass for inoperable ICA aneurysms¹. Additionally, Lin et al. identified ACA aneurysms with recurrences after microsurgical clipping that were subsequently treated with flow diversion using PED, resulting in complete occlusion in 83% of cases. Therefore, flow diversion with PED is a viable option for treating recurrences after surgical clipping⁴. However, the use of a combined technique requires careful evaluation, individualized treatment planning, and close follow-up.

The selection of the explicit operative surgical technique is influenced by the aneurysm's anatomy, its relationship to adjacent brain structures, and the perioperative neurological state. The chosen operative approach aims to optimize field exposure with minimal brain retraction and increased bone removal, allowing for direct and unobstructed access to the aneurysm. The widely utilized orbitozygomatic-pterional trans-sylvian approach is commonly employed for treating anterior circulation and certain posterior circulation giant aneurysms^{28,30,39,43}. This approach offers multiple access points in various directions within the sylvian fissure, and in specific cases, additional bone removal, such as clinoidectomy, may be necessary to enhance exposure of the ICA giant aneurysms. The removal of the zygomatic arch provides a reasonable subtemporal access, with inferior retraction of the temporalis muscle. For addressing the tip of the basilar aneurysm and its branches, the subtemporal approach can provide appropriate access^{8,45}.

For lower basilar, vertebral artery, and PICA giant aneurysms, the far lateral suboccipital approach is considered a standard technique^{31,44}. This procedure requires removal of the rim of the foramen magnum, a portion of the occipital condyle, and the C1 arch up to the sulcus arteriosus to achieve ideal access.

Intraoperative cerebral protection plays a crucial role in the management of these complex lesions and can be facilitated prior to the commencement of the operation. Proper head positioning, extensive bone exposure, and minimal brain retraction can be achieved through CSF drainage and administration of Mannitol. Additional protection during the procedure involves maintaining mildly elevated blood pressure to enhance cerebral perfusion pressure, administering barbiturates, and implementing patient cooling techniques^{26,46}.

Barbiturate-induced EEG burst suppression is an effective method during cross-clamping. Barbiturates have the ability to reduce neuronal metabolic rate and increase brain tolerance to focal ischemia by mitigating substrate supply shortages. However, their role in global ischemia is not well-established^{26,38,45,47}. Administering barbiturates before applying the clip has proven to be particularly beneficial, as evidenced by Spetzler et al.'s animal model study, which confirmed adequate brain protection from focal ischemia^{45,47}.

Hypothermia serves as an adjunctive therapy with significant value in preventing ischemia by reducing cerebral metabolic rate of oxygen consumption through decreased cellular oxygen requirement for

enzymatic reactions. Attention must be given to the patient's clotting factors^{22,40}. The author believes that maintaining a hypothermia level of 32 degrees Celsius is efficient in neural protection with fewer complications. Furthermore, minimizing temporary clamping time and preserving existing collaterals from the external carotid artery are crucial for preventing ischemic insults²⁶.

The surgical outcomes for the management of large and giant aneurysms exhibit variability across different series^{1,4,6,7,9,10,21,24,34}. However, with meticulous perioperative planning, excellent to good outcomes can be achieved in the majority of patients, surpassing 80% success rates, and with a relatively low mortality rate ranging from approximately 5–15%^{1,5,6,11,34,41}. When comparing these surgical outcomes to the natural course of untreated giant aneurysms, it becomes evident that surgery offers superior results and significantly impacts the prognosis.

We firmly believe that the chosen technique for aneurysm occlusion, along with its location and the preoperative neurological status of the patient, greatly influences the overall outcome. It is our conviction that the best results can be achieved when a suitable surgical technique is selected for effectively isolating the giant aneurysm, accompanied by comprehensive perioperative planning.

Conclusion

The management of giant intracranial aneurysms necessitates an individualized approach, as there is no universally effective technique to address the complexity of these lesions. It is crucial that intervention, either directly or indirectly, is carried out at the earliest opportunity by a skilled vascular neurosurgeon who possesses knowledge of various surgical techniques. The primary goal should be to isolate the aneurysm neck from hemodynamic changes and decompress the sac while preserving the parent vessels. It is important to note that leaving behind the adherent wall does not pose any harm, as these lesions do not exhibit regrowth.

With the recent advancements in the field and comprehensive preoperative planning, coupled with the selection of an appropriate surgical technique and the implementation of adequate intraoperative brain protection measures, a favorable outcome can be achieved. By adhering to these principles, the overall prognosis for patients with giant intracranial aneurysms can be significantly improved. However, further research and advancements in surgical techniques are necessary to continue enhancing the management and outcomes of these challenging lesions.

Declarations

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have

agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Declaration of interest

None.

Acknowledgments

None.

Data Availability

The data used in this study is available upon request from the corresponding author.

Conflict of interest

The authors of this article have carefully considered any potential conflicts of interest and have found none to report. They have no relevant financial or non-financial interests that could impact the article's content, and they have no affiliations or involvement with any organizations with a financial or proprietary interest in the material discussed. The authors declare that they have no competing interests related to the manuscript's subject matter and certify that they have no ties to any entity that could present a conflict.

References

1. Lawton MT, Spetzler RF. Surgical management of giant intracranial aneurysms: experience with 171 patients. *Clin Neurosurg.* 1995;42:245-266.
2. Locksley HB, Sahs AL, Sandler R. SECTION III: Subarachnoid Hemorrhage Unrelated to Intracranial Aneurysm and A–V Malformation. *J Neurosurg.* 1966;24(6):1034-1056. doi:10.3171/jns.1966.24.6.1034
3. Morley TP, Barr HWK. Giant Intracranial Aneurysms: Diagnosis, Course, and Management. *Neurosurgery.* 1969;16(Supplement 1):73-94. doi:10.1093/neurosurgery/16.CN_suppl_1.73
4. Campos JK, Ball BZ, Cheaney II B, et al. Multimodal management of giant cerebral aneurysms: review of literature and case presentation. *Stroke Vasc Neurol.* 2020;5(1):22-28. doi:10.1136/svn-2019-000304
5. Guglielmi G. History of Endovascular Endosaccular Occlusion of Brain Aneurysms: 1965–1990. *Interventional Neuroradiology.* 2007;13(3):217-224. doi:10.1177/159101990701300301
6. Lawton MT, Hamilton MG, Morcos JJ, Spetzler RF. Revascularization and Aneurysm Surgery: Current Techniques, Indications, and Outcome. *Neurosurgery.* 1996;38(1):83-94. doi:10.1097/00006123-199601000-00020

7. Østergaard JR. Risk factors in intracranial saccular aneurysms Aspects on the formation and rupture of aneurysms, and development of cerebral vasospasm. *Acta Neurol Scand.* 1989;80(2):81-98. doi:10.1111/j.1600-0404.1989.tb03847.x
8. Sano K. Temporo-polar Approach to Aneurysms of the Basilar Artery at and Around the Distal Bifurcation: Technical Note. *Neurol Res.* 1980;2(1):361-367. doi:10.1080/01616412.1980.11739588
9. Taki W, Nishi S, Yamashita K, et al. Selection and combination of various endovascular techniques in the treatment of giant aneurysms. *J Neurosurg.* 1992;77(1):37-42. doi:10.3171/jns.1992.77.1.0037
10. Drake CG. Giant Intracranial Aneurysms: Experience with Surgical Treatment in 174 Patients. *Neurosurgery.* 1979;26(Supplement 1):12-95. doi:10.1093/neurosurgery/26.CN_suppl_1.12
11. Hosobuchi Y. Direct surgical treatment of giant intracranial aneurysms. *J Neurosurg.* 1979;51(6):743-756. doi:10.3171/jns.1979.51.6.0743
12. Sughrue ME, Saloner D, Rayz VL, Lawton MT. Giant intracranial aneurysms: evolution of management in a contemporary surgical series. *Neurosurgery.* 2011;69(6):1261-1270; discussion 1270-1. doi:10.1227/NEU.0b013e31822bb8a6
13. Song J, Mao Y. Giant Aneurysm Management. In: ; 2022:133-160. doi:10.1007/978-3-030-87649-4_7
14. Drake CG, Peerless SJ. Giant fusiform intracranial aneurysms: review of 120 patients treated surgically from 1965 to 1992. *J Neurosurg.* 1997;87(2):141-162. doi:10.3171/jns.1997.87.2.0141
15. Nagahiro S, Takada A, Goto S, Kai Y, Ushio Y. Thrombosed growing giant aneurysms of the vertebral artery: growth mechanism and management. *J Neurosurg.* 1995;82(5):796-801. doi:10.3171/jns.1995.82.5.0796
16. Yaşargil MG. *Microsurgery Applied to Neurosurgery.* Academic Press; 1969.
17. Sundt TM, Sundt TM. Principles of preparation of vein bypass grafts to maximize patency. *J Neurosurg.* 1987;66(2):172-180. doi:10.3171/jns.1987.66.2.0172
18. Suzuki J, Ohara H. Clinicopathological study of cerebral aneurysms. *J Neurosurg.* 1978;48(4):505-514. doi:10.3171/jns.1978.48.4.0505
19. Yong-Zhong G, August H, Alphen M van. Pathogenesis and histopathology of saccular aneurysms: review of the literature. *Neurol Res.* 1990;12(4):249-255. doi:10.1080/01616412.1990.11739952
20. Ferguson GG. Physical factors in the initiation, growth, and rupture of human intracranial saccular aneurysms. *J Neurosurg.* 1972;37(6):666-677. doi:10.3171/jns.1972.37.6.0666
21. Drake CG, Peerless SJ, Ferguson GG. Hunterian proximal arterial occlusion for giant aneurysms of the carotid circulation. *J Neurosurg.* 1994;81(5):656-665. doi:10.3171/jns.1994.81.5.0656
22. Spetzler RF, Hadley MN, Rigamonti D, et al. Aneurysms of the basilar artery treated with circulatory arrest, hypothermia, and barbiturate cerebral protection. *J Neurosurg.* 1988;68(6):868-879. doi:10.3171/jns.1988.68.6.0868
23. Spetzler RF, Riina HA, Lemole GM. Giant Aneurysms. *Neurosurgery.* 2001;49(4):902-908. doi:10.1097/00006123-200110000-00022

24. Whittle IR, Dorsch NW, Besser M. Giant intracranial aneurysms: diagnosis, management, and outcome. *Surg Neurol.* 1984;21(3):218-230. doi:10.1016/0090-3019(84)90191-5
25. Giannotta SL, Litofsky NS. Reoperative management of intracranial aneurysms. *J Neurosurg.* 1995;83(3):387-393. doi:10.3171/jns.1995.83.3.0387
26. Lougheed WM, Marshall BM, Hunter M, Michel ER, Sandwith-Smyth H. Common carotid to intracranial internal carotid bypass venous graft. *J Neurosurg.* 1971;34(1):114-118. doi:10.3171/jns.1971.34.1.0114
27. Wiebers DO, Whisnant JP, Sundt TM, O'Fallon WM. The significance of unruptured intracranial saccular aneurysms. *J Neurosurg.* 1987;66(1):23-29. doi:10.3171/jns.1987.66.1.0023
28. Serbinenko FA. Balloon catheterization and occlusion of major cerebral vessels. *J Neurosurg.* 1974;41(2):125-145. doi:10.3171/jns.1974.41.2.0125
29. Guglielmi G, Viñuela F, Dion J, Duckwiler G. Electrothrombosis of saccular aneurysms via endovascular approach. *J Neurosurg.* 1991;75(1):8-14. doi:10.3171/jns.1991.75.1.0008
30. Guglielmi G, Viñuela F, Sepetka I, Macellari V. Electrothrombosis of saccular aneurysms via endovascular approach. *J Neurosurg.* 1991;75(1):1-7. doi:10.3171/jns.1991.75.1.0001
31. Heros RC. Lateral suboccipital approach for vertebral and vertebrobasilar artery lesions. *J Neurosurg.* 1986;64(4):559-562. doi:10.3171/jns.1986.64.4.0559
32. Khurana VG, Piepgras DG, Whisnant JP. Ruptured giant intracranial aneurysms. Part I. A study of rebleeding. *J Neurosurg.* 1998;88(3):425-429. doi:10.3171/jns.1998.88.3.0425
33. Knuckey NW, Haas R, Jenkins R, Epstein MH. Thrombosis of difficult intracranial aneurysms by the endovascular placement of platinum-Dacron microcoils. *J Neurosurg.* 1992;77(1):43-50. doi:10.3171/jns.1992.77.1.0043
34. Sundt TM, Piepgras DG. Surgical approach to giant intracranial aneurysms. *J Neurosurg.* 1979;51(6):731-742. doi:10.3171/jns.1979.51.6.0731
35. Khayata MH, Spetzler RF, Mooy JJA, Herman JM, Rekate HL. Combined surgical and endovascular treatment of a giant vertebral artery aneurysm in a child. *J Neurosurg.* 1994;81(2):304-307. doi:10.3171/jns.1994.81.2.0304
36. Piepgras DG, Khurana VG, Whisnant JP. Ruptured giant intracranial aneurysms. Part II. A retrospective analysis of timing and outcome of surgical treatment. *J Neurosurg.* 1998;88(3):430-435. doi:10.3171/jns.1998.88.3.0430
37. Regli L, Piepgras DG, Hansen KK. Late patency of long saphenous vein bypass grafts to the anterior and posterior cerebral circulation. *J Neurosurg.* 1995;83(5):806-811. doi:10.3171/jns.1995.83.5.0806
38. Roski RA, Spetzler RF, Nulsen FE. Late complications of carotid ligation in the treatment of intracranial aneurysms. *J Neurosurg.* 1981;54(5):583-587. doi:10.3171/jns.1981.54.5.0583
39. Smith RR, Al-Mefty O, Middleton TH. An Orbitocranial Approach to Complex Aneurysms of the Anterior Circulation. *Neurosurgery.* 1989;24(3):385-391. doi:10.1227/00006123-198903000-00013

40. Spetzler R, Chater N. Microvascular bypass surgery. *J Neurosurg.* 1976;45(5):508-513. doi:10.3171/jns.1976.45.5.0508
41. Sundt TM, Piepgras DG, Fode NC, Meyer FB. Giant intracranial aneurysms. *Clin Neurosurg.* 1991;37:116-154.
42. Sundt TM, Piepgras DG, Marsh WR, Fode NC. Saphenous vein bypass grafts for giant aneurysms and intracranial occlusive disease. *J Neurosurg.* 1986;65(4):439-450. doi:10.3171/jns.1986.65.4.0439
43. Hakuba A, Liu S shan, Shuro N. The orbitozygomatic infratemporal approach: A new surgical technique. *Surg Neurol.* 1986;26(3):271-276. doi:10.1016/0090-3019(86)90161-8
44. Herkes GK, Morgan M, Grinnell V, et al. EEG monitoring during angiographic balloon test carotid occlusion: experience in sixteen cases. *Clin Exp Neurol.* 1993;30:98-103.
45. Selman WR, Spetzler RF, Roski RA, Roessmann U, Crumrine R, Macko R. Barbiturate coma in focal cerebral ischemia. *J Neurosurg.* 1982;56(5):685-690. doi:10.3171/jns.1982.56.5.0685
46. Streefkerk HJN, Van der Zwan A, Verdaasdonk RM, Mansvelt Beck HJ, Tulleken CAF. Cerebral Revascularization. In: ; 2003:145-225. doi:10.1007/978-3-7091-0641-9_3
47. SHAPIRO HM. BARBITURATES IN BRAIN ISCHAEMIA. *Br J Anaesth.* 1985;57(1):82-95. doi:10.1093/bja/57.1.82