The strategy of coping with stress and selected health related behavioural patterns of demobilized battle participants

Dorota Ortenburger

Jan Długosz University in Czeszochowa

Jacek Wąsik (✉ jwasik@konto.pl)

Jan Długosz University <https://orcid.org/0000-0002-6285-7283>

Olga Andriejczuk

Lesi Ukrainki University in Luck

Research article

Keywords: stress, trauma, coping strategies, pain, rehabilitation, psychology.

Posted Date: August 15th, 2019

DOI: <https://doi.org/10.21203/rs.2.12944/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: The aim of the herein paper is to express knowledge of the role of ties (relations) between mental factors such as trust in your own possibilities of coping with pain and availing of the active strategies of coping with stress, while also the degree of readiness to limit stress with the aid of the use of alcohol. **Methods:** The research covered the group of 38 demobilized ATO participants (21-46 years) undergoing treatment and rehabilitation in the Hospital. A measurement was conducted with the aid of certified psychological questionnaires and from medical documentation. **Results:** A correlation occurred between the active strategy of coping with the stress felt and the strategy of coping with pain of the ICS type – Internal control of pain ($r=0.71$; $p<0.05$), as well as the negative correlation between the readiness to utilize stimulants with the aim of easing stress and the strategy of coping with pain of the ICS type ($r=-0.58$; $p<0.05$). **Conclusion:** The ways of coping with stress and pain by the hospitalized participants of combat action may be significant factors (of a psychological nature) that have an impact on their readiness to undertake behaviour that is significant for their health (such as alcohol and others) with the aim of reducing the symptoms of nervous tension felt.

Background

The participants of combat action with relation to the tasks executed are exposed to experiencing traumatic events. This is first and foremost connected with the possibility of endangering health, while also confrontation with their own deaths and the deaths of other people. Such people face extraordinarily difficult challenges such as continuing actions aimed at the realization of the stipulated aims in situations of intensive stress. This type of events bring multiple negative consequences in the sphere of mental and physical health. They may be the source of anxiety, unease, insomnia, irritability and anger, symptoms of depression, or becoming involved in behaviour that is not beneficial to their health, such as among other things, the intensive use of alcohol [1].

Several types of strategies of coping with strong nervous tension and its repercussions are distinguished. These are as follows: active coping with stress (planning possible actions, searching for various possibilities of help, positive re-evaluation), helplessness (use of psychoactive substances, alcohol, ceasing of remedial action), searching for support (emotional and instrumental), avoidance behaviour (including: dealing with something else, denial, venting emotion) [2].

The ability to cope with stress is a significant part of the processes of self-regulation and the skill of adjusting the strategies of procedure to the requirements of the situation which a person is facing. This is also strongly individualized, which is illustrated by both scientific research and colloquial observations that highlight how people differ from each other [3].

Using alcohol (or other stimulants) with the aim of easing the effects of experiences of a traumatic nature constitutes a problem for a multitude of people who are involved in actions that endanger their safety, to a large extent soldiers and non-professional veterans of combat action [4]. Searching for relief

and improvement of self-esteem by means of alcohol is treated as a non-constructive strategy in easing the psycho-physiological effects of stress, which improves self-esteem only for a brief period, after which it deepens the existing health, financial and social problems [5]. It is the source of problems over a long-term perspective as these phenomena may occur in interactions based on feedback, e.g. the abuse of alcohol may lead to conflicts – and conflicts increase exasperation, irritability and anger.

Some researchers note that not only soldiers, but other groups also face the consequences of post-traumatic stress. The occurrence of similar mental consequences have been noted in a group of medical rescuers, which is confirmed by research conducted in several countries [6]. Approximately half of British rescuers experienced intrusive thoughts, feelings of irritation, problems with sleeping and distancing themselves from other people [2]. Research conducted by Polish researchers indicates that almost half of them felt the impact of traumatic experiences in their professional lives and family lives. Simultaneously, the majority of medical rescuers analysed admit that extreme stress during rescue missions does not overcome their possibilities of adaptation and does not cause a reduction in their readiness to participate in future rescue missions [6].

The participants of combat action are frequently exposed to pain. Coping with pain and its consequences are of significance for our state of health [7]. Pain belongs to the strong stressors, in which the following components permeate: sensory, affective, motivational, cognitive [1,12]. Research illustrates that in terms of people who want to reduce the unpleasant experiences, there is a risk of developing an addiction [4,6]. The relation between health and coping with long-term stress and pain is widely analysed and widely documented in psychoneuroimmunology [8,9].

In accordance with the holistic paradigm that underlies psychoneuroimmunology, pain connects the physical and mental experiences that have a reciprocal impact on each other. People who are unable to deal with stress feel the acute effects of chronic pain and those who as a result of psycho-physical overload develop symptoms of depression [10]. Hence, the abuse of, for instance alcohol during feelings of helplessness may constitute a way to change the state of awareness with the aim of reducing the feeling of pain which ultimately makes depression more profound.

The differences in terms of attitudes towards pain (from complete helplessness to various types of feelings of effectiveness) are visible in the form of convictions – W (trust in personal management), L (faith in medical care), P (conviction about the case) [3]. These differences are particularly visible in situations of experiencing pressure, whose extreme example is participation in combat action.

Participation in this type of action, as well as the injuries incurred lead to the fact that the adjustment of the soldier to normal everyday life constitutes a challenge. A person in such situations instigates behaviour that is directed at coping with the consequences of traumatic events. Hence, it is important to become familiar with the relations and dependencies of the strategies of coping with stress and pain with the aim of preparing therapeutic programs. With relation to this fact, an attempt at an analysis was undertaken as to whether the specified type of strategies of coping with stress and pain are associated

with the use of stimulants (primarily alcohol) among the group of hospitalized participants of combat action.

With regard to this fact, we concentrated on becoming familiar with the relations between the following:

1. strategy of coping with pain type “W”, while releasing stress by means of various stimulants.
2. availing of active constructive remedial strategies that are aimed at reducing the symptoms of nervous tension and the frequency of using alcohol.
3. Strategies of coping with pain in patients and their diagnosis in the sphere of the basis of the ailments experienced.

The following hypothesis was put forward: trust in your own possibilities to cope with stress and the use of active strategies to cope with stress are associated with a lower consumption of alcohol.

Methods

Subject

The research covered the group of 38 demobilized ATO participants undergoing treatment and rehabilitation in the Hospital aged between 21 and 46 years. A significant age differentiation (age span between the oldest and the youngest participant of defence action) was revealed from the specifics of the analysed group. The cause of this case was the fact that some participants were called to arms “ad hoc”, or at once and the vast majority of these will not become regular soldiers. Thus, there was no regulation from top-down in terms of the age as in typical military formations. A measurement was conducted with the aid of certified psychological questionnaires (Beliefs About Pain Control Questionnaire and Mini-Cope Inventory). Additional, medical data relating to the analysed people, including the state of health of the patients was acquired from medical documentation.

Protocol

A measurement was conducted with the aid of certified psychological questionnaires: (Beliefs About Pain Control Questionnaire – BPCQ (13); Data relating to mental functioning was acquired by the application of the Mini-Cope Inventory, which constitutes a fully-fledged standardized version [14]. A shorter version of Cope (The Coping Orientations to Problems Experienced) was applied due to the reactions of those under analysis, who displayed exasperation and a reluctance towards time-consuming completion of expanded questionnaires. Additional information referring to the analysed people in the state of health of patients was acquired via medical documentation. This data encompassed a diagnosis of the type of pain that occurs in patients with particular attention paid to whether the pain experienced had features of neuropathic pain.

Statistical analysis

Analysis was conducted on the regression between the use of stimulants by soldiers (dependent variable) and two dependent variables (active strategy of coping with stress and the type of conviction defined as: "Internal control of pain". For the statistical analyses, the value of $\alpha = 0.05$ was considered as significant.

Ethics

Research includes human data, which have been performed in accordance with the Declaration of Helsinki and have been approved by the Lesya Ukrainka Eastern European National University, Lutsk, Ukraine Bioethics Commission (no.2/2018). Before interviews, the nature and the purpose of the study were explained, and full confidentiality was assured to all participants. All participants were informed about their right not to participate in the study and gave their oral consent before the study.

Results

In Fig. 1, a graphic interpretation has been presented of the dependencies between three chosen variables: the application of the active strategy of coping with stress (ACS), coping with pain of the type ICP (Internal control of pain), as well as the readiness to consume alcohol as an antidote for nervous tension (ACS). In Table 1, the values of the coefficient correlations of the same variables have been illustrated. Table 2 contains the average values of the distribution of data relating to the intensity of the particular types of convictions regarding the control of pain; according to the criteria: 1. the existence of neuropathic pain, 2. pain of a mixed nature– probably (not fully confirmed) damage to the nerve underlying chronic pain, 3. lack of neuropathic pain.

Table 1. Table of correlations between chosen variables $p < 0.05$.

Variable	ASS	ICP	ACS
ASS	-	0.71	-0.37
ICP	0.71	-	-0.58
ACS	-0.37	-0.58	-

ASS –Active strategy coping with stress, ICP – Internal control of pain, ACS –Alcohol Coping strategy

Table 2. Average values of distribution of data relating to the intensification of particular types of convictions with regard to controlling pain (in scores).

No.	Nature (type) of pain	ICP		DCP		PCE	
		M	SD	M	SD	M	SD
1	Neuropathic pain. Confirmed damage to the nerve underlying chronic pain	16.17	2.31	12.72	2.91	13.17	3.50
2	Pain of a mixed nature. Not fully confirmed (probable) damage to nerve underlying chronic pain	20.79	2.35	12.74	4.53	13.58	3.12
3	Lack of pain of a neuropathic nature (the lack of confirmation of damage to nerve)	24.91	3.75	12.91	6.47	14.36	3.98
Total number of patients		20.00	4.31	12.77	4.45	13.60	3.43

Discussion

The acquired results indicate that the group of hospitalized participants of anti-terrorist action encompassed within the research, who are people of a lower value of the ICP indicator and who do not avail of the active strategies may be more susceptible to "escaping to alcohol" (Fig. 1).

It was revealed that a correlation ($r=0.71$; $p<0.01$) exists between the active strategy of coping with the stress experienced (ASS) and the strategy of coping with pain of the ICP type.

This provides the argument which is supportive of the view that the analysed people who learnt to cope with various forms of stress themselves are "vaccinated" against other factors that evoke a stressful reaction, i.e. pain. At the basis of this covariance there may be a more profound neurophysiological mechanism that is described on the basis of the theory of the Gate Control Theory.

In accordance with the concept entitled "*Gate Control Theory*," the response to the unpleasant stimulants does not only depend on the nature of the stimulant, but also to a large extent on the individual features of the nervous system of the particular person [1]. At a psychological level, this may signify that the basis of coping with stress and pain may involve a common feature which is defined as "proactivity" in the sphere of individual responses to situations that arouse anxieties in a person, which may be indicated by the results of previous research [15].

Active coping with stress is indicated by a moderate correlation ($r=-0.37$; $p<0.01$) with the willingness to consume alcohol (and use of other stimulants) with the aim of easing stress. However, we do not know how many people indicated a tendency to consume alcohol prior to conscription and how many developed this tendency during military service, or in the aftermath of defence action. On the basis of the herein research, we may merely state that we have noted covariance (negative sign of correlation) between active strategies and readiness to reach out for alcohol as a way to reduce the unpleasant symptoms of excessive nervous tension.

The lower tendency towards the consumption of alcohol co-exists with the greater conviction that it is possible to restrict the experienced pain effectively by yourself ($r=-0.58$, $p<0.01$, as well as a higher level of activity of coping with stress ($r=-0.37$, $p<0.01$). It is worth noting that in the case whereby the acquired results illustrated a contrasting tendency, namely, if the increase in the readiness to consume alcohol as a remedial action for stress is accompanied by growth in the value of the variable of "active coping with stress", this could certify to the fact that behaviour involving the consumption of alcohol is treated by the analysed people as their own "actions" (alternate). This would also find clarification in the light of subject-related literature, in which there is mention of the phenomenon that exists in people, namely, "the illusions that enable them to live peacefully". Health psychology describes these types of mechanisms involving self-deceit in broader terms [16].

The data presented in Table 2 illustrates that patients differ with regard to diagnosis with relation to the presence of neuropathic pain. This indicates that the level of trust in your own possibilities in the sphere of coping in the case of the feeling of pain turns out to be an issue that requires greater attention in the

situation whereby among the hospitalized soldiers where pain occurred, this accompanies injuries associated with damage to nerves (neuropathic pain). Neuropathic pain leads to the case whereby the therapy is long-lasting and less efficient than the treatment of other types of pain [14]. This results from its specifics and the time of duration.

Pain of a neuropathic nature, which emerges as a result of damage to a nerve or its sheath, may involve many years of healing in terms of the tissue which evokes negative emotions [17,18]. Due to the increasing difficulty in the active constructive coping in terms of the situation of pressure, in this group there may be an increased level of risk of the use of stimulants with the aim of reducing the unpleasant symptoms of excessive nervous tension (irritability, frustration) [17,19]. The results presented are encouraging in terms of continuing the research in its current form.

The acquired data illustrates that the applied method may have a practical application in terms of screening and prophylactic tests, as well as the evaluation of the effectiveness of the therapy [13]. We are of the opinion that following the conclusion of the therapy for the hospitalized participants of combat action, it is justifiable to conduct similar research with the aid of the same methods. This would provide information about the effectiveness of the therapeutic impact. The acquired data provides arguments in order to take account of the psychological aspect in a broader sense in terms of the possible therapeutic effects during the course of medical rehabilitation for veterans of combat action. This is illustrated by the research results of a wide section of the research program which encompassed the participants of ATO actions [20,21].

Conclusions

The ways of coping with stress and pain on the part of the hospitalized participants of combat action may be factors (psychological independent variables) that potentially influence the readiness to undertake behaviour that is significant for their health (such as use of alcohol, etc.) with the aim of reducing the symptoms of nervous tension experienced.

This is backed up by the acquired results that indicate covariance (negative) between the level of trust in their own possibilities of coping with pain and the readiness to use alcohol as a strategy for easing the effects of stress. A negative correlation was also noted between the degree of using active strategies of coping with stress and the readiness to use alcohol.

With regard to the fact that undertaking constructive ways of coping with the mental effects of traumatic experiences increases the chances of recovering, it is worth taking these results into account, particularly during the course of preparing programs for patients who have incurred damage to their health during combat action and who do not indicate psychologically active remedial action.

Abbreviations

ACS: Alcohol Coping strategy; ASS: Active strategy coping with stress; ATO: Anti-Terrorist Operation; BPCQ: About Pain Control Questionnaire; Cope: Coping Orientations to Problems Experienced; Type W: trust in personal management; Type L: faith in medical care; Type P: conviction about the case.; ICP: Internal control of pain

Declarations

Ethics approval and consent to participate

Research includes human data, which have been performed in accordance with the Declaration of Helsinki and have been approved by the Lesya Ukrainka Eastern European National University, Lutsk, Ukraine Bioethics Commission (no.2/2018). Before interviews, the nature and the purpose of the study were explained, and full confidentiality was assured to all participants. All participants were informed about their right not to participate in the study and gave their oral consent before the study.

Consent for publication

Not applicable.

Availability of data and material

The datasets analyzed during the current study are not publicly available but are available from the corresponding author on reasonable request.

Competing interests

All other authors declare that they have no competing interests.

Funding

This research received no external funding.

Authors' contributions

DO formulated the research questions, designed the study, interpreted the results and wrote the first draft of manuscript. JW revised the manuscript and interpreted the results and managed the whole project and contributed in all steps. OA collected the data.

Acknowledgements

We acknowledge the time and effort of all participants in this study.

References

1. Melzack R. Pain and the neuromatrix in the brain. J Dent Educ 2001; 65(12):1378-82.

2. Carver C, Scheier M, Weintraub J. Assessing coping strategies: a theoretically based approach. *J Pers Soc Psychol* 1989;56(2):267-83.
3. Strong J, Ashton R, Chant D. The measurement of attitudes towards and beliefs about pain. *Pain* 1992;48:227-36.
4. Williams J, McDevitt-Murphy M, Murphy J, Crouse E. Deployment Risk Factors and Postdeployment Health Profiles Associated With Traumatic Brain Injury in Heavy Drinking Veterans. *Military Med* 2012;177(7):789–96.
5. Sheridan C, Radmacher S. *Health Psychology: Challenging the Biomedical Model*, International Edition, John Wiley and Sons, United Kingdom, 1992
6. Oginska-Bulik N. Negative and positive effects of traumatic experiences in a group of emergency workers – The role of personal and social resources. *Medycyna Pracy* 2013;64(4):463–72.
7. Villemure C, Bushnell *Cognitive modulation of pain: how do attention and emotion influence pain processing? Pain* 2002; 95:195– 99.
8. Dunn A, Wang J, Ando T. Effects of cytokines on cerebral neurotransmission. Comparison with the effects of stress. *Adv Exp Med Biol* 1999;461:117–27. doi: 10.1007/978-0-585-37970-8
9. Schleifer S, Marbach J, Keller S. Psychoneuroimmunology: Potential Relevance to Chronic Orofacial Pain. *Anesthesia Progress* 1990;37(2-3):93–8.
10. Brian L. Stress, Depression and the Activation of the Immune System, *World J Biol Psych* 2000;1(1):17-25.
11. Schedlowski M, Schmidt R. Stress and the immune system. *Naturwissenschaften*. 1996;83(5):214-20.
12. Strong J, Ashton R, Chant D. The measurement of attitudes towards and beliefs about pain. *Pain* 1999;48:227-36.
13. Skevington S. A standardized scale to measure beliefs about controlling pain (BPCQ): a preliminary study. *Psychology and Health* 1990;4:221-32.
14. Carver C. You want to measure coping but your protocol's too long: Consider the Brief COPE. *Inter J Behav Med* 1997;4:92-100.
15. Proios I, Batsiou S, Proios M. Volition profile of people with physical disability. *Phys Activ Rev* 2018;6:151-60.
16. Taylor S, Armor D. Positive illusions and coping with adversity. *J Pers* 1996;(64)4:873-98.
17. Egli M, Koob G, Edwards S. Alcohol dependence as a chronic pain disorder. *Neuroscience and Biobehavioral Rev* 2012;36(10):2179–92.
18. Zale E, Maisto S, Ditre J. Interrelations between Pain and Alcohol: An Integrative Review. *Clinical Psych Rev* 2015, 37, 57–71.
19. Ortenburger D, Rodziewicz-Gruhn J, Wąsik J, Marfina O, Polina N. Selected problems of the relation between pain-immunity and depression, *Phys Activ Rev* 2017;5:74-7

20. Ortenburger D, Wąsik J, Tsos A, Bielikowa N, Andrijchuk O, Indyka Forms of expressing anger in hospitalised Ukrainian post-deployed service members. *Ann Agric Env Med* 2018;25(1):172–5.
21. Grygus I. The role of physical activity in the rehabilitation of patients suffering from mild persistent bronchial asthma. *Phys Activ Rev* 2017;5:155-66

Figures

Figure 1

Graphic presentation of dependency between three variables analysed: ASS – Active strategy coping with stress, ICP – Internal control of pain, ACS – Alcohol coping strategy