Survival and prognostic factors analyses in malignant giant cell tumor of bone

Jin Zhang

Tianjin Tumor Hospital

Xin Wang

Southern Medical University Nanfang Hospital

Feng Lin

Tianjin Tumor Hospital

Guijun Xu

Tianjin Tumor Hospital

Haixiao Wu

Tianjin Tumor Hospital

Jincai Duan

Tianjin Tumor Hospital

Min Mao

First affiliated hospital, Army Medical University

Chao Zhang (✉ drzhangchao@tmu.edu.cn)

Tianjin Tumor Hospital <https://orcid.org/0000-0001-7096-8488>

Research article

Keywords: giant tumor cancer of bone, malignant, survival, prognosis, SEER

Posted Date: August 15th, 2019

DOI: <https://doi.org/10.21203/rs.2.12960/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background The characteristics and survival in patients with malignant giant tumor cancer of bone (GCTB) were not investigated thoroughly due to the limited population. We evaluated the issues based on a large cohort in the National Cancer Institute's Surveillance, Epidemiology, and End Results (SEER) database. **Methods** Patients who were diagnosed with malignant GCTB from 1975 to 2016 were extracted from the SEER database. The overall survival (OS) was calculated by Kaplan–Meier analysis and the inter-group difference was tested by log-rank test. Univariate and multivariate Cox proportional hazard regression were conducted to identify the independent survival factors. **Results** A total of 325 patients with malignant GCTB were included. The overall 1-, 5-, and 10-year survival rates were 94.3% (95% CI: 91.7-96.8), 82.3% (95% CI: 77.9-86.6), and 80.1% (95% CI: 75.4-84.7), respectively. In the univariate analysis, age older than 34 years, grade IV, T2/3 stage, M1, distant and surgery of the primary site. Multivariate Cox regression showed the poor survival in patients with age older than 34 years (hazard ratio (HR) =3.68, 95% CI: 2.06-6.57, P<0.001), T2 stage (HR=4.96, 95% CI: 1.57-15.63, P=0.006), distant tumor (HR=3.02, 95% CI: 1.32-6.92, P=0.009), and the extra-skeletal sites (HR=8.84, 95% CI: 2.89-27.07, P<0.001), respectively. **Conclusions** This large population-based series described the clinical characteristics of the malignant GCTB. Age, T2, distant tumor and extra-skeletal sites were determinant survival factors of the patients with malignant GCTB.

Background

Giant cell tumor of bone (GCTB) was rare. It was reported that GCTB accounted for approximately 5% to 6% of primary bone tumors [1]. The annual incidence was reported to be 1.7 (1.4-1.9) per million [2]. Metaphyseal of long bones were the major affected sites and many studies were conducted to investigate both the clinical characteristics [3,4] and treatments[5].

Most of the GCTB were thought to be pathologically benign cases. Some tumors showed aggressive features such as cortical and subchondral involvement, reactive bony zone infiltration, and soft tissue mass [6,7]. Recently, a study including four large series identified 92 cases with malignant GCTB in 2,315 patients, the incidence was from 1.1% to 11.3% and the cumulative incidence was 4.0% [8]. As to the malignant GCTB, both survival and prognostic prediction were important for individualize treatment generation. But the issues were not intensively studied due to the lack of long-term follow-up clinical data [9]. Itkin and his colleagues retrospectively retrieved the data of 242 patients from 26 studies with a median follow-up of 6.9 years [10]. They reported an estimated overall survival rate of 86.9%. Few studies investigated the prognostic factors of patients with malignant GCTB.

The Surveillance, Epidemiology, and End Results (SEER) database is one of the largest cancer databases in USA. Based on this database, a previous study estimated the survival among 117 malignant GCTB from 1975 to 2004. The prognostic factors for poor survival included older patients and the distant metastasis at diagnosis [11].

This database has been extended and some new factors were added including the TNM stages, new version of AJCC grade, insurance status, and records of some specific metastatic organs. Update of the new data should be re-evaluated to provide further understanding on the malignant GCTB. In the present study, based on the updated data from the database, we attempted to evaluate the survival of patients with malignant GCTB and to identify the prognostic factors.

Patients And Methods

Data Source

The SEER database was used to extract information of patients with malignant GCTB by the SEER*Stat Software version 8.3.5 (<https://seer.cancer.gov/seerstat/>) (Information Management Service, Inc. Calverton, MD, USA). According to the ICD-O-3/WHO2008, patients who were diagnosed as malignant GCTB (Code: 9250/3: giant cell tumor of bone, malignant) between 1975 and 2016 were collected. Patients were excluded if they were diagnosed at autopsy or via death certificate. The malignant GCTB located in extra-skeletal sites such as kidney, lung, pancreas and thyroid were excluded.

Statistical analysis

According to the records and distribution of patients in the SEER database, all the patient-related variables were selected as following: age (<34 or ≥ 34 years); gender (male and female); marital status (unmarried or married); race (white, black, Indian/Alaska Native (IA) and Asian or Pacific Islander (API)); primary site (long bones of lower limb with associated joints, long bones of upper limb with scapula and associated joints, pelvis and spine, irregular bone and other site out of bone); tumor grade (I, II, III, or IV); T stage (T1, T2, T3); N stage (N0 and N1); M stage (M0 and M1); SEER historic stage (localized, regional, and distant); years interval of diagnosis (1975-1984, 1985-1994, 1995-2004, and 2005-2016); and the surgical treatments of primary site (none or yes). The patients without clear records were classified as the unknown group for the corresponding parameter.

Continuous data was described as mean $\pm$ standard deviation. Differences between groups were analyzed with univariate analysis. Categorical data was presented as the number and percentage, and the difference was evaluated by Pearson chi-square test. The 1-, 5-, and 10-year overall survival (OS) were calculated by the Kaplan–Meier method and the log-rank test was used to evaluate the difference. The prognostic factors ($P < 0.05$) in the univariate analysis were collected for multivariate Cox regression model. To evaluate the influence of the extra-skeletal sites on survival, a sensitivity analysis was performed by including or excluding these cases. All statistical analyses were performed using SPSS 23.0 (IBM Corporation, Armonk, NY, USA) and MedCalc 15.2.2. Two-sided P -values < 0.05 were considered statistically significant.

Results

Patient characteristics

The selection process of the patients was illustrated in Figure 1. According to the inclusion criteria, 334 patients with malignant GCTB were initially selected from 1975 to 2016. After excluding one case which was diagnosed via death certificate and eight patients with extra-skeletal GCTB, 325 patients were finally included. Table 1 showed the detail clinicopathological features of the patients. The median age of the patients was 34 years and a female predomination (52.3%) was observed. More tumors located in the long bones and associated joints (64.6%), especially in the lower limb (48.6%). More than 50% of patients had no clear records in tumor grade, T stage, N stage and M stage. Twenty-two patients showed distant metastasis including lung metastasis (10/22, 45.45%), bone metastasis (5/22, 22.73%) and the other metastasis (7/22, 31.82%). As to SEER historic stage, localized and regional stages were the main types occupying 61.2% of all the patients while 13.5% of patients were at distant stage. Surgery was performed on 53.85% of all the patients.

There eight patients were diagnosed with extra-skeletal GCTB. They were recorded from 1987 to 2016 and five patients were female cases. Two of the patients were diagnosed around 20 years old, while the others were older than 50 years. As to the primary site, three cases were found in pancreas, one in kidney, one in thyroid, one in mediastinum, one in lung and one subcutaneous case. There was not clear record of tumor grade in four patients, one at grade I, two at grade III and one at grade IV. Two patients had long survival (135 and 136 months), one died at 24 months after diagnosis, and five patients died within 7 months.

Survival analysis and prognostic factors identification

The overall survival of the entire cohort was illustrated in Figure 2. The overall 1-, 5-, and 10-year survival rates were 94.3% (95% CI: 91.7-96.8), 82.3% (95% CI: 77.9-86.6), and 80.1% (95% CI: 75.4-84.7), respectively. Detail OS for patients with malignant GCTB originated from bone in each characteristic was summarized in Table 2.

The results of univariate and multivariate Cox hazard regression were summarized in Table 3. Lower OS was noticed in age older than 34 years ($P<0.001$), grade IV ($P=0.041$), T2/3 ($P<0.001$), unknown N stage ($P=0.019$), M1 stage ($P<0.001$), the SEER historic distant stage and surgery of the primary ($P=0.05$) under univariate analysis. Multivariate Cox regression showed the poor survival in patients with age older than 34 years (hazard ratio (HR) =3.68, 95% CI: 2.06-6.57, $P<0.001$), T2 stage (HR=4.96, 95% CI: 1.57-15.63, $P=0.006$), and distant tumor (HR=3.02, 95% CI: 1.32-6.92, $P=0.009$), respectively. The sensitivity analysis suggested that the primary malignant GCTB located in the extra-skeletal sites was associated with poor survival (HR=8.84, 95% CI: 2.89-27.07, $P<0.001$). Analysis including the extra-skeletal cases did not significantly affect the results (supplementary table).

Discussion

In the present study, with the large cohort of malignant GCTB, the 5- year OS was 82.3% (95% CI: 77.9-86.6). This rate was consistent with the reported rate of 86.9% by Itkin et al[10] and 80% by Domovitev et

al[12]. The prognostic factors were revealed, and poor survival was correlated with the following factors including age older than 34 years, higher T stage, distant tumor and the extra-skeletal sites.

Similar with the survival trend from previous SEER cohort analysis [11], in the present study, poor survival can be found in older patients and those with SEER historic distant stage. The records of metastatic sites were added in SEER dataset since 2010. Compared with the 5-year OS of 81.3% for patients without metastasis, it significantly decreased to 53.9 % for those with metastasis. Previous studies reported that the incidence of lung metastasis in benign giant cell tumor was around 3.5% [13,14] while approximately 8% in the malignant cases [15,16]. In our study, 14 out of 89 cases extracted from 2010 to 2016 were recorded as M1 status and lung was the major metastatic site (10 out of 89, 11.2%). The higher incidence of lung metastasis may be caused by regular screening. More lung metastases were reported to develop in the early time after initial diagnosis of GCTB and after recurrence [16,17]. Screening of possible metastasis should be especially performed at these situations.

The classification of TNM was added in the SEER database since 2004 and the results showed a trend of poor survival in higher T stage or higher N stage. The multivariate Cox regression suggested that higher T stage was one of the prognostic factors. Tumor size as the independent risk factor for lung metastasis was previously found [18]. Based on the Campanacci classification, which has been widely accepted in evaluating giant cell tumor, grade III represented the aggressive type. A series of studies reported Campanacci grade III as the risk factor for tumor recurrence [19,20] and lung metastasis [21,22]. The recurrence was also reported to be a risk factor for resulting in metastasis [21,22]. Therefore, T stage and/or Campanacci classification played important role in evaluating the survival of patients with giant cell tumor.

To our knowledge, it was the first time to study the extra-skeletal malignant GCTB. Based on our results, the extra-skeletal site was associated with the poor survival. Only two patients lived longer than 11 years while five patients died less than seven months. Therefore, attention should be paid on the patients with the lesion in the extra-skeletal site. Limited by few available cases, further analysis on revealing potential prognostic factors in such population cannot be conducted. More cases from multiple centers will be needed.

To overcome giant cell tumor, either resection with reconstruction or intralesional curettage has been widely accepted [23-25]. The surgery with/without radiotherapy did not significantly improve the survival in the previous SEER study [11]. Similarly, the surgery was not an independent survival factor of the patients with malignant GCTB in the present study. The effect of surgery on the survival of malignant GCTB was different from other malignant cancers such as colorectal cancer [26] and ovarian cancer[27]. The difference may be caused by significantly higher survival in the giant cell tumor than other malignant cancers.

Undoubtedly, there are some limitations in the present study. More than 50% of patients had no clear records in tumor grade, T stage, N stage and M stage. More clear data was needed to further reveal the effect of AJCC grade and TNM stage on survival. Limited by the absent records of surgery in the SEER

database, further analysis of the correlation between surgery and recurrence or metastasis cannot be performed. The results of the present study need further external validation.

Conclusions

In conclusion, based on the large population from the SEER database, the clinical characteristics and prognostic factors among the patients with malignant GCTB were analyzed. Age older than 34 years, T2 stage, SEER historic distant stage and the extra-skeletal sites were negatively associated with survival.

Abbreviations

GCTB: giant cell tumor of bone; SEER: Surveillance, Epidemiology, and End Results database; AJCC: American Joint Committee on Cancer; OS: overall survival; HR: hazard ratio; CI: confidence interval.

Declarations

Acknowledgements

Not Applicable

Authors' contributions

JZ, MM and CZ made contributions to the conception of this study. GX, HW and JD made contributed to acquisition of data and interpretation of results. XW and FL performed statistical analysis. FL, HW and JD prepared the tables and figures. JZ and MM mainly prepared the manuscript. GX and CZ reviewed and reedited the manuscript. All authors read and approved the final manuscript.

Funding

The present study was sponsored by Natural Science Foundation of China (81702161, 81802508), Natural Science Foundation of Tianjin Science and Technology Committee China (17JCQNJC11000, 15JCQNJC12400), Top talent training program of the first affiliated hospital of PLA Army Medical University (SWH2018BJKJ-12), the Doctor Start-up Grant of Tianjin Medical University Cancer Institute and Hospital (B1711), Laboratory of Tumor Immunology and Pathology (Army Medical University), Ministry of Education (2017jszl01).

The Natural Science Foundation of China, Natural Science Foundation of Tianjin Science and Technology Committee China and Doctor Start-up Grant of Tianjin Medical University Cancer Institute and Hospital mainly provided sponsor on data collection and manuscript writing. The top talent training program of the first affiliated hospital of PLA Army Medical University and Laboratory of Tumor Immunology and Pathology (Army Medical University) mainly sponsored the data analysis.

Availability of data and materials

The datasets analyzed during the current study are derived from the Surveillance, Epidemiology, and End Results (SEER) database in USA.

Ethics approval and consent to participate

The patients' consent for the data in the public SEER database was not required due to the data because cancer is a reportable disease in every state of the United States. The present study complied with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Consent for publication

Not applicable.

Conflict of interest

All the authors report no conflicts of interest in this work.

References

1. Lopez-Pousa A, Martin BJ, Garrido T, Vazquez J. Giant cell tumour of bone: new treatments in development. *Clin Transl Oncol*. 2015;17(6):419-30.
2. Verschoor AJ, Bovee J, Mastboom M, Sander DP, Van De Sande M, Gelderblom H. Incidence and demographics of giant cell tumor of bone in The Netherlands: First nationwide Pathology Registry Study. *Acta Orthop*. 2018;89(5):570-74.
3. Errani C, Ruggieri P, Asenzio MA, Toscano A, Colangeli S, Rimondi E, et al. Giant cell tumor of the extremity: A review of 349 cases from a single institution. *Cancer Treat Rev*. 2010;36(1):1-07.
4. Zou C, Lin T, Wang B, Zhao Z, Li B, Xie X, et al. Managements of giant cell tumor within the distal radius: A retrospective study of 58 cases from a single center. *J Bone Oncol*. 2019;14(100211).
5. van der Heijden L, Dijkstra PD, Campanacci DA, Gibbons CL, van de Sande MA. Giant cell tumor with pathologic fracture: should we curette or resect? *Clin Orthop Relat Res*. 2013;471(3):820-29.
6. Present D, Bertoni F, Hudson T, Enneking WF. The correlation between the radiologic staging studies and histopathologic findings in aggressive stage 3 giant cell tumor of bone. *Cancer-Am Cancer Soc*. 1986;57(2):237-44.
7. Chakarun CJ, Forrester DM, Gottsegen CJ, Patel DB, White EA, Matcuk GJ. Giant cell tumor of bone: review, mimics, and new developments in treatment. *Radiographics*. 2013;33(1):197-211.
8. Palmerini E, Picci P, Reichardt P, Downey G. Malignancy in Giant Cell Tumor of Bone: A Review of the Literature. *Technol Cancer Res Treat*. 2019;18(1078107648).
9. Gong L, Liu W, Sun X, Sajdik C, Tian X, Niu X, et al. Histological and clinical characteristics of malignant giant cell tumor of bone. *Virchows Arch*. 2012;460(3):327-34.
10. Itkin B, Straminsky S, De Ronato G, Lewi D, Marantz A, Bardach A. Prognosis of metastatic giant cell tumor of bone in the pre-denosumab era. A systematic review and a meta-analysis. *Jpn J Clin Oncol*.

2018;48(7):640-52.

11. Beebe-Dimmer JL, Cetin K, Fryzek JP, Schuetze SM, Schwartz K. The epidemiology of malignant giant cell tumors of bone: an analysis of data from the Surveillance, Epidemiology and End Results Program (1975-2004). *Rare Tumors*. 2009;1(2):e52.
12. Domovitev SV, Healey JH. Primary malignant giant-cell tumor of bone has high survival rate. *Ann Surg Oncol*. 2010;17(3):694-701.
13. Junior RC, Pereira MG, Garcia PB, Santos PA, Cavalcanti AS, Meohas W. Epidemiological study on giant cell tumor recurrence at the Brazilian National Institute of Traumatology and Orthopedics. *Rev Bras Ortop*. 2016;51(4):459-65.
14. Niu X, Zhang Q, Hao L, Ding Y, Li Y, Xu H, et al. Giant cell tumor of the extremity: retrospective analysis of 621 Chinese patients from one institution. *J Bone Joint Surg Am*. 2012;94(5):461-67.
15. Jiang N, Qin CH, Tan CX, Wen SF, Ma YF, Dong F, et al. A retrospective analysis of 140 patients with giant cell tumor in the extremity: a multicenter study based on four hospitals in South China. *Cancer Epidemiol*. 2013;37(3):294-99.
16. Rosario M, Kim HS, Yun JY, Han I. Surveillance for lung metastasis from giant cell tumor of bone. *J Surg Oncol*. 2017;116(7):907-13.
17. Dominkus M, Ruggieri P, Bertoni F, Briccoli A, Picci P, Rocca M, et al. Histologically verified lung metastases in benign giant cell tumours—14 cases from a single institution. *Int Orthop*. 2006;30(6):499-504.
18. Wang B, Chen W, Xie X, Tu J, Huang G, Zou C, et al. Development and validation of a prognostic index to predict pulmonary metastasis of giant cell tumor of bone. *Oncotarget*. 2017;8(64):108054-63.
19. Urakawa H, Yonemoto T, Matsumoto S, Takagi T, Asanuma K, Watanuki M, et al. Clinical outcome of primary giant cell tumor of bone after curettage with or without perioperative denosumab in Japan: from a questionnaire for JCOG 1610 study. *World J Surg Oncol*. 2018;16(1):160.
20. Cheng DD, Hu T, Zhang HZ, Huang J, Yang QC. Factors Affecting the Recurrence of Giant Cell Tumor of Bone After Surgery: A Clinicopathological Study of 80 Cases from a Single Center. *Cell Physiol Biochem*. 2015;36(5):1961-70.
21. Yang Y, Huang Z, Niu X, Xu H, Li Y, Liu W. Clinical characteristics and risk factors analysis of lung metastasis of benign giant cell tumor of bone. *J Bone Oncol*. 2017;7(23-28).
22. Kito M, Matusmoto S, Ae K, Tanizawa T, Gokita T, Kobayashi H, et al. Pulmonary metastasis from giant cell tumor of bone: clinical outcome prior to the introduction of molecular target therapy. *Jpn J Clin Oncol*. 2017;47(6):529-34.
23. Pazionis TJ, Alradwan H, Deheshi BM, Turcotte R, Farrokhyar F, Ghert M. A Systematic Review and Meta-Analysis of En-Bloc vs Intralesional Resection for Giant Cell Tumor of Bone of the Distal Radius. *Open Orthop J*. 2013;7(103-08).
24. Zhang S, Zhang J, Wang X. Comparison of tumor curettage and resection for treatment of giant cell tumor of the bone around the knee joint. *Pak J Med Sci*. 2016;32(3):662-66.

25. Charest-Morin R, Fisher CG, Varga PP, Gokaslan ZL, Rhines LD, Reynolds JJ, et al. En Bloc Resection Versus Intralesional Surgery in the Treatment of Giant Cell Tumor of the Spine. *Spine (Phila Pa 1976)*. 2017;42(18):1383-90.

26. Guo X, Zhang C, Ma W, Tian F, Xu G, Han X, et al. Patterns of bone metastases in newly diagnosed colorectal cancer: a real-world analysis in the SEER database. *Int J Colorectal Dis*. 2019;

27. Zhang C, Guo X, Peltzer K, Ma W, Qi L, Zhang Y, et al. The prevalence, associated factors for bone metastases development and prognosis in newly diagnosed ovarian cancer: a large population based real-world study. *J Cancer*. 2019;10(14):3133-39.

Tables

Table 1. Description of the patients with malignant giant cell tumor of bone.

Subject characteristics	Number	Present (%)
Sex		
Male	155	47.69
Female	170	52.31
Age		
<34	159	48.92
≥34	166	51.08
Marital status		
Unmarried	136	41.85
Married	162	49.85
Unknown	27	8.31
Insurance recode		
Uninsured	113	34.77
Insured	8	2.46
Unknown	204	62.77
Race		
White	241	74.15
Black	41	12.62
AI	8	2.46
API	28	8.62
Unknown	7	2.15
Year of diagnosis		
1975-1984	31	9.54
1985-1994	42	12.92
1995-2004	90	27.69
2005-2016	162	49.85
Primary site		
Long bones of lower limb and associated joints	158	48.62
Long bones of upper limb and associated joints	52	16.00
Pelvis and spine	59	18.15
Irregular bone, short bone, and flat bone	47	14.46
Unknown	9	2.77
Grade		
Grade I	9	2.77
Grade II	18	5.54
Grade III	12	3.69
Grade IV	31	9.54
Unknown	255	78.46
T stage 2004+		
T1	61	18.77
T2	41	12.62
T3	4	1.23
Unknown	219	67.38
N stage 2004+		
N0	149	45.85
N1	1	0.31
Unknown	175	53.85
M stage 2004+		
M0	142	43.69

M1	22	6.77
Unknown	161	49.54
SEER historic stage		
Localized	111	34.15
Regional	88	27.08
Distant	44	13.54
Unknown	82	25.23
Surg (prim) 1998+		
None	59	18.15
Yes	175	53.85
Unknown	91	28.00

Note: IA: Indian/Alaska Native; API: Asian or Pacific Islander

Table 2. The overall survival (OS) in patients with malignant giant cell tumor of bone diagnosed between 1975 and 2016.

Factors	1-year OS, % (95% CI)	5-year OS, % (95% CI)	10-year OS, % (95% CI)	<i>P</i> <i>value</i>
Sex				0.397
Male	95.4 (92.0-98.7)	83.1 (76.7-89.3)	79.2 (72.1-86.2)	
Female	93.3 (89.5-97.0)	81.6 (75.5-87.5)	80.8 (74.6-86.9)	
Age				< 0.001
<34	98.7 (96.9-100.0)	92.4 (88.1-96.6)	NA	
≥34	89.9 (85.2-94.5)	72.4 (65.1-79.5)	67.9 (60.1-75.6)	
Marital status				0.307
Unmarried	92.2 (87.6-96.8)	79.5 (72.3-86.5)	75.5 (67.7-83.2)	
Married	95.6 (92.4-98.7)	82.9 (76.7-88.9)	NA	
Unknown	96.2 (88.8-100.0)		87.0 (73.1-100.0)	
Insurance recode				0.251
Uninsured	92.5 (87.5-97.4)	76.7 (67.7-85.5)	74.0 (63.9-83.9)	
Insured		NA	NA	
Unknown	95.0 (92.0-97.9)	84.7 (79.7-89.6)	82.6 (77.3-87.8)	
Race				0.560
White	93.5 (90.3-96.6)	82.5 (77.4-87.5)	80.1 (74.6-85.5)	
Black	97.5 (92.7-100.0)	81.9 (69.7-93.8)	NA	
AI		75.0 (45.0-100.0)	NA	
API	89.0 (77.2-100.0)	80.3 (64.7-95.6)	73.0 (53.4-92.2)	
Unknown				
Year of diagnosis				0.124
1975-1984	90.3 (79.9-100.0)	87.1 (75.3-98.7)	NA	
1985-1994	97.4 (92.4-100.0)	92.3 (83.9-100.0)	87.2 (76.7-97.5)	
1995-2004	94.4 (89.6-99.1)	85.1 (77.6-92.4)	82.7 (74.7-90.5)	
2005-2016	93.5 (89.6-97.3)	76.3 (69.0-83.5)	74.9 (67.2-82.4)	
Primary site				0.333
Long bones of lower limb and associated joints	94.7 (91.1-98.2)	84.0 (78.0-89.9)	81.3 (74.7-87.7)	
Long bones of upper limb and associated joints	98.0 (94.0-100.0)	86.9 (77.1-96.5)	84.3 (73.5-94.8)	
Pelvis and spine	94.8 (89.1-100.0)	74.9 (62.9-86.6)	NA	
Irregular, short, and flat bone	91.2 (82.9-99.3)	83.9 (72.9-94.7)	81.0 (69.0-92.7)	
Unknown	77.8 (50.6-100.0)	64.8 (32.5-96.5)	NA	
Grade				< 0.001
Grade I		NA	NA	
Grade II	94.4 (83.8-100.0)	81.9 (63.2-100.0)	NA	
Grade III	91.7 (76.1-100.0)	60.0 (27.3-92.1)	40.0 (1.2-78.0)	

Grade IV	71.8 (55.0-88.3)	53.8 (34.0-73.2)	47.1 (25.9-67.8)	
Unknown	96.7 (94.5-98.9)	86.1 (81.6-90.5)	84.4 (79.6-89.1)	
T stage 2004+				< 0.001
T1	96.7 (92.2-100.0)	92.8 (86.0-99.5)	NA	
T2	84.4 (72.9-95.7)	47.0 (26.2-67.4)	NA	
T3		25.0 (-17.5-66.7)	NA	
Unknown	95.3 (92.5-98.1)	85.5 (80.7-90.2)	82.8 (77.6-87.9)	
N stage 2004+				0.050
N0	93.7 (89.7-97.6)	76.2 (68.6-83.6)	74.8 (66.8-82.6)	
N1	NA	NA	NA	
Unknown	94.7 (91.3-98.0)	86.8 (81.6-91.8)	84.2 (78.6-89.7)	
M stage 2004+				< 0.001
M0	97.2 (94.5-99.9)	81.3 (74.1-88.3)	79.8 (72.2-87.3)	
M1	75.4 (56.6-93.8)	53.9 (31.6-75.8)	NA	
Unknown	94.3 (90.7-97.9)	86.5 (81.1-91.8)	83.8 (78.0-89.5)	
SEER historic stage				< 0.001
Localized	97.3 (94.3-100.0)	87.4 (81.0-93.7)	86.1 (79.2-92.8)	
Regional	94.2 (89.3-99.0)	80.0 (71.1-88.7)	78.1 (68.7-87.3)	
Distant	84.0 (73.1-94.6)	65.2 (50.9-79.2)	59.4 (44.3-74.2)	
Unknown	96.0 (91.5-100.0)	87.9 (80.0-95.6)	86.2 (77.8-94.5)	
Surg (prim) 1998+				0.024
None	90.9 (83.3-98.3)	70.3 (56.9-83.4)	67.0 (52.7-81.0)	
Yes	94.7 (91.3-98.0)	82.3 (76.3-88.2)	80.2 (73.7-86.6)	
Unknown	95.5 (91.2-99.7)	88.7 (82.1-95.2)	86.4 (79.2-93.4)	

Abbreviations: OS=overall survival; CI=Confidence Interval; Surg (prim) – surgical treatment of primary site; NA=Not available.

Table 3. Univariate and multivariate Cox regression analyzing the prognostic factors for survival of patients with malignant giant cell tumor of bone (1975 and 2016).

Subject characteristics	Univariate		Multivariate	
	HR (95%CI)	<i>P</i> -value	HR (95%CI)	<i>P</i> -value
Sex				
Male	1.00(Reference)		1.00(Reference)	
Female	0.82 (0.51-1.31)	0.399	0.88 (0.53-1.47)	0.635
Age				
<34	1.00(Reference)		1.00(Reference)	
≥34	4.28 (2.44-7.51)	<0.001	3.68 (2.06-6.57)	<0.001
Marital status				
Unmarried	1.00(Reference)		NA	NA
Married	0.68 (0.41-1.12)	0.127	NA	NA
Unknown	0.84 (0.37-1.90)	0.669	NA	NA
Insurance recode				
Uninsured	1.00(Reference)			
Insured	0.74 (0.10-5.47)	0.765	NA	NA
Unknown	0.63 (0.37-1.09)	0.100	NA	NA
Race				
White	1.00(Reference)			
Black	0.96 (0.47-1.94)	0.902	NA	NA
AI	1.07 (0.26-4.42)	0.920	NA	NA
API	1.54 (0.73-3.24)	0.258	NA	NA
Unknown				
Year of diagnosis				
1975-1984	1.00(Reference)		1.00(Reference)	
1985-1994	1.09 (0.42-2.79)	0.864	1.18 (0.43-3.26)	0.754
1995-2004	1.56 (0.63-3.89)	0.341	2.24 (0.63-7.94)	0.211
2005-2016	2.37 (0.97-5.81)	0.059	3.56 (0.65-19.50)	0.144
Primary site				
Long bones of lower limb and associated joints	1.00(Reference)		NA	NA
Long bones of upper limb and associated joints	0.67 (0.31-1.46)	0.317	NA	NA
Pelvis and spine	1.42 (0.77-2.62)	0.263	NA	NA
Irregular bone, short bone, and flat bone	1.25 (0.63-2.47)	0.529	NA	NA
Unknown	2.02 (0.62-6.59)	0.245	NA	NA
Grade				
Grade I	1.00(Reference)		NA	NA
Grade II	2.14 (0.25-18.45)	0.488	NA	NA
Grade III	5.47 (0.64-47.04)	0.121	NA	NA
Grade IV	8.37 (1.09-64.19)	0.041	NA	NA
Unknown	1.70 (0.23-12.38)	0.599	NA	NA
T stage				
T1	1.00(Reference)		1.00(Reference)	
T2	8.10 (2.68-24.48)	<0.001	4.96 (1.57-15.63)	0.006

T3	16.07 (3.55-72.68)	<0.001	2.44 (0.42-14.03)	0.317
Unknown	2.12 (0.76-5.97)	0.153	2.09 (0.61-7.20)	0.243
N stage				
N0	1.00(Reference)		NA	
N1	NA	NA	NA	NA
Unknown	0.53 (0.32-0.90)	0.019	NA	NA
M stage				
M0	1.00(Reference)		1.00(Reference)	
M1	3.86 (1.83-8.11)	<0.001	1.14 (0.44-2.99)	0.786
Unknown	0.77 (0.44-1.36)	0.374	1.11 (0.33-3.78)	0.869
SEER historic stage				
Localized	1.00(Reference)		1.00(Reference)	
Regional	1.52 (0.79-2.93)	0.209	1.18 (0.59-2.37)	0.638
Distant	3.97 (2.08-7.60)	<0.001	3.02 (1.32-6.92)	0.009
Unknown	1.19 (0.58-2.41)	0.635	1.21 (0.56-2.61)	0.633
Surg (prim)				
None	1.00(Reference)		1.00(Reference)	
Yes	0.55 (0.30-1.00)	0.050	0.62 (0.31-1.21)	0.162
Unknown	0.39 (0.19-0.79)	0.009	1.54 (0.48-4.99)	0.471

Abbreviations: Met=Metastases.

Figures

Figure 1

Process of the patient selection for analyzing both the survival and prognostic factors of patients with malignant giant cell tumor of bone.

Figure 2

The overall survival of the patients with malignant giant cell tumor of bone from the SEER database.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplement1.pdf](#)