Association between non-motor symptoms and health-related quality of life for patients with Parkinson's disease: Evidence from a smartphone-based survey

Xiaojing Fan

Xi'an Jiaotong University <https://orcid.org/0000-0002-6704-0678>

Yaxin Zhao

Xi'an Jiaotong University

Huanyuan Luo

Liverpool School of Tropical Medicine

Ben James

uMotif Limited

Duolao Wang (✉ duolao.wang@lstmed.ac.uk)

Liverpool School of Tropical Medicine

Bruce Hellman

uMotif Limited

Research

Keywords: Health-Related Quality of Life, PDQ-8, Parkinson's disease, Non-Motor Symptom

Posted Date: June 24th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-37037/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background

This study aims to explore the association between the total number of non-motor symptoms and the Parkinson's disease (PD) patients' health-related quality of life quantitatively, using data from one of the largest smartphone-powered studies and a simplified version of 8-item Parkinson's disease Quality of life Questionnaire (PDQ-8).

Methods

The data used for analysis constitutes one part of a dataset derived from the project named '100 for Parkinson's'. 1246 patients were included in the baseline survey. The 30-item Non-Motor Symptom (NMS-30) Questionnaire and PDQ-8 Summary Index (PDQ-8 SI) were used to measure health-related quality of life and a generalized linear model was used to analyze the association of the non-motor symptoms and quality of life.

Results

The mean number of NMS per patient was 11.81 ± 5.53 . For patients with limited ability to work: the key symptoms were urgency (77.4%), sad or blue (71.05%) and difficulty getting to sleep or staying asleep (70.90%). After controlling for the life-style determinants and psychological symptoms prior PD diagnosis determinants, GLM presented the total sum of NMS-30 questions had the strongest positive influence on health-related quality of life as measured by the PDQ-8 SI. Where the sum of NMS scores increases by 1 point, the PDQ-8 SI will increase by 1.33 points (95%CI: 1.13, 1.52; $P < 0.001$). For patients without limited ability to work: urgency (62.45%), difficulty getting to sleep or staying asleep (59.81%) and getting up regularly at night to pass urine (55.09%) were the most frequent symptoms. GLM showed that where the total sum of NMS scores increases by 1 point, the PDQ-8 SI will increase by 1.56 points (95%CI: 1.37, 1.76; $P < 0.001$) after controlling for other confounders.

Conclusions

This smartphone-based study finds a higher total number of non-motor symptoms is related to a lower health-related quality of life for PD patients. It provides useful evidence for the PDQ-8 instrument and is helpful for a multidisciplinary approach to patient care and facilitate the provision of more comprehensive education for patients and caregivers.

Background

Parkinson's disease (PD) is a neurological disease, characterized by slow progress, and can be clinically manifested as motor and non-motor symptoms [1]. It is the second most common neurodegenerative disease in the world, and its incidence and prevalence are shifted by demographic changes [2]. Compared with 2.5 million in 1990, there are 6.1 million people with PD globally in 2016 [3]. In addition to typical motor symptoms, patients also suffer from various non-motor symptoms, such as psychosis, falls, depression, anxiety and sleep disturbance that seriously affect quality of life [1, 4–6]. Non-motor symptoms are related to age and disease severity and often under recognized and inadequately treated [7]. The non-motor symptoms mainly included six aspects: neuropsychiatric symptoms, sleep disorders, autonomic symptoms, gastrointestinal symptoms, sensory symptoms and other symptoms including fatigue, diplopia, blurred vision and so on [7, 8].

Health-related Quality of Life (HRQL) has become an important endpoint in the evaluation of public policy. The World Health Organization's Quality of Life Group defines quality of life as individual's perception of his or her life status within the context of the culture and value system in which individual lives and concerns[9]. Quality of life is one of the most important concepts and indicators for studying management of chronic disease such as PD [10]. Most studies evaluated HRQL for people with PD through the 39-item Parkinson's disease Quality of life Questionnaire (PDQ-39) [11–13]. PDQ-39 includes 39 items in a single questionnaire, which can be time consuming for respondents to complete. An 8-item Parkinson's Disease Quality of life Questionnaire (PDQ-8) is a shorter version of PDQ-39 and has been proven to be a valid and reliable instrument to assess quality of life in a PD population, while reducing respondent burden[14].

Loading [MathJax]/jax/output/CommonHTML/jax.js

According to previous studies, non-motor symptoms such as cognitive impairment, autonomic dysfunction and sleep disorders have become major factors of health-related quality of life for PD patients [7, 15]. Fewer population-based researches report specific data on the magnitude of association between total number of non-motor symptoms and the health-related quality of life measured by the PDQ-8. In addition, although it turns out that smartphone-based study can provide the same reliable results as field study, and can reduce the burden and errors of data entry [16, 17], few scholars in the UK use smartphone-based way to capture data. In this study, we use one of the largest smartphone-supported study named '100 for Parkinson's' to capture data and measure the specific magnitude of association between total number of non-motor symptoms and the health-related quality of life for PD patients quantitatively.

Methods

Participants

The data used for analysis came from the project named '100 for Parkinson's', which has been detailed introduced in the previous study [18]. Patients who had been diagnosed with PD before the survey, used the uMotif application on their own mobile phones and/or tablets to record their demographic characteristics, PDQ-8, NMS-30, medications and so on. There were no exclusion criteria applied in this study. 1246 patients with PD consented to take part in this project and completed both the NMS-30 questionnaire and PDQ-8 questionnaire.

Main outcome and predictor variable

The Non-Motor Symptom Questionnaire is used to screen and evaluate non-motor symptoms of Parkinson's [19]. It is a 30-item self-completed questionnaire consisting of neuropsychiatric symptoms, sleep disorders, autonomic symptoms, gastrointestinal symptoms, sensory symptoms and other symptoms including unexplained change in weight, falling, and so on. Every symptom was listed in the questionnaire and patients can easily understand and provide answers (including "yes," "no," and "don't know") to each question [20]. The total number of non-motor symptoms means the number of "yes" for items of the NMS-30. In this study, we divided NMS into 6 kinds according to previous study [7], the detailed symptoms were showed in Additional file 1.

The health-related quality of life of PD patients was assessed by the 8-item Parkinson's disease Questionnaire (PDQ-8). The scale has been validated and measured by eight dimensions of quality of life related specifically to PD [21]. The questions were derived from the 7th, 12th, 25th, 27th, 31st, 35th and 37th question of PDQ-39 [22, 23]. The 8 questions are scored by using the following codes: it would be scored 0 when the answer is "Never"; it would be scored 1 when the answer is "Occasionally"; it would be scored 2 when the answer is "Sometimes"; it would be scored 3 when the answer is "Often"; it would be scored 4 when the answer is "Always". Formula for the PDQ-8 summary index (PDQ-8 SI) is as following:

$$PDQ-8SI = \frac{T_s}{4 \times 8} \times 100$$

1

T_s means sum of scores of each questionnaire for each patients with PD, 4 means the maximum score of per question, 8 means the total number of questions. The PDQ-8 SI is ranged from 0 to 100 [21, 24]. The total score of the PDQ-8 SI is used to report outcomes, health-related quality of life gets worse when the PDQ-8 SI is higher.

Statistical Analysis

Considering the fact that motor characteristic of patients with PD (with and without limited ability to work) having different quality of life, data analysis was divided into two groups of participants: patients with limited ability to work and without limited ability to work. Differences in variables between PDQ-8 questions and motor characteristic were compared using Chi-square tests. The generalized linear model (GLM) was employed to explore the association between the total number of non-motor symptoms and the health-related quality of life for patients with PD and measure the potential determinants of quality of life quantitatively when controlling for confounding factors (including socio-economic factors, lifestyle factors and psychological symptoms prior PD diagnosis). The PDQ-8 SI score is the outcome variable; the sum of NMS-30 questions is the main determinant. The statistical analyses were performed using STATA statistical software version 12.0 (StataCorp LP, College station 77845, USA), related figures were made using Excel 2016. All p values were 2-sided and $p < 0.05$ were considered statistically significant.

Results

Basic characteristics

The basic demographic features of 1246 patients with PD were displayed in Table 1. As for socio-economic characteristics, there were 577 females and 615 males. The majority of patients' education were university/college/equivalent (72.64%). For the Life-style characteristics, 62.87% of patients never smoked tobacco and 15.33% of patients never consume drinks containing alcohol. 87.67% of patients live with others (including their wife, partners, children, relatives and hired career). In the psychological symptoms prior PD diagnosis characteristics, from the sample, 261 (21.62%) patients suffered from depression prior to PD diagnosis, 225 (18.6%) had anxiety prior to PD diagnosis, and 677 (56.09%) had limited ability to work.

Table 1
Basic characteristics of patients with Parkinson's disease (n = 1246)

Variables	n	%
Socio-economic characteristics		
Gender		
Female	577	48.41
Male	615	51.59
Age of participants (years)		
≤59	561	45.03
60–69	446	35.79
≥70	239	19.18
Education		
Up to 16 years old	192	16.21
Up to 18 years old	132	11.15
University / College / Equivalent	860	72.64
Life-style characteristics		
Do you smoke tobacco?		
No, never	750	62.87
No, but used to	371	31.10
Yes	72	6.03
How often do you have a drink containing alcohol?		
Never	182	15.33
2–3 times per week	291	24.52
2–4 times per month	246	20.72
4 or more times per week	251	21.15
Monthly or less	217	18.28
Live style		
Live with others	1031	87.67
Live alone	145	12.33
Psychological symptoms prior PD diagnosis		
Depression prior PD diagnosis		
No	946	78.38
Yes	261	21.62
Anxiety prior PD diagnosis		
No	985	81.40
Yes	225	18.60
Motor characteristic		

Variables	n	%
Without limited ability to work	530	43.91
With limited ability to work	677	56.09

Distribution of NMS-30 and PDQ-8 questions

1.28% of patients had no NMS and 98.72% presented more than one symptom. The mean number of NMS per patient was 11.81 ± 5.53 . Figure 1 showed neuropsychiatric symptoms, sleep disorders, autonomic symptoms and gastrointestinal symptoms were less frequent in the patients without limitations to work than the patients with limited ability to work. The key symptoms were urgency (77.4%), feeling sad or blue (71.05%) and difficulty getting to sleep or staying asleep (70.90%) respectively for the PD patients with limited ability to work, whereas the urgency (62.45%), difficulty getting to sleep or staying asleep (59.81%) and getting up regularly at night to pass urine (55.09%) were the most frequent symptoms for the PD patients without limited ability to work. The distribution of PDQ-8 questions between patients with and without limit ability to work were shown in Table 2. Compared to those without limited ability to work, patients with limited ability to work reported more problems in always difficulty getting around (3.69% vs 0.19%), dressing (1.18% vs 0.19%), always feeling depressed (2.07% vs 0.75%), always problems with close personal relationships (2.66% vs 1.13%), always feeling unable to communicate properly (1.62% vs 0.57%), always painful muscle cramps (2.66% vs 0.75%) and always feeling embarrassed in public (2.51% vs 0.19%) except problems with concentration (13.96% vs 7.39%).

Table 2
Distribution of PDQ-8 questions between patients with and without limited ability to work (n = 1207)

PDQ-8 questions	Limited ability to work				χ^2	<i>P</i>
	Without (n = 530)		With (n = 677)			
	n	%	n	%		
Difficulty getting around					324.12	< 0.001
Always	1	0.19	25	3.69		
Often	343	64.72	114	16.84		
Sometimes	131	24.72	247	36.48		
Occasionally	16	3.01	141	20.83		
Never	39	7.36	150	22.16		
Difficulty dressing					193.12	< 0.001
Always	1	0.19	8	1.18		
Often	334	64.91	196	28.95		
Sometimes	160	30.19	281	41.51		
Occasionally	7	1.32	74	10.93		
Never	18	3.40	118	17.43		
Felt depressed					104.81	< 0.001
Always	4	0.75	14	2.07		
Often	239	45.09	154	22.75		
Sometimes	204	38.49	260	38.40		
Occasionally	22	4.15	115	16.99		
Never	61	11.51	134	19.79		
Problems with close personal relationships					127.65	< 0.001
Always	6	1.13	18	2.66		
Often	329	62.08	219	32.35		
Sometimes	147	27.74	240	35.45		
Occasionally	18	3.40	70	10.34		
Never	30	5.66	130	19.20		
Problems with concentration					120.02	< 0.001
Always	74	13.96	50	7.39		
Often	237	44.72	162	23.93		
Sometimes	144	27.17	199	29.39		
Occasionally	32	6.04	140	20.68		
Never	43	8.11	126	18.61		
Felt unable to communicate properly					200.90	< 0.001
Always	3	0.57	11	1.62		

PDQ-8 questions	Limited ability to work				χ^2	<i>P</i>
	Without (n = 530)		With (n = 677)			
	n	%	n	%		
Often	257	48.49	120	17.73		
Sometimes	214	40.38	267	39.44		
Occasionally	11	2.08	127	18.76		
Never	45	8.49	152	22.45		
Painful muscle cramps or spasm					75.45	< 0.001
Always	4	0.75	18	2.66		
Often	148	27.92	108	15.95		
Sometimes	226	42.64	214	31.61		
Occasionally	69	13.02	199	29.39		
Never	83	15.66	138	20.38		
Felt embarrassed in public					107.02	< 0.001
Always	1	0.19	17	2.51		
Often	272	51.32	188	27.77		
Sometimes	188	35.47	245	36.19		
Occasionally	23	4.34	101	14.92		
Never	46	8.68	126	18.61		

The association of NMS-30 and PDQ-8 SI

For patients with limited ability to work (Table 3), GLM presented the sum of NMS-30 questions had the strongest positive influence on the health-related quality of life as measured by the PDQ-8 SI score, among the factors analyzed in this study. When controlling socioeconomic characteristic in model 1, the estimates of the association of NMS and PDQ-8 SI is 1.33 (95%CI: 1.13, 1.52; $P < 0.001$). After controlling the life-style determinants and psychological symptoms prior to PD diagnosis determinants, the results in model 3 are still statistically significant ($P < 0.001$), indicating that where the sum of NMS increases by 1 point, the PDQ-8 SI would increase by 1.33 points, meaning the quality of life will decrease. Besides, the GLM indicated that depression prior to PD diagnosis ($\beta = 3.08$, 95%CI: 0.14, 6.02; $P = 0.040$) is also the potential determinant of worse quality of life of PD patients. However, age of 60–69 ($\beta = -2.87$, 95%CI: -5.20, -0.54; $P = 0.016$) and education of up to 18 years old ($\beta = -4.19$, 95%CI: -8.18, -0.21; $P = 0.039$) were the potential determinants of better health-related quality of life, compared to age ≤ 59 and education of up to 16 years old, respectively.

Table 3

The multivariate analysis of the association between total number of Non-Motor Symptoms and PDQ-8 Single index by generalized linear model analysis for patients with limited ability to work (n = 677)

Variables	Model 1				Model 2				Model 3			
	Estimates	<i>P</i>	95%CI		Estimates	<i>P</i>	95%CI		Estimates	<i>P</i>	95%CI	
Main determinants												
Sum of NMS-30 questions	1.33	< 0.001	1.13	1.52	1.30	< 0.001	1.10	1.50	1.33	< 0.001	1.12	1.54
Socio-economic determinants												
Gender												
Female	Reference				Reference				Reference			
Male	-0.15	0.886	-2.18	1.88	-0.001	0.999	-2.16	2.16	-0.21	0.854	-2.41	2.00
Age of participants (years)												
≤59	Reference				Reference				Reference			
60–69	-3.44	0.002	-5.65	-1.24	-2.83	0.016	-5.13	-0.54	-2.87	0.016	-5.20	-0.54
≥70	-2.42	0.121	-5.48	0.64	-1.73	0.283	-4.89	1.43	-2.52	0.129	-5.79	0.74
Education												
Up to 16 years old	Reference				Reference				Reference			
Up to 18 years old	-3.46	0.080	-7.34	0.42	-3.90	0.052	-7.85	0.04	-4.19	0.039	-8.18	-0.21
University / College / Equivalent	-1.33	0.353	-4.14	1.48	-1.59	0.280	-4.46	1.29	-1.83	0.221	-4.76	1.10
Life-style determinants												
Do you smoke tobacco?												
No, never					Reference				Reference			
No, but used to					-0.35	0.764	-2.65	1.94	-0.59	0.624	-2.93	1.76
Yes					3.62	0.086	-0.52	7.76	3.35	0.117	-0.84	7.55
How often do you have a drink containing alcohol?												
Never					Reference				Reference			
2–3 times per week					-0.02	0.991	-3.41	3.37	0.14	0.938	-3.31	3.58

Model 1 is a model controlling the main determinants and socio-economic determinants including gender, age and education; model 2 is a model adding life-style characteristics (e.g. smoking status, drinking status and live style) as covariates into model 1; model 3 is a model controlling the main determinants, socio-economic determinants, life-style characteristics and psychological Loading [MathJax]/jax/output/CommonHTML/jax.js sion prior PD diagnosis and anxiety prior PD diagnosis).

Variables	Model 1			Model 2			Model 3		
	Estimates	<i>P</i>	95%CI	Estimates	<i>P</i>	95%CI	Estimates	<i>P</i>	95%CI
2–4 times per month				1.73	0.312	-1.63 5.09	1.59	0.365	-1.85 5.04
4 or more times per week				-0.81	0.641	-4.22 2.60	-0.33	0.852	-3.84 3.17
Monthly or less				-0.65	0.694	-3.91 2.61	-0.87	0.610	-4.20 2.47
Live style									
Live with others				Reference			Reference		
Live alone				0.92	0.566	-2.22 4.06	1.11	0.496	-2.09 4.32
Psychological symptoms prior PD diagnosis									
Depression prior PD diagnosis									
No							Reference		
Yes							3.08	0.040	0.14 6.02
Anxiety prior PD diagnosis									
No							Reference		
Yes							-2.26	0.157	-5.39 0.87
Model 1 is a model controlling the main determinants and socio-economic determinants including gender, age and education; model 2 is a model adding life-style characteristics (e.g. smoking status, drinking status and live style) as covariates into model 1; model 3 is a model controlling the main determinants, socio-economic determinants, life-style characteristics and psychological symptoms prior PD diagnosis (e.g. depression prior PD diagnosis and anxiety prior PD diagnosis).									

For patients without limited ability to work - after controlling for the socio-economic determinants (Table 4), life-style and psychological symptoms prior to PD diagnosis determinants in model 3 – where the total sum of NMS increases by 1 point, the PDQ-8 SI score will increase by 1.56 points (95%CI: 1.37, 1.76; $P < 0.001$). In other words, where the non-motor symptoms are more, the health-related quality of life of PD patients without limited ability to work was worse. The GLM also indicated that depression prior to PD diagnosis ($\beta = 4.06$, 95%CI: 1.03, 7.09; $P = 0.009$) was also the potential determinant of worse quality of life than no depression prior to PD diagnosis; while age of 60–69 ($\beta = -2.84$, 95%CI: -4.85, -0.82; $P = 0.006$) was the potential determinant of better quality of life, compared to age ≤ 59 and education of up to 16 years old, respectively.

Table 4

The multivariate analysis of the association between total number of Non-Motor Symptoms and PDQ-8 Single index by generalized linear model analysis for patients without limited ability to work (n = 530)

Variables	Model 1				Model 2				Model 3			
	Estimates	<i>P</i>	95%CI		Estimates	<i>P</i>	95%CI		Estimates	<i>P</i>	95%CI	
Main determinants												
Sum of NMS-30 questions	1.59	< 0.001	1.41	1.78	1.60	< 0.001	1.41	1.79	1.56	< 0.001	1.37	1.76
Socio-economic determinants												
Gender												
Female	Reference				Reference				Reference			
Male	-0.35	0.696	-2.09	1.40	-0.15	0.871	-1.99	1.68	0.10	0.918	-1.75	1.94
Age of participants (years)												
≤59	Reference				Reference				Reference			
60–69	-3.02	0.002	-4.94	-1.10	-2.75	0.007	-4.76	-0.74	-2.84	0.006	-4.85	-0.82
≥70	-1.52	0.225	-3.98	0.94	-1.27	0.335	-3.84	1.31	-1.42	0.282	-4.00	1.17
Education												
Up to 16 years old	Reference				Reference				Reference			
Up to 18 years old	0.35	0.845	-3.12	3.81	0.41	0.821	-3.16	3.98	0.37	0.837	-3.19	3.94
University / College / Equivalent	-2.06	0.092	-4.45	0.34	-2.13	0.088	-4.57	0.31	-2.24	0.075	-4.71	0.22
Life-style determinants												
Do you smoke tobacco?												
No, never					Reference				Reference			
No, but used to					-0.44	0.672	-2.45	1.58	-0.31	0.764	-2.33	1.71
Yes					3.77	0.084	-0.51	8.05	3.50	0.107	-0.75	7.74
How often do you have a drink containing alcohol?												
Never					Reference				Reference			
2–3 times per week					-2.02	0.222	-5.26	1.22	-1.43	0.395	-4.71	1.86

Model 1 is a model controlling the main determinants and socio-economic determinants including gender, age and education; model 2 is a model adding life-style characteristics (e.g. smoking status, drinking status and live style) as covariates into model 1; model 3 is a model controlling the main determinants, socio-economic determinants, life-style characteristics and psychological Loading [MathJax]/jax/output/CommonHTML/jax.js sion prior PD diagnosis and anxiety prior PD diagnosis).

Variables	Model 1			Model 2			Model 3		
	Estimates	<i>P</i>	95%CI	Estimates	<i>P</i>	95%CI	Estimates	<i>P</i>	95%CI
2–4 times per month				-2.37	0.166	-5.72 0.98	-1.66	0.339	-5.05 1.74
4 or more times per week				-3.40	0.048	-6.78 -0.03	-2.19	0.213	-5.63 1.25
Monthly or less				-2.29	0.218	-5.94 1.35	-1.72	0.361	-5.42 1.97
Live style									
Live with others				Reference			Reference		
Live alone				-0.28	0.843	-3.02 2.47	0.06	0.966	-2.72 2.84
Psychological symptoms prior PD diagnosis									
Depression prior PD diagnosis									
No							Reference		
Yes							4.06	0.009	1.03 7.09
Anxiety prior PD diagnosis									
No							Reference		
Yes							-1.10	0.479	-4.16 1.95
Model 1 is a model controlling the main determinants and socio-economic determinants including gender, age and education; model 2 is a model adding life-style characteristics (e.g. smoking status, drinking status and live style) as covariates into model 1; model 3 is a model controlling the main determinants, socio-economic determinants, life-style characteristics and psychological symptoms prior PD diagnosis (e.g. depression prior PD diagnosis and anxiety prior PD diagnosis).									

Discussion

This study is a comprehensive exploration of the association of total number of non-motor symptoms and health-related quality of life for PD patients quantitatively using a large smartphone-based way to capture data in the UK. For patients with PD, NMS have been considered as a significant content, and appropriate therapy can considerably improve quality of life [25]. Many researchers have explored how some items of the NMS independently have effect on the health-related quality of life, such as depression, sleep, mood and attention [13, 26], less researchers focused on the association between the total number of NMS and health-related quality of life through PDQ-8. This study explored the association between total number of NMS and health-related quality of life for PD patients and found that the higher the sum of NMS-30 questions was, the worse the health-related quality of life became. The NMS could be related to degeneration of different brain structures, may reduce the PD patients' health-related quality of life, and may cause severe disability and shorten life expectancy, which should be considered in the treatment plan [7].

Our study reveals that almost every patient with PD presented with more than one symptom. The relative NMS occurrence, as evaluated by NMS-30, is consistent with several other studies, showing sleep (difficulty getting to sleep), mood/cognition (sad or blue), memory/attention and urination (urgency, getting up regularly at night to pass urine) as the most prevalent symptoms both for PD patients with and without limited ability to work [27–29]. However, the absolute occurrence of the most frequent symptoms are lower than Paulo Bugalho's study (Sleep/fatigue, affect/cognition, attention/memory were all above 80%) [30], but higher than other studies [28, 29], in which above 60% are rare, while in our study "Difficulty getting to sleep" is 70.90% for patients with limited ability to work and is 59.81% for patients without limited ability to work; "Urgency to pass urine makes you rush to the toilet" is 77.40% for patients

Loading [MathJax]/jax/output/CommonHTML/jax.js patients without limited ability to work, further investigation is needed to verify.

Depression is reported to be one of the common complication of PD ranging in prevalence between 2.7% and 70% but is under self-reported[31]. Some studies showed depression can occur before the onset of PD, and it may have a significant effect on the patient's quality of life [31–34]. In our study, depression (Feeling sad low or blue) is one of the most prevalent symptoms both for patients with and without limited ability to work. Depression prior PD diagnosis was also a potential determinant of worse quality of life.

Age and education were found to be correlated with health-related quality of life of PD patients significantly. It is well known that age is the greatest risk factor for the development of PD. A previous study showed age-specific incidence rates of PD increased sharply beginning at age 60 years and peaking after 80 years [35]. In our study, patients aged 60–69 years had better quality of life than patients aged less than 59 years, which is inconsistent with previous research, further investigation is needed to verify. Compared with patients educated up to 16 years old, patients educated up to 18 years old is a potential determinant of better quality of life, which is accordance with the conclusion that higher education is associated with better health-related quality of life [36, 37].

Limitations

Our study presents some limitations. Firstly, due to the limitations of question setting in the questionnaire, some potential factors that affect health-related quality of life were not controlled for in this study, such as individual exercise, occupation, income level or marital status. Secondly, this study conducted a brief question on whether patients' motor symptoms limited their ability to work and divided the characteristic of motor symptoms into with or without limited ability to work, but lacked a detailed evaluation of Parkinson's patients' motor symptoms. More detailed study are needed to verify the association between non-motor symptoms and health-related quality of life for patients with Parkinson's disease after controlling for their motor symptoms. Thirdly, this is a quantitative study that reports the association between non-motor symptoms of PD patients and health-related quality of life. In the future, more qualitative research will be needed to verify the association.

Conclusions

In this study, we find a reversed association between total numbers of non-motor symptoms and health-related quality of life for PD patients; the more of the total number of non-motor symptoms, the worse of the health-related quality of life. Improved awareness and understanding of the association will encourage to provide reference for more comprehensive care for PD patients and their caregivers.

List of abbreviations

Abbreviations

Health-related Quality of Life	HRQL
Generalized Linear Model	GLM
30-item Non-Motor Symptom	NMS-30
Parkinson's disease	PD
8-item Parkinson's Disease Quality of life Questionnaire	PDQ-8
39-item Parkinson's Disease Quality of life Questionnaire	PDQ-39
PDQ-8 Summary Index	PDQ-8 SI
Participant Centered Consent	PCC

Declarations

Ethics approval and consent to participate

The ethics of this study was approved by the Liverpool School of Tropical Medicine (No.15-050) and the New England IRB for participants from the USA (No.16-042). In addition, this study was conducted in accordance with the Declaration of Helsinki. E-consent was obtained by projects from each participant before investigation.

Consent for publication

Not applicable.

Availability of data and materials

Researchers who want to use these data should contact Bruce Hellman (bruce@umotif.com).

Competing interests

The authors declare that they have no competing interests.

Funding

This study was funded by the 65th batch of Postdoctoral Research Foundation of China (CN) (No.2019M653684) and the Central University Basic Scientific Research Business Expenses Special Funds (CN) (No. SK2020047). The '100 for Parkinson's' study was funded by uMotif Limited.

Authors' contributions

DW and BH was responsible for overall design of this study. BH and BJ were responsible for the field working including data collection and management. XF, YZ and DW were responsible for the statistical analysis plan. XF, YZ and DW performed the statistical analysis. The manuscript was prepared by XF, YZ, HL, BJ, BH and DW. All authors critically reviewed and approved the final manuscript.

Acknowledgements

We express our appreciation to the Nesta/Cabinet Office Centre for Social Action Innovation Fund, a growing number of patient groups (the Cure Parkinson's Trust and Parkinson's UK, the European Parkinson's Disease Association), uMotif Limited and global sponsor partners for their support.

References

1. Kalia LV, Lang AE. Parkinson's disease. *Lancet*. 2015;386(9996):896–912.
2. Poewe W, Seppi K, Tanner CM, Halliday GM, Brundin P, Volkmann J, Schrag AE, Lang AE. Parkinson disease. *Nat Rev Dis Primers* 2017, 23(3):17013. doi: 17010.11038/nrdp.12017.17013.
3. GBD 2016 Parkinson's Disease Collaborators. Global, regional, and national burden of Parkinson's disease, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet Neurol*. 2018;17(11):939–53.
4. Prasuhn J, Piskol L, Vollstedt EJ, Graf J, Schmidt A, Tadic V, Tunc S, Hampf J, Warlich E, Bibergeil C. Non-motor symptoms and quality of life in subjects with mild parkinsonian signs. *Acta Neurologica Scandinavica* 2017, 136(5).
5. Hurt CS, Rixon L, Chaudhuri KR, Mossmorris R, Samuel M, Brown RG. Identifying barriers to help-seeking for non-motor symptoms in people with Parkinson's disease. *Journal of Health Psychology*. 2016;24(5):561–71.
6. Bartolomei L, Pastore A, Meligrana L, Sanson E, Bonetto N, Minicuci GM, Marsala SZ, Mesiano T, Bragagnolo L, Antonini A. Relevance of sleep quality on caregiver burden in Parkinson's disease. *Neurol Sci*. 2018;39(5):1–5.
7. Chaudhuri KR, Healy Dg Fau - Schapira AHV, Schapira AH. Non-motor symptoms of Parkinson's disease: diagnosis and management. *Lancet Neurol*. 2006;5(3):235–45.
8. Wang G, Wan Y, Cheng Q, Xiao Q, Wang Y, Zhang J, Ma J-F, Wang X-J, Zhou H-Y, Chen S-D. Malnutrition and associated factors in Chinese patients with Parkinson's disease: Results from a pilot investigation. *Parkinsonism Relat Disord*. 2010;16(2):119–23.
9. Group TW. The World Health Organization Quality of Life assessment (WHOQOL): position paper from the World Health Organization. *Soc Sci Med*. 1995;41(10):1403–9.
10. Martinez-Martin P. What is quality of life and how do we measure it? Relevance to Parkinson's disease and movement disorders. *Mov Disord*. 2017;32(3):382–92.
11. Kim SR, Kim JY, Kim HK, Lim KE, Kim MS, Chung SJ. Association among type D personality, non-motor symptoms, and quality of life in Parkinson's disease: A cross-sectional study. *Geriatric Nursing* 2017, 38(5).

12. Zhang L, Cao B, Ou R, Wei Q, Zhao B, Yang J, Wu Y, Shang H. Non-motor symptoms and the quality of life in multiple system atrophy with different subtypes. *Parkinsonism Relat Disord.* 2017;35:63–8.
13. Prakash KM, Nadkarni NV, Lye WK, Yong MH, Tan EK. The impact of non-motor symptoms on the quality of life of Parkinson's disease patients: a longitudinal study. *Eur J Neurol.* 2016;23(5):854.
14. Fereshtehnejad SM, Naderi N, Rahmani A, Shahidi GA, Delbari A, Lökk J. Psychometric study of the Persian short-form eight-item Parkinson's disease questionnaire (PDQ-8) to evaluate health related quality of life (HRQoL). *Health Quality of Life Outcomes.* 2014;12(1):78.
15. Marras C, Chaudhuri KR. Nonmotor features of Parkinson's disease subtypes. *Mov Disord.* 2016;31(8):1095–102.
16. Bot BM, Suver C, Neto EC, Kellen M, Klein A, Bare C, Doerr M, Pratap A, Wilbanks J, Dorsey ER, et al. The mPower study, Parkinson disease mobile data collected using ResearchKit. *Scientific Data.* 2016;3:160011.
17. Trister AD, Dorsey ER, Friend SH. Smartphones as new tools in the management and understanding of Parkinson's disease. *Npj Parkinsons Disease.* 2016;2(1):16006.
18. Fan X, Wang D, Hellman B, Janssen MF, Bakker G, Coghlan R, Hursey A, Matthews H, Whetstone I. Assessment of Health-Related Quality of Life between People with Parkinson's Disease and Non-Parkinson's: Using Data Drawn from the '100 for Parkinson's' Smartphone-Based Prospective Study. *Int J Environ Res Public Health* 2018, 15(11):pii: E2538. doi: 2510.3390/ijerph15112538.
19. Chaudhuri KR, Martinez-Martin P, Brown RG, Sethi K, Stocchi F, Odin P, Ondo W, Abe K, Macphee G, Macmahon D. The metric properties of a novel non-motor symptoms scale for Parkinson's disease: Results from an international pilot study. *Movement Disorders Official Journal of the Movement Disorder Society.* 2007;22(13):1901.
20. Chaudhuri KR, Martinezmartin P, Schapira AH, Stocchi F, Sethi K, Odin P, Brown RG, Koller W, Barone P, Macphee G. International multicenter pilot study of the first comprehensive self-completed nonmotor symptoms questionnaire for Parkinson's disease: the NMSQuest study. *Mov Disord.* 2006;21(7):916–23.
21. Jenkinson C, Fitzpatrick R, Peto V, Greenhall R, Hyman N. The PDQ-8: Development and validation of a short-form parkinson's disease questionnaire. *Psychol Health.* 1997;12(6):805–14.
22. Franchignoni F, Giordano A, Ferriero G. Rasch analysis of the short form 8-item Parkinson's Disease Questionnaire (PDQ-8). *Qual Life Res.* 2008;17(4):541–8.
23. Jenkinson C, Fitzpatrick R, Peto V, Dummett S, Morley D, Saunders P. The Parkinson's disease questionnaires user manual Health Services Research Unit. University of Oxford: University of Oxford; 2012.
24. Tan LCS, Lau P-N, Au W-L, Luo N. Validation of PDQ-8 as an independent instrument in English and Chinese. *J Neurol Sci.* 2007;255(1–2):77–80.
25. Martinez-Martin P, Schapira AHV, Stocchi FF, Sethi K, Odin P, MacPhee G, Brown RG, Naidu Y, Clayton L, Abe K, et al. Prevalence of nonmotor symptoms in Parkinson's disease in an international setting; study using nonmotor symptoms questionnaire in 545 patients. *Mov Disord.* 2007;22(11):1623–9.
26. Berganzo K, Tijero B, Gonzalez-Eizaguirre A, Somme J, Lezcano E, Gabilondo I, Fernandez M, Zarranz JJ, Gomez-Esteban JC. Motor and non-motor symptoms of Parkinson's disease and their impact on quality of life and on different clinical subgroups. *Neurologia.* 2016;31(9):585–91.
27. Gallagher DA, Lees AJ, Schrag A. What are the most important nonmotor symptoms in patients with Parkinson's disease and are we missing them? *Mov Disord.* 2010;25(15):2493–500.
28. Zis P, Martinez-Martin P, Sauerbier A, Rizos A, Sharma JC, Worth PF, Sophia R, Silverdale M, Chaudhuri KR. Non-motor symptoms burden in treated and untreated early Parkinson's disease patients: argument for non-motor subtypes. *Eur J Neurol.* 2015;22(8):1145–50.
29. Duncan GW, Khoo TK, Yarnall AJ, O'Brien JT, Coleman SY, Brooks DJ, Barker RA, Burn DJ. Health-related quality of life in early Parkinson's disease: the impact of nonmotor symptoms. *Mov Disord.* 2014;29(2):195–202.
30. Bugalho P, Lampreia T, Miguel R, Mendonça MD, Caetano A, Barbosa R. Non-Motor symptoms in Portuguese Parkinson's Disease patients: correlation and impact on Quality of Life and Activities of Daily Living. *Sci Rep.* 2016;6:32267.
31. Burn DJ. Beyond the iron mask: towards better recognition and treatment of depression associated with Parkinson's disease. *Mov Disord.* 2002;17(3):445–54.
32. Ray S, Agarwal P. Depression and Anxiety in Parkinson's Disease. *Clin Geriatr Med.* 2020;36(1):93–104.

33. Marsh L. Depression and Parkinson's disease: current knowledge. *Curr Neurol Neurosci Rep* 2013, 13(12):409. doi: 410.1007/s11910-11013-10409-11915.

34. Timmer MHM, van Beek M, Bloem BR, Esselink RAJ. What a neurologist should know about depression in Parkinson's disease. *Pract Neurol*. 2017;17(5):359–68.

35. Driver JA, Logroschino G, Gaziano JM, Kurth T. Incidence and remaining lifetime risk of Parkinson disease in advanced age. *Neurology*. 2009;72(5):432–8.

36. Chrobak-Bien J, Gawor A, Paplaczky M, Malecka-Panas E, Gasiorowska A. Analysis of factors affecting the quality of life of those suffering from Crohn's disease. *Pol Przegl Chir*. 2017;89(4):16–22.

37. Su M, Zhou Z, Si Y, Wei X, Xu Y, Fan X, Chen G. Comparing the effects of China's three basic health insurance schemes on the equity of health-related quality of life: using the method of coarsened exact matching. *Health Qual Life Outcomes*. 2018;16(1):41. doi:10.1186/s12955-12018-10868-12950.

Figures

Figure 1

Estimates association of PDQ-8 Single index with NMS-30 and each domain of NMS-30 for PD patients with and without limited ability to work (n=1207)

Figure 2

The distribution of NMS-30 questions between patients with and without limited ability to work (n=1207)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Responseletter.docx](#)
- [Additionalfile1.docx](#)