

Microscopic extra-laminar sequestrectomy (MELS) for the treatment of hidden zone lumbar disc herniation: report of the surgical technique, patient selection, and clinical outcomes

Chunxiao Wang

Southern Medical University Nanfang Hospital

Yao Zhang

Yantai Affiliated Hospital of Binzhou Medical University

Xiaojie Tang

Yantai Affiliated Hospital of Binzhou Medical University

Haifei Cao

Yantai Affiliated Hospital of Binzhou Medical University

Qinyong Song

Yantai Affiliated Hospital of Binzhou Medical University

Jiangwei Tan (✉ drzal0221@163.com)

Binzhou Medical University - Yantai Campus <https://orcid.org/0000-0001-7830-5427>

Chengzhen Jin

Yantai Affiliated Hospital of Binzhou Medical University

Hongkai Song

Yantai Affiliated Hospital of Binzhou Medical University

Research article

Keywords: Hidden Zone, Lumbar Disc Herniation, Microscopic Sequestrectomy, Extra-Laminar Approach

Posted Date: June 25th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-37318/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Surgical management of lumbar disc herniation in the hidden zone is technically challenging due to its difficult surgical exposure. The conventional interlaminar approach harbors the potential risk of post-surgical instability, while other approaches consist of complicated procedures with a steep learning curve and prolonged operation time.

Objective

To introduce a safe and effective technique named microscopic extra-laminar sequestrectomy (MELS) for treatment of hidden zone lumbar disc herniation and present clinical outcomes within a two year follow-up period.

Methods

Between Jan 2016 to Jan 2018, twenty one patients (13 males) with hidden zone lumbar disc herniation were enrolled in this study. All patients underwent MELS (19 patients underwent sequestrectomy only, 2 patients underwent an additional inferior discectomy). The nerve root and herniated fragment were visually exposed using this extra-laminar approach. The operation duration, blood loss, intra- and postoperative complications, and recurrences were recorded. The Visual Analog Scale (VAS), Oswestry Disability Index (ODI), and the modified MacNab criteria were used to evaluate clinical outcomes. Postoperative stability was evaluated both radiologically and clinically.

Results

The mean follow-up period was 20.95 ± 2.09 months, ranging from 18 to 24 months. The mean operation time was 32.43 ± 7.19 min and the mean blood loss was 25.52 ± 5.37 ml. All patients showed complete neurological symptom relief after surgery. The VAS and ODI score were significantly improved at the final follow-up compared to those before operation (7.88 ± 0.70 vs 0.10 ± 0.30 , 59.24 ± 10.83 vs 11.29 ± 3.59 , respectively, $p < 0.05$). Seventeen patients (81%) obtained an “excellent” outcome and the remaining four (19%) patients obtained a “good” outcome based the MacNab criteria. One patient suffered reherniation at the same level one year after the initial surgery and underwent a transforaminal endoscopic discectomy. No major complications and postoperative instability were observed.

Conclusions

Our observation suggest that MELS is a safe and effective method in the management of hidden zone lumbar disc herniation. Due to its relative simplicity, it comprises a flat surgical learning curve and shorter operation duration, and overall results in reduced disturbance to lumbar stability.

1. Introduction

Lumbar disc herniation is the most common diagnosis among degenerative disorders of the lumbar spine, affecting 2 to 3% of the population, and is the main cause of spinal surgery in the adult population [1–3]. Wiltse *et al.* [4] reported that the lateral lumbar spinal canal can be subdivided into three regions: the subarticular (lateral recess), foraminal (pedicle), and extraforaminal (far lateral) zone. MacNab [5] described the subarticular and foraminal region as the “hidden zone” due to difficult surgical exposure. Surgical management of “hidden zone” lumbar disc herniation (HZLDH) using a standard interlaminar approach requires hemilaminotomy or hemilaminectomy of the upper vertebra and is often associated with partial or complete resection of facet joints, resulting in an increased risk of lumbar segmental instability [6–9]. Di Lorenzo *et al.* [10] introduced a more direct approach to access the hidden zone through a fenestration on the hemilamina. Soldner *et al.* [11] further described this approach and named it translaminar fenestration. However, due to segment-dependent vertebral anatomy, the fenestration must be performed very laterally in the upper lumbar levels to reach the medial hidden zone. Disruption of the lateral hemilamina (pars interarticularis) has been linked to an increased risk of stress fracture and instability [12]. Dezawa *et al.* [13] described an endoscopic translaminar approach, in which trauma to the pars interarticularis is smaller. However this approach is more technically demanding and the surgical hand-eye coordination learning curve is steep. Therefore, while many options for the surgical management of hidden zone lumbar disc herniation exist, most of them are challenging or may result in post-operative spinal instability. In this study, we therefore developed a novel safe and effective technique called microscopic extra-laminar sequestrectomy (MELS) which we believe will be easy to implement for most surgeons. The purpose of this study is to describe this novel surgical technique, and outline patient selection and preliminary clinical outcomes used for its evaluation.

2. Methods

2.1 Patient recruitment

We undertook a non-randomized prospective study, which was approved by the ethics committee of our institution (Spine surgery department, Yantai Affiliated Hospital of Binzhou Medical University). All patients provided informed consent. Between Jan 2016 to Jan 2018, twenty one patients (13 males and 8 females) who were diagnosed with HZLDH were enrolled in this study and underwent MELS. The mean age was 59.58 ± 7.67 years, ranging from 48 to 77 years. All patients included in this study failed 4 weeks of conservative treatment before surgery. Magnetic resonance imaging (MRI) and computerized tomography (CT) were performed preoperatively and the imaging manifestations showed consistency with spinal symptoms in all patients. The inclusion criteria were as follows: 1) age > 18 years; 2) no previous history of spinal surgery; 3) MRI and CT identified the fragment was located in the hidden zone;

4) good general condition, no severe cardiopulmonary or hepatorenal dysfunction. Exclusion criteria were: 1) central stenosis (less than 10 mm) or lateral recess stenosis (less than 3 mm) confirmed by MR imaging and CT scans; 2) concomitant diseases involving systematic infection or malignant tumor; 3) segmental instability confirmed by dynamic radiographs.

2.2 Surgical procedure

All the surgeries were performed by the same senior surgeon (the corresponding author). After endotracheal general anesthesia, the patient was positioned prone on the operative table, the hip and knee were flexed to widen the interlaminar space, the abdomen was kept free-hanging to reduce the intraoperative bleeding. After the spinal level of interest was identified by fluoroscopy, the Quadrant system (Medtronic, USA) was assembled and fixed to the operative table. A longitudinal incision of 2–3 cm length was made 2 cm lateral to the midline on the pathological side and a guide needle was docked on the lateral border of the lamina through the deep fascia. Sequential dilators were introduced and a suitable blade was appropriately anchored on the target lamina. After the final position of the retractor was reconfirmed by the C-arm, the lateral side of the lamina and the pars interarticularis were exposed. A blunt sublaminar dissection was performed from the lateral margin with a curette to detach the ligamentum flavum. A 2–3 mm crescent-shaped lateral lamina was excised and ligamentum flavum was removed to expose the exiting nerve root and the ganglion, around which hemostasis with a bipolar was potentially needed. This exploration was performed in a caudal to cranial direction along the nerve root with a hook probe. The nerve root was cranially retracted and appropriately protected, the sequestered nucleus pulposus was explored and removed by tracking along its path. In some cases where the routine hook probe (10 mm) was not able to reach the disc fragment, the lateral margin of the pars interarticularis would have to be removed partially to reach closer to the more medially located sequestration, while sometimes a longer hook (15 mm) was required to drag far fragments out (Fig. 1). The clearance of the intervertebral disc material depends on both preoperative MRI and intraoperative findings. The discectomy was performed in three patients as the migrated fragments were obviously linked with the inferior disc. After a small part of the superior articular process was removed, the intervertebral space can be approached and the discectomy was easily performed (Fig. 2, 3). After adequate hemostasis, the retractors were taken out and a latex strip drainage was placed in position before the surgical wound was sutured. The drainage was removed 24 to 48 hours after surgery, and the stitches were 10 to 12 days later.

2.3 Evaluation of outcomes

The clinical outcomes were assessed before surgery and at 3 days, 1 month, 6 months, 1 year, and 18 to 24 months after surgery. We used the Visual Analog Scale (VAS) to assess back and leg pain, the Oswestry Disability Index (ODI) to assess functional capacity, and the modified MacNab criteria to evaluate patient satisfaction. Operating times, blood loss, intra- and perioperative complications, the length of hospital stays and recurrences were recorded. Immediate postoperative CT and MRI images were taken for each patient to evaluate the destruction of pars and completeness of decompression. The

occurrence of postoperative instability was evaluated both radiologically and clinically. A dynamic X-ray was taken at least 1 year after surgery. An anterior translation of more than 3 mm between the operated vertebra and the inferior vertebra was considered as radiological instability. New onset or significantly exacerbated back pain which induced disability was considered clinical instability.

2.4 Statistical analysis

All data are represented as mean \pm standard deviation ($\bar{x} \pm s$). The VAS and ODI before and after surgery, and at the last follow-up were compared using repeated-measures ANOVA. $P < 0.05$ was considered to be statistically significant. All analyses were conducted using SPSS 19.0 software (SPSS, Inc., Chicago, IL, USA).

3. Results

The location of the sequestration were L2 in 2 (9.5%) patients, L3 in 8 (38.0%) patients, L4 in 9 (43.0%) patients, L5 in 2 (9.5%) patients (baseline patient characteristics are shown in Table 1). All patients had HZLDH and successfully underwent the MELS approach. The mean operation duration was 32.43 ± 7.19 min and the mean blood loss was 25.52 ± 5.37 ml. The mean length of the hospital stay was 4.19 ± 0.87 days. We observed no intra- or perioperative complications.

Table 1
Baseline characteristics (N = 21)

Age (range)	58.66 ± 7.77 (47–77)
Gender	
Male	13 (61.9%)
Female	8 (38.1%)
Body Mass Index (range)	25.81 ± 2.15 (22–30)
Smoking history	4 (19.0%)
Chronic comorbidity	
Hypertension	4 (19.0%)
Diabetes mellitus	2 (9.5%)
Location of sequestration	
L2	2 (9.5%)
L3	8 (38.0%)
L4	9 (43.0%)
L5	2 (9.5%)
Total	21 (100%)

3.1 Clinical outcomes

The mean follow-up period was 20.95 ± 2.09 months (range: 18–24 months). The VAS and ODI were significantly improved at all follow-up intervals ($p < 0.01$, Table 2). As measured using the modified MacNab criteria, 17 patients (81%) obtained an “excellent” outcome and the remaining 4 (19%) patients obtained a “good” outcome (Table 3). Postoperative lower back pain occurred in 5 patients (23.8%), but could be significantly relieved with the use of painkillers. One patient suffered severe leg pain one year after the initial surgery and was diagnosed with contralateral lumbar disc re-herniation of the same level that was initially operated on. The dynamic lumbar X-Ray before the second surgery showed that no instability occurred. After performance of an endoscopic transforaminal discectomy, the patient underwent complete recovery.

Table 2
The changes of ODI and VAS after surgery

	Pre-op	Post-op	1 month	6 months	1 year	Last follow-up	p ¹	p ²	p ³
VAS ¹	7.88 ± 0.70	1.14 ± 0.79	0.43 ± 0.51	0.14 ± 0.36	0.43 ± 1.75	0.10 ± 0.30	< 0.05	< 0.05	< 0.05
VAS ²	3.71 ± 1.38	1.62 ± 0.92	0.76 ± 0.62	0.48 ± 0.51	0.57 ± 0.93	0.19 ± 0.40	< 0.05	< 0.05	< 0.05
ODI	59.24 ± 10.83	35.81 ± 6.04	25.71 ± 7.40	14.52 ± 7.45	14.29 ± 5.63	11.29 ± 3.59	< 0.05	< 0.05	< 0.05

Pre-op: preoperative; Post-op: 3 days after surgery; VAS¹ indicates the leg pain score, VAS² indicates the lower back pain score; P¹: comparison between Pre-op and Post-op; P²: comparison between Post-op and 1 year; P³: comparison between Post-op and last follow-up.

Table 3
Outcomes of the modified MacNab criteria

Grade	Definition	n (%)
Excellent	complete relief of symptoms, back to normal work and life	17 (81%)
Good	mild discomfort and still able to participate in all activities;	4 (19%)
Fair	symptoms alleviated, limited mobility, influence the normal activities	0
Poor	no better than preoperative status, even worse	0

3.2 Radiography outcomes

All patients showed complete removal of the fragments and no patient exhibited pars fracture as observed in the immediate postoperative CT and MRI. The dynamic X-Ray taken at least one year after surgery revealed that no patient showed postoperative instability.

4. Discussion

Definition of HZLDH

It has been estimated that roughly 10–20% of all disc herniations migrate in a cranio-lateral direction and may hence be located in the preforaminal and foraminal regions of the “hidden zone” [6]. Some authors

specifically describe the hidden zone as an area bordered laterally by the pedicle, ventrally by the dorsal part of the vertebral body and covered dorsally by the pars interarticularis of the hemilamina [5, 6]. However, the description of this type of disc herniation varies from author to author. Soldner *et al.* [11] use “canalicular” instead of “foraminal” and termed the HZLDH as “canalicular and cranio-posterolateral” lumbar disc herniation. Schulz *et al.* [14] termed sequestration in the hidden zone as “craniolateral lumbar disc herniation”. Papavero *et al.* [15] described this pathology as “fragment extruded cephalad into the spinal or root canal impinged the exiting root”. Despite the wide variety of descriptions, all authors mentioned above recognize the difficult surgical exposure in this clinical scenario. (Fig. 4)

Current surgical strategies

The standard surgical procedure for HZLDH is the interlaminar approach, during which removal of a major portion of the pars interarticularis is necessary [7, 9]. Donaldson *et al.* [16] reported that the resection of more than 50% of the facet joint is required in approximately two thirds of cases using the microsurgical interlaminar approach, which is likely to cause postoperative instability. For this reason, the interlaminar approach has gradually lost its popularity [8]. Since Di Lorenzo introduced the microscopic translaminar approach, the concept of preserving the bony borders of the lamina and sparing facet joints has been widely accepted in the management of HZLDH and has since been further described by several authors [11–14]. Several further novel approaches have been described recently. Wang *et al.* [17] described an approach called endoscopic transpedicle fenestration, with a bony hole drilled on the pedicle, sparing the lamina and facet joints. Reinshagen *et al.* [18] introduced the microscopic translaminar crossover approach. An angled fenestration is created in the contralateral hemilamina, whereby the medial portion of the hemilamina, just at the base of the spinous, is targeted to the hidden zone on the symptomatic (ipsilateral) side. This approach offers advantages when treating recurrent patients who previously underwent laminotomy. Here, we introduced the MELS for the first time to provide a novel option for the HZLDH.

Considerations of surgical anatomy

Papavero *et al.* [15] compared a disc fragment that is extruded cephalad underneath the lamina to a fish underneath the surface of a frozen lake, with there being two methods to hook the “fish”. The first is to cross the surface with an icebreaker and to catch the fish. The second option is to cut a small hole in the ice surface targeting the fish and to cast the rod. The translaminar approach remains popular for the treatment of HZLDH, however some authors have argued that this technique has its limitations [6]. Daghighi *et al.* [19] reported that disc fragments cranially migrated into the hidden zone are more commonly seen in higher lumbar levels. Due to segment-dependent changes of vertebral anatomy, the translaminar approach must be located very laterally in the upper lumbar levels in order to reach the medial hidden zone [11, 15, 20] (Fig. 5). Disruption of the lateral hemilamina and pars interarticularis have furthermore been correlated to an increased risk of stress fracture and instability [6, 12].

In our technique, the sequestered nuclear pulposus impinging the exiting nerve root can be compared to a melon on a vine, thus the “MELS” can be described using the Chinese saying “follow the vine to get the

melon" (tracking along the stem). Along the lateral border of the lamina, the vine can be easily found, thus the melon near the vine can be explored and dragged out with a hook. When managing the higher level with relative slender lamina, an undercut of the lateral hemilamina is enough to find the nerve root and the fragment. When managing fragments at the hidden zone of the L5 level, as the above-mentioned segment-related anatomy features, the wide and short lamina of L5 lead to a relatively small operating space, thus small parts of antero-superior S1 articular process were sometimes needed to explore the deep located L5 nerve root. Ivanov *et al.* [12] reported that the lateral half of the pars has the largest thickness and removing one fourth of the lateral aspect of the isthmus has minimal influence on the stresses in the remaining neural arches. In the MELS, we removed only 2–3 mm lateral margin of lamina and the isthmus was maintained almost intact. No radiological instability was observed at the final follow-up in all patients and no patient suffered severe postoperative low back pain-induced disability.

Fragmentectomy without discectomy

There is still a dispute regarding whether to deal with the intervertebral space or not in the treatment of HZLDH. Faulhauer *et al.* [21] proposed that fragment excision is superior to conventional disc removal due to a lower rate of postoperative spinal instability complications, while Kotil *et al.* [22] reported an increased recurrence rate for fragmentectomy compared with discectomy. Ebeling *et al.* [23] reported that cranio-lateral disc herniations commonly appear as a complete sequestration, thus management of intervertebral space is seldom needed. Moreover, Barth *et al.* [24] reported that discectomy did not result in additional benefits in the treatment of lumbar disc herniation, and the sequestrectomy demonstrated significantly less postoperative disc degeneration than standard microdiscectomy. In view of these considerations, we chose not to manage the intervertebral space in most of our patients. However, we performed disc removal in two cases because we found that the migrated fragments connected to the inferior disc closely, which indicated a tendency that more fragments would come out along the path. The recurrence rate of 4.8% (1/21) within the 2 years follow-up was relatively low compared with the study by Papavero *et al.* (7%) and was comparable with the study by Soldner *et al.* (3.3%) [11, 15, respectively]. However, larger patient groups and longer follow-up periods are needed to clarify the necessity of discectomy after fragmentectomy.

Limitations

There are several limitations in this study. Firstly, our sample size was small, involving only 21 patients. Although the main purpose of this study was to introduce this novel approach, and a satisfactory outcome was obtained in the present study, larger sample sizes will be needed to provide stronger evidence for our conclusions. Secondly, management of L5 may be slightly more challenging, as the operation space is relative small. Finally, it is difficult to access the superior intervertebral space via this extralaminar approach, though the inferior discectomy is feasible. However, if superior discectomy is required, we would recommend the use of other approaches.

5. Conclusion

Our results suggest that MELS is a safe and effective surgical treatment of HZLDH. This approach possesses several advantages: 1) easy to grasp for most spine surgeons, with a flat learning curve; 2) simple approach for sequestration and more reliable decompression; and 3) less disruption to the pars interarticularis and less risk of iatrogenic lumbar instability. Thus, we conclude that MELS represents a good surgical option for the treatment of HZLDH.

Abbreviations

List of abbreviations	
MELS	microscopic extra-laminar sequestrectomy
HZLDH	"hidden zone" lumbar disc herniation
VAS	visual analog scale
ODI	Oswestry disability index
MRI	magnetic resonance imaging
CT	computerized tomography

Declarations

Ethics approval and consent to participate

The Ethics Committee of Yantai affiliated hospital of Binzhou medical university give the approval for the study. Written informed consent to participate in this study was obtained from all patients.

Consent to publication

All participants gave written informed consent for their personal or clinical details along with any identifying images to be published in this study.

Availability of data and materials

All data used by or generated in this study is available from the corresponding author upon reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

No funding was received for this research.

Author's contributions

1. T designed this study and perform all the surgeries as the senior surgeon.
2. W participated in the surgery and the study design.
3. Z collected the data and wrote the manuscript.
4. Tang participated in the proof of this manuscript.

Other authors participated in the follow-up and data collection.

All authors read and approved the final manuscript.

Acknowledgements

Not applicable.

References

1. Luis RV, Emiliano NV, Juan ESH, Gustavo G. Lumbar Disc Herniation. *Rev Bras Ortop.* 2010;45(1):17–22.
2. Chen BL, Guo JB, Zhang HW, Zhang YJ, Zhu Y, Zhang J, et al. Surgical versus non-operative treatment for lumbar disc herniation: a systematic review and meta-analysis. *Clin Rehabil.* 2018;32:146–60.
3. Lurie JD, Tosteson TD, Tosteson AN, Zhao W, Morgan TS, Abdu WA, et al. Surgical versus nonoperative treatment for lumbar disc herniation: eight-year results for the spine patient outcomes research trial. *Spine.* 2014;39:3–16.
4. Wiltse LL, Berger PE, McCulloch JA. A system for reporting the size and location of lesions in the spine. *Spine (Phila Pa 1976).* 1997;22:1534–7.
5. Macnab I. Negative disc exploration. An analysis of the causes of nerve-root involvement in sixty-eight patients. *J Bone Joint Surg Am.* 1971;53:891–903.
6. Clemens R, Navid R, Marek M, Bernhard R. Surgical Approaches to the Lumbar Hidden Zone: Current Strategies and Future Directions. *EBioMedicine.* 2015;2:1005–7.
7. Caspar W, Campbell B, Barbier DD, Kretschmmer R, Gotfried Y. The Caspar microsurgical discectomy and comparison with a conventional standard lumbar disc procedure. *Neurosurgery.* 1991;28:78–86.
8. Abumi K, Panjabi MM, Kramer KM, Duranceau J, Oxland T, Crisco JJ. 1990. Biomechanical evaluation of lumbar spinal stability after graded facetectomies. *Spine.* 1990;15:1142–1147.
9. Schulz C, Kunz U, Mauer UM, Mathieu R. Early postoperative results after removal of cranially migrated lumbar disc prolapse: retrospective comparison of three different surgical strategies. *Adv Orthop.* 2014;702163.
10. Di Lorenzo N, Porta F, Onnis G, Cannas A, Arbau G, Maleci A. Pars interarticularis fenestration in the treatment of foraminal lumbar disc herniation: a further surgical approach. *Neurosurgery.*

- 1998;42:87–89; discussion:89–90.
11. Soldner F, Hoelper BM, Wallenfang T, Behr R. The translaminar approach to canalicular and cranio-dorsolateral lumbar disc herniations. *Acta Neurochir (Wien)*. 2002;144:315–20.
 12. Ivanov AA, Faizan A, Ebraheim NA, Yeasting R, Goel VK. The effect of removing the lateral part of the pars interarticularis on stress distribution at the neural arch in lumbar foraminal microdecompression at L3–L4 and L4–L5: anatomic and finite element investigations. *Spine*. 2007;32:2462–6.
 13. Dezawa A, Mikami H, Sairyo K. Percutaneous endoscopic translaminar approach for herniated nucleus pulposus in the hidden zone of the lumbar spine. *Asian J Endosc Surg*. 2012;5:200–3.
 14. Schulz C, Abdeltawab A, Mauer UM. Translaminar approach to cranio-laterally herniated lumbar disc prolapse. *Acta Neurochir (Wien)*. 2012;154:711–4.
 15. Papavero L, Langer N, Fritzsche E, Emami P, Westphal M, Kothe R. The translaminar approach to lumbar disc herniations impinging the exiting root. *Neurosurgery*. 2008;62:173–7. discussion 177-8.
 16. Donaldson WF, Star MJ, Thorne RP. Surgical treatment for the far lateral herniated lumbar disc. *Spine*. 1993;18:1263–7.
 17. Wang D, Pan H, Hu Q, Zhu H, Zhu L, He Y, et al. Percutaneous endoscopic transpedicle approach for herniated nucleus pulposus in the lumbar hidden zone. *Asian J Endosc Surg*. 2017;10:87–91.
 18. Reinshagen C, Ruess D, Molcanyi M, Redjal N, Walcott BP, Goldbrunner R, et al. A novel translaminar crossover approach for pathologies in the lumbar hidden zone. *J Clin Neurosci*. 2015;22:1030–5.
 19. Daghighi MH, Pouriesa M, Maleki M, Fouladi DF, Pezeshki MZ, Mazaheri KR, et al. Migration patterns of herniated disc fragments: a study on 1,020 patients with extruded lumbar disc herniation. *Spine J*. 2014;14:1970–7.
 20. Reulen HJ, Müller A, Ebeling U. Microsurgical anatomy of the lateral approach to extraforaminal lumbar disc herniations. *Neurosurgery*. 1996;39:345 – 50; discussion 350–351..
 21. Faulhauer K, Manicke C. Fragment excision versus conventional disc removal in the microsurgical treatment of herniated lumbar disc. *Acta Neurochir (Wien)*. 1995;133:107–11.
 22. Kotil K, Köksal NS, Kayaci S. Long term results of lumbar sequestrectomy versus aggressive microdiscectomy. *J Clin Neurosci*. 2014;21:1714–8.
 23. Ebeling U, Reulen HJ. Are there typical localisations of lumbar disc herniations? A prospective study. *Acta Neurochirurg (Wein)*. 1992;117:143–8.
 24. Barth M, Diepers M, Weiss C, Thomé C. Two-year outcome after lumbar microdiscectomy versus microscopic sequestrectomy: part 2: radiographic evaluation and correlation with clinical outcome. *Spine*. 2008;33:273–9.

Figures




Figure 1

Illustration of the anatomical parameters in different levels. The solid black line indicates the width of lamina (from L5-L3: 22 mm, 18.2 mm, 15.4 mm); the dotted black line indicates the height of lamina (from L5-L3: 17.3 mm, 21.2 mm, 23.1 mm); the solid white line indicates the distance from lateral margin of pars to lateral border of vertebral body (from L5-L3: 2.8mm, 4.8mm, 5.3mm).




Figure 2


Spinal canal area classification. The red area indicates the foraminal zone (canalicular zone); the green area indicates the preforaminal zone or subarticular zone; the blue area indicates the central canal zone, and the transparent square zone indicates the hidden zone.



A




B



C


Figure 3

a,b: The preoperative MRI showed the migrated fragments were obviously linked with the inferior disc. c: the intervertebral space can also be approached after removing a small part of the IAP (white arrow: exiting nerve root; black arrow: posterior wall of vertebral body; yellow arrow: intervertebral space).



A

B



C




D




E

Figure 4

a, b: The preoperative MRI showed a complete sequestered fragment. c: The intra-operative view: the exiting nerve root was exposed after the “crescent-shaped” excision of the lamina (white circle: exiting nerve root; white star: lateral margin of the lamina; SAP: superior articular process; IAP: inferior articular process). d,e: immediate postoperative CT shows the crescent-shaped excision of the lateral lamina (white arrow).



A



B

Figure 5

Schematic illustration of the extralaminar approach. a: After crescent-shape excision of the lamina, the sequestered fragments can be explored with a hook along the exiting nerve root. b: Postoperative three-dimensional reconstruction detailing the shape of excision (highlighted with the white arrow).