Synthetic hormone boosts diabetic wound healing in rats

Yi-Fan Shen

Jing-Huan Huang

Kai-Yang Wang

Jin Zheng

Lin Cai

Hong Gao

Xiao-Lin Li

Jing-Feng Li

Video Byte

Keywords: Cell Communication and Signaling, PTH, multifunctional factor, diabetic chronic wound, exosomes, synergistic effect, hormone, PTHrP-2, epithelial cell, endothelial cell, angiogenesis, HUVEC, exosomes, fibrogenesis, epithelialization, wound healing, diabetes, hyperglycemia, PI3K-Akt, fibroblast

Posted Date: June 23rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-37679/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Diabetic wounds are a growing problem worldwide. One solution could lie in a hormone secreted by the parathyroid. PTH is critical for re-growing bone. But evidence suggests that it could also help regenerate skin and blood vessels. Researchers recently tested this hypothesis on rats. Applying a synthetic version of PTH to diabetic wounds significantly improved wound healing. But not as they believed it would. Synthetic PTH did not appear to directly activate the cells it repaired. Instead, separate experiments showed that the PTH derivative indirectly led to repair, using exosomes, tiny sacs ejected by cells to communicate with other cells. If replicated in more realistic models of diabetes, these findings could lead to a powerful new way to accelerate wound healing.