Negative Regulation of Ptch1 by miR-342-5p and FoxO3 Induced Autophagy Involved in Renal Fibrosis

Simin Tang

First People's Hospital of Foshan

Yi Wang

The Third Affiliated Hospital of Southern Medical University

Guiling Xie

The Third Affiliated Hospital of Southern Medical University

Wenjun Li

The Third Affiliated Hospital of Southern Medical University

Yanna Chen

The Third Affiliated Hospital of Southern Medical University

Jinshu Liang

The Third Affiliated Hospital of Southern Medical University

Pei Liu

The Third Affiliated Hospital of Southern Medical University

Fuhu Song

The Third Affiliated Hospital of Southern Medical University

Jun Zhou (✉ zhoujun7843@126.com)

The Third Affiliated Hospital of Southern Medical University <https://orcid.org/0000-0003-1545-8092>

Research

Keywords: Renal fibrosis, miR-352-5p, Ptch1, FoxO3, Bioinformatics, Autophagy

Posted Date: June 26th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-37890/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Frontiers in Bioengineering and Biotechnology on October 29th, 2020. See the published version at <https://doi.org/10.3389/fbioe.2020.583318>.

Abstract

Background: The pathogenesis of renal fibrosis (RF) is not well understood. Here, we performed an integrative database analysis of miRNAs and mRNAs to discover the major regulatory pathway in RF.

Methods: Putative miRNAs and mRNAs involved in RF in unilateral ureteral obstruction (UUO) model mice were extracted and analyzed using the Gene Expression Omnibus (GEO) database. The bioinformatics analysis suggested that *Ptch1* expression is negatively regulated by miR-342-5p and FoxO3. Then real-time PCR, Western blot, Fluorescence in situ hybridization were done to confirm the hypothesis.

Results: Sixty-three differentially expressed miRNAs (DE-miRNAs) in GSE118340, 141 DE-miRNAs in GSE42716, and 183 DE-mRNAs in GSE69101 were identified. Various bioinformatic analyses revealed miR-342-5p as a strong candidate regulator in RF. We also predicted that miR-342-5p targets *Ptch1* and that FoxO3 is the transcription factor of *Ptch1*. We also observed that TGF- β 1 upregulated the expression of miR-342-5p and inhibited the expression of FoxO3 and *Ptch1* in TCMK-1 cells. Furthermore, downregulation of miR-342-5p or FoxO3 reversed the inhibitory effect of TGF- β 1 on the expression of *Ptch1* in TCMK-1 cells. Additionally, downregulation of *Ptch1* increased TGF- β 1-induced autophagy, as evidenced by an increase in the number of GFP-LC3 puncta and the increased protein expression of SOSTM1/p62 and LC3II/LC3I.

Conclusion: Our findings showed that downregulation of *Ptch1* induced autophagy in TGF- β 1-stimulated TCMK-1 cells, and *Ptch1* expression is negatively regulated by miR-342-5p and FoxO3. These findings will further our understanding of the molecular mechanisms of RF and provide useful novel therapeutic targets for RF.

1. Background

As the terminal pathological alteration of almost all progressive kidney diseases, renal fibrosis (RF) is characterized by hyperproliferation, activation of fibroblasts, and increased deposition of extracellular matrix (ECM), thereby distorting normal kidney tissue architecture and causing loss of kidney function [1]. There is an urgent unmet clinical need for effective therapies and understanding of the molecular pathogenesis of RF. MicroRNAs (miRNAs) are endogenous small non-coding 21-25 nucleotide single-stranded RNAs. miRNAs regulate gene expression by promoting the degradation of mRNAs or inhibiting the translation of mRNAs. miRNAs are involved in the development of various diseases, and several miRNAs are closely associated with RF, including miR-184, miR-34a, and miR-199a-3p [2-4]. Our review showed the recent progress, biological functions, and clinical applications of miRNAs in the progression of RF [5]. However, the role of most miRNAs in RF is still unclear. Autophagy is a highly evolutionarily conserved lysosomal degradation pathway that plays a vital role in maintaining cellular homeostasis [6]. Autophagy dysfunction is implicated in the pathogenesis of acute kidney injury and RF diseases [7-9]. In the past few years, miRNA-mediated autophagy has been reported to be involved in RF [10,7,11]. However, the specific mechanism by which miRNA-mediated autophagy regulates RF has not been fully elucidated.

Obstructive ureteral mechanical damage is the primary clinical etiology of RF. Exploring the genetic changes in the obstructive kidneys helps to understand the molecular and cellular mechanisms involved in the pathogenesis of RF. Unilateral ureteral obstruction (UUO) is a classic animal model that can simulate obstructive nephropathy [12]. Recently, with the rapid development of microarray and high-throughput sequencing technologies, the mechanisms of RF at the genomic level have been increasingly explored [13-15]. We analyzed two comprehensive and integrated miRNA datasets (GSE118340 and GSE42716) and one mRNA dataset (GSE69101) from the Gene Expression Omnibus (GEO) database. Two representative DE-miRNAs (miR-342-5p and miR-466a-3p) were shared and extracted for further analysis. Previous studies have reported that miR-342-5p was strictly related to the deposition of extracellular matrix (ECM), inflammatory

cytokine release, and the TGF- β signaling pathway [16-18], which are involved in the mechanisms of RF. However, the mechanisms by which miRNAs mediate the specific molecular changes observed in RF have not been studied. There are no studies on the role of miR-342-5p in the development of RF. Thus, we further focused on the mechanism by which miR-342-5p is involved in RF progression.

Combined with in-depth bioinformatics analysis, we predicted that miR-342-5p directly targets Ptch1, which is an endogenous inhibitor of pro-fibrotic signaling pathways, and is regulated by the transcription factor (TF) FoxO3. Herein, we verified the relationship between miR-342-5p, FoxO3, and Ptch1 in vitro. Our results indicated that miR-342-5p expression was upregulated in TCMK-1 cells stimulated with TGF- β 1, while FoxO3 and Ptch1 were downregulated. Rescue experiments showed that miR-342-5p promoted the fibrotic process by mediating Ptch1 and FoxO3. In addition, autophagy was increased in response to TGF- β 1, and knockdown of Ptch1 in TCMK-1 cells increased autophagy markers. Collectively, these data suggest that miR-342-5p and FoxO3 regulate Ptch1, which mediates autophagy involved in the RF process. These findings further the understanding of the molecular mechanisms of RF and provide therapeutic targets for RF diseases.

2. Methods

2.1 Microarray dataset

Two miRNA datasets, GSE118340 and GSE42716, were downloaded from the GEO database (<https://www.ncbi.nlm.nih.gov/geo/>). GSE11840 was based on the GPL19057 (Illumina NextSeq 500), and GSE42716 was based on the GPL10384 [Agilent-021828 Unrestricted Mouse miRNA Microarray (V2)]. In total, four kidney samples of the UUO model (GSM3325596, GSM3325597, GSM3325598, GSM3325599) and three kidney samples of the sham control (GSM3325589, GSM3325590, GSM3325591) from GSE118340 were selected for analysis. Four kidney samples of the UUO model (GSM1048441, GSM1048442, GSM1048443, GSM1048444) and four kidney samples of the sham control (GSM1048437, GSM1048438, GSM1048439, GSM1048440) from GSE42716 were also selected. The mRNA dataset GSE96101 was also downloaded from the GEO dataset, which is based on the GPL4134 (Agilent-014868 Whole Mouse Genome Microarray 4x44K G4122F). Six kidney samples of the UUO model (GSM2533412, GSM2533413, GSM2533414, GSM2533415, GSM2533416, GSM2533417) and five kidney samples of the sham control (GSM2533473, GSM2533474, GSM2533475, GSM2533476, GSM2533477) were selected for analysis. The above kidney samples were all harvested from mice seven days after UUO surgery and then were subjected to microarray analysis of miRNAs and mRNAs. Contralateral kidneys without ureteral obstruction were used as controls. Subsequently, data quality from these datasets was assessed using R software.

2.2 Identification of differentially expressed miRNAs (DE-miRNAs) and mRNAs (DE-mRNAs)

The R software “DESeq2” package was used to identify DE-miRNAs in kidney tissues between UUO and sham control [19]. DE mRNAs were identified using the online tool GEO2R (<http://www.ncbi.nlm.nih.gov/geo/geo2r>), which allows users to compare two or more datasets in the GEO series to determine DEGs under experimental conditions [20]. Genes with more than one probe set were removed. In the present study, DE-miRNAs and DE-mRNAs were combined using the cut-off point of adj.P-value < 0.05, and $|\log FC| > 1.0$. Then, the R heatmap and ggplot2 package were applied to paint the heatmap and Volcano. adj.P < 0.05 means statistical significance.

2.3. GO annotation and KEGG pathway enrichment analysis

The selected DE-miRNAs were then uploaded to miRpath v3.0 (<http://www.microrna.gr/miRPathv3/>), an online website that integrates miRNA target gene databases, including TarBase, TargetScan, and microT-CDS [21]. It only needs to input the miRNA Id of interest and obtain the results of Gene Ontology (GO) and Kyoto Encyclopedia of Gene and

Genome (KEGG) enrichment analysis. The selected DE-mRNAs were then uploaded to DAVID (<https://david.ncifcrf.gov>), a database that conducts functional annotation bioinformatics microarray analysis [22]. The top 10 GO annotations and KEGG pathways were visualized using GraphPad Prism.

2.4. miRWalk and Targetscan

miRWalk (<http://mirwalk.umm.uni-heidelberg.de/>) can predict the mRNAs targeted by specific miRNAs using a machine learning algorithm. Thus, this database was used to predict the targeted mRNAs of miR-342-5p. Targetscan 7.2 (http://www.targetscan.org/mmu_72/), an online database that is currently used to predict the location of miRNA binding sites with an accuracy of up to 90%, was applied to revalidate the binding sites of miRNA and mRNA.

2.5 Comparative Toxicogenomics Database (CTD)

CTD (<http://ctdbase.org>) is a powerful, publicly available database designed to advance understanding of how environmental exposures affect human health and to clarify the relationship among genes, drugs, and diseases. We used this database to evaluate the inference score of selected mRNAs in chronic kidney diseases (CKDs).

2.6 The iRegulon

The iRegulon, a Cytoscape App that consists of a transcription factor (TF) and its direct transcriptional targets, which contain common TF binding sites in their cis-regulatory control elements. It predicts TFs by calculating motif enrichment analysis. The motif enrichment analysis applies multiple position weight matrices (PWM), and finally sorts each motif. The preferred motif was used to predict the final TFs [23].

2.7 Cell culture and treatment

Mouse kidney epithelial cell line TCMK-1 was cultured in RPMI-1640 (HycloneCat.No.SH30809.01B) supplemented with 10% fetal bovine serum (Hyclone, Cat.No.SH30087.01) and 1% penicillin-streptomycin (Hyclone, Cat.No. SH30010) at 37°C under saturated humidity and 5% CO₂ conditions. The cells were starved for 12 h; on reaching 70% confluence, TGF-β1 (eBioscience, 14-8342-82) was added to the medium at a final concentration of 10 ng/ ml for 0, 1, 2, 4, 8, 12, and 24 h, to choose suitable time points for the follow-up experiment.

2.8 Cell transfection

TCMK-1 cells were passaged one day before transfection to obtain 30%-50% confluence. Transfection was performed using Lipofectamine™ RNAiMAX (Invitrogen, Cat. 13778075), and the miRNA / siRNA working concentration was 50 nM using Opti-MEM (Invitrogen, Cat. No. 31985070). In LC3 plasmid cell transfection, TCMK-1 cells were passaged one day before transfection to obtain 70%-80% confluence. Transfection was performed using Lipofectamine 2000 (Invitrogen, Cat. 11668019), Premo™ Autophagy Sensor LC3B-RFP (BacMam 2.0 Invitrogen™, P36236), and Opti-MEM (Invitrogen, Cat. No. 31985070). Approximately four h after transfection, the transfection mixture was replaced with DMEM-10% FBS. The punctate distribution of GFP-LC3 was observed using fluorescence microscopy. The remaining cells were then harvested for RNA and protein preparation.

2.9 Quantitative real-time PCR

Total RNA of TCMK-1 cells was extracted with TRIzol® Reagent (Invitrogen, 15596026) according to the manufacturer's protocol. DNA was removed by treating with DNase (Fermentas, EN0521) and the purity of the RNA was assessed by determining the absorbance ratio at OD260/OD280. Total RNA was reverse transcribed into cDNA using TaKaRa PrimeScript II 1st Strand cDNA Synthesis Kit (TaKaRa, D6210A), Oligo dT Primer, dNTP Mixture, total RNA, and RNase

Free dH₂O. Real-time PCR was performed using the TaKaRa SYBR® Premix Ex Taq™ II kit (Perfect Real Time), and ABI PRISM® 7500 Sequence Detection System Real-Time PCR System. Primer sequences for the TCMK-1 cell samples were as follows:

FoxO3-F: 5' - AACCGGCTCCTTCAACAGTA

FoxO3-R: 5' GAAGCAAGCAGGTCTTGGA

Ptch1-F: 5' - TCTGCTTCGGTGAAGTGTG

Ptch1-R: 5' - CCACGTCCTGTAGCTCTATG

β-actin-F: 5' GCTTCTAGGCGGACTGTTAC

β-actin-R: 5' CCATGCCAATGTTGTCTCTT

miR-342-5p-F: 5' AACTCCAGCTGGGAGGGGTGCTATCTGTGAT

miR-342-5p-R: 5' CTCAACTGGT GTCGTGGA

miR-342-5p-RT:5' CTCAACTGGTGTCTGGAGTCGGCAATTCAGTTGAGCTCAATCA

U6-F:5' - CTCGCTTCGGCAGCACA

U6-R:5' AACGCTTCACGAATTTGCGT

U6-RT: 5' AACGCTTCAC GAATTTGCGT

Reactions to determine the levels of Ptch1 and FoxO3 were carried out in a total volume of 20 µL, including 10 µL SYBR® Premix Ex Taq™ (2×), 2 µL cDNA, 6.4 dH₂O, and 0.8 µL of each specific primer (10 µM), while the reaction for miR-342-5p was carried out in a total volume of 8 µL, including 2 µL 10 mM dNTP (Promega), 0.5 µL RNase inhibitor (Promega), 0.5 µL miR-342-5 primer, 0.5 µL U6 primer, 4 µL 5x buffer, and 0.5 µL M-MLV (Promega). The cycling conditions were set at 94°C for 3-4 min; 30 cycles at 94°C for 30 s, 60°C for 30 s, and 72°C for 30 s; the last cycle (1x) was set at 72°C for 10 min and 16°C for ∞. β-actin was utilized for standardization of Ptch1 and FoxO3, and U6 was used as the endogenous control for miR-342-5p. The levels of mRNAs and miRNAs were calculated using the 2^{-ΔΔCt} method with the ABI PRISM® 7500 Sequence Detection System SDS Shell Software 1.6.

2.10 Western blot

Protein samples were extracted with RIPA buffer. The extracted total protein was separated by SDS-PAGE and transferred to PVDF membranes. Following incubation in blocking buffer for 1 h at room temperature, the membrane was incubated at 4°C overnight with Ptch1 antibody (Abcam, ab109096), Foxo3 antibody (Thermo, FA1-14171), SQSTM1 / p62 antibody (Abcam, ab56416), and LC3B (D11) XP antibody (Cell Signaling Technology, 3868S). The results were visualized using Beyo ECL Plus substrate and exposed to X-ray film. β-actin was used for normalization.

2.11 Fluorescence in situ hybridization

Fluorescence in situ hybridization (FISH) was performed to detect the location of miR-342-5p, FoxO3, and Ptch1. We used a miR-342-5p probe labeled with FAM (green) and FoxO3 and Ptch1 probes labeled with cy3 (red). The cells were then visualized using a scanning laser confocal microscope (Leica, TCS SP2 AOBs).

2.12. Statistical Analyses

All data are shown as the mean $\pm$ standard deviation (s.d) and visualized using GraphPad Prism 8. Statistical analysis was conducted using SPSS 24.0. $P < 0.05$ was considered significant. All experiments were performed at least three times.

3. Results

3.1 Identification of candidate DE-miRNAs

To identify potential miRNA-mRNA regulatory networks in RF, the datasets GSE118340 and GSE42716 were selected to screen DE-miRNAs between UUO samples and normal samples. Briefly, the distribution of gene expression data in each group was revealed in the heatmap and unsupervised clustering analysis (Figure 2A and Figure 2B). Sixty three DE-miRNAs and 141 DE-miRNAs (adj.P < 0.05 , $|\log FC| > 1$) were detected from GSE118340 and GSE42716 using DESeq2 package analysis. A volcano plot was used to visualize the DE-miRNAs from the two datasets (Figure 2C and Figure 2D). The Venn diagram showed that only two DE-miRNAs (miR-342-5p and miR-466a-3p) were shared in both the GSE118340 and GSE42716 datasets and were extracted for further analysis (Figure 2E).

3.2 GO annotation and KEGG pathway enrichment analysis

miR-342-5p and miR-466a-3p relevant GO annotation and KEGG pathways were identified using the mirPath v3.0 web server, and the results showed that 39 GO and 13 KEGG pathways were significantly ($P < 0.05$) regulated by miR-342-5p, while 36 GO and 13 KEGG pathways (Supplementary Materials Table S1 and Table S2) were significantly regulated by miR-466a-3p. The top 10 GO annotations and KEGG pathways of miR-342-5p and miR-466a-3p are shown in Figures 3A, Figure 3B, 3C, and 3D. All the above identifications were based on merging and meta-analysis algorithms according to experimentally validated miRNA target interactions. The pathways regulated by the miR-342-5p closely related to the pathophysiology of RF should be involved in ECM-receptor interaction; whereas, miR-466a-3p is closely associated with the TGF- β signaling pathway.

3.3 Identification of DE-mRNAs

Using the GEO2R online tool, 183 DEGs (adj.P < 0.01 , $|\log FC| > 1$) were detected in the GSE96101, including 177 upregulated and 6 downregulated DEGs. In addition, a volcano plot and a heatmap of all DE-mRNAs were generated using the R ggplot2 package (Figure 4A and **Figure 4B**). Then, mRNAs obtained from miRwalk and DE-mRNAs extracted from GSE96101 were subjected to Venn analysis. Venn diagram showing 48 mRNAs were overlapped and are listed below in detail (Figure 4C). GO annotation and KEGG pathways were obtained from DAVID and are summarized in Supplementary Materials (Table S3). We found that Ptch1 was mainly involved in intracellular protein transport, protein localization, and protein processing.

3.4 Prediction of miR-342-5p and FoxO3 correlation with Ptch1

Next, we used Targetscan7.2 to revalidate the binding sites of miRNAs and mRNAs, further narrowing the scope of the interaction of miRNA-mRNA. We found that only 8 DE-mRNAs were interacted with to miR-342-5p. Among them, Vsx2 and Ptch1 had the higher context score (Table 1). Then, the CTD database was used to assess the inference score of these 8 DE-mRNAs in CKDs. The results showed that Dhcr24 and Carhsp1 had higher inference scores (Table 2). Combining the results obtained from these two databases, we used regression to calculate and select the most suitable target genes of miR-342-5p. The results showed that Ptch1 is the most promising target mRNA for miR-342-5p (Figure

5A). Furthermore, we used the iRegulon to predict the TFs of the network. The results showed that FoxO3 is a potential TF of Ptch1 (Figure 5B).

Table 1 Targetscan7.2 predicts the mRNA targets of miR-342-5p

Target gene	Representative transcript	3P-seq tags + 5	Total sites	8mer sites	7mer-m8 sites	7mer-A1 sites	6mer sites	Cumulative weighted context score	Total context score
Vsx2	ENSMUST000000021665.6	5	2	2	0	0	1	-0.61	-0.61
Ptch1	ENSMUST000000021921.5	4095	1	1	0	0	1	-0.36	-0.36
Gcnt2	ENSMUST000000110191.3	49	1	0	0	1	1	-0.13	-0.13
Six4	ENSMUST000000043208.7	266	1	0	0	1	1	-0.09	-0.09
Dhcr24	ENSMUST000000047973.3	9729	2	0	0	2	1	-0.08	-0.16
Vti1a	ENSMUST000000095950.2	693	1	0	0	1	0	-0.02	-0.03
Ap2m1	ENSMUST000000007216.8	2635	1	0	0	1	0	0	-0.08
Carhsp1	ENSMUST000000008537.8	1872	1	0	1	0	0	0	-0.19

Table 2 CTD database assesses the inference score of targeting mRNAs in chronic renal disease

Gene Symbol	Disease Name	Direct Evidence	Inference Score
Dhcr24	Kidney Failure, Chronic	0	33.74
Carhsp1	Kidney Failure, Chronic	0	25.27
Gcnt2	Kidney Failure, Chronic	0	17.3
Six4	Kidney Failure, Chronic	0	17.1
Vti1a	Kidney Failure, Chronic	0	16.9
Ap2m1	Kidney Failure, Chronic	0	16.4
Ptch1	Kidney Failure, Chronic	0	12.45
Vsx2	Kidney Failure, Chronic	0	2.65

3.5 TGF- β 1 induces upregulation of miR-342-5p and downregulation of FoxO3 and Ptch1 in TCMK-1 cells

To determine whether miR-342-5p, FoxO3, and Ptch1 correlate with the progression of RF, we measured the expression levels of miR-342-5p, FoxO3, and Ptch1 in TCMK-1 cells treated with TFG- β 1. The results showed that miR-342-5p expression levels were gradually increased, while those of FoxO3 and Ptch1 decreased in a time-dependent manner (Figure 5C, Figure 5D, and Figure 5E). Of note, FoxO3 and Ptch1 expression levels were the most significantly reduced at the 4-h time point. As demonstrated by FISH, miR-342-5p was mainly distributed in the cytoplasm of TCMK-1 cells, as was FoxO3 and Ptch1. The merged images showed that miR-342-5p was partially co-localized with Ptch1 (Figure 5F), indicating a potential interaction between them. These data indicated that miRNA-342-5p, FoxO3, and Ptch1 may be directly involved in the development of RF.

3.6 Ptch1 expression is regulated by miR-342-5p and FoxO3 in TCMK-1 cells treated with TGF- β 1

Real-time PCR and western blotting analyses showed that the expression levels of Ptch1 were regulated by miR-342-5p and FoxO3 in TCMK-1 cells treated with TGF- β 1. Downregulation of miR-342-5p reversed the inhibitory effect of TGF- β 1 treatment on Ptch1 expression in TCMK-1 cells, while the knockdown of *FoxO3* gene exacerbated the TGF- β 1-mediated inhibition of Ptch1 expression in TCMK-1 cells (Figure 6A, Figure 6B and 6C). These data suggest that miR-342-5p negatively regulates Ptch1, which is transcriptionally regulated by FoxO3 during the progression of RF.

3.7 Downregulation of Ptch1 aggravated TGF- β 1-induced autophagy in TCMK-1 cells

Autophagy mediates the development of RF, and a previous study showed that Ptch1 could inhibit autophagy in Hedgehog (Hh)-dependent cancers [24]. However, whether Ptch1 regulates autophagy and participates in the pathophysiological process of RF is not yet clear. Western blot analysis showed that the miR-342-5p inhibitor could reverse TGF- β 1-induced downregulation of Ptch1 expression, while FoxO3 siRNA promoted the effect of TGF- β 1 (Figure 6D and Figure 6E). In addition, inhibition of Ptch1 promoted the TGF- β 1-induced increase in LC3 II/I expression levels and also promote the TGF- β 1-induced decrease in SQSTM1/p62 expression levels in TCMK-1 cells. In addition, we found that autophagy indicated by GP-LC3 spots increased after TGF- β 1 stimulation. Downregulation of Ptch1 increased the number of GFP-LC3 spots (Figure 6F and Figure 6G), indicating that inhibition of Ptch1 could exacerbate TGF- β 1-induced autophagy in TCMK-1 cells. A schematic diagram showing the putative involvement of miR-342-5p, FoxO3, Ptch1, and autophagy in the progression of RF is shown in Figure 6H.

4. Discussion

RF is the typical final pathological phenotype of all CKDs [1]. Research on the mechanism of RF will contribute to a breakthrough in the treatment of CKD. In the past few years, even though the contribution of miRNAs to the pathogenesis of RF has been widely studied, the role of most miRNAs is not well understood. With the development of high-throughput sequencing analyses, an increasing number of studies have explored putative miRNAs that cause renal injury. Here, we evaluated an integrative network of miRNA and mRNA data to discover possible regulators of RF. In the present study, miR-342-5p was found to be a promising candidate miRNA affecting the progression of RF. KEGG pathway analysis of miR-342-5p showed that ECM receptor interaction is closely related to the pathophysiology of RF. As has been previously reported, the production of ECM proteins provided tissue scaffolds for normal repair events after renal injury [25,26]. Furthermore, researchers have found that miR-342-5p is upregulated more than sevenfold in paired samples of renal cell carcinoma tissue and adjacent non-tumorous renal parenchyma [27].

During the past few years, many studies have shown that the expression levels of miRNAs and their downstream target genes are closely associated with the progression of RF [3,4,2,10]. After applying various bioinformatics technologies as shown above, we found that Ptch1 is the most promising target mRNA of miR-342-5p in RF. Ptch1 is mainly expressed in epithelial cells around the renal tubules. Extracellular hedgehog (Hh) ligands can bind to the Ptch1 receptor, allowing transmembrane protein Smoothed (SMO) to signal downstream and activate Glioma (Gli) TFs [28,29]. It has been shown that hypermethylation of Ptch1 is associated with the perpetuation of fibroblast activation and fibrosis in the liver [30]. Similarly, inhibition of Ptch1 may have pro-fibrotic effects in kidney tissue by activating Hh signaling [31,32]. Furthermore, Ptch1 gene mutations have been suggested to play an essential role in the pathogenesis of nephroblastoma [33]. Bioinformatics analysis predicted that miR-342-5p interacts with Ptch1 in the progression of RF. Our in vitro experiments confirmed that TGF- β 1 induces up-regulation of miR-342-5p and down-regulation of Ptch1 in TCMK-1 cells. The rescue experiment implied that Ptch1 expression was regulated by miR-342-5p in TCMK-1 cells treated with TGF- β 1. Thus, miR-342-5p is most likely to play a crucial role in the development of RF by regulating Ptch1.

Our bioinformatics results showed that FoxO3 is the TF of Ptch1. FoxO3 plays a critical role in various biological processes including development, proliferation, apoptosis, metabolism, and differentiation, by regulating expression of a wide spectrum of genes. It has been shown that FoxO3 may attenuate the idiopathic pulmonary fibrosis myofibroblast phenotype in vitro and bleomycin-induced lung fibrosis in vivo [34], prevent the progression of kidney diseases [35,36] and promote kidney injury after UUO[37]. Hence, FoxO3 may play a role in promoting the fibrosis process by regulating Ptch1. Our results indicated that TGF- β 1 increases miR-342-5p and decreases FoxO3 and Ptch1 in TCMK-1 cells. FISH experiments indicated that miR-342-5p, FoxO3, and Ptch1 were mainly distributed in the cytoplasm of TCMK-1 cells. The merged images showed that miR-342-5p was partially colocalized with Ptch1. Thus, miRNA-342-5p might activate fibrotic responses in the kidney by interacting with Ptch1, which is directly regulated by FoxO3. As reported previously, Ptch1 has therapeutic implications in Hh-dependent cancers by inhibiting autophagy [29]. Autophagy can maintain cell homeostasis and energy under different physiological states [6]. Quantification of immunofluorescent LC3-GFP puncta showed that inhibiting Ptch1 can enhance the effect of TGF- β 1 on the promotion of autophagy. Therefore, the inhibition of Ptch1 could promote RF by activating TGF- β 1 induced autophagy.

5. Conclusion

Through bioinformatics analysis tools and in vitro experiments, we found that Ptch1 regulated by miR-342-5p and FoxO3 could promote autophagy, which plays an essential role in the progression of RF. Further study is required to fully elucidate the role of miR-342-5p, Ptch1, and FoxO3 in RF. Nevertheless, our novel findings provide new insights into the pathogenesis and therapeutic targets of RF.

List Of Abbreviations

Renal fibrosis: RF

Extracellular matrix: ECM

MicroRNAs: miRNAs

Unilateral ureteral obstruction: UUO

Gene Expression Omnibus: GEO

Transcription factor: TF

Differentially expressed: DE

Gene Ontology: GO

Kyoto Encyclopedia of Gene and Genome: KEGG

Comparative Toxicogenomics Database: CTD

chronic kidney diseases: CKDs

Fluorescence in situ hybridization: FISH

Hedgehog: Hh

SMO: Smoothen

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The data that support the findings of this study are downloaded from the GEO database (<https://www.ncbi.nlm.nih.gov/geo/>; GEO accession: GSE118340, GSE42716 and GSE121190). All in vitro experiments data generated or analyzed are included in this published article [and its supplementary information files].

Competing interests

The authors have no conflict of interest to declare.

Funding

This work was supported by Natural Science Foundation of Guangdong Province [2019A1515011087] (to Jun Zhou).

Authors' contributions

ST and YW contributed equally to this work. ST carried out the conception and the derivation of the study, carried out the bioinformatics analysis and drafted the manuscript. YW generate the data from the GEO and analyze the data preliminarily, and then help to performed real-time PCR, western blotting. WL, GX and JL conduct the cell culture and cell transfection. PL, YC and FS collect and analyze the numerical data. JZ guided the general research strategy and gave the critical revision of this manuscript. All authors read and approved this manuscript.

Acknowledgements

Authors would like to thank all members for their kind comments.

References

1. Boor P, Ostendorf T, Floege J (2010) Renal fibrosis: novel insights into mechanisms and therapeutic targets. *Nat Rev Nephrol* 6 (11):643-656. doi:10.1038/nrneph.2010.120
2. Liu Y, Bi X, Xiong J, Han W, Xiao T, Xu X, Yang K, Liu C, Jiang W, He T, Yu Y, Li Y, Zhang J, Zhang B, Zhao J (2019) MicroRNA-34a Promotes Renal Fibrosis by Downregulation of Klotho in Tubular Epithelial Cells. *Mol Ther* 27 (5):1051-1065. doi:10.1016/j.ymthe.2019.02.009
3. Chen B (2019) The miRNA-184 drives renal fibrosis by targeting HIF1AN in vitro and in vivo. *Int Urol Nephrol* 51 (3):543-550. doi:10.1007/s11255-018-2025-4
4. Yang R, Xu X, Li H, Chen J, Xiang X, Dong Z, Zhang D (2017) p53 induces miR199a-3p to suppress SOCS7 for STAT3 activation and renal fibrosis in UUO. *Sci Rep* 7:43409. doi:10.1038/srep43409
5. Fan Y, Chen H, Huang Z, Zheng H, Zhou J (2020) Emerging role of miRNAs in renal fibrosis. *RNA Biol* 17 (1):1-12. doi:10.1080/15476286.2019.1667215

6. Klionsky DJ (2007) Autophagy: from phenomenology to molecular understanding in less than a decade. *Nat Rev Mol Cell Biol* 8(11):931-937. doi:10.1038/nrm2245
7. Nam SA, Kim WY, Kim JW, Park SH, Kim HL, Lee MS, Komatsu M, Ha H, Lim JH, Park CW, Yang CW, Kim J, Kim YK (2019) Autophagy attenuates tubulointerstitial fibrosis through regulating transforming growth factor-beta and NLRP3 inflammasome signaling pathway. *Cell Death Dis* 10 (2):78. doi:10.1038/s41419-019-1356-0
8. Hartleben B, Godel M, Meyer-Schwesinger C, Liu S, Ulrich T, Kobler S, Wiech T, Grahmmer F, Arnold SJ, Lindenmeyer MT, Cohen CD, Pavenstadt H, Kerjaschki D, Mizushima N, Shaw AS, Walz G, Huber TB (2010) Autophagy influences glomerular disease susceptibility and maintains podocyte homeostasis in aging mice. *J Clin Invest* 120 (4):1084-1096. doi:10.1172/JCI39492
9. Jiang M, Wei Q, Dong G, Komatsu M, Su Y, Dong Z (2012) Autophagy in proximal tubules protects against acute kidney injury. *Kidney Int* 82 (12):1271-1283. doi:10.1038/ki.2012.261
10. Li XY, Wang SS, Han Z, Han F, Chang YP, Yang Y, Xue M, Sun B, Chen LM (2017) Triptolide Restores Autophagy to Alleviate Diabetic Renal Fibrosis through the miR-141-3p/PTEN/Akt/mTOR Pathway. *Mol Ther Nucleic Acids* 9:48-56. doi:10.1016/j.omtn.2017.08.011
11. Yang X, Wang H, Tu Y, Li Y, Zou Y, Li G, Wang L, Zhong X (2020) WNT1-inducible signaling protein-1 mediates TGF-beta1-induced renal fibrosis in tubular epithelial cells and unilateral ureteral obstruction mouse models via autophagy. *J Cell Physiol* 235 (3):2009-2022. doi:10.1002/jcp.29187
12. Chevalier RL, Forbes MS, Thornhill BA (2009) Ureteral obstruction as a model of renal interstitial fibrosis and obstructive nephropathy. *Kidney Int* 75 (11):1145-1152. doi:10.1038/ki.2009.86
13. Higashi AY, Aronow BJ, Dressler GR (2019) Expression Profiling of Fibroblasts in Chronic and Acute Disease Models Reveals Novel Pathways in Kidney Fibrosis. *J Am Soc Nephrol* 30 (1):80-94. doi:10.1681/ASN.2018060644
14. Lee CG, Kim JG, Kim HJ, Kwon HK, Cho IJ, Choi DW, Lee WH, Kim WD, Hwang SJ, Choi S, Kim SG (2014) Discovery of an integrative network of microRNAs and transcriptomics changes for acute kidney injury. *Kidney Int* 86 (5):943-953. doi:10.1038/ki.2014.117
15. Zhou LT, Qiu S, Lv LL, Li ZL, Liu H, Tang RN, Ma KL, Liu BC (2018) Integrative Bioinformatics Analysis Provides Insight into the Molecular Mechanisms of Chronic Kidney Disease. *Kidney Blood Press Res* 43 (2):568-581. doi:10.1159/000488830
16. Qu Y, Liu D, Jia H, Yang Z (2019) CircularRNA_{circ_0004002} regulates cell proliferation, apoptosis, and epithelial-mesenchymal transition through targeting miR-342-5p and Wnt3a in anorectal malformations. *Journal of Biological Chemistry* 120:15483-15493. doi:10.1002/jcb.28814
17. Wei Y, Nazari-Jahantigh M, Chan L, Zhu M, Heyll K, Corbalán-Campos J, Hartmann P, Thiemann A, Weber C, Schober A (2013) The microRNA-342-5p Fosters Inflammatory Macrophage Activation Through an Akt1- and microRNA-155-Dependent Pathway During Atherosclerosis. *Vascular Medicine* (129):1610-1619. doi:10.1161/CIRCULATIONAHA.1161/CIRCULATIONAHA.112.000736
18. Yan X-C, Cao J, Liang L, Wang L, Gao F, Yang Z-Y, Duan J-L, Chang T-F, Deng S-M, Liu Y, Dou G-R, Zhang J, Zheng Q-J, Zhang P, Han H (2016) miR-342-5p Is a Notch Downstream Molecule and Regulates Multiple Angiogenic Pathways Including Notch, Vascular Endothelial Growth Factor and Transforming Growth Factor b Signaling. *J Am Heart Assoc* (5):e003042. doi:10.1161/JAHA.115.003042
19. Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15 (12):550. doi:10.1186/s13059-014-0550-8
20. Davis S, Meltzer PS (2007) GEOquery: a bridge between the Gene Expression Omnibus (GEO) and BioConductor. *Bioinformatics* 23 (14):1846-1847. doi:10.1093/bioinformatics/btm254

21. Vlachos IS, Zagganas K, Paraskevopoulou MD, Georgakilas G, Karagkouni D, Vergoulis T, Dalamagas T, Hatzigeorgiou AG (2015) DIANA-miRPath v3.0: deciphering microRNA function with experimental support. *Nucleic Acids Res* 43 (W1):W460-466. doi:10.1093/nar/gkv403
22. Huang da W, Sherman BT, Lempicki RA (2009) Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 4 (1):44-57. doi:10.1038/nprot.2008.211
23. Janky R, Verfaillie A, Imrichova H, Van de Sande B, Standaert L, Christiaens V, Hulselmans G, Herten K, Naval Sanchez M, Potier D, Svetlichnyy D, Kalender Atak Z, Fiers M, Marine JC, Aerts S (2014) iRegulon: from a gene list to a gene regulatory network using large motif and track collections. *PLoS Comput Biol* 10 (7):e1003731. doi:10.1371/journal.pcbi.1003731
24. Chen X, Morales-Alcala CC, Riobo-Del Galdo NA (2018) Autophagic Flux Is Regulated by Interaction Between the C-terminal Domain of PATCHED1 and ATG101. *Mol Cancer Res* 16 (5):909-919. doi:10.1158/1541-7786.MCR-17-0597
25. Shen Y, Miao N, Xu J, Gan X, Xu D, Zhou L, Xue H, Zhang W, Lu L (2016) Metformin Prevents Renal Fibrosis in Mice with Unilateral Ureteral Obstruction and Inhibits Ang II-Induced ECM Production in Renal Fibroblasts. *Int J Mol Sci* 17 (2). doi:10.3390/ijms17020146
26. LORENA D, UCHIO K, COSTA AMA, DESMOULIÈRE A (2002) Normal scarring: importance of myofibroblasts. *Wound repair and regeneration* (1067-1967). doi:<https://doi.org/10.1046/j.1524-475X.2002.00201.x>
27. Redova M, Poprach A, Besse A, Iliev R, Nekvindova J, Lakomy R, Radova L, Svoboda M, Dolezel J, Vyzula R, Slaby O (2013) MiR-210 expression in tumor tissue and in vitro effects of its silencing in renal cell carcinoma. *Tumour Biol* 34 (1):481-491. doi:10.1007/s13277-012-0573-2
28. A RiA (1998) Combinatorial Gli gene function in floor plate and neuronal inductions by Sonic hedgehog. *Development* (125):2203–2212
29. Alexandre C, Jacinto A, Ingham PW (1996) Transcriptional activation of hedgehog target genes in *Drosophila* is mediated directly by the Cubitus interruptus protein, a member of the GLI family of zinc finger DNA-binding proteins. *Genes Dev* (10):2003–2013. doi:10.1101/gad.10.16.2003
30. Yang JJ, Tao H, Huang C, Shi KH, Ma TT, Bian EB, Zhang L, Liu LP, Hu W, Lv XW, Li J (2013) DNA methylation and MeCP2 regulation of PTCH1 expression during rats hepatic fibrosis. *Cell Signal* 25 (5):1202-1211. doi:10.1016/j.cellsig.2013.01.005
31. Bai Y, Lu H, Wu C, Liang Y, Wang S, Lin C, Chen B, Xia P (2014) Resveratrol inhibits epithelial-mesenchymal transition and renal fibrosis by antagonizing the hedgehog signaling pathway. *Biochem Pharmacol* 92 (3):484-493. doi:10.1016/j.bcp.2014.09.002
32. Bai Y, Lu H, Lin C, Xu Y, Hu D, Liang Y, Hong W, Chen B (2016) Sonic hedgehog-mediated epithelial-mesenchymal transition in renal tubulointerstitial fibrosis. *Int J Mol Med* 37 (5):1317-1327. doi:10.3892/ijmm.2016.2546
33. Isidor B, Bourdeaut F, Lafon D, Plessis G, Lacaze E, Kannengiesser C, Rossignol S, Pichon O, Briand A, Martin-Coignard D, Piccione M, David A, Delattre O, Jeanpierre C, Sévenet N, Le Caignec C (2012) Wilms' tumor in patients with 9q22.3 microdeletion syndrome suggests a role for PTCH1 in nephroblastomas. *European Journal of Human Genetics* 21 (7):784-787. doi:10.1038/ejhg.2012.252
34. Al-Tamari HM, Dabral S, Schmall A, Sarvari P, Ruppert C, Paik J, DePinho RA, Grimminger F, Eickelberg O, Guenther A, Seeger W, Savai R, Pullamsetti SS (2018) FoxO3 an important player in fibrogenesis and therapeutic target for idiopathic pulmonary fibrosis. *EMBO Mol Med* 10 (2):276-293. doi:10.15252/emmm.201606261
35. Li L, Kang H, Zhang Q, D'Agati VD, Al-Awqati Q, Lin F (2019) FoxO3 activation in hypoxic tubules prevents chronic kidney disease. *J Clin Invest* 129 (6):2374-2389. doi:10.1172/JCI122256

36. Lim SW, Shin YJ, Luo K, Quan Y, Cui S, Ko EJ, Chung BH, Yang CW (2019) Ginseng increases Klotho expression by FoxO3-mediated manganese superoxide dismutase in a mouse model of tacrolimus-induced renal injury. Aging 11:1-22

37. Li L, Zviti R, Ha C, Wang ZV, Hill JA, Lin F (2017) Forkhead box O3 (FoxO3) regulates kidney tubular autophagy following urinary tract obstruction. Journal of Biological Chemistry 292 (33):13774-13783.
doi:10.1074/jbc.M117.791483

Figures

Figure 1

Figure 1

Simple flow chart of bioinformatics analysis and experimental design

Figure 2

Figure 2

Candidate DE-miRNAs were obtained from GSE118340 and GSE42718. A and B) Heatmap showing the expression levels of each gene in different groups from GSE118340 and GSE42718; C and D) Volcano plot showing the DE-miRNAs in GSE118340 and GSE42718; E) Venn diagram showing miR-342-5p and miR-466a-3p are common DE-miRNAs in both GSE118340 and GSE42718.

Figure 3

Figure 3

GO annotation and KEGG pathway analysis of miR-342-5p and miR-466a-3p. A and B) GO annotation and KEGG pathway analysis of miR-342-5p obtained from miRPath V3.0. C and D) GO annotation and KEGG pathway analysis of miR-466a-3p obtained from miRPath V3.0.

Figure 4

Figure 4

Candidate DE-mRNAs obtained from GSE96101. A) Heatmap showing the the expression levels of each gene in different groups from GSE96101; B) Volcano plot showing the DE-mRNAs in GSE96101; E) Venn diagram showing 48 common DE mRNAs obtained from GSE96101 and predicted mRNAs from miRWalk.

Figure 5

Figure 5

Ptch1 is the most promising target mRNA of miR-342-5p. A) Regression analysis showing Ptch1 as the most promising target mRNA of miR-342-5p; B) The iRegulon prediction showing FoxO3 as potential transcription factor of Ptch1; C, D, and E) Real-time PCR showing that TGF- β 1 increases miR-342-5p and decreases FoxO3 and Ptch1 in TCMK-1 cells; F) FISH experiments showing that miR-342-5p is mainly distributed in the cytoplasm of TCMK-1 cells, as is FoxO3 and Ptch1. The merged images show that miR-342-5p is partially colocalized with FoxO3 and Ptch1.

Figure 6

Figure 6

Downregulation of Ptch1 induced autophagy in TCMK-1 cells. A, B, and C) Real-time PCR and western blot showing that downregulation of miR-342-5p reverses TGF-β1-mediated inhibition of the expression of Ptch1 in TCMK-1 cells, while the downregulation of FoxO3 promotes TGF-β1-mediated inhibition of the expression of Ptch1 in TCMK-1 cells; D and E) Western blot showing that knockdown of Ptch1 in TCMK-1 cells increased expression of SOSTM1/p62 and LC3II/LC3I; F and G) Quantification of immunofluorescent LC3-GFP puncta showing that Ptch1 knockdown enhanced autophagy, as evidenced by an increase in the number of GFP-LC3 spots. H) A simple schematic diagram showing the relationship between miR-342-5p, FoxO3, and Ptch1 in the progression of RF, and the role of autophagy in RF.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [renamedf0e6c.xlsx](#)
- [renameda9fd3.xlsx](#)
- [renamed5424c.xlsx](#)
- [renamed53f42.xls](#)