Overexpression of the dystrophins Dp40 and Dp40L170P modifies neurite outgrowth and the protein expression profile of PC12 cells

César García-Cruz

Centro de Investigacion y de Estudios Avanzados Unidad Zacatenco: Centro de Investigacion y de Estudios Avanzados del Instituto Politecnico Nacional

Candelaria Merino-Jiménez

Centro de Investigacion y de Estudios Avanzados Unidad Zacatenco: Centro de Investigacion y de Estudios Avanzados del Instituto Politecnico Nacional

Jorge Aragón

Centro de Investigacion y de Estudios Avanzados Unidad Zacatenco: Centro de Investigacion y de Estudios Avanzados del Instituto Politecnico Nacional

Víctor Ceja

Centro de Investigacion y de Estudios Avanzados Unidad Zacatenco: Centro de Investigacion y de Estudios Avanzados del Instituto Politecnico Nacional

Brenda González-Assad

Centro de Investigacion y de Estudios Avanzados Unidad Zacatenco: Centro de Investigacion y de Estudios Avanzados del Instituto Politecnico Nacional

Juan Pablo Reyes-Grajeda

Instituto Nacional de Medicina Genómica: Instituto Nacional de Medicina Genómica

Cecilia Montanez (✉ cecim@cinvestav.mx)

Centro de Investigacion y de Estudios Avanzados del Instituto Politecnico Nacional <https://orcid.org/0000-0002-7186-9016>

Research Article

Keywords: Dystrophin, Dp40, Neurite outgrowth, PC12 cells, α -internexin, VAMP

Posted Date: April 5th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-381163/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Dp40 is ubiquitously expressed, including in the central nervous system. Dp40 mRNA and protein are detected in the early stages and postnatal stages of the mouse brain, respectively. In addition to being present in the nucleus, membrane, and cytoplasm, Dp40 is detected in neurites and postsynaptic spines in hippocampal neurons. Although Dp40 is expressed from the same promoter as Dp71, its role in the cognitive impairment present in Duchenne muscular dystrophy patients is still unknown. Here, we studied the effects of overexpression of Dp40 and Dp40_{L170P} (a mutant of Dp40) during the neuronal differentiation process of PC12 Tet-On cells. We found that Dp40 overexpression increased the percentage of PC12 cells with neurites and neurite length, while Dp40_{L170P} overexpression decreased them compared to Dp40 overexpression. Two-dimensional gel electrophoresis analysis carried out in nerve growth factor-differentiated PC12-Dp40_{L170P} cells showed that the protein expression profile was modified compared to that of the control cells (PC12 Tet-On). The proteins with the highest upregulated expression were α -internexin and S100a6, which are involved in cytoskeletal structure. The expression of vesicle-associated membrane proteins increased in differentiated PC12-Dp40 cells, in contrast to PC12-Dp40_{L170P} cells, while neurofilament light-chain was decreased in both differentiated cells. HspB1 was absent in undifferentiated cells and weakly detected in all differentiated cells. These results suggest that the subcellular distribution and expression of Dp40 has an important role in the neurite outgrowth of PC12 cells through the regulation of proteins involved in neurofilaments and exocytosis of synaptic vesicles, functions that might be affected in PC12-Dp40_{L170P}.

Introduction

Dp40 is the smallest dystrophin reported to date and is transcribed from intron 62 to exon 70 [1]. Moreover, the Dp40 transcript is expressed in several human foetal tissues, such as muscle, lung, liver, and brain, embryonic stem cells, adult muscle and the schwannoma cell line [1]. In addition, its expression has been reported in different regions of the brain, such as the cortex, cerebellum and hippocampus of mice [1, 2]. Dp40 mRNA is expressed in undifferentiated and nerve growth factor- (NGF) differentiated PC12 cells [3]. Unlike other Dp71 isoforms, Dp40 lacks the C-terminal end and therefore the motifs and domains that interact with syntrophins and dystrobrevins [4]. Similar to Dp71, Dp40 contains part of the WW domain, which provides the main binding site to β -dystroglycan (β -DG), a component of the dystrophin-associated protein complex (DAPC), as well as the EF-hand motifs and the ZZ domain involved in Ca^{2+} detection and transport to the nucleus, respectively [5, 6]. The EF-hand motifs and ZZ domain are necessary for the WW domain to interact efficiently with β -DG [7, 8]. It has been reported that Dp40 interacts with syntaxin 1A (STX1A), vesicle-associated membrane protein 2 (VAMP2) and synaptosome-associated protein 25 (SNAP25), a group of presynaptic proteins involved in exocytosis of synaptic vesicles of the hippocampus and cortex in the mouse brain [9]. In addition, Dp40 protein shows high expression in neuronal cells and a decrease in non-neuronal cells in primary culture of the mouse hippocampus. Moreover, the Dp40 protein is only expressed in postnatal stages, not in the embryonic stages of the mouse brain [2]. In PC12 cells, transient expression of Dp40 showed that it is located in the membrane and cytoplasm of undifferentiated cells, while its subcellular distribution changes in NGF-differentiated PC12 cells at day 3 post-treatment, when it is located in the membrane and cytoplasm with a significant increase in the nucleus. In addition, the mutant Dp40_{L170P} has a change in residue 170 from leucine to proline, promoting exclusive nuclear localization in PC12 cells [3]. Furthermore, Dp40 colocalized with β -DG opposite to Dp40_{L170P} in NGF-differentiated PC12 cells [3]. Additionally, primary culture of mouse brain neurons showed that Dp40 is located in the membrane, nucleus and excitatory dendritic spines [2].

One-third of patients with Duchenne muscular dystrophy (DMD) have different degrees of cognitive deficits coupled with progressive muscular degeneration that characterize the disease. Cognitive impairment in DMD patients has been mostly associated with alterations in dystrophin Dp71 expression [10], and the disruption of Dp71 was shown to alter DAPC [11]. Although Dp40 is ubiquitously expressed from the same promoter as Dp71 [1], its participation in DAPC and therefore in this disease is still unknown. Interestingly, a report showed that six patients with different degrees of cognitive deficit had a deletion of three base pairs at positions 9711–9714 in the *dmd* gene. [12]. Importantly, this deletion is located in the same residue where the punctual change from leucine to proline is located in the mutant of Dp40 (Dp40_{L170P}). Therefore, in this work, to contribute to the knowledge of Dp40 function, we created PC12 Tet-On cells that overexpress dystrophin Dp40 or Dp40_{L170P} in an inducible and stable manner to analyse the effect of overexpression of these proteins on the neurite outgrowth process through morphometric and proteomic analyses. The results obtained showed that Dp40 overexpression stimulates neurite outgrowth in the opposite manner as Dp40_{L170P}, which its expression caused a reduction in neurite number and length. In addition, we carried out a proteomic analysis using two-dimensional gel electrophoresis (2-DE) to compare the Dp40_{L170P} expression profile with that of the control. We identified proteins related to alteration of the neurite outgrowth process. Dp40_{L170P} increased the expression levels of α -internexin and S100a6, proteins involved in intermediate filaments and reorganization of the cytoskeleton, respectively. Additionally, we evaluated the expression of VAMP, NF-L and HspB1, which are related to secretory processes, neurofilaments and cytoskeletal remodelling.

The results of this study provide valuable information about the role of Dp40 in neurite outgrowth during neural differentiation and its participation in cognitive deficits when the Dp40 isoform is disrupted.

Materials And Methods

Vector construction

The cDNA fragments of Dp40 and Dp40_{L170P} were obtained from the vectors pcDNA4/HisMax-TOPO/Dp40 and pcDNA4/HisMax-TOPO/Dp40_{L170P} [3] through amplification by PCR using the primers 5'TAGATCACGCGTACATGAGGGAACACCTCAAAGGC3' (pTRE-MluI) and 5'GATCTAGCGGCCGCTCACGTTTCCATGTTGTCCCCCTCTAAC3' (Dp40-NotI), which add the restriction sites MluI and NotI, respectively. The DNA sequence of Dp40 corresponds to the *Rattus norvegicus* sequence reported in GenBank (KF154977.1). With T4 DNA ligase, the Dp40 and Dp40_{L170P} fragments were cloned into pGEM-T Easy (Promega, Madison, WI, USA) as a transient vector and then sequenced. After Dp40 and Dp40_{L170P} fragment excision with restriction enzymes and cloning in the vector pTRE2pur-Myc (Clontech, Mountain View, CA, USA.), which adds a Myc flag tag to the N-terminal end of Dp40 and Dp40_{L170P}, an inducible expression system Tet-On was generated using the vectors pTRE2pur-Myc/Dp40 and pTRE2pur-Myc/Dp40_{L170P}.

Cell culture and NGF differentiation

pTRE2pur-Myc/Dp40 and pTRE2pur-Myc/Dp40_{L170P} vectors were used to stably transfect PC12 Tet-On cells (PC12 cells that express a regulator protein from the Tet-On system) (Clontech). Then, we generated a pool of puromycin resistant PC12-Dp40 and PC12-Dp40_{L170P} cells to isolate clones that express the recombinant proteins Myc-Dp40 and Myc-Dp40_{L170P} in an inducible manner. As a control, PC12 Tet-On cells were stably transfected with an empty pTRE2pur-Myc vector to obtain PC12 control cells as previously described [13]. All cell lines were cultured in Dulbecco's modified Eagle's medium (Gibco, Rockville, Maryland, USA) supplemented with 10% heat-inactivated horse serum, 5% Tet System Approved foetal bovine serum, 100 U/ml penicillin, 1 mg/ml streptomycin, 250 ng/ml mycostatin, and 100 µg/ml geneticin (G418). The expression of Myc-Dp40 and Myc-Dp40_{L170P} was induced with 100 ng/ml doxycycline for 24 h to obtain undifferentiated PC12 cells. For NGF-differentiated PC12 cells, induction was maintained throughout the differentiation process. Under differentiation conditions, the medium supplemented with 100 ng/ml doxycycline and 50 ng/ml NGF was changed every three days for nine days [14].

Morphometric analysis of neurite outgrowth

PC12-Dp40, PC12-Dp40_{L170P} and PC12 control cells were plated at low confluence on collagen-coated plastic dishes and cultured in the presence of NGF for nine days. Ten micrographs from three independent experiments were taken with an inverted microscope (Axiovert. A1 Zeiss) with 10X objective. To quantify the neurite outgrowth ratio, we considered all cells that produced a neurite greater than a cell body. For neurite length, all cells with neurites greater than two cell bodies were considered. Axiovision 4.8 software was used to count the neurites and to measure their length.

Protein extraction and 2-DE

Total protein extracts of the NGF-differentiated PC12 control and PC12-Dp40_{L170P} cells were obtained and run in 2-DE gels as previously described [15]. Protein concentrations were determined by the Bradford method. For PC12 control and PC12-Dp40_{L170P} cell lines, three protein samples were obtained from three independent experiments. The 2-DE gels obtained were stained using Bio-Safe Coomassie Stain (Bio-Rad, Hercules, CA, USA) according to the manufacturer's instructions.

Scanning and image analysis

The stained 2-DE gels were scanned with an Image Quant 4000 instrument (GE Healthcare, Chicago, IL, USA). For determination of the protein abundance, the percent volume (% volume) of each spot was calculated using Image Master 2D Platinum 7.0 software (GE Healthcare). Spots were manually examined to eliminate artefacts. The nonparametric Mann-Whitney test was used to compare data from each group and detect spots with different expression levels. Spots with *P* values < 0.05 and changes of 1.1-fold or more were considered statistically significant. Spots of interest were excised from the 2-DE gels for identification using mass spectrometry (MS).

In-gel trypsin digestion and protein identification by MS

The selected protein spots were excised and placed in Eppendorf tubes with 50 µl of destaining solution (50% v/v methanol and 5% v/v acetic acid), followed by washes with Milli-Q H₂O. Gel fragments were dehydrated by incubation in 100 µl of acetonitrile (ACN) for 10 min,

and the supernatant was then removed; this step was repeated once. Dry gel pieces were rehydrated with 200 ng of trypsin (Promega V528A) in 50 mM NH_4HCO_3 and 5% ACN and incubated overnight at 37 °C. The resulting peptides were extracted with 40 μl of 50% ACN and 5% formic acid, and the solution volume was reduced in a concentrator (Eppendorf 5301). Peptides from each sample were desalted on C18 columns (ZipTipC18). A 1:1 mixture of peptide solution and matrix solution (5 mg/ml CHCA 50% v/v and TFA 0.1% v/v) was analysed using a 4800 Plus MALDI TOF/TOF mass spectrometer (Sciex). The search was performed with the enzyme specificity of trypsin, and one missed cleavage was allowed. The detected protein threshold was 66%, and the precursor mass tolerance was 0.5 to 1 Da. The MS data were compared with the *Rattus norvegicus* database (downloaded in September 2016) using Protein Pilot™ software (version 2.0.1) and the Mascot algorithm [16].

Western blotting

Western blotting (WB) was performed as previously described [14]. The mouse monoclonal antibodies anti- β -actin (1:500), anti-c-Myc (1:200) and anti-VAMP1/2 (1:200) were purchased from Santa Cruz Biotechnology (Dallas, TX, USA). Rabbit monoclonal anti- α -internexin (1:10,000), mouse monoclonal NF-L (1:500) and rabbit polyclonal HspB1 (1:500) antibodies were purchased from Abcam (Burlingame, CA, USA).

Indirect immunofluorescence microscopy

Indirect immunofluorescence (ImmF) was carried out in undifferentiated and NGF-differentiated PC12 Tet-On cells as previously described [3]. Alexa Fluor 488- and Alexa Fluor 594-conjugated secondary antibodies (Invitrogen, Life Technologies, NY, USA) were used to detect the primary antibody signal. Images were captured using a Leica confocal microscope (Leica TCS SP8) with a 40X objective at zoom 2. The fluorescence intensity and colocalization ratio were quantified from equatorial Z-sections obtained from three independent experiments.

Statistical analysis

Data are shown as the mean $\pm$ SD of three independent experiments. Statistical analyses were performed using Student's t-test with GraphPad Prism 5 software. P-values < 0.05 were considered statistically significant.

Results

Overexpression of Myc-Dp40 and Myc-Dp40_{L170P} showed different effects on the neurite outgrowth of PC12-Tet-On cells.

PC12 Tet-On cells were stably transfected with the pTRE2pur-Myc/Dp40 or pTRE2pur-Myc/Dp40_{L170P} vector, and vector integration was tested by genomic DNA PCR (Fig. S1). To characterize the isolated clones PC12-Dp40 and PC12-Dp40_{L170P}, we determined the minimum concentration of doxycycline to induce the overexpression of the recombinant proteins through WB using an anti-c-Myc antibody. Myc-Dp40 and Myc-Dp40_{L170P} were overexpressed using 50 to 1000 ng/ml doxycycline in undifferentiated PC12 Tet-On cells (Fig. S2). Because higher concentrations of doxycycline did not result in an increase in recombinant protein expression, we used 100 ng/ml doxycycline to overexpress Dp40 proteins to eliminate the cytotoxic effect of doxycycline reported at 200 ng/ml [17]. With this doxycycline concentration, undifferentiated and NGF-differentiated PC12 Tet-On cells were analysed to evaluate the morphological effect of Myc-Dp40 and Myc-Dp40_{L170P} overexpression on neurite outgrowth processes compared with those of the PC12 control cells. Myc-Dp40 and Myc-Dp40_{L170P} overexpression did not affect the morphology of the undifferentiated PC12 cells (Fig. 1a-c), in contrast to the NGF-differentiated PC12 cells (Fig. 1d-f). The differentiation ratio (Fig. 1g) showed that the PC12-Dp40 cells ($40.6\% \pm 2.90$; $P = 0.0003$) had a greater number of cells with neurites than the PC12 control cells ($4.96\% \pm 0.72$), while PC12-Dp40_{L170P} did not result in a significant difference compared with the PC12 control cells. However, a decrease in the differentiation ratio of the NGF-differentiated Dp40_{L170P} cells ($4.50\% \pm 1.82$; $P = 0.0005$) was observed compared with that of the PC12-Dp40 cells. Additionally, neurite length quantification (Fig. 1h) showed that the NGF-differentiated PC12-Dp40 cells ($17.3 \mu\text{m} \pm 0.30$; $P = 0.0086$) had longer neurites than the PC12 control cells ($14.0 \mu\text{m} \pm 0.62$). However, the PC12 control cells presented longer neurites than the PC12-Dp40_{L170P} cells ($11.5 \mu\text{m} \pm 0.052$; $P = 0.0359$). Therefore, the differentiated PC12-Dp40_{L170P} cells ($11.5 \mu\text{m} \pm 0.52$; $P = 0.006$) showed lower neurite outgrowth than the PC12-Dp40 cells ($17.3 \mu\text{m} \pm 0.30$). Based on these results, Myc-Dp40 expression stimulates neurite outgrowth, whereas Myc-Dp40_{L170P} inhibits this process.

Protein expression profile of differentiated PC12-Dp40_{L170P} cells

To understand the inhibitory mechanism of neurite outgrowth produced by Dp40_{L170P}, we analysed the total protein extracts of the PC12 control and PC12-Dp40_{L170P} NGF-differentiated cells for nine days by 2-DE (Fig. 2a and b). Differentially expressed protein spots with at least a 1.1-fold change were considered differentially expressed. Among the 344 protein spots detected, 14 (spots ID: 1-14) showed differential expression (Fig. 2c), 13 proteins showed upregulated expression and one showed downregulated expression, and magnified

views of the differentially expressed protein spots are presented in Fig. 2c. Thus, Dp40_{L170P} modified the protein expression profile of PC12 Tet-On cells. The 14 spots selected were excised from the 2-DE gels and analysed by MS. The details of each identified protein are summarized in Table 1. The proteins with highly upregulated expression were S100a6 (2.1-fold, p/5.2) and α -internexin (2.0-fold, p/5.3), which are involved in the reorganization of cytoskeletal structure and neurofilaments present in immature neurons, respectively. Only the expression of ectonucleotide pyrophosphatase (1.5-fold, p/5.5), a protein related to cellular communication, was downregulated. The other proteins identified were related to chaperone-like activity (T-complex protein 1 subunit zeta and the endoplasmic reticulum chaperone BIP) and metabolism (phosphoglycerate mutase 1, L-lactate dehydrogenase A chain and alpha enolase). This last group of proteins usually shows upregulated expression in several proteomic analyses, probably due to cellular stress responses [18].

Myc-Dp40 and Myc-Dp40_{L170P} overexpression modified the expression of α -internexin, VAMP, HspB1 and NF-L during the neuronal differentiation process

After MS identification, we validated the differential expression of α -internexin, one of the top two proteins with upregulated expression determined by WB in undifferentiated (Fig. 3) and differentiated (Fig. 4) PC12 control cells, PC12-Dp40 cells and PC12-Dp40_{L170P} cells. First, we evaluated the expression of the recombinant proteins: the Myc-Dp40 and Myc-Dp40_{L170P} proteins were overexpressed 13.3- and 7-fold in the undifferentiated PC12-Dp40 and PC12-Dp40_{L170P} cells, respectively, showing a significant increase compared those of the PC12 control cells. In addition, Myc-Dp40 (1.9-fold) was more highly expressed than Myc-Dp40_{L170P} in the PC12-Dp40 and PC12-Dp40_{L170P} cells. α -Internexin, which is involved in the expression and assembly of neurofilaments in the central nervous system [19, 20], presented a 5.25-fold increase in the PC12-Dp40_{L170P} cells compared with the PC12 control cells. However, α -internexin expression was not significantly different in the undifferentiated PC12-Dp40 cells compared with the PC12 control and PC12-Dp40_{L170P} cells. Since a previous report [9] showed that VAMP interacts with the Dp40 protein, we decided to validate its expression. The results showed that VAMP is expressed at low levels and without a significant difference in all undifferentiated cells. Recently, it was observed that Dp71e _{Δ 71} and mutant Dp71 _{Δ 78-79} overexpression increased the HspB1, a remodeler of the cytoskeleton, during the neuronal differentiation process of PC12 cells [15, 21]. Thus, we analysed the expression of the HspB1 protein; however, we did not detect this protein under undifferentiated conditions. In addition, NF-L expression was evaluated because it is a neuronal differentiation marker that is expressed in postmitotic neurons as part of the cytoskeletal structure. However, we did not observe a significant difference in NF-L expression in any undifferentiated PC12 cells.

In the NGF-differentiated PC12 cells (Fig. 4), Myc-Dp40 and Myc-Dp40_{L170P} expression showed an increase of 9.8- and 4.3-fold in the PC12-Dp40 and PC12-Dp40_{L170P} cells, respectively compared with the PC12 control cells. However, there was no significant difference between Myc-Dp40 and Myc-Dp40_{L170P}. α -Internexin expression in the PC12-Dp40 cells did not show a significant difference compared with that in the PC12 control cells. However, its expression in the PC12-Dp40_{L170P} cells was increased 44- and 3-fold compared with that in the PC12 control and PC12-Dp40 cells, respectively. This increase is in accordance with what we observed in the 2-DE gels in the differentiated PC12 control cells and the PC12-Dp40_{L170P} cells. VAMP expression was increased 5.4- and 1.6-fold in the NGF-differentiated PC12-Dp40 and PC12-Dp40_{L170P} cells, respectively, compared with that in the PC12 control cells. However, the PC12-Dp40 cells showed an expression increase of 3.3-fold compared to the PC12-Dp40_{L170P} cells. HspB1 presented low expression and no significant difference in the differentiated cell lines (Fig. 4b). NF-L expression was not significantly different between the differentiated PC12 control and PC12-Dp40 cells. In contrast, the PC12 control cells showed an increase of 2-fold compared with the PC12-Dp40_{L170P} cells but no significant difference compared with the PC12-Dp40 cells. All these results showed that the disruption of dystrophin Dp40 through the expression of the mutant Dp40_{L170P} could affect the expression of neurofilaments such as α -internexin and NF-L during the neuronal differentiation process of PC12 cells.

Differential distribution of the Myc-Dp40, Myc-Dp40_{L170P} and α -internexin proteins in PC12- Tet-On cells

Differences in the subcellular distribution of Myc-Dp40, Myc-Dp40_{L170P} and c-Myc epitopes were evaluated through ImmF assays in the undifferentiated and NGF-differentiated PC12 Tet-On cells (Figs. 5 and 6, respectively). Myc-Dp40 protein localization did not present a significant difference between the cytoplasm (49.08% $\pm$ 2.20) and nucleus (50.92% $\pm$ 2.20; P = NS) in the undifferentiated cells, while Myc-Dp40_{L170P} showed an increase in the nucleus (75.06% $\pm$ 1.69) compared to the cytoplasm (24.94% $\pm$ 1.69; P < 0.0001), unlike the c-Myc peptide in the PC12 control cells, which presented an increase in immunoreactivity in the cytoplasm (66.53% $\pm$ 1.58) compared to the nucleus (33.47% $\pm$ 1.580; P < 0.0001). Under NGF-differentiated conditions, in the PC12 control cells, the c-Myc-peptide was localized to the same extent in the cytoplasm (52.68% $\pm$ 2.98) and nucleus (47.32% $\pm$ 2.989; P = NS). However, Myc-Dp40 was mainly located in the cytoplasm (61.58% $\pm$ 3.00) compared to the nucleus (38.42% $\pm$ 3.00; P < 0.0001), while Myc-Dp40_{L170P} showed higher immunoreactivity in the nucleus (62.20% $\pm$ 2.19) than in the cytoplasm (37.80% $\pm$ 2.19; P < 0.0001).

α -Internexin has been detected in the cytoplasm of postmitotic neurons of the periphery and central nervous system [22, 23]. In the undifferentiated PC12 control, PC12-Dp40 and PC12-Dp40_{L170P} cells, α -internexin was observed in the cytoplasm. We found that the immunoreactivity of this protein was similar in the undifferentiated PC12 control ($0.73\% \pm 0.19$), PC12-Dp40 ($0.83\% \pm 0.08$; $P = \text{NS}$) and PC12-Dp40_{L170P} cells ($1.42\% \pm 0.37$; $P = \text{NS}$). However, the NGF-differentiated PC12-Dp40_{L170P} cells ($3.03\% \pm 0.10$) showed a higher percentage of immunoreactivity than the PC12-Dp40 ($1.76\% \pm 0.28$; $P = 0.013$) and PC12 control cells ($1.52\% \pm 0.21$; $P = 0.003$), but we did not observe a significant difference between the NGF-differentiated PC12-Dp40 ($0.44\% \pm 0.157$; $P = \text{NS}$) and PC12 control cells ($0.36\% \pm 0.003$). In the NGF-differentiated PC12-Dp40 cells, α -internexin was observed in the cytoplasm and along the length of neurites, while in the PC12 control and PC12-Dp40_{L170P} cells, it was mostly distributed in the cytoplasm. Based on these results, the subcellular distributions of Myc-Dp40 and Myc-Dp40_{L170P} were mostly cytoplasmic and nuclear, respectively. In addition, the overexpression of Myc-Dp40_{L170P} increases α -internexin expression in the undifferentiated and NGF-differentiated PC12-Dp40_{L170P} cells.

Discussion

PC12 cells are a neuronal differentiation model widely used to study the reorganization of the cytoskeleton, electrical excitability and secretory vesicle dynamics during the neuronal differentiation process due to their similarities to sympathetic neurons [24–26]. Previous studies have shown that PC12 expresses short dystrophins, including Dp71 isoforms from Dp71d, Dp71f, Dp71e groups and Dp40 [3, 27, 28] and some of them show increased expression during the neuronal differentiation process [27]. Interestingly, through antisense technology against the N-terminus of short dystrophin mRNAs (Dp71/Dp40), it has been shown that these dystrophins are essential for neurite outgrowth [29]. In addition, one study revealed that Dp40 accumulates in the nucleus during the neuronal differentiation process of PC12 cells [3]. Moreover, the exchange of leucine to proline in residue 170 of Dp40 (named Dp40_{L170P}) promotes exclusive nuclear localization of Dp40, probably because it disrupts a nuclear export signal, which decreases the colocalization percentage between Dp40 and β -DG in PC12 cells [3, 30].

To obtain a better understanding of the role of dystrophin Dp40 in the neurite outgrowth process, we evaluated the effect of Dp40 and Dp40_{L170P} overexpression during neuronal differentiation of PC12 Tet-On cells. As has been reported for Dp71 $_{\Delta 78-79}$ and Dp71e $_{\Delta 71}$ [21, 31], Dp40 overexpression also stimulated neurite outgrowth by increasing the ratio of cells with neurites and the neurite length in PC12 cells. However, Dp40_{L170P} decreased the neurite length compared with those of the PC12-Dp40 and PC12 control cells (Fig. 1). This result suggests that Dp40 overexpression could stimulate the neurite outgrowth process through the subcellular distribution of Dp40. Moreover, Dp40 has been detected in neurites and dendritic spines in PC12 cells and hippocampal neurons in primary culture [2, 3]. In addition, in this study, we observed that Dp40 was present in the neurites and cytoplasm and to a lesser extent in the nucleus of the NGF-differentiated PC12-Dp40 cells (Fig. 5). However, unlike what was previously reported [3], where Dp40_{L170P} was located exclusively in the nucleus, in this work, Dp40_{L170P} not only increased its presence in the nucleus but was also observed in the cytoplasm of the undifferentiated and NGF-differentiated PC12-Dp40_{L170P} cells. We speculated that these differences in the subcellular distribution compared with the previous findings [3] are because recombinant protein overexpression (Myc-Dp40 and Myc-Dp40_{L170P}) was mediated through stable transfection in the Tet-On system and not through transient transfection, as in the previous report.

The differentially expressed proteins identified through MS were related to the reorganization of the cytoskeleton (S100a6, α -internexin), cellular communication (ectonucleotide pyrophosphatase), chaperone-like activity (T-complex protein 1 subunit zeta and the endoplasmic reticulum chaperone BIP) and metabolism (phosphoglycerate mutase 1, L-lactate dehydrogenase A chain and α -enolase) according to the Protein Data Bank. Additionally, it has been reported that one-third of the differentially regulated proteins in proteomic studies are involved in metabolism, possibly because of the treatment to obtain the protein extract [18]. The protein with the highest differential expression was S100a6, with an increase of 2.1-fold (Table 1). The S100a6 protein belongs to the calcium binding protein family, which has been associated with several processes, such as proliferation, apoptosis cytoskeletal dynamics, and cellular response to stress factors [32], including stimulating neurite outgrowth in PC12 cells [33]. However, the expression of this protein was upregulated in the PC12-Dp40_{L170P} cells, which showed lower neurite outgrowth than the PC12-Dp40 cells. Therefore, it is likely that S100a6 overexpression is not the only element required to promote neurite outgrowth. Thus, we wanted to focus on the α -internexin protein, which could explain the disruption of neurite outgrowth in PC12-Dp40_{L170P} cells.

α -Internexin was the second protein with the greatest upregulation in expression, with an increase of 2.0-fold. This protein belongs to intermediate filament type IV and is expressed early during brain development in most neurons [22]. After validation of the α -internexin expression, we observed that this protein showed decreased expression from undifferentiated to differentiated PC12 control cells (Figs. 3 and 4), in which it was practically absent, consistent with the high α -internexin expression detected in immature myenteric neurons and its decrease with age [34]. However, α -internexin expression was highly expressed in the NGF-differentiated PC12-Dp40_{L170P} cells compared

with the PC12 control and PC12-Dp40 cells, while NF-L presented low expression in the PC12-Dp40_{L170P} cells, in contrast to the PC12 control cells, which showed high expression (Fig. 4). This finding contrasts with the fact that it has been reported that the later neuronal marker NF-L increases its expression during neuronal differentiation and that α -internexin decreases its expression in myenteric neurons during development [22, 35], suggesting that Dp40 does not promote neuronal differentiation by increasing neurofilaments such as NF-L. A study reported that α -internexin expression is upregulated after peripheral nerve injury in facial motor neurons of rats [36], possibly as a compensatory mechanism to reassemble other neurofilaments and promote neurite outgrowth. It was reported that α -internexin is the only neurofilament capable of coassembling itself [37], with other neurofilaments increasing interfilament spacing [38], probably to allow the assembly of other neurofilaments and promote neurite outgrowth. In addition, α -internexin has been identified as a candidate involved in early stages of brain regeneration in lesion models in the cerebellum [39], a brain structure that has been suggested to participate in integrating learning processes [40]. However, the overexpression of α -internexin in transgenic mice induced abnormal swelling of Purkinje cell axons in the cerebellum, promoting neuronal dysfunction [41], which could be related to the disruption of neurite outgrowth in the PC12-Dp40_{L170P} cells. Interestingly, the increase in α -internexin detected in the differentiated PC12-Dp40 and PC12-Dp40_{L170P} cells was not proportional to the amount of recombinant proteins produced (Figs. 3 and 4) because Myc-Dp40 is more highly expressed than Myc-Dp40_{L170P} in the undifferentiated and differentiated PC12 cells. Therefore, differences in the expression levels of Myc-Dp40 and Myc-Dp40_{L170P} do not influence α -internexin expression. Thus, it is possible that the increase in α -internexin expression in the differentiated PC12-Dp40_{L170P} cells is due to the nuclear distribution of Myc-Dp40_{L170P} instead of protein production.

Dp40 overexpression stimulated the neurite outgrowth during neuronal differentiation, which could be related to the increase in VAMP in the differentiated PC12-Dp40 cells and its decrease in the PC12-Dp40_{L170P} cells (Fig. 4), where neurite outgrowth is disrupted. VAMP2 overexpression in PC12 cells stimulates neurite outgrowth [42], and Dp40 interacts with VAMP2 in synaptic vesicle fractions in the adult mouse brain and colocalizes with VAMP in neurites of primary cultured hippocampal neurons [9]. Thus, we suggest that Dp40 overexpression promoted neuritic outgrowth by increasing VAMP expression in the differentiated PC12-Dp40 cells (Fig. 4) and decreasing it in the PC12-Dp40_{L170P} cells that express the mutant Dp40_{L170P}. The low expression of NF-L in the differentiated PC12-Dp40_{L170P} suggests that the disruption of dystrophin Dp40 could alter the expression and function of neurofilaments affecting neurite outgrowth because the coordination between membrane trafficking and cytoskeletal remodelling are critical requirements for axonal growth [43]. The lack of long neurites observed in the PC12-Dp40_{L170P} cells could be associated with the nuclear distribution of the mutant Dp40_{L170P} protein (Fig. 5) by regulating VAMP expression through its decrease and preventing the Dp40 interaction with VAMP and with other synaptic vesicle proteins (SNAP25 and STX1A), disrupting the exocytosis cycle and therefore its participation in neurite outgrowth [43]. Considering the facts mentioned, it is possible that the disruption of VAMP as a consequence of the nuclear localization of Dp40_{L170P} in the differentiated PC12-Dp40_{L170P} cells promotes the α -internexin overexpression as a scaffold neurofilament to restore the reassembly of neurofilaments and therefore neurite outgrowth.

However, previous reports have shown that overexpression of Dp71 $_{\Delta 78-79}$ and Dp71e $_{\Delta 71}$ stimulates neurite outgrowth in NGF-differentiated PC12 cells, and both cell lines presented a high expression of HspB1 [15, 21]. Because Myc-Dp40 overexpression promotes neuronal differentiation (Fig. 1), we tested the expression level of HspB1 in the differentiated PC12 cells. Interestingly, the expression of this protein was very low in the differentiated PC12 cells (Fig. 4). This finding indicates that Dp71 and Dp40 promote neurite outgrowth through a different pathway. Thus, our results suggest that Dp40 also plays an important role in the neuronal differentiation process through the regulation of the expression of proteins related to synaptic vesicles such as VAMP and neurofilaments such as α -internexin, possibly through its differential subcellular distribution in PC12 cells. Analysis of the mechanisms that stimulate the disruption of neuronal differentiation as a consequence of the subcellular distribution of Dp40 should be the next step. Taken together, these results suggested that the neurite outgrowth promoted by Dp40 overexpression could be carried out through a different strategy than the increase in HspB1, as was previously reported in PC12 cells that overexpress dystrophin Dp71 [15, 21], possibly through VAMP expression, which is increased in the differentiated PC12-Dp40 cells but decreased in the PC12-Dp40_{L170P} cells.

Conclusion

In this work, we reported the effect of Dp40 and the mutant Dp40_{L170P} on the neuronal differentiation process of PC12 cells. Dp40 expression promotes an increase in the percentage of cells with neurites as well as in neurite length, while Dp40_{L170P} expression decreases neurite length. Dp40_{L170P} modifies the protein expression profile of PC12 Tet-On cells, upregulating the expression of proteins involved in the cytoskeletal reorganization and structural proteins such as α -internexin and S100a6. Additionally, during neuronal differentiation, Dp40 overexpression increased the expression of VAMP, in contrast to Dp40_{L170P}, which decreased it. However, the low expression of HspB1 in differentiated PC12-Dp40 cells suggests that Dp40 promotes neurite outgrowth through a different pathway than of dystrophin Dp71e $_{\Delta 71}$.

and Dp71_{Δ78–79}. These data support the hypothesis that the disruption of Dp40 alters neurite outgrowth and could contribute to the cognitive deficit present in DMD patients with mutations in residue 170 of dystrophin Dp40.

Declarations

Acknowledgement

We thank the proteomics service unit from the Instituto Nacional de Medicina Genómica for identifying the peptides, Ivan Galván, MSc, for his assistance with confocal microscopy and Clemencia Salas for providing technical assistance.

Funding

This work was supported by the following grants from the Consejo Nacional de Ciencia y Tecnología (CONACyT): C. García-Cruz Postdoctoral Fellowship 29931, C. Merino-Jiménez Postdoctoral Fellowship 24868 and C. Montañez Grant CB-2017-2018-A1-S-24868-M, ECOS-NORD CONACYT grant number 276330.

Conflicts of interest/Competing interests

The authors have no conflicts of interest to declare.

Availability of data and material

All data are available in the paper and electronic supplementary material.

Code availability (Not applicable)

Authors' contributions

CPG designed the study, analysed the data and wrote the paper. CMJ performed the proteomic analysis. JPRG designed and performed the MS analysis. JA, VC and BGA performed the cell cultures and participated in constructing the vectors, provided technical assistance and contributed to the revision of the paper. CM coordinated the study, analysed the results, and wrote the paper. All the authors reviewed the results and approved the final version of the manuscript.

Ethics approval(Not applicable)

Consent to participate(Not applicable)

Consent for publication(Not applicable)

References

1. Tinsley JM, Blake DJ, Davies KE (1993) Apo-dystrophin-3: a 2.2kb transcript from the DMD locus encoding the dystrophin glycoprotein binding site. *Human Mole Genet* 2(5):521–524. <https://doi.org/10.1093/hmg/2.5.521>
2. Fujimoto T, Itoh K, Yaoi T, Fushiki S (2014) Somatodendritic and excitatory postsynaptic distribution of neuron-type dystrophin isoform, Dp40, in hippocampal neurons. *Biochem Biophys Res Commun* 452(1):79–84. <https://doi.org/10.1016/j.bbrc.2014.08.064>
3. Aragón J, Martínez-Herrera A, Bermúdez-Cruz RM, Bazán ML, Soid-Raggi G, Ceja V, Coy-Arechavaleta AS, Alemán V, Depardón F, Montañez C (2015) EF-hand domains are involved in the differential cellular distribution of dystrophin Dp40. *Neurosci Lett* 600:115–120. <https://doi.org/10.1016/j.neulet.2015.05.038>
4. Blake DJ, Weir A, Newey SE, Davies KE (2002) Function and genetics of dystrophin and dystrophin-related proteins in muscle. *Physiol Rev* 82(2):291–329. <https://doi.org/10.1152/physrev.00028.2001>
5. Suárez-Sánchez R, Aguilar A, Wagstaff K, Velez G, Azuara-Medina P, Gomez P, Vásquez-Limeta A, Hernández-Hernández O, Lieu K, Jans D (2014) Nucleocytoplasmic shuttling of the Duchenne muscular dystrophy gene product dystrophin Dp71d is dependent on the importin α/β and CRM1 nuclear transporters and microtubule motor dynein. *Biochim Biophys Acta* 1843(5):985–1001. <https://doi.org/10.1016/j.bbamcr.2014.01.027>
6. Chen Y, Xue S, Zhou Y, Yang JJ (2010) Calciomics: prediction and analysis of EF-hand calcium binding proteins by protein engineering. *Sci China Chem* 53(1):52–60. <https://doi.org/10.1007/s11426-010-0011-5>

7. Rentschler S, Linn H, Deininger K, Bedford MT, Espanel X, Sudol M (1999) The WW domain of dystrophin requires EF-hands region to interact with β -dystroglycan. *Biol Chem* 380(4):431–442. <https://doi.org/10.1515/BC.1999.057>
8. Hnia K, Zouiten D, Cantel S, Chazalette D, Hugon G, Fehrentz J-A, Masmoudi A, Diment A, Bramham J, Mornet D (2007) ZZ domain of dystrophin and utrophin: topology and mapping of a β -dystroglycan interaction site. *Biochem J* 401(3):667–677. <https://doi.org/10.1042/BJ20061051>
9. Tozawa T, Itoh K, Yaoi T, Tando S, Umekage M, Dai H, Hosoi H, Fushiki S (2012) The shortest isoform of dystrophin (Dp40) interacts with a group of presynaptic proteins to form a presumptive novel complex in the mouse brain. *Mol Neurobiol* 45(2):287–297. <https://doi.org/10.1007/s12035-012-8233-5>
10. Daoud F, Candelario-Martínez A, Billard J-M, Avital A, Khelfaoui M, Rozenvald Y, Guegan M, Mornet D, Jaillard D, Nudel U (2009) Role of mental retardation-associated dystrophin-gene product Dp71 in excitatory synapse organization, synaptic plasticity and behavioral functions. *PLoS One* 4(8):e6574. <https://doi.org/10.1371/journal.pone.0006574>
11. Benabdesselam R, Dorbani-Mamine L, Benmessaoud-Mesbah O, Rendon A, Mhaouty-Kodja S, Hardin-Pouzet H (2012) Dp71 gene disruption alters the composition of the dystrophin-associated protein complex and neuronal nitric oxide synthase expression in the hypothalamic supraoptic and paraventricular nuclei. *J Endocrinol* 213(3):239–249. <https://doi.org/10.1530/JOE-12-0066>
12. de Brouwer AP, Nabuurs SB, Verhaart IE, Oudakker AR, Hordijk R, Yntema HG, Hordijk-Hos JM, Voeselek K, De Vries BB, Van Essen T (2014) A 3-base pair deletion, c. 9711_9713del, in DMD results in intellectual disability without muscular dystrophy. *Eur J Hum Genet* 22(4):480–485. <https://doi.org/10.1038/ejhg.2013.169>
13. Herrera-Salazar A, García-Villegas R, Aragón J, Sánchez-Trujillo A, Ceja V, Martínez-Herrera A, Merino-Jiménez C, Montañez C (2016) Overexpression of mutant dystrophin Dp71 [INCREMENT] 78–79 stimulates cell proliferation. *Neuroreport* 27(1):6–11. <https://doi.org/10.1097/WNR.0000000000000475>
14. Aragón J, Martínez-Herrera A, Romo-Yáñez J, Ceja V, Azotla-Vilchis C, Siqueiros-Márquez L, Soid-Raggi G, Herrera-Salazar A, Montañez C (2016) Identification of Dp71 isoforms expressed in PC12 cells: subcellular localization and colocalization with β -dystroglycan and α 1-syntrophin. *J Mol Neurosci* 58(2):201–209. <https://doi.org/10.1007/s12031-015-0657-8>
15. Merino-Jiménez C, Aragón J, Ceja V, Rodríguez-Martínez G, Cázares-Raga FE, Chardonnet S, Pionneau C, Rendon A, Montañez C (2016) Dp71 Δ 78–79 dystrophin mutant stimulates neurite outgrowth in PC12 cells via upregulation and phosphorylation of HspB1. *Proteomics* 16(9):1331–1340. <https://doi.org/10.1002/pmic.201500211>
16. Perkins DN, Pappin DJ, Creasy DM, Cottrell JS (1999) Probability-based protein identification by searching sequence databases using mass spectrometry data. *Electrophoresis* 20(18):3551–3567. [https://doi.org/10.1002/\(SICI\)1522-2683\(19991201\)20:18<3551::AID-ELPS3551>3.0.CO;2-2](https://doi.org/10.1002/(SICI)1522-2683(19991201)20:18<3551::AID-ELPS3551>3.0.CO;2-2)
17. Ermak G, Cancasci VJ, Davies KJ (2003) Cytotoxic effect of doxycycline and its implications for tet-on gene expression systems. *Anal Biochem* 318(1):152–154. [https://doi.org/10.1016/S0003-2697\(03\)00166-0](https://doi.org/10.1016/S0003-2697(03)00166-0)
18. Petrak J, Ivanek R, Toman O, Cmejla R, Cmejlova J, Vyoral D, Zivny J, Vulpe CD (2008) Deja vu in proteomics. A hit parade of repeatedly identified differentially expressed proteins. *Proteomics* 8(9):1744–1749. <https://doi.org/10.1002/pmic.200700919>
19. Yuan A, Rao MV, Sasaki T, Chen Y, Kumar A, Liem RK, Eyer J, Peterson AC, Julien J-P, Nixon RA (2006) α -Internexin is structurally and functionally associated with the neurofilament triplet proteins in the mature CNS. *J Neurosci* 26(39):10006–10019. <https://doi.org/10.1523/JNEUROSCI.2580-06.2006>
20. Omary MB, Liem RK (2016) *Methods in Enzymology. Intermediate filament proteins*, vol 568. Academic Press, Elsevier Inc.
21. García-Cruz C, Merino-Jiménez C, Ceja V, Aragón J, Siqueiros-Márquez L, Reyes-Grajeda JP, Montañez C (2019) The dystrophin isoform Dp71e Δ 71 is involved in neurite outgrowth and neuronal differentiation of PC12 cells. *J Proteomics* 191:80–87. <https://doi.org/10.1016/j.jprot.2018.03.027>
22. Kaplan MP, Chin S, Fliegner KH, Liem R (1990) Alpha-internexin, a novel neuronal intermediate filament protein, precedes the low molecular weight neurofilament protein (NF-L) in the developing rat brain. *J Neurosci* 10(8):2735–2748. <https://doi.org/10.1523/JNEUROSCI.10-08-02735.1990>
23. Chiu F-C, Barnes E, Das K, Haley J, Socolow P, Macaluso FP, Fant J (1989) Characterization of a novel 66 kd subunit of mammalian neurofilaments. *Neuron* 2(5):1435–1445. [https://doi.org/10.1016/0896-6273\(89\)90189-x](https://doi.org/10.1016/0896-6273(89)90189-x)
24. Greene LA, Tischler AS (1976) Establishment of a noradrenergic clonal line of rat adrenal pheochromocytoma cells which respond to nerve growth factor. *Proc Natl Acad Sci U S A* 73(7):2424–2428. <https://doi.org/10.1073/pnas.73.7.2424>
25. Pollock JD, Krempin M, Rudy B (1990) Differential effects of NGF, FGF, EGF, cAMP, and dexamethasone on neurite outgrowth and sodium channel expression in PC12 cells. *J Neurosci* 10(8):2626–2637. <https://doi.org/10.1523/JNEUROSCI.10-08-02626.1990>

26. Martin T, Grishanin R (2003) PC12 cells as a model for studies of regulated secretion in neuronal and endocrine cells. *Methods Cell Biol* 71(34):267–286. [https://doi.org/10.1016/S0091-679X\(03\)01012-4](https://doi.org/10.1016/S0091-679X(03)01012-4)
27. Marquez FG, Cisneros B, Garcia F, Ceja V, Velazquez F, Depardon F, Cervantes L, Rendon A, Mornet D, Rosas-vargas H, Mustre M, Montanez C (2003) Differential expression and subcellular distribution of dystrophin Dp71 isoforms during differentiation process. *Neuroscience* 118(4):957–966. [https://doi.org/10.1016/S0306-4522\(03\)00063-0](https://doi.org/10.1016/S0306-4522(03)00063-0)
28. Saint Martin A, Aragon J, Depardon-Benitez F, Sanchez-Trujillo A, Mendoza-Hernandez G, Ceja V, Montanez C (2012) Identification of Dp71e, a new dystrophin with a novel carboxy-terminal end. *FEBS J* 279(1):66–77. <https://doi.org/10.1111/j.1742-4658.2011.08399.x>
29. Acosta R, Montañez C, Fuentes-Mera L, Gonzalez E, Gómez P, Quintero-Mora L, Mornet D, Alvarez-Salas LM, Cisneros B (2004) Dystrophin Dp71 is required for neurite outgrowth in PC12 cells. *Exp Cell Res* 296(2):265–275. <https://doi.org/10.1016/j.yexcr.2004.01.015>
30. Martínez-Herrera A, Aragón J, Bermúdez-Cruz RM, Bazán ML, Soid-Raggi G, Ceja V, Coy-Arechavaleta AS, Alemán V, Depardón F, Montañez C (2015) In silico analyses of dystrophin Dp40 cellular distribution, nuclear export signals and structure modeling. *Data Brief* 4:349–352. <https://doi.org/10.1016/j.dib.2015.06.007>
31. Aragón J, Romo-Yáñez J, Martínez-Herrera A, Ceja V, Rendon A, Montañez C (2011) Characterization of Dp71Δ78-79, a novel dystrophin mutant that stimulates PC12 cell differentiation. *J Neurochem* 119(4):697–707. <https://doi.org/10.1111/j.1471-4159.2011.07347.x>
32. Donato R (2003) Intracellular and extracellular roles of S100 proteins. *Microsc Res Tech* 60(6):540–551. <https://doi.org/10.1002/jemt.10296>
33. Masiakowski P, Shooter E (1990) Changes in PC12 cell morphology induced by transfection with 42C cDNA, coding for a member of the S-100 protein family. *J Neurosci Res* 27(3):264–269. <https://doi.org/10.1002/jnr.490270304>
34. Fausson-Pellegrini M-S, Matini P, DeFelici M (1999) The cytoskeleton of the myenteric neurons during murine embryonic life. *Anat Embryol* 199(5):459–469. <https://doi.org/10.1007/s004290050244>
35. Fliegner KH, Kaplan MP, Wood TL, Pintar JE, Liem RK (1994) Expression of the gene for the neuronal intermediate filament protein α -internexin coincides with the onset of neuronal differentiation in the developing rat nervous system. *J Comp Neurol* 342(2):161–173. <https://doi.org/10.1002/cne.903420202>
36. McGraw TS, Mickle JP, Shaw G, Streit WJ (2002) Axonally transported peripheral signals regulate α -internexin expression in regenerating motoneurons. *J Neurosci* 22(12):4955–4963. <https://doi.org/10.1523/JNEUROSCI.22-12-04955.2002>
37. Ching GY, Liem R (1993) Assembly of type IV neuronal intermediate filaments in nonneuronal cells in the absence of preexisting cytoplasmic intermediate filaments. *J Cell Biol* 122(6):1323–1335. <https://doi.org/10.1083/jcb.122.6.1323>
38. Kornreich M, Malka-Gibor E, Laser-Azogui A, Doron O, Herrmann H, Beck R (2015) Composite bottlebrush mechanics: α -internexin fine-tunes neurofilament network properties. *Soft Matter* 11(29):5839–5849. <https://doi.org/10.1039/C5SM00662G>
39. Ilieș I, Zupanc M, Zupanc G (2012) Proteome analysis reveals protein candidates involved in early stages of brain regeneration of teleost fish. *Neuroscience* 219:302–313. <https://doi.org/10.1016/j.neuroscience.2012.05.028>
40. Caligiore D, Arbib MA, Miall RC, Baldassarre G (2019) The super-learning hypothesis: Integrating learning processes across cortex, cerebellum and basal ganglia. *Neurosci Biobehav Rev* 100:19–34. <https://doi.org/10.1016/j.neubiorev.2019.02.008>
41. Ching GY, Chien C-L, Flores R, Liem RK (1999) Overexpression of α -internexin causes abnormal neurofilamentous accumulations and motor coordination deficits in transgenic mice. *J Neurosci* 19(8):2974–2986. <https://doi.org/10.1523/JNEUROSCI.19-08-02974.1999>
42. Shirasu M, Kimura K, Kataoka M, Takahashi M, Okajima S, Kawaguchi S, Hirasawa Y, Ide C, Mizoguchi A (2000) VAMP-2 promotes neurite elongation and SNAP-25A increases neurite sprouting in PC12 cells. *Neurosci Res* 37(4):265–275. [https://doi.org/10.1016/S0168-0102\(00\)00125-5](https://doi.org/10.1016/S0168-0102(00)00125-5)
43. Bloom OE, Morgan JR (2011) Membrane trafficking events underlying axon repair, growth, and regeneration. *Mol Cell Neurosci* 48(4):339–348. <https://doi.org/10.1016/j.mcn.2011.04.003>

Tables

Proteins identified by mass spectrometry											
Spot ID	Variation	Fold change in spot	MS	Accession no. ^a	Protein	Gene	MW [kDa]	pI	Match pept.	SC (%) ^b	
1	+	2	MALDI	P23565	Alpha internexin	Ina	56.11	5.20	14	29.8	
2	+	2.1	MALDI	P05964	S100A6 isoform	S100a6	10.03	5.30	1	58.4	
5	+	1.15	MALDI	P25113	Phosphoglycerate mutase 1	Pgam1	28.83	6.67	4	29.1	
6	+	1.47	MALDI	P04642	L-lactate dehydrogenase A chain	Ldha	36.45	8.45	2	21.4	
7	+	1.41	MALDI	P04764	Alpha enolase	Eno1	47.12	6.12	12	45.3	
8	-	1.52	MALDI	P84039	Ectonucleotide pyrophosphatase/phosphodiesterase	Enpp5	54.38	5.52	1	16.4	
9	+	1.56	MALDI	Q3MHS9	T-complex protein 1 subunit zeta	Cct6a	58.01	6.46	2	10.5	
10	+	1.20	MALDI	P06761	Endoplasmic reticulum chaperone BIP	Hspa5	72.33	5.07	5	17.6	
^a Accession number according to the swiss-prot <i>Rattus norvegicus</i> database											
^b SC %= % of the sequence identified											

Figures

Figure 1

Morphometric analysis of undifferentiated and NGF-differentiated PC12-Dp40 and PC12-Dp40L170P cells. PC12 control, PC12-Dp40 and PC12-Dp40L170P cells were differentiated with NGF for nine days. (a) Morphometric analysis was performed using clear-field micrographs of the undifferentiated (top panel) and differentiated PC12 cells (bottom panel). (b) Quantification of the differentiation ratio (g) and neurite length (h) of the NGF-differentiated PC12-Dp40 and PC12-Dp40L170P cells compared to the PC12 control cells is shown. The scale bar represents 10 μm

Figure 2

Two-dimensional gel electrophoresis images of proteins from the NGF-differentiated PC12 control and PC12-Dp40L170P cells. Total protein extracts of the NGF-differentiated PC12 control and PC12-Dp40L170P cells were collected after differentiation for nine days and separated by isoelectric focusing using immobilized pH nonlinear strips (7 cm), followed by SDS-PAGE. Gels were stained with coomassie colloidal blue. (a) Protein profile of the PC12 control cells and (b) the PC12-Dp40L170P cells. Numbers 1-14 indicate the IDs of the differentially expressed protein spots. (c) Magnified views of the differentially expressed protein spots. Arrows represent the nonlinear immobilized pH gradient used for IEF. The positions of standard markers for the second dimension are shown on the left side of each image

Figure 3

Expression of the proteins α-interneixin, VAMP, HspB1, NF-L, Myc-Dp40 and Myc-Dp40L170P in the undifferentiated PC12-Dp40 and PC12-Dp40L170P cells. Protein extracts were obtained from the undifferentiated PC12 control, PC12-Dp40 and Dp40L170P cells and analysed by WB. (a) Expression of α-interneixin, VAMP, HspB1, NF-L, HspB1, Myc-Dp40 and Myc-Dp40L170P. (b) Relative expression of α-interneixin, VAMP, and NF-L was determined as indicated in the Materials and Methods. The graph represents the mean ± SD from three independent experiments. *P ≤ 0.034 and 0.016 for α-interneixin and NF-L comparing the PC12 control cells and the PC12-Dp40L170P cells. ***P ≤ 0.0006 and **P ≤ 0.0018 for Myc-Dp40 and Myc-Dp40L170P comparing the PC12-Dp40 and PC12-Dp40L170P with PC12 control cells. *P ≤ 0.0124 for Myc-Dp40 comparing the PC12-Dp40 and PC12-Dp40L170P cells. β-actin was used as a loading control. MW is indicated in kDa

Figure 4

Page 13/15

Differential expression of α -internexin, VAMP, HspB1, NF-L, Myc-Dp40 and Myc-Dp40L170P in the NGF-differentiated PC12-Dp40 and PC12-Dp40L170P cells. Protein extracts were obtained from the NGF-differentiated PC12 control, PC12-Dp40 and PC12-Dp40L170P cells and analysed by WB. (a) Expression of α -internexin, VAMP, HspB1, NF-L, Myc-Dp40 and Myc-Dp40L170P. (b) Relative expression of α -internexin, VAMP, HspB1, NF-L, Myc-Dp40 and Myc-Dp40L170P was determined as indicated in the Materials and Methods. The graph presents the mean $\pm$ SD from three independent experiments. **P $\leq$ 0.002 and *P $\leq$ 0.021 for α -internexin comparing the PC12-Dp40L170P with PC12 control and PC12-Dp40 cells, respectively. **P $\leq$ 0.008 and **P $\leq$ 0.007 for VAMP comparing the PC12 control with PC12-Dp40 and PC12-Dp40L170P cells, respectively, and *P $\leq$ 0.015 comparing PC12-Dp40 and PC12-Dp40L170P. *P $\leq$ 0.040 for NF-L comparing the PC12 control and PC12-Dp40L170P cells. *P $\leq$ 0.010 and *P $\leq$ 0.040 for Myc-Dp40 and Myc-Dp40L170P comparing the PC12-Dp40 and PC12-Dp40L170P with PC12 control cells. β -actin was used as a loading control. MW is indicated in kDa

Figure 5

Cellular distribution of α -internexin in undifferentiated and NGF-differentiated PC12-Dp40 and PC12-Dp40L170P cells. ImmF was performed on the PC12-Dp40, PC12-Dp40L170P and PC12 control cells after the induction of differentiation. ImmF staining for the Myc-Dp40 and Myc-Dp40L170P proteins was performed using the anti-c-Myc antibody (green) and α -internexin (red). Nuclei were stained with DAPI (blue). Images represent a single equatorial Z-section from confocal images to show the localization of each protein. (a) Undifferentiated and (b) differentiated PC12 control, PC12-Dp40 and PC12-Dp40L170P cells. Merged images correspond to the overlap of Myc-Dp40 and Myc-Dp40L170P with the α -internexin protein. The figure is a representative result of three independent experiments. The scale bar represents 20 μ m

This is a list of supplementary files associated with this preprint. Click to download.

- [Electronicsupplementarymaterial.docx](#)