

Acircularity and Circularity Indexes of the Foveal Avascular Zone in High Myopia

Helong Piao

Chonnam National University Medical School and Hospital

Yue Guo

Wenzhou Medical University

Haowei Zhang

Chonnam National University Medical School and Hospital

Mi Sun Sung

Chonnam National University Medical School and Hospital

Sang Woo Park (✉ exo70@naver.com)

Chonnam National University Medical School and Hospital

Research Article

Keywords: foveal avascular zone (FAZ), Myopization, Acircularity and circularity indexes, high myopia

Posted Date: April 6th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-385310/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on August 19th, 2021.
See the published version at <https://doi.org/10.1038/s41598-021-96304-9>.

Abstract

We explored the association between foveal avascular zone (FAZ) parameters and high myopia using optical coherence tomography angiography (OCTA). To evaluate the specific influence of high myopia on the FAZ, we divided 106 eyes of 106 patients into quartiles based on the axial length. The upper quartile was then defined as the high myopia group ($n = 27$), while the lower quartile was the non-high myopia group ($n = 26$). The FAZ parameters of these two groups were then retrospectively compared to investigate variations in FAZ size and morphology. The mean axial length and spherical equivalent were 23.96 ± 0.94 mm and -2.27 ± 1.58 diopters in the non-high myopia group and 27.42 ± 0.36 mm and -8.12 ± 2.14 diopters in the high myopia group, respectively. The areas of superficial and deep FAZ, perimeters of deep FAZ, major axis lengths of deep FAZ, minor axis lengths of superficial and deep FAZ were significantly larger in eyes with high myopia than in eyes with non-high myopia ($P < 0.05$). Linear regression analyses showed that no significant correlation was observed between FAZ areas and acircularity and circularity index of FAZ in non-high myopia group, conversely, FAZ areas strongly correlated with acircularity and circularity index of FAZ in high myopia group. We found that An increase in the FAZ area in highly myopic eyes was accompanied by a significant variation in FAZ acircularity and circularity indexes. Further research should address whether these findings are associated with future disease development in highly myopic eyes.

Introduction

Myopization, especially high myopization, is a complex pathological process involving the whole eyeball. It is characterized by axial elongation, which can cause retinal detachment, choroid neovascularization, macular hemorrhage, and retinal ischemia, all of which lead to visual dysfunction.¹⁻³ One meta-analysis suggested that nearly half of the world's population will be myopic by 2050, with up to 10% showing high myopia.⁴ With the increased incidence of myopia, more knowledge about myopia progression is necessary. For several decades, many studies have focused on the pathological changes of myopia, including the effects of axial elongation on the chorioretinal thickness and retinal vasculature.⁵⁻⁹ However, the pathological processes underlying myopia development remain unclear.

Recently, optical coherence tomography angiography (OCTA) has been commonly used in the clinical practice. It allows clinicians to view the various layers of the retinal and choroidal microvasculature non-invasively. Since many eye diseases involve the loss of retinal vessels, ophthalmologists could diagnose and manage patients using OCTA monitoring. One recent study showed that even when the OCT-measured parameters reach floor level, OCTA-measured vessel density is still present a promising indicator for monitoring progress in advanced glaucoma.¹⁰ It follows that OCTA may possess better discriminatory power than OCT.

The foveal avascular zone (FAZ) is an important measurement parameter in OCTA. It is a capillary dropout zone in the macula and has the highest density of cone photoreceptors.⁵ The FAZ is surrounded by an interconnecting capillary network derived from branches of the central retinal artery (CRA), which

branches out on the retinal nerve fiber layer (RNFL) level to form the superficial plexus within the ganglion cell layer. Deeper branches reaching into the inner nuclear layer form the deep plexus.¹¹ Therefore, the FAZ reflects alterations in the central retinal microvasculature and is related to central visual function.¹² Several clinical elements can influence FAZ parameters, including diabetic retinopathy (DR), axial length, sex, and glaucoma.^{12–16} Kwon et al. reported an increase in FAZ area and a decrease in the circularity index in glaucomatous eyes.¹² Krawitz et al. used the acircularity index and axial ratio of the FAZ to demonstrate that the FAZ shape was deformed in patients with DR.¹³ He et al. and Cheng et al. showed that increased FAZ area and decreased vessel density were correlated with high myopia.^{14,15} Interestingly, FAZ enlargement occurs in both glaucomatous and myopic eyes. Given that glaucoma is closely related to myopia, further research should address how the FAZ varies among myopic individuals.

Since myopia progression mainly affects the posterior pole of the retina through axial elongation, highly myopic eyes may show some characteristic changes in the FAZ. Accordingly, in the present study, we measured the acircularity index, circularity index, and other biomarkers of the FAZ to quantify the shape of the FAZ and determine whether FAZ deformation was significantly correlated with high myopia.

Methods

Subjects

This retrospective, cross-sectional study was conducted according to the tenets of the Declaration of Helsinki and was approved by the Institutional Review Board of Chonnam National University Hospital. The participants were informed of the study objectives and written informed consent was obtained from all participants.

Healthy myopic volunteers were recruited consecutively from July 1, 2016 to September 31, 2016. A detailed medical history was recorded for each subject. All subjects underwent complete ophthalmic examination, including best-corrected visual acuity (BCVA), intraocular pressure (IOP) using Goldmann applanation tonometry, slit-lamp examination, refractive error assessment, fundus examination using color stereoscopic disc photography and red-free RNFL fundus photography, and Swedish Interactive Threshold Algorithm standard 30 – 2 perimetry using a Humphrey Field Analyzer (Carl Zeiss Meditec Inc., Dublin, California, USA). All IOP measurements were recorded between 4:00 PM and 7:00 PM. Axial length and central corneal thickness were measured using optical low-coherence reflectometry (Lenstar; Haag-Streit AG, Koeniz, Switzerland). Avanti spectral-domain OCT (RTVue-XR Avanti; Optovue, Fremont, California, USA) was used to measure the chorioretinal thickness.

All volunteers were required to meet the following inclusion criteria: (1) age between 18 and 30 years; (2) astigmatism within ± 2 diopters; (3) BCVA of 20/25 or better; (4) IOP ≤ 21 mmHg; (5) normal optic nerve head with non-glaucomatous changes on stereoscopic photographs, including an intact neuroretinal rim without thinning, pit, or localized pallor; (6) absence of RNFL abnormalities on red-free fundus photographs; and (7) normal visual field results in both eyes. The exclusion criteria included (1) history of

glaucoma or intraocular surgery; (2) IOP > 21 mmHg; (3) myopic degeneration, including peripheral retinal tears, intrachoroidal cavitations, or choroidal neovascularization; (4) history of systemic disease that could affect the eyes, such as diabetes mellitus, (5) other evidence of retinal pathology or opaque media. For cases in which both eyes of a subject met the inclusion criteria, one eye was chosen randomly.

Spectral-Domain Optical Coherence Tomography Angiography

The spectral-domain OCT instrument used was the RTVue Avanti with prototype AngioVue® OCTA software (Optovue, Inc., Fremont, CA). This instrument operates at ~ 840 nm wavelength and conducts 70,000 A-scans per second to acquire OCTA volumes consisting of two repeated B-scans from 304 sequential, uniformly spaced locations. Each B-scan consists of 304 A-scans, giving a total of $2 \times 304 \times 304$ A-scans per acquisition, with a total acquisition time of approximately 3 seconds and an axial optical resolution of ~ 5 μm . Macular 3×3 -mm region scans centered on the fovea were captured from each subject. OCTA perfusion maps were generated using the split-spectrum amplitude decorrelation angiography (SSADA) algorithm, as previously described.¹⁷ The Optovue software built into the Avanti OCT automatically segmented the inner retinal blood vessels into the superficial and deep capillary plexus layers. The boundaries of the superficial capillary plexus extended from 3 mm below the inner limiting membrane to 15 mm below the inner plexiform layer (IPL). The deep capillary plexus extended from 15 mm to 70 mm below the IPL.

FAZ Measurements

The following measurements were made, as described in detail elsewhere: FAZ area, FAZ perimeter, acircularity index, circularity index, axial ratio, major axis length, and minor axis length.^{13,18} Briefly, the area, perimeter length, acircularity index, and circularity index of FAZ were measured automatically, and the axial ratio, major axis length, and minor axis length of the FAZ were measured manually using ImageJ software 1.52a (National Institutes of Health, USA) (Fig. 1). Littman and the modified Bennett formulas were used to correct ocular magnification induced by axial variation on the area of FAZ.¹⁹ A perfectly circular FAZ has an acircularity index equal to 1.0, with deviations from a circular shape leading to an increase in this metric. Acircularity is defined by the following equation:

Acircularity = perimeter of FAZ / perimeter of the circle with equal area

Circularity is a shape descriptor that can mathematically indicate the degree of similarity of the FAZ to a perfect circle. A value of 1.0 designates a perfect circle; as the circularity value decreases, the shape is increasingly less circular. Circularity is defined by the following equation:

Circularity = $4\pi \times \text{area} / \text{perimeter}^2$

The following parameters were determined from a best-fit ellipse of the FAZ: the length of the major and minor axes, and the axial ratio, which is calculated using the following equation:

Axial ratio = major axis length of FAZ / minor axis length of FAZ

To decrease the measurement error, the average data from two independent examiners (H.P. and H.Z.) were used statistically in this study.

Statistical Analysis

IBM SPSS Statistics for Windows, Version 25.0 (IBM Corp., Armonk, NY, USA) was used for statistical analysis. The inter-observer agreement of the FAZ parameters was evaluated by calculating the intraclass correlation coefficients. The Mann–Whitney U test was used to compare the FAZ parameters between non-high myopia and high myopia groups. Linear regression analysis was used to investigate whether the areas of the superficial and deep FAZ were correlated with other FAZ parameters in each group. *P*-values < 0.05 were considered statistically significant.

Results

A total of 106 eyes of 106 subjects with myopia were enrolled. The general characteristics of the study subjects are listed in Table 1. The mean axial length was 25.9 ± 1.4 mm (range: 22.0–28.1 mm); mean spherical equivalent, -5.4 ± 2.9 diopters (range: -12.125 to -0.25 diopters); mean foveal thickness, 251.0 ± 19.8 μ m (range: 200–297 μ m); and subfoveal choroidal thickness, 258.4 ± 17.4 μ m (range: 132 to 373 μ m). Inter-observer agreement of the following FAZ measurements showed excellent reproducibility, as estimated using intraclass correlation coefficients: area, perimeter, circularity index, acircularity index, axial ratio, and major and minor axis length (data not shown).

Table 1
Demographic and ocular factors of study subjects

	Number of subjects (n = 106)		
	Mean	SD	Range
Age (years)	23.8	4.2	18–34
Sex, male/female	74/32		
Axial length (mm)	25.9	1.4	22.1–28.1
IOP (mm Hg)	14.2	2.6	10–21
Spherical equivalent (diopter)	-5.4	2.9	-12.125--0.25
Central corneal thickness (μm)	555.5	31.0	487–632
Average keratometry (diopter)	42.4	2.3	38.2–47.2
Foveal thickness (μm)	251.0	19.8	200–297
Parafoveal thickness (μm)	313.4	15.2	268–356
Subfoveal choroidal thickness (μm)	258.4	17.8	132–373
IOP = intraocular pressure.			

To investigate the morphological changes in the FAZ that were correlated with high myopia, we grouped all subjects based on the method recommended by Joarder and Firozzaman.²⁰ We first defined the upper and lower quartiles (upper quartile: n = 27, cut-off value = 26.84 mm; lower quartile: n = 26, cut-off value = 25.02 mm). The remaining 53 subjects belonged to the second and third quartiles. After grouping, only those with high myopia were included in the upper quartile group, while none of those with high myopia were in the lower quartile group, in which all participants had low to moderate myopia. The results of the grouping and a comparison of FAZ parameters between the two groups are shown in Table 2. In the non-high myopia group, the mean axial length was 23.96 ± 0.94 mm (range: 22.09–25.02 mm) and the spherical equivalent was -2.27 ± 1.58 diopters (range: -5.25 to -0.25 diopters), while in the high myopia group, the mean axial length was 27.42 ± 0.36 mm (range: 26.86–28.11 mm) and the spherical equivalent was -8.12 ± 2.14 diopters (range: -12.125 to -6.25 diopters). The following measurements were significantly larger in eyes with high myopia than in those with non-high myopia: the areas of the superficial and deep FAZ, the perimeters of the deep FAZ, the major axis lengths of the deep FAZ, and the minor axis lengths of the superficial and deep FAZ ($P < 0.05$ in all cases). No other parameter showed any significant difference between the groups (acircularity index, circularity index, and axial ratio of the superficial and deep FAZ). These results showed that significant differences occurred more often with the deep FAZ.

Table 2

Ocular factors and FAZ parameters of OCTA in non-high myopia (lower quartile) and high myopia (upper quartile) groups.

Variables	Non-high myopia (n = 26)	High myopia (n = 27)	P Value
Axial length (mm)	24.0 ± 0.9 (22.09–25.02)	27.4 ± 0.3 (26.86–28.11)	< 0.001
Spherical equivalent (diopter)	-2.3 ± 1.6 (-5.25--0.25)	-8.1 ± 2.1 (-12.125--6.25)	< 0.001
Area of superficial FAZ (mm ²)	0.3 ± 0.08 (0.18–0.52)	0.43 ± 0.22 (0.20–1.15)	0.024
Area of deep FAZ (mm ²)	0.45 ± 0.12 (0.23–0.68)	0.66 ± 0.25 (0.33–1.36)	< 0.001
Perimeter of superficial FAZ (mm)	2.9 ± 0.46 (2.05–4.08)	3.47 ± 1.41 (2.06–8.60)	0.122
Perimeter of deep FAZ (mm)	3.96 ± 0.71 (2.79–5.62)	4.87 ± 1.41 (2.82–8.93)	0.005
Circularity index of superficial FAZ	0.48 ± 0.10 (0.29–0.67)	0.48 ± 0.12 (0.20–0.67)	0.873
Circularity index of deep FAZ	0.37 ± 0.06 (0.23–0.46)	0.36 ± 0.09 (0.20–0.56)	0.423
Acircularity index of superficial FAZ	1.47 ± 0.17 (1.22–1.85)	1.49 ± 0.24 (1.22–2.27)	0.873
Acircularity index of deep FAZ	1.67 ± 0.17 (1.47–2.11)	1.69 ± 0.21 (1.33–2.26)	0.423
Axial ratio of superficial FAZ	1.24 ± 0.19 (1.03–1.92)	1.21 ± 0.20 (1.01–1.70)	0.310
Axial ratio of deep FAZ	1.18 ± 0.13 (1.01–1.48)	1.24 ± 0.17 (1.02–1.65)	0.179
Major axis length of superficial FAZ (mm)	0.78 ± 0.11 (0.55–1.01)	0.93 ± 0.34 (0.55–2.09)	0.065
Major axis length of deep FAZ (mm)	0.93 ± 0.15 (0.67–1.31)	1.13 ± 0.21 (0.82–1.71)	< 0.001
Minor axis length of superficial FAZ (mm)	0.64 ± 0.10 (0.44–0.87)	0.77 ± 0.26 (0.43–1.72)	0.012

FAZ = foveal avascular zone.

Data are mean ± standard deviation unless otherwise indicated.

Factors with statistical significance are shown in boldface

Variables	Non-high myopia (n = 26)	High myopia (n = 27)	<i>P</i> Value
Minor axis length of deep FAZ (mm)	0.79 ± 0.11 (0.60–1.02)	0.92 ± 0.15 (0.72–1.34)	0.003
FAZ = foveal avascular zone.			
Data are mean ± standard deviation unless otherwise indicated.			
Factors with statistical significance are shown in boldface			

Linear regression analysis adjusted for age and sex was conducted to examine whether the FAZ parameters of OCTA were correlated with FAZ area in the non-high myopia and high myopia groups. As shown in Table 3, the areas of the superficial and deep FAZ were not significantly correlated with the circularity index of the FAZ in the non-high myopia group. Conversely, the areas of the superficial and deep FAZ were significantly negatively correlated with circularity index ($P = 0.001$ in both layers) of the FAZ in the high myopia group. Likewise, the areas of the superficial and deep FAZ were not significantly correlated with the acircularity index of the FAZ in the non-high myopia group (Fig. 2A, B), but the areas of the superficial and deep FAZ were significantly positively correlated with acircularity index of the FAZ in the high myopia group ($P < 0.001$ in both layers) (Fig. 2C, D). In both groups, the areas of the superficial and deep FAZ were significantly positively correlated with the perimeters of the FAZ ($P < 0.001$ in both layers). Only the area of the superficial FAZ in the non-high myopia group was significantly correlated with the axial ratio of the FAZ ($P = 0.011$). In addition, the areas of the superficial FAZ in the non-high myopia group were weakly correlated with the major axis lengths of the FAZ ($P = 0.024$), while other major and minor axis lengths of the FAZ were strongly associated with the FAZ areas ($P < 0.001$ in both groups).

Table 3

Associations of the FAZ parameters of OCTA with FAZ area in non-high myopia (lower quartile) and high myopia (upper quartile) groups

	Area of Superficial FAZ			Area of deep FAZ		
	R ²	Coefficient	P Value*	R ²	Coefficient	P Value*
Non-high myopia						
Circularity index of FAZ	0.027	0.190	0.536	0.069	0.001	0.997
Acircularity index of FAZ	0.053	-0.478	0.347	0.079	-0.040	0.951
Perimeter of FAZ	0.534	3.787	< 0.001	0.704	4.553	< 0.001
Axial ratio of FAZ	0.269	-1.339	0.011	0.013	0.129	0.659
Major axis length of FAZ	0.406	0.580	0.024	0.697	1.074	< 0.001
Minor axis length of FAZ	0.818	1.086	< 0.001	0.754	0.853	< 0.001
High myopia						
Circularity index of FAZ	0.427	-0.352	0.001	0.409	-0.223	0.001
Acircularity index of FAZ	0.588	0.814	< 0.001	0.494	0.595	< 0.001
Perimeter of FAZ	0.931	6.243	< 0.001	0.915	5.516	< 0.001
Axial ratio of FAZ	0.094	0.186	0.336	0.254	0.065	0.600
Major axis length of FAZ	0.892	1.458	< 0.001	0.708	0.707	< 0.001
Minor axis length of FAZ	0.764	1.030	< 0.001	0.805	0.520	< 0.001
FAZ = foveal avascular zone.						
*P value adjusted for age and sex.						
Factors with statistical significance are shown in boldface.						

Discussion

In this study, we utilized OCTA to analyze the macular microvasculature and further explore the morphological changes of FAZ. We found that several FAZ parameters were larger in highly myopic eyes than in non-highly myopic eyes, and that the differences were even more pronounced in the deep FAZ layer. Moreover, the acircularity and circularity indexes of the FAZ were significantly correlated with FAZ enlargement in highly myopic eyes, whereas no significant relationships were found between these parameters in the non-highly myopic group. To our knowledge, this was the first report to determine FAZ deformation in high myopia.

Several reports have investigated the FAZ in myopic eyes.^{5, 14–16, 21, 22} Sung et al. reported lower peripapillary vessel density and larger superficial and deep FAZ areas in highly myopic eyes, as elucidated using OCTA.⁵ He et al. also demonstrated that reduced radial peripapillary capillary, deep parafoveal vessel density, and enlarged FAZ area occurred in high myopia.¹⁴ Similarly, Cheng et al. showed that increased FAZ area and decreased vessel density in the superficial and deep layers were correlated with axial elongation.¹⁵ All these studies reported that FAZ enlargement was accompanied by a reduction of retinal vessel density in the macular region, indicating that the size of the FAZ indirectly reflects variations in retinal perfusion. Although we did not measure vessel density in the present study, our results are in accordance with those of the previous reports.

The mechanism of FAZ enlargement in myopic eyes remains unclear, although some hypotheses have been offered. First, retinal vascular trunk dragging during myopia progression may decrease retinal blood flow and further result in variation in the FAZ size. Two previous studies reported that the FAZ areas were significantly associated with optic nerve head tilt, which occurs during axial elongation.^{5, 14} Second, in eyes with axial elongation, overall macular thinning may cause reduced oxygen consumption, resulting in decreased retinal blood flow and increased FAZ area.^{23, 24} In the current study, the FAZ areas in highly myopic eyes were significantly larger than those in non-highly myopic eyes, and the difference was more significant in the deep FAZ layers, suggesting that the deep capillary plexus is more susceptible to myopia-related changes, possible because the arterial blood supply differs between the superficial and deep capillary plexuses. The oxygen and nutrition demand of the superficial retina is met by the central retinal artery, while that of the deep retina is met by the choroidal vascular system.²⁵ Since choroidal thinning is more prominent in highly myopic eyes,²⁶ FAZ enlargement might be more pronounced in the deep retinal layer.

Numerous studies have researched FAZ deformation, but none have explored the morphological changes in the FAZ in high myopia. In the present study, we speculated that either or both of the following mechanisms influence FAZ morphology: First, the retinal vascular trunk in the ONH was dragged, resulting in irregular reduction in macular blood perfusion, with consequent FAZ deformation and enlargement during axial elongation. Second, despite individual variations, the FAZ morphology in non-highly myopic eyes may be relatively circular in most cases. Moreover, as the retinal vascular trunk was dragged nasally in highly myopic eyes, the macular microvasculature may have been distorted towards the ONH, leading to FAZ deformation. One prospective observational study on progressive myopia indicated that the retinal vasculature at the posterior pole was unchanged, and that only the central vascular trunk in the lamina cribrosa was dragged nasally.²⁷ However, that study did not measure the retinal vasculature of the macular region using OCTA; from a macroscopic point of view, it demonstrated that the posterior polar blood vessels did not change, but it did not analyze the macular microvasculature. As such, the results were insufficient to explain the lack of variation in the FAZ capillaries. Therefore, depending on geometrical change in the FAZ, it may be possible to confirm myopia severity and macular perfusion status. Future longitudinal study should verify this.

Of note, we found that the FAZ areas were not significantly associated with the acircularity or circularity index in non-highly myopic eyes. However, they were positively correlated with acircularity index and negatively correlated with circularity index in highly myopic eyes, suggesting that the FAZ area becomes larger and more acircular as myopia progresses. Variations in the FAZ acircularity/circularity indexes and FAZ size may have multiple clinical significances. Kwon et al. showed that a decreased FAZ circularity index was significantly associated with central visual field defect, and that the FAZ size was significantly associated with the severity of central visual field defect in glaucomatous eyes.²⁸ Choi et al. reported that decreased FAZ circularity index may represent a disruption of the parafoveal capillary network in patients with glaucoma.²⁹ Kim et al. also showed that decreased FAZ circularity is a good indicator of vascular dropout, and that it is associated with disease progression in vascular maculopathy.³⁰ Another study found a significant difference in acircularity index between controls and non-proliferative DR. This result could inform objective staging of the disease.¹³ Before the disease has progressed significantly, the retinal microvascular system in myopic eyes has already suffered structural damage, and discernible microvascular changes are visible in the FAZ. Given that FAZ acircularity is associated with visual field defects and macular vascular dropout, our results might explain the reason for the high prevalence of central scotoma in myopic glaucoma.

Recently, Krawitz et al. reported that, unlike other indicators such as area, perimeter, and length, the acircularity index can quantify FAZ geometry without the need for axial length measurements to correct for retinal magnification.¹³ It follows that the acircularity index (including circularity index) could serve as indicators to predict pathological myopic changes in advance. Long-term monitoring of FAZ evolution using the acircularity and circularity indexes may be necessary to substantiate the clinical value of these metrics, which are not influenced by ocular magnification correction.

The limitations of the current study were as follows: The study was retrospective, and to evaluate the specific influence of axial myopia, we set further criteria, which resulted in a small sample size. A prospective big data study is needed to improve data accuracy. Furthermore, we did not measure the retinal thickness in detail. One previous study demonstrated that early foveal microcirculatory alterations in diabetic eyes were related to macular ganglion cell/inner plexiform layer thickness, regardless of the presence of DR.³⁰ Intraretinal thickness may therefore be related to FAZ deformation in myopic eyes. Future studies should investigate this relationship further to deepen the current understanding of myopia progression.

In conclusion, we observed FAZ enlargement in highly myopic eyes, especially in the deep retinal layer. This enlargement accompanied changes in the acircularity and circularity indexes of the FAZ in highly myopic eyes. Further research should address whether these findings are associated with future disease development in highly myopic eyes.

Declarations

Data Availability

The datasets generated during the current study are available from the corresponding author on reasonable request.

ACKNOWLEDGEMENTS

This research was supported by a Patient-Centered Clinical Research Coordinating Center grant funded by the Ministry of Health & Welfare, Republic of Korea (HI19C0481, HC19C0276). The funding organizations had no role in the design or conduct of this research.

Author Contributions Statement

The design and conduct of the study (H.P., M.S.S., S.W.P.); collection of data (H.P., H.Z., M.S.S., Y.G.); management, analysis, and interpretation of data (H.P., Y.G., H.Z., M.S.S., S.W.P.); and preparation of the manuscript (H.P., M.S.S., S.W.P.). All authors reviewed the manuscript.

Additional information

The authors have no conflicts of interests to disclose regarding any material discussed in this article.

References

1. Cho, B. J., Shin, J. Y. & Yu, H. G. Complications of Pathologic M *Eye & contact lens* **42**, 9-15 (2016).
2. Costanzo, E. *et al.* Type 1 Choroidal Neovascularization Lesion Size: Indocyanine Green Angiography Versus Optical Coherence Tomography Angiography. *Investigative ophthalmology & visual science* **57**, OCT307-OCT313 (2016)
3. Kaneko, Y., Moriyama, M., Hirahara, S., Ogura, Y. & Ohno-Matsui, K. Areas of nonperfusion in peripheral retina of eyes with pathologic myopia detected by ultra-widefield fluorescein angiography. *Investigative ophthalmology & visual science* **55**, 1432-1439 (2014).
4. Holden, B. A. *et al.* Global Prevalence of Myopia and High Myopia and Temporal Trends from 2000 through 2050. *Ophthalmology* **123**, 1036-1042 (2016).
5. Sung, M. S., Lee, T. H., Heo, H. & Park, S. W. Association Between Optic Nerve Head Deformation and Retinal Microvasculature in High Myopia. *Am J Ophthalmol* **188**, 81-90 (2018).
6. Jonas, J. B. *et al.* Retinal Thickness and Axial Length. *Investigative ophthalmology & visual science* **57**, 1791-1797 (2016).
7. Ucak, T. *et al.* Alterations in optical coherence tomography angiography findings in patients with high myopia. *Eye* **34**, 1129-1135 (2020).
8. Wu, H. *et al.* Assessment of Choroidal Vascularity and Choriocapillaris Blood Perfusion in Anisomyopic Adults by SS-OCT/OCTA. *Investigative ophthalmology & visual science* **62**, 8 (2021).
9. Shen, L. *et al.* Scleral and Choroidal Thickness in Secondary High Axial Myopia. *Retina* **36**, 1579-1585 (2016).

10. Moghimi, S. *et al.* Measurement Floors and Dynamic Ranges of OCT and OCT Angiography in Glaucoma. *Ophthalmology* **126**, 980-988 (2019).
11. Zivkovic, M. *et al.* Foveal Avascular Zone in Normal Tension Glaucoma Measured by Optical Coherence Tomography Angiography. *BioMed research international* **2017**, 3079141 (2017).
12. Kwon, J., Choi, J., Shin, J. W., Lee, J. & Kook, M. S. Glaucoma Diagnostic Capabilities of Foveal Avascular Zone Parameters Using Optical Coherence Tomography Angiography According to Visual Field Defect Location. *J Glaucoma* **26**, 1120-1129 (2017).
13. Krawitz, B. D. *et al.* Acircularity index and axis ratio of the foveal avascular zone in diabetic eyes and healthy controls measured by optical coherence tomography angiography. *Vision Res* **139**, 177-186 (2017).
14. He, J. N. *et al.* Association between retinal microvasculature and optic disc alterations in high myopia. *Eye* **33**, 1494-1503 (2019).
15. Cheng, D. *et al.* Deep perifoveal vessel density as an indicator of capillary loss in high myopia. *Eye* **33**, 1961-1968 (2019).
16. Tan, C. S. *et al.* Optical Coherence Tomography Angiography Evaluation of the Parafoveal Vasculature and Its Relationship With Ocular Factors. *Investigative ophthalmology & visual science* **57**, Oct224-Oct234 (2016).
17. Jia, Y. L. *et al.* Split-spectrum amplitude-decorrelation angiography with optical coherence tomography. *Opt Express* **20**, 4710-4725 (2012).
18. Shiihara, H. *et al.* Objective evaluation of size and shape of superficial foveal avascular zone in normal subjects by optical coherence tomography angiography. *Sci Rep* **8**, 10143 (2018).
19. Sampson, D. M. *et al.* Axial Length Variation Impacts on Superficial Retinal Vessel Density and Foveal Avascular Zone Area Measurements Using Optical Coherence Tomography Angiography. *Investigative ophthalmology & visual science* **58**, 3065-3072 (2017).
20. Joarder, A.H. & Firozzaman, Quartiles for discrete data. *Teaching Statistics*. **23**, 86-89 (2001).
21. Min, C.H. *et al.* Macular microvasculature in high myopia without pathologic changes: an optical coherence tomography angiography s *Korean J Ophthalmol*. **34**, 106-112 (2020).
22. Golebiewska, J., Biala-Gosek, K., Czeszyk, A. & Hautz, W. Optical coherence tomography angiography of superficial retinal vessel density and foveal avascular zone in myopic children. *Plos One* **14**, e0219785 (2019).
23. Song, W. K., Lee, S. C., Lee, E. S., Kim, C. Y. & Kim, S. S. Macular Thickness Variations with Sex, Age, and Axial Length in Healthy Subjects: A Spectral Domain-Optical Coherence Tomography Study. *Investigative ophthalmology & visual science* **51**, 3913-3918 (2010).
24. La Spina, C., Corvi, F., Bandello, F. & Querques, G. Static characteristics and dynamic functionality of retinal vessels in longer eyes with or without pathologic myopia. *Graef Arch Clin Exp* **254**, 827-834 (2016).

25. Kur, J., Newman, E. A. & Chan-Ling, T. Cellular and physiological mechanisms underlying blood flow regulation in the retina and choroid in health and disease. *Prog Retin Eye Res* **31**, 377-406 (2012).
26. Al-Sheikh, M. *et al.* Quantitative OCT Angiography of the Retinal Microvasculature and the Choriocapillaris in Myopic Eyes. *Investigative ophthalmology & visual science* **58**, 2063-2069 (2017).
27. Lee, K. M., Choung, H. K., Kim, M., Oh, S. & Kim, S. H. Positional Change of Optic Nerve Head Vasculature during Axial Elongation as Evidence of Lamina Cribrosa Shifting Boramae Myopia Cohort Study Report 2. *Ophthalmology* **125**, 1224-1233 (2018).
28. Kwon, J., Choi, J., Shin, J. W., Lee, J. Y. & Kook, M. S. Alterations of the Foveal Avascular Zone Measured by Optical Coherence Tomography Angiography in Glaucoma Patients With Central Visual Field Defects. *Investigative ophthalmology & visual science* **58**, 1637-1645 (2017).
29. Choi, J. *et al.* Quantitative optical coherence tomography angiography of macular vascular structure and foveal avascular zone in glaucoma. *Plos One* **12**, e0184948 (2017).
30. Kim, K., Kim, E. S. & Yu, S. Y. Optical coherence tomography angiography analysis of foveal microvascular changes and inner retinal layer thinning in patients with diabetes. *Brit J Ophthalmol* **102**, 1226-1231 (2018).

Figures




Figure 1

(A) Optical coherence tomography angiogram of the FAZ. (B) The FAZ was separated from the background image and the area (A) and perimeter (P) were automatically measured using the ImageJ software. (C) An irregular outline was manually fit using the ImageJ software. (D) The major axis (black line) and minor axis (yellow line) lengths were also manually calculated. FAZ = foveal avascular zone.




Figure 2

Scatter plots showing the association between the FAZ areas and acircularity index of FAZ. (A, B) The areas of the superficial and deep FAZ were not significantly correlated with the acircularity index in the non-high myopia group. (C, D) The areas of the superficial and deep FAZ were strongly associated with the acircularity index in the high myopia group. FAZ = foveal avascular zone.