

# Development of Integration Indexes to Determine the Extent of Family Planning and Child Immunization Services Integration in Health Facilities in Urban Areas of Nigeria

**Kate L. Sheahan** (✉ [kate.sheahan@va.gov](mailto:kate.sheahan@va.gov))

University of North Carolina at Chapel Hill <https://orcid.org/0000-0001-5473-3646>

**Jennifer Orgill-Meyer**

Franklin and Marshall College

**Ilene Speizer**

University of North Carolina at Chapel Hill

**Sian Curtis**

University of North Carolina at Chapel Hill

**John Paul**

University of North Carolina at Chapel Hill

**Morris Weinberger**

University of North Carolina at Chapel Hill

**Antonia V. Bennett**

University of North Carolina at Chapel Hill

---

## Research

**Keywords:** Family planning, immunization, integration, measurement, Nigeria

**Posted Date:** December 31st, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-38714/v2>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

**Version of Record:** A version of this preprint was published on February 23rd, 2021. See the published version at <https://doi.org/10.1186/s12978-021-01105-y>.

## Abstract

**Background:** Integrating family planning into child immunization services may address unmet need for contraception by offering family planning information and services to postpartum women during routine child immunization visits. However, policies and programs promoting integration are often based on insubstantial or conflicting evidence about its effects on service delivery and health outcomes. While integration models vary, many studies measure integration as binary (a facility is integrated or not) rather than a multidimensional and varying continuum. It is thus challenging to ascertain the determinants and effects of integrated service delivery. This study creates Facility and Provider Integration Indexes, which measure capacity to support integrated family planning and child immunization services and applies them to analyze the extent of integration across 400 health facilities.

**Methods:** This study utilizes cross-sectional health facility (N= 400; 58% hospitals, 42% primary healthcare centers) and healthcare provider (N= 1,479) survey data that were collected in six urban areas of Nigeria for the impact evaluation of the Nigerian Urban Reproductive Health Initiative. Principal Components Analysis was used to develop Provider and Facility Integration Indexes that estimate the extent of integration in these health facilities. The Provider Integration Index measures provider skills and practices that support integrated service delivery while the Facility Integration Index measures facility norms that support integrated service delivery. Index scores range from zero (low) to ten (high).

**Results:** Mean Provider Integration Index score is 5.42 (SD 3.10), and mean Facility Integration Index score is 6.22 (SD 2.72). Twenty-three percent of facilities were classified as having low Provider Integration scores, 32% as medium, and 45% as high. Fourteen percent of facilities were classified as having low Facility Integration scores, 38% as medium, and 48% as high.

**Conclusion:** Many facilities in our sample have achieved high levels of integration, while many others have not. Results suggest that measuring integration as a binary variable does not (a) reflect the true variation in integration within and across health facilities, (b) enable nuanced measurement of the determinants or effects of integration, or (c) provide tailored, actionable information about how best to improve integration. Overall, results reinforce the importance of utilizing more nuanced measures of facility-level integration.

## Plain English Summary

Inadequate spacing between pregnancies can lead to adverse health outcomes among women and babies. The WHO recommends 24-months between pregnancies. Access to postpartum contraception is critical, yet many women wishing to space their pregnancies do not use it. Integrating family planning (FP) services into child immunization (CI) services may increase access to postpartum contraception. While previous research shows that integration is acceptable to women, evidence about its effects on service delivery and health outcomes is scarce and inconsistent. This is due, in part, to challenges surrounding the measurement of integration. One such challenge is that integration within health facilities is often measured as binary (i.e., a facility is designated as integrated or not), though numerous factors influence the extent of integration within facilities over time. Without capturing variation in integration, analyzing the effects of integration is difficult. This study addresses that challenge by developing Provider (i.e., nurse/midwife) and Facility Integration Indexes, which measure capacity to support integrated FP and CI services. Utilizing cross-sectional data collected for the Nigerian Urban Reproductive Health Initiative, we apply the indexes to describe the extent of integration across 400 facilities: 23% of facilities have low Provider Integration scores, 32% have medium, and 45% have high. Fourteen percent of facilities have low

Facility Integration scores, 38% have medium, and 48% have high. These results suggest that nuanced measures of integration, like those described in this paper, may enable more accurate evaluation of integration's effects, and provide more specific information about whether and how to support integration.

## Introduction

Nigeria has among the highest maternal and infant mortality rates in the world. In 2015, Nigeria had a maternal mortality ratio (MMR) of 814/100,000 live births, which was the fourth highest in the world (1). At the same time, Nigeria's infant mortality rate (IMR) was 67/1,000 live births (2). Nigeria signed on, and recently reconfirmed its resolve, to achieve Sustainable Development Goal 3 to reduce MMR to below 70/100,000 live births and to end preventable deaths of newborns and children under age 5 years by 2030 (3).

Family planning use has the potential to drastically reduce newborn and maternal deaths globally (4). Yet, unmet need for contraception remains high among Nigerian women; 24.8% of women of reproductive age (15-49 years) want to stop or delay childbearing but are not using contraception (5). The Federal Government of Nigeria set a target to increase the modern contraceptive prevalence rate from 10% in 2012 to 27% by 2020 (6). By 2018, the modern contraceptive prevalence rate among all women had increased to 13.8%, though this remains well below the target (5). Thus, increasing access to, and utilization of, family planning methods remains an urgent priority in Nigeria.

## Contraception Among Postpartum Women: Critical to Maternal and Infant Health

Postpartum contraceptive use is particularly critical to the health of women and babies. Interpregnancy intervals of less than 18 months are associated with increased risk of infant and maternal morbidity and mortality (7). Further, under-five mortality is significantly higher among children born less than 24-months after the preceding birth. One in 4 children in Nigeria are born less than two years after a sibling. The under-five mortality is 183/1000 among these children; this declines to 93/1000 among children born 3 years after the previous birth (2). The World Health Organization (WHO) recommends an interval of at least 24 months after a live birth prior to attempting the next pregnancy (8). In Nigeria, 65% of women who have given birth within the last 23 months may have an unmet need for contraception (9). Meeting this postpartum need for contraception would contribute substantially to contraceptive prevalence, which would in turn decrease MMR and IMR.

## Integration of Family Planning and Child Immunization Services

Integrating family planning into routine child immunization services has the potential to address unmet need for contraception among postpartum women by leveraging repeated immunization consultations to offer family planning information and services. Although modern contraceptive use in the postpartum period may be low in many contexts, immunization coverage is generally high. Immunization services may thus provide a solid platform for the integration of family planning services (10). In 2019, 85% of infants globally received the recommended 3 doses of the diphtheria-tetanus-pertussis (DPT-3) vaccine, although coverage in Nigeria is lower at around 50% (11, 12). While integration models vary in purpose and design, two primary models are commonly implemented (13, 14). The first, combined service provision, entails offering both family planning and immunization services on the same day at the same location. The second, single service provision plus referral, entails offering either family

planning or immunization services along with education, screening, and/or referral for the other service at a different place or time. Previous research has shown that integration can be feasible and acceptable to providers and clients (15-18). However, numerous factors and challenges influence the extent of integration attained within a facility, including health system characteristics, provider and client characteristics, staffing and space constraints, and the cultural context (19-21).

Within health facilities, family planning services may be integrated into child immunization services at the primary and secondary levels. The Nigerian Ministry of Health promotes the integration of family planning into immunization services as an important approach to increasing family planning availability and accessibility (22). The Minimum Standards for Primary Healthcare (MSPH) in Nigeria identify both family planning and immunization services as minimum components of primary healthcare and require provision of these services at all public primary healthcare facilities while advocating that privately owned facilities align with these standards (23). Both immunization and family planning services form part of a standard package of primary healthcare services commonly provided within hospitals.

## Evidence and Measurement Gaps

Policy and programming recommendations that promote integrating family planning and child immunization services are often based on insubstantial or conflicting evidence about its impact on health services delivery and patient outcomes (24-27). Given the scarcity of evidence and the resource and planning implications of integration, it is critical to develop context specific evidence that captures the nature, extent, and effects of integration in order to inform policy and program design (13).

The complexity of implementing integrated service delivery combined with the varied integration models and definitions prompt questions related to how integration and its effects should be measured (28-30). Many studies measure integration as a binary variable (a facility is integrated or not) rather than a multidimensional continuum that varies across time and place (17, 18, 28). A few studies have developed more nuanced measures of integration and assessed associations between these levels and health and service delivery outcomes. For example, the Integra Initiative developed indexes that measure the extent and type of HIV and reproductive health integration, and Church (2015) analyzed whether degree of HIV and reproductive health integration was associated with client demand for services and unmet need for family planning (20, 30). Mackenzie (2018) identified that the level of maternal, newborn, and child health and family planning integration varied widely across facilities as well as across clinical areas (31). Multi-dimensional, continuous measures of integration are valuable because they enable analyses related to the extent of integration within and across facilities and the effects of varying degrees of integration on service delivery and health outcomes. No studies, to our knowledge, have measured the extent of family planning and child immunization services integration (32, 33). This study aims to address this gap by developing integration indexes that quantify the extent of facility-level family planning and child immunization services integration as a varying and multifaceted outcome. Indexes like these may be used to analyze associations between the extent of integration and outcomes such as receipt of services, client satisfaction, and quality of care. The methodological process demonstrated in this study may be adapted by those aiming to better understand the extent of integration within facilities and over time.

## Methods

# Setting and Data Source

The data for this study were collected by the Measurement, Learning & Evaluation (MLE) project for the impact evaluation of the Nigerian Urban Reproductive Health Initiative (NURHI). NURHI, funded by the Bill & Melinda Gates Foundation, aimed to increase modern family planning use among the urban poor through a multi-pronged approach that included improving the quality of family planning services in high-volume urban health facilities.

The intervention supported contraceptive supply chains and logistics, training for family planning counseling and provision, and facility level management systems. This study utilizes health facility (N= 400, 58% hospitals, 42% primary healthcare centers) and healthcare provider (N= 1,479) baseline survey data that were collected in 2011 from Abuja (Nigeria's capital), Benin City, Ibadan, Ilorin, Kaduna and Zaria (34). NURHI selected these cities because they include both northern and southern regions of the country and each has a population of approximately, or more than, one million. The northern and southern regions of Nigeria differ in their cultural, economic, and religious characteristics; the north is poorer and predominantly Muslim while the south is more affluent and predominantly Christian.

## Study Sample

Two categories of healthcare facilities are included in the sample: high-volume facilities (HVF) (n = 112) and preferred-provider facilities (PPF) (n = 288). HVF and PPF can be either public or private, and they can be either primary or secondary facilities. The sample includes all public facilities in the study cities. HVF, generally the top service delivery sites by client load, offered both antenatal care and immunization services; these facilities served more than 1,000 antenatal clients per year. The NURHI program supported all HVF, and all of these facilities are included in the sample (34). PPF were identified from a baseline household survey conducted by MLE that contained a representative sample of 16,144 women aged 15-49. Women were asked the name of each facility where they went for family planning, maternal health, and child health services. Using this information, MLE created a list of facilities that women reported by study cluster (primary sampling unit). The most commonly mentioned facility in each primary sampling unit was categorized as a PPF. If the PPF was already included in the sample as a public facility or an HVF, then the second most-commonly mentioned facility was included. If the second most-commonly stated facility was already in the sample, no additional facility was included. Including the PPFs along with the public facilities and HVFs ensures that the sample includes facilities that women in these urban areas actually visit.

## Survey Instruments

This study utilizes instruments developed for the NURHI impact evaluation, which draws upon validated tools selected from the Quick Investigation of Quality (35). Facility and provider surveys were conducted in each facility by trained interviewers hired by Data Research and Mapping Corporation; the MLE project provided technical assistance for training of interviewers. The surveys collected information on the readiness of facilities and providers to offer integrated services, normal family planning service provision practices, gaps in commodities, equipment, training and resources, the extent of family planning integration into maternal, newborn and child health services, and other health facility characteristics. One facility audit was conducted per facility by asking questions of a manager or another administrator. In larger facilities, four providers were selected through simple random sampling to complete the provider survey; in facilities with four or fewer providers, all were approached for interview.

Providers eligible for inclusion were medically qualified to provide clinical services and assigned to provide direct family planning and/or maternal, newborn and child health services to clients at that facility; their responses are analyzed in this study.

## Statistical Method

This study employs Principal Components Analysis (PCA) to create two family planning and child immunization services integration indexes: a Provider Integration Index and a Facility Integration Index. All analyses were conducted using Stata version 13.1 (Stata Corp, LP, College Station, Texas).

## Constructing and Interpreting the Indexes

### *Selection of Variables for Inclusion in the PCA*

Drawing on Mayhew (2016), we posit that numerous characteristics and processes interact within a health facility to result in varying degrees of integrated service delivery (30). While the Nigerian Ministry of Health does not provide a specific definition of integrated family planning and immunization services, their 2008 National Guidelines for the Integration of Reproductive Health and HIV Programmes offers this explanation:

Integration in the health sector has been defined by offering two or more services at the same facility during the same operating hour, with the provider of one service actively encouraging clients to consider using the other services during the same visit, in order to make those services more convenient and efficient. Integrated services should be offered at the same point but where that is not possible, strong referral systems are required to ensure that clients receive high quality service (36).

NURHI's Strategy for Integrating Family Planning into Maternal, Newborn, Child Health and HIV/AIDS Services references this guidance (37). This study also refers to this guidance to inform the attributes measured in the indexes. Additionally, we reviewed the integration literature to identify facility-level attributes that support service integration. Several critical attributes emerged, including a) facility norms that support concurrent service provision (e.g., operational management standards and procedures that support the availability of both child immunization and family planning services at the same consultation or on the same day), and b) provider capacity to offer multiple services (e.g., provider(s) has the skills and willingness to offer family planning information or services during a child immunization visit) (26, 30, 38-41).

Because this study was conceptualized after data collection, we leveraged the available data and selected eight indicators for inclusion in the indexes (Table 1). Table 2 describes these and other facility characteristics. A few of these indicators warrant additional explanation. The indicators used to develop the integration indexes focus primarily on family planning information and services that are provided during the child immunization visits. The indexes are thus most appropriate for use within that context, though they do also measure a range of referral scenarios. Improving outcomes through integration relies upon both high coverage and quality of integrated services. A substantial body of research links higher quality family planning services with increased contraceptive adoption, prevalence, and continuation; poor family planning service quality can hinder use (42). Therefore, the

level of quality provided and the absence of barriers that limit coverage and quality are essential indicators of effective integration (40, 43). We analyzed quality of integrated family planning services by measuring the range and breadth of family planning topics that providers discuss with a client during child health service visits.

Because the extent of integration can be influenced by provider bias (44-46), we include social norm-based service barriers by measuring the extent to which providers at a facility require spousal consent prior to provision of a family planning method during an integrated visit. In Nigeria, spousal consent is a tenacious barrier to contraceptive use that may be mitigated by training providers that standard service provision guidelines do not include a requirement for spousal consent, providing supportive supervision on guideline implementation, and utilizing more comprehensive behavior change approaches with providers (47). While numerous such barriers exist and could have been employed in the indexes this is the only variable in our dataset that captures such barriers to family planning specifically during immunization visits.

Several variables refer to child immunization, child growth monitoring, or child health service visits. Child health services visits include either immunization or growth monitoring visits, but not sick child visits. In variables referring to child health services, it was not possible to differentiate data pertaining only to child immunizations from data pertaining only to child growth monitoring. However, child immunization visits comprise the vast majority of all child health services visits. In the concurrent health facility client exit interview, 1,714 attended the facility for a child health service visit. Of these, only 90 (5%) report that child growth monitoring was the primary purpose of their visit while the remaining 95% reported immunization as the primary purpose of their visit. Facility-level variables are based on a summary of provider responses. Means were imputed for missing data.

## PCA Application

PCA was applied following the selection and transformation of variables. Input variables were standardized to a mean of zero and a standard deviation of one prior to the analysis to prevent variables with greater variance from dominating each component. The Kaiser-Meyer-Olkin (KMO) test of sampling adequacy was used to ascertain the suitability of the data for use in a PCA. Our KMO test yielded a score of 0.8, indicating sampling adequacy for each variable and the complete model. Based on evaluation of the eigenvalues (Table 3) and the scree plot (Figure 1) we retained two components. The factor loading scores (see factor loadings column in Table 4), which show the correlation coefficient between each variable and component, were examined to determine which dimensions of integration are represented by the components. The scores confirmed the anticipated dimensions: provider integration and facility integration.

## Creating the Indexes

We constructed the Provider Integration Index and Facility Integration Index using weights calculated for each of the variables by dividing its factor loading by the sum of the factor loadings of all variables in that component (see weights column in Table 4). Next, we multiplied the variables included in each component by their associated weights and summed the values. Finally, we calculated the Provider Integration Index score and the Facility Integration Index score for each facility by multiplying these values by ten. The indexes thus range in value from zero to ten, with a higher score indicating a higher level of integration. A sensitivity analysis was conducted to identify the effects of excluding from the sample those facilities that do not offer child immunization (n=61); there were no statistically significant differences between the indexes that include all facilities versus those with the restricted set of facilities. We retained these facilities in the sample because one goal of the paper is to assess

integration across the range of facilities and circumstances represented by our sample. Excluding these facilities would prevent us from knowing the full extent of integration across our sample. Additionally, one key benefit of developing these indexes is the ability to apply them to understand the effects of integration on health and service delivery outcomes. Having a score for facilities that do not offer child immunization allows future research to better identify correlations between level of integration (even very low level) and other outcomes.

As a robustness check, we also created the indexes using Exploratory Factor Analysis. The indexes created using EFA correlate strongly with the indexes created using PCA (Spearman rank correlation 0.99 for the Provider Indexes and 0.89 for the Facility Indexes), indicating that the results are robust to the use of either method.

## Results

Eighty-five percent of facilities offer either child immunization or child growth monitoring services while 77% of facilities provide both family planning and child immunization services (Table 2). Private facilities are less likely than public facilities to provide both child immunization and family planning services; 93% of public facilities provide both child immunization and family planning services, while 66% of private facilities offer both services. Among the 93 facilities (81= private, 12= public) that do not offer both child immunization and family planning services, it is generally child immunization services that are not offered. Among the private facilities that do not offer both child immunization and family planning services, 89% do not offer child immunization services while 9% do not offer family planning services. Among the public facilities that do not offer both child immunization and family planning services, 50% do not offer child immunization services while 50% do not offer family planning services (data not shown). Among facilities that provide family planning services, 84% offer them either five or seven days a week. Immunization provision is less frequent; 51% of the facilities that provide them do so once or twice per week while 26% provide them every day. On average, 58% of providers in each facility report offering family planning information during child health visits while 56% of providers in each facility have been trained to provide both child immunization and family planning services. Providers, on average, addressed fewer than two of the seven elements of family planning service provision during child health visits. Fifty percent of providers report requesting partner consent prior to providing family planning at a child health visit, though this varies by method.

Index scores vary across facilities (Figures 2 and 3). Table 4 provides Provider and Facility Integration Index scores and their classifications. Please refer to the “Creating the Indexes” section for a description of how values in Table 4 were calculated and how these values link to the index scores. The mean Provider Integration Index score is 5.4 (standard deviation: 3.1, range: 0 - 10) and the mean Facility Integration Index score is 6.2 (standard deviation: 2.7, range: 0 - 9.9) (Table 4). Each facility was classified as having “low integration” (index score 0 - 3.29), “medium integration” (3.30 - 6.59) or “high integration” (6.60 -10.00). These classifications were determined by dividing raw scores equally into thirds along the score continuum of zero to ten. Twenty-three percent of facilities are classified as having low Provider Integration Index scores, 32% are classified as having medium scores and 45% are classified as having high scores. Mean Provider Integration Index scores are 0.75, 4.99, and 8.23 for the low, medium, and high groups respectively. Fourteen percent of facilities are classified as having low Facility Integration Index scores, 38% have medium scores and 48% have high scores. The mean Facility Integration Index scores are 0.50, 6.00, and 8.10 for the low, medium, and high groups respectively. The mean value of each component within each classification group is presented within the Integration Index Score Classification Group Means columns. For example, on average, 4% of providers in facilities classified as having low integration index scores routinely offer family planning information during child immunization visits, 49% of providers in facilities with medium scores do, and 93% of providers in facilities with high integration index scores do.

# Index Coherence and Robustness

Following Filmer and Pritchett (2001), we examined the internal coherence of the indexes by comparing facility characteristics and index scores across low (0-3.29), medium (3.30-6.59), and high integration groups (6.60-10) (48). We note significant differences in facility characteristics and overall index scores across all groups for both indexes, meaning that the integration variables were significantly different between the low and medium score groups and the medium and high score groups. These differences indicate that both indexes have strong internal coherence (see p-values in Table 4).

We assessed the robustness of the indexes by examining how the classifications of facilities having high Integration Index scores changed when different sub-sets of variables were entered into the PCA (48). To assess the robustness of the Provider Integration Index, we ran 6 variations of the PCA. The first variation (“base case”) included all variables. Each subsequent model omitted one of the provider integration variables. Similarly, to assess the robustness of the Facility Integration Index, we ran 4 variations of the PCA. The first variation (the “base case”) included all variables. Each subsequent model omitted one of the facility integration variables. Looking only at the sites classified as “high integration” in the base case, we examine the impact on classification when we omit one variable at a time from the PCA. Table 5 shows the percentage of facilities classified as “high integration” in the base case compared with the percentage of facilities classified as “high integration” in the subsequent models. The Provider Integration Index is highly robust to the inclusion of different sub-sets of variables in the model. This is evidenced by the very similar classification results across models.

Table 6 presents the same results for the Facility Integration Index. Unlike the Provider Integration Index, the Facility Integration Index shows considerable sensitivity to the different sub-sets of variables included in the PCA models. When the variables indicating normal practices at the facility are omitted, the Facility Integration Index score shifts towards the mean value of the score of operational days when both child immunization and family planning services are offered. Based on these assessments, we retained all base case variables in the model in order to reflect more characteristics of provider and facility integration (49).

## Discussion

The purpose of this paper is to demonstrate the development of novel indexes that may be adapted and applied to measure integration along a continuum, and to apply these indexes to describe the extent of family planning and child immunization integration across a large sample of primary and secondary health facilities. The index scores allow for individual facility scoring and ranking; the components within the indexes enable even more nuanced analyses to inform policy and program strengthening. These integration indexes may be adapted and employed to enrich understanding of integration levels within and across health facilities, identify facilities for intervention, inform and monitor the effectiveness of interventions, and investigate the effect of integration on a range of service delivery and health outcomes. The NURHI survey instruments are publicly available and provide a valuable reference for the adaptation or development of indicators for future programs and research (50). Some widely used surveys offer the opportunity to adapt this methodology to analyze integration of family planning into other services. For example, the DHS Program’s Service Provision Assessment (SPA) and the PMA surveys include indicators that assess service availability, provider training and/or scope of practice, and client receipt of family planning information and services during other consultations, though not immunization services. Future revisions of these surveys may offer the opportunity to include questions related to immunization services, which could facilitate analyses about the integration of family planning and child immunization services. With these other

survey tools, a similar non-binary index of integration can be created to assess family planning integration into other reproductive health services.

Our results suggest that facility norms and provider capacity to support integrated family planning and child immunization service delivery varies considerably across our sample. The identification of two distinct dimensions of integration, the heterogeneity of the scores and the substantial percentage of facilities within each integration index classification level, suggests that measuring integration as a binary variable does not reflect the true variation in integration within and across health facilities in urban areas of Nigeria. This is consistent with the findings of Mayhew (2016) and Pfitzer (2020) and underscores the importance of employing ordinal or continuous measures of integration in research that examines the determinants and effects of integration.

The development of the integration indexes required us to take a detailed look at provider and facility characteristics in each index. Several characteristics warrant particular attention. First, the majority of facilities with low Provider Integration Index scores (n= 94) do not provide both child immunization and family planning services, which is essential to integration. While we considered excluding these facilities from the sample, we opted to retain them because doing so enables description of the full spectrum of integration across our sample. Additionally, one benefit of these indexes is the ability to apply them to analyze the evolution of integration and its effects on a variety of outcomes. Having the complete range of scores allows future research to track integration in all of the facilities and identify correlations between level of integration (even very low level) and other outcomes. This is particularly important when considering that low levels of integration may impart some benefits, such as increased privacy, that may positively influence outcomes. Lastly, we retained these facilities in the sample because the Minimum Standards for Primary Healthcare in Nigeria stipulate that primary healthcare centers should provide both services, and both services form part of a standard package of primary health services provided by hospitals. Our results show that it is primarily private facilities that do not provide both family planning and immunization services. In Nigeria, routine immunization has been funded primarily by the government and Gavi (51). Private facilities may not qualify for either source of funding and may thus be less likely to offer immunization services. The private sector provides a substantial amount of health services in developing countries, including in urban areas of Nigeria (52). Thus, it is important to understand its capacity to provide integrated services and consider whether and how private sector facilities might implement integration of family planning into other services. Secondly, the median score of the operational days when both child immunization and family planning services are offered (median = 0.14, interquartile range = 0.38 [0.02-0.40]) indicates that many facilities do not offer integrated services on a daily basis. Immunization services are provided on fewer days per week than family planning services, likely due to logistic issues such as reliance upon cold boxes that are delivered on specific days. Policies and programs supporting the integration of family planning and child immunization services could ensure that staffing and management procedures enable a full range of high-quality family planning services on immunization days.

Another factor we observed is the high percentage of providers requiring partner consent prior to providing a method of contraception. Our results show that 50% of providers report requiring partner consent during integrated visits; the percentage varies by contraceptive method selected. This substantial barrier to family planning service during integrated visits warrants additional investigation. Finally, it is important to consider the quality of family planning services when they are integrated into immunization services. Our results indicate that providers discuss a limited range of information with women during integrated visits. While provision of family planning information and services within the immunization setting may offer benefits, such as decreased need for repeat visits, it also presents challenges, such as potentially insufficient staffing, consultation time, and privacy to allow high quality

information provision and discussion about women's reproductive goals and needs (28). This is an important consideration for the design of integrated service provision; integration approaches that establish easily accessible and efficient linkages between immunization and family planning services may prove more beneficial and be more likely to offer women a full range of methods than those which offer limited family planning information and services within the immunization setting. This is an important area for continuing research. Policies and programs promoting integration should ensure sufficient support to individual providers and facility systems to ensure that integration approaches enhance family planning service quality.

This study has several limitations. First, while the Nigerian Ministry of Health promotes the integration of family planning and child immunization services, it does not provide a precise definition of integration. We therefore relied upon Nigeria's 2008 National Guidelines for the Integration of Reproductive Health and HIV Programmes (this, to our knowledge, is the most detailed and comprehensive definition of integration promoted by the Nigerian Ministry of Health), the NURHI integration strategy, and the broader integration literature to shape the attributes included in the indexes. This issue has challenged previous research; the lack of precise definitions of integration has long complicated its measurement (53). Also, because this was a secondary analysis, we could not get precise information about some factors that are important to integration, such as service overlap. While we know the number of days per week immunization and family planning services are offered, we do not know *which* days of the week, so although family planning services are provided 5-7 days per week in the majority (approximately 85%) of facilities that provide family planning services, it is possible that we misrepresented the overlap in some instances (e.g., a facility that offers immunization on Monday and family planning on Monday, Wednesday and Friday would score the same as a facility that offers immunization on Tuesday and family planning on Monday, Wednesday and Friday). Future research could address this by collecting more specific information about the days and hours when these services are offered. Similarly, while we assume that providers are aware of normal practices in the facility, it is possible that they lack completely accurate knowledge. Either of these situations may lead to inaccuracies that would affect the index scores. Additionally, the provider and facility survey responses may be subject to social desirability bias. Future studies may address this issue by triangulating information between clients and providers. Our sampling strategy ensured inclusion of public facilities, which was NURHI's focus, as well as private facilities that women actually attended. This provided an opportunity to collect relevant information about a variety of well-attended facilities across the six cities. However, the high number of private facilities precluded a census. Thus, our results are specific to our sample and are not generalizable to all urban facilities in Nigeria or other contexts. For example, because of NURHI's focus on high-volume facilities, the sample includes 58% hospitals and thus over represents secondary facilities, which comprise 12% of total health facilities in Nigeria (54). Last, this data was collected in 2011 and our results should be interpreted as a description of the extent of integration within these facilities at the time of data collection. Indeed, whether, how, and why the extent of integration changes within facilities over time is an important area for future research.

Notwithstanding these limitations, this research advances the conversation about how to measure integration by describing the development of more nuanced measures of integration that identify facility and provider attributes that support integrated service delivery. Such indexes are valuable because they allow for more detailed measurement of the determinants and effects of integration over time. This research also describes the nature and extent of family planning and child immunization services integration in six cities of Nigeria. This is important information in light of the Nigerian government's goal to reduce MMR and IMR by increasing contraceptive prevalence, in part, by reaching more postpartum women through integration of family planning and child immunization services.

## Conclusion

Integration of family planning and immunization services is complex and challenging, and evidence about its effects on service delivery and health outcomes has been inconsistent. Previous research on the integration of family planning and child immunization has designated health facilities as integrated or not based on whether the facility received an intervention intended to increase integration. This study takes a novel approach by developing indexes that offer continuous measures of facility-level family planning and child immunization services integration and using the indexes to identify the extent of integration within a sample of 400 health facilities in six urban sites of Nigeria. Measuring the degree of integration is valuable as a means of monitoring health facility and system performance. Over time, such measurement will enable clearer understanding of the extent, patterns and adoption of integration as well the effect of programs on affecting integration on service delivery and health outcomes.

This research underscores the need for policies and programs seeking to promote or improve integration to start with a clear, context-specific definition and approach that recognizes the dual provider and facility dimensions. The definition should align with specific Ministry of Health and program objectives and strategies and should be captured in data collection initiatives from the outset. While the method outlined in this paper is most appropriate for programs that include a large number of facilities and have the necessary data collection and analytic expertise, smaller programs may incorporate some of the approaches highlighted in this paper. Such programs can articulate a precise and context-specific definition of integration and ensure that project implementation and routine monitoring and evaluation activities capture this definition with a variety of measures. Future research should analyze the determinants of integration and the effects of varying degrees of integration on service delivery and health outcomes. While this research focuses specifically on service delivery within health facilities, further research should examine how health systems pillars such as governance, human resources, logistics, financing, and information management systems influence integrated service delivery. A more robust evidence base is essential to the development of integration policies and programs that will increase contraceptive prevalence among postpartum women to promote healthy birth spacing and, ultimately, reduce maternal and infant morbidity and mortality globally.

## Abbreviations

AIDS: Acquired Immune Deficiency Syndrome; CGM: Child Growth Monitoring; CI: Child Immunization; EFA: Exploratory Factor Analysis; FP: Family Planning; HIV: Human Immunodeficiency Syndrome; HVF: High-volume Facility; IMR: Infant Mortality Rate; KMO: Kaiser-Meyer-Olkin; MLE: Measurement, Learning & Evaluation; MMR: Maternal Mortality Rate; MSPH: Minimum Standards for Primary Healthcare; NURHI: Nigerian Urban Reproductive Health Initiative; PCA: Principal Components Analysis; PPF: Preferred Provider Facility; SD: Standard Deviation; WHO: World Health Organization

## Declarations

### **Ethics approval and consent to participate:**

The study protocol and all consent procedures and consent forms were approved by the Institutional Review Board at the University of North Carolina at Chapel Hill and by the National Health Research Ethics Committee of Nigeria in Nigeria. All facility respondents were asked to consent to participate in the study. Prior to being surveyed, health providers were asked to sign consent forms that included details on the purpose of the study, potential benefits and

potential risk, and clarified that the information provided would not be identifiable. Facility administrators who were asked questions about the services offered at the facility through a facility audit were asked for verbal consent to participate.

**Consent for publication:**

Not applicable

**Data availability:**

Data from this study and all documentation are available upon request through the MLE Dataverse website at: <https://dataverse.unc.edu/dataverse/mle>.

**Competing Interests:**

The authors declare that they have no competing interests.

**Funding:**

The Bill & Melinda Gates Foundation provided funding for the data collection of the data used in this study. The Bill & Melinda Gates Foundation did not play a role in the design of this study or in the analysis or interpretation of data.

**Authors' contributions:**

KS led the design and implementation of research, data analysis, interpretation of results and manuscript writing. JOM contributed to data analysis and interpretation of results. IS provided critical contextual, theoretical, and methodological guidance and expertise. SC provided theoretical and methodological expertise. MW and JP contributed to conceptualization and presentation of the research. AVB contributed to conceptualization and presentation of the research and provided overall guidance and direction. All authors provided critical feedback and helped shape the research, interpretation of results and manuscript. All authors read and approved the final manuscript.

**Acknowledgements:**

Not applicable

## References

1. World Health Organization. World health statistics 2016: monitoring health for the SDGs, sustainable development goals. Geneva: WHO Press; 2016.
2. National Population Commission (NPC) and ICF. Nigeria Demographic and Health Survey 2018. Abuja and Rockville, Maryland: NPC; 2019.
3. Government of the Federal Republic of Nigeria. Integration of the SDGs into National Development Planning: A Second Voluntary Review. Abuja; 2020.

4. Bhutta ZA, Das JK, Bahl R, Lawn JE, Salam RA, Paul VK, et al. Can available interventions end preventable deaths in mothers, newborn babies, and stillbirths, and at what cost? *Lancet* (London, England). 2014;384(9940):347-70.
5. Family Planning 2020. Nigeria 2018 Core Indicators 1-9 Country Fact Sheet 2018 Available from: <http://www.familyplanning2020.org/sites/default/files/Nigeria%202018%201-9%20Handout.pdf>
6. Nigeria Family Planning 2020 Commitment Update [press release]. 2017.
7. Conde-Agudelo A, Rosas-Bermudez A, Kafury-Goeta AC. Birth spacing and risk of adverse perinatal outcomes: a meta-analysis. *Jama*. 2006;295(15):1809-23.
8. World Health Organization. Report of a WHO Technical Consultation on Birth Spacing. Geneva: World Health Organisation; 2006.
9. Moore Z, Pfitzer A, Gubin R, Charurat E, Elliott L, Croft T. Missed opportunities for family planning: an analysis of pregnancy risk and contraceptive method use among postpartum women in 21 low- and middle-income countries. *Contraception*. 2015;92(1):31-9.
10. World Health Organization. Working together: an integration resource guide for immunization services throughout the life course. Geneva: 2018.
11. World Health Organization. Global Immunization Coverage 1980 to 2019. 2019. Available from: <https://www.who.int/data/gho/data/themes/immunization>.
12. World Health Organization. WHO Vaccine Preventable Diseases 2020 Nigeria Summary 2020 Available from: [https://www.who.int/immunization/monitoring\\_surveillance/data/nga.pdf](https://www.who.int/immunization/monitoring_surveillance/data/nga.pdf)
13. Wallace AS, Ryman TK, Dietz V. Experiences integrating delivery of maternal and child health services with childhood immunization programs: systematic review update. *The Journal of infectious diseases*. 2012;205 Suppl 1:S6-19.
14. High-Impact Practices in Family Planning (HIP). Family Planning and Immunization Integration: Reaching postpartum women with family planning services. July 2013: United States Agency for International Development; 2013.
15. Cooper CM, Fields R, Mazzeo CI, Taylor N, Pfitzer A, Momolu M, et al. Successful proof of concept of family planning and immunization integration in Liberia. *Global health, science and practice*. 2015;3(1):71-84.
16. Dulli L, Green M, Katz K. Increasing postpartum family planning services in Madagascar: Assessing the feasibility and acceptability of immunization services as an entry point to family planning. 2010.
17. Dulli LS, Eichleay M, Rademacher K, Sortijas S, Nsengiyumva T. Meeting Postpartum Women's Family Planning Needs Through Integrated Family Planning and Immunization Services: Results of a Cluster-Randomized Controlled Trial in Rwanda. *Global health, science and practice*. 2016;4(1):73-86.
18. Huntington D, Aplogan A. The integration of family planning and childhood immunization services in Togo. *Studies in family planning*. 1994;25(3):176-83.

19. Cleland J, Shah IH, Daniele M. Interventions to Improve Postpartum Family Planning in Low- and Middle-Income Countries: Program Implications and Research Priorities. *Studies in family planning*. 2015;46(4):423-41.
20. Church K, Wringe A, Lewin S, Ploubidis GB, Fakudze P, Mayhew SH. Exploring the Feasibility of Service Integration in a Low-Income Setting: A Mixed Methods Investigation into Different Models of Reproductive Health and HIV Care in Swaziland. *PloS one*. 2015;10(5):e0126144.
21. Ahumuza SE, Rujumba J, Nkoyooyo A, Byaruhanga R, Wanyenze RK. Challenges encountered in providing integrated HIV, antenatal and postnatal care services: a case study of Katakwi and Mubende districts in Uganda. *Reproductive health*. 2016;13:41.
22. Nigerian Federal Ministry of Health. Nigeria Family Planning Blueprint (Scale Up Plan). In: Health NFMo, editor. Abuja:
23. Nigerian National Primary Healthcare Development Agency. Minimum Standards for Primary Healthcare in Nigeria. Abuja: 2012.
24. Kuhlmann AS, Gavin L, Galavotti C. The integration of family planning with other health services: a literature review. *International perspectives on sexual and reproductive health*. 2010;36(4):189-96.
25. Speizer IS, Fotso JC, Okigbo C, Faye CM, Seck C. Influence of integrated services on postpartum family planning use: a cross-sectional survey from urban Senegal. *BMC public health*. 2013;13:752.
26. Vance G, Janowitz B, Chen M, Boyer B, Kasonde P, Asare G, et al. Integrating family planning messages into immunization services: a cluster-randomized trial in Ghana and Zambia. *Health policy and planning*. 2014;29(3):359-66.
27. Blazer C, Prata N. Postpartum family planning: current evidence on successful interventions. *Open Access J Contracept*. 2016;7:53-67.
28. Nelson AR, Cooper CM, Kamara S, Taylor ND, Zikeh T, Kanneh-Kesselly C, et al. Operationalizing Integrated Immunization and Family Planning Services in Rural Liberia: Lessons Learned From Evaluating Service Quality and Utilization. *Global health, science and practice*. 2019;7(3):418-34.
29. Mounier-Jack S, Mayhew SH, Mays N. Integrated care: learning between high-income, and low- and middle-income country health systems. *Health policy and planning*. 2017;32(suppl\_4):iv6-iv12.
30. Mayhew SH, Ploubidis GB, Sloggett A, Church K, Obure CD, Birdthistle I, et al. Innovation in Evaluating the Impact of Integrated Service-Delivery: The Integra Indexes of HIV and Reproductive Health Integration. *PloS one*. 2016;11(1):e0146694.
31. Baumgartner JN, Green M, Weaver MA, Mpangile G, Kohi TW, Mujaya SN, et al. Integrating family planning services into HIV care and treatment clinics in Tanzania: evaluation of a facilitated referral model. *Health policy and planning*. 2014;29(5):570-9.
32. Hoang T, Goetz MB, Yano EM, Rossman B, Anaya HD, Knapp H, et al. The impact of integrated HIV care on patient health outcomes. *Medical care*. 2009;47(5):560-7.

33. Mattocks KM, Kroll-Desrosiers A, Kinney R, Singer S. Understanding Maternity Care Coordination for Women Veterans Using an Integrated Care Model Approach. *J Gen Intern Med.* 2019;34(Suppl 1):50-7.
34. MLE N, & National Population Commission 2010-2011 Nigeria Baseline Survey for the Urban Reproductive Health Initiative. JHU CPC, USA; Abuja, Nigeria: Measurement, Learning and Evaluation (MLE) [UNC, USA; Abuja, Nigeria]; Nigerian Urban Reproductive Health Initiative(NURHI) [JHU CCP,USA;Abuja, Nigeria]; and National Population Commission [Abuja, Nigeria]; 2011.
35. MEASURE Evaluation. Quick investigation of quality (QIQ): a user's guide for monitoring quality of care in family planning. Chapel Hill, North Carolina: MEASURE Evaluation, University of North Carolina; 2016.
36. Nigeria Federal Ministry of Health. National Guidelines for the Integration of Reproductive Health and HIV Programmes in Nigeria. Abuja: 2008.
37. Strategy for Integrating Family Planning into Maternal, Newborn, Child Health and HIV Services Abuja: 2009 <https://www.nurhitoolkit.org/document-library/fp-integration-strategy#.X9fddC9h1R4>
38. Dudley L, Garner P. Strategies for integrating primary health services in low- and middle-income countries at the point of delivery. *The Cochrane database of systematic reviews.* 2011(7):Cd003318.
39. Watt N, Sigfrid L, Legido-Quigley H, Hogarth S, Maimaris W, Otero-Garcia L, et al. Health systems facilitators and barriers to the integration of HIV and chronic disease services: a systematic review. *Health policy and planning.* 2017;32(suppl\_4):iv13-iv26.
40. Kerber KJ, de Graft-Johnson JE, Bhutta ZA, Okong P, Starrs A, Lawn JE. Continuum of care for maternal, newborn, and child health: from slogan to service delivery. *Lancet (London, England).* 2007;370(9595):1358-69.
41. Mayhew SH, Sweeney S, Warren CE, Collumbien M, Ndwiga C, Mutemwa R, et al. Numbers, systems, people: how interactions influence integration. Insights from case studies of HIV and reproductive health services delivery in Kenya. *Health policy and planning.* 2017;32(suppl\_4):iv67-iv81.
42. RamaRao S, et al. The link between quality of care and contraceptive use. *International Family Planning Perspectives.* 2003;29(2):76-83.
43. Curry N., Ham C. Clinical and service integration: the route to improved outcomes. London: The King's Fund; 2010.
44. de Jongh TE, Gurol-Urganci I, Allen E, Jiayue Zhu N, Atun R. Barriers and enablers to integrating maternal and child health services to antenatal care in low and middle income countries. *BJOG : an international journal of obstetrics and gynaecology.* 2016;123(4):549-57.
45. Dynes MM, Bernstein E, Morof D, Kelly L, Ruiz A, Mongo W, et al. Client and provider factors associated with integration of family planning services among maternal and reproductive health clients in Kigoma Region, Tanzania: a cross-sectional study, April-July 2016. *Reproductive health.* 2018;15(1):152.
46. Pfitzer A, Maly C, Tappis H, Kabue M, Mackenzie D, Healy S, et al. Characteristics of successful integrated family planning and maternal and child health services: Findings from a mixed-method, descriptive evaluation. *F1000Res.* 2019;8:229.

47. Solo J, Festin M. Provider Bias in Family Planning Services: A Review of Its Meaning and Manifestations. *Global health, science and practice*. 2019;7(3):371-85.
48. Filmer D, Pritchett LH. Estimating wealth effects without expenditure data—or tears: an application to educational enrollments in states of India. *Demography*. 2001;38(1):115-32.
49. Vyas S, Kumaranayake L. Constructing socio-economic status indices: how to use principal components analysis. *Health policy and planning*. 2006;21(6):459-68.
50. Measurement, Learning, and Evaluation Project, NURHI, & Data Research and Mapping Consult. Report of the Baseline Facility Survey for the Nigerian Urban Reproductive Health Initiative (NURHI). 2011. Available from: <http://www.nurhitoolkit.org/document-library/nurhi-baseline-facility-report#.WYx9LIKZN8c>
51. Olumide F, Chidiabere O, Akinyede O, Botchway P, Soyemi K. Vaccine financing in Nigeria: are we making progress towards self-financing/sustenance? *The Pan African Medical Journal*. 2017;27(Suppl 3).
52. Levy JK, Curtis S, Zimmer C, Speizer IS. Assessing gaps and poverty-related inequalities in the public and private sector family planning supply environment of urban Nigeria. *Journal of Urban Health*. 2014;91(1):186-210.
53. Atun R, de Jongh T, Secci F, Ohiri K, Adeyi O. A systematic review of the evidence on integration of targeted health interventions into health systems. *Health policy and planning*. 2010;25(1):1-14.
54. Makinde OA, Sule A, Ayankogbe O, Boone D. Distribution of health facilities in Nigeria: implications and options for universal health coverage. *The International journal of health planning and management*. 2018;33(4):e1179-e92.

## Tables

**Table 1: Description of PCA Input Variables**

| <b><i>Input Variable Description</i></b>   | <b><i>Type</i></b>  | <b><i>Survey Source</i></b> |
|--|---|-----------------------------|
| <i>What proportion of providers at facility offer both CI and FP services?*</i>  | Continuous between 0 and 1  | Provider                    |
| <i>What proportion of providers at facility routinely offers FP information during CI or CGM visits*?</i>  | Continuous between 0 and 1  | Provider                    |
| <i>What is the average count of FP items that providers at facility tell client during CHS visits?***</i>  | Ordinal between 0 and 7<br>FP items include: 1) Identify reproductive goals 2) Provide information about different FP methods 3) Discuss the client's FP preferences 4) Help women select a suitable method 5) Educate women to use the selected method 6) Explain side effects 7) Explain specific medical reasons to return   | Provider                    |
| <i>What proportion of providers at facility do not request partner consent prior to woman's receipt of FP services during CHS visit?*</i>  | Continuous between 0 and 1  | Provider                    |
| <i>Does the facility provide both child immunization and family planning services?</i>   | Binary (0= no, 1= yes)  | Facility                    |
| <i>What is the normal practice at this facility if client wants FP information during CHS visit?***</i>  | Ordinal between 0 and 7<br>Responses include: 0) Facility does not provide child health services 1) Facility does provide child health services but does not provide family planning services 2) Client is given no information or referral 3) Client is given referral to another facility 4) No appointment made, client told to return on a different day 5) Appointment made for different day 6) Client sometimes receives information on same day 7) Client always receives information on same day | Facility                    |
| <i>What is the normal practice at this facility if client wants hormonal method of FP during CHS visit?***</i>   | Ordinal between 0 and 7<br>Responses include: 0) Facility does not provide child health services 1) Facility does provide child health services but does not provide family planning services 2) Client is given no information or referral 3) Client is given referral to another facility 4) No appointment made, client told to return on a different day 5) Appointment made for different day 6) Client sometimes receives method on same day 7) Client always receives method on same day           | Facility                    |
| <i>What is the score of operational days when both CI and FP services are offered?****</i>   | Continuous between 0 and 1<br>Defined as: (Proportion of operational days that child immunization services are provided) multiplied by (Proportion of operational days that family planning services are provided)  | Facility                    |
| <p><i>Notes: CI: Child Immunization FP: Family Planning CGM: Child Growth Monitoring CHS: Child Health Service CHS visits include either CI or CGM visits, but not sick child visits. In variables referring to CHS visits, it was not possible to differentiate data pertaining only to CI from data pertaining only to CGM. Client exit interviews show that CI visits comprise the 95% of all CHS visits. *Proportion of providers was obtained from the provider survey by taking an average of provider responses to dichotomous survey questions (0= no, 1= yes). For example, if two providers responded that they did not routinely offer FP information during a CI or CGM visit and two responded that they did then the facility would score a 0.5 on this item. ** Facility score calculated by adding one point for affirmative responses to items 1-7. ***Facility scores reflect the response, ranked from 0 (low) to 7 (high).**** The score for a facility that is open 7 days/week and offers immunization 1 day/week and family planning services 7 days/week would be calculated: <math>(1/7) \times (7/7) = 0.14</math></i></p> |   |                             |

**Table 2: Facility Characteristics**

| Characteristic   | Total (N=400), n (%) or Mean Value (SD) |
|--|---|
| <i>Facility Type</i>   |   |
| High Volume  | 112 (28)                                |
| Preferred Provider   | 288 (72)                                |
| <i>Ownership</i>   |   |
| Publicly Owned   | 164 (41)                                |
| Privately Owned  | 236 (59)                                |
| <i>Level</i>   |   |
| Primary  | 166 (42)                                |
| Secondary  | 234 (58)                                |
| <i>Location</i>  |   |
| Abuja  | 48 (12)                                 |
| Benin  | 71 (18)                                 |
| Ibadan   | 62 (15)                                 |
| Ilorin   | 72 (18)                                 |
| Kaduna   | 92 (23)                                 |
| Zaria  | 55 (14)                                 |
| Facilities that offer CI and/or CGM  | 339 (85)                                |
| Facilities that provide CI and FP services   | 307 (77)                                |
| Normal practice if FP info wanted during CH visit  | 5.82<br>(2.23)                          |
| Normal practice if hormonal FP wanted during CH visit  | 4.63<br>(1.97)                          |
| Score of days where both CI and FP are offered*  | 0.14<br>(0.38, 0.02-0.4)                |
| Proportion providers at facility who offer CI and at least 1 modern FP method  | 0.56<br>(0.40)                          |
| Proportion providers at facility who routinely offer FP info during CI/CGM   | 0.58<br>(0.39)                          |
| Average FP items that a provider at a facility tells client during CHS   | 1.67<br>(1.52)                          |
| Proportion providers at facility that do not request partner consent for FP during CHS   | 0.50<br>(0.37)                          |
| <i>Notes:</i> See Table 1 for complete variable definitions. CI: Child Immunization FP: Family Planning CGM: Child Growth Monitoring CHS: Child Health Service * The median and interquartile range are presented. |   |

**Table 3: Main PCA Results from Analysis of Health Facility Data from Six Cities in Nigeria**

| Component | Eigenvalue | Proportion of Explained Variance | Proportion of Cumulative Explained Variance |
|-----------|------------|----------------------------------|---|
| Comp1     | 4.456      | 0.557                            | 0.557                                       |
| Comp2     | 1.532      | 0.191                            | 0.748                                       |
| Comp3     | 0.795      | 0.099                            | 0.848                                       |
| Comp4     | 0.450      | 0.056                            | 0.904                                       |
| Comp5     | 0.315      | 0.039                            | 0.943                                       |
| Comp6     | 0.256      | 0.032                            | 0.975                                       |
| Comp7     | 0.107      | 0.013                            | 0.989                                       |
| Comp8     | 0.089      | 0.011                            | 0.999                                       |

**Table 4: Provider and Facility Integration Index Scores and Variable Means by Group**

| Integration Variable Description   | Factor Loading | Sum of Factor Loadings | Weight | Overall Mean (SD)  | Integration Index Score Classification Group Means |                                 |                                 | p-values |             |
|--|----------------|------------------------|--------|--------------------|--|---------------------------------|---------------------------------|----------|-------------|
|  |                |                        |        |                    | Low Score (0-3.29)                                 | Medium Score (3.30-6.59)        | High Score (6.60-10)            | Low-Med. | Med. - High |
| Integration of providers at who offer CI and at modern FP method                       | 0.39           | 1.87                   | 0.21   | 0.56 (0.4)         | 0.05   | 0.47                            | 0.90                            | 0.00     | 0.00        |
| Integration of providers at who routinely offer during CI/CGM                          | 0.41           | 1.87                   | 0.22   | 0.58 (0.39)        | 0.04   | 0.49                            | 0.93                            | 0.00     | 0.00        |
| Integration of providers at who offer FP items that a r at a facility tells during CHS | 0.33           | 1.87                   | 0.18   | 1.67 (1.52)        | 0.14   | 1.76                            | 3.38                            | 0.00     | 0.00        |
| Integration of providers at that do not request CI during CHS                          | 0.37           | 1.87                   | 0.20   | 0.5 (0.37)         | 0.04   | 0.41                            | 0.81                            | 0.00     | 0.00        |
| Integration of providers at who provides CI and FP during CHS                          | 0.38           | 1.87                   | 0.20   | 0.77 (0.42)        | 0.23   | 0.88                            | 0.97                            | 0.00     | 0.00        |
| <b>Provider Integration Index</b>  |                |                        |        | <b>5.42 (3.10)</b> | <b>0.75 (23% of facilities)</b>                    | <b>4.99 (32% of facilities)</b> | <b>8.23 (45% of facilities)</b> | 0.00     | 0.00        |
| <b>Integration Variable Description</b>  |                |                        |        |                    |  |                                 |                                 |          |             |
| Integration of providers at who practice if FP info during CH visit                    | 0.38           | 1.21                   | 0.32   | 5.82 (2.23)        | 0.09   | 0.91                            | 0.98                            | 0.00     | 0.00        |
| Integration of providers at who practice if hormonal methods during CH visit           | 0.41           | 1.21                   | 0.34   | 4.63 (1.97)        | 0.05   | 0.80                            | 0.96                            | 0.00     | 0.00        |
| Integration of providers at who offer FP if days where both CI and FP are offered      | 0.42           | 1.21                   | 0.34   | 0.29 (0.35)        | 0.01   | 0.11                            | 0.49                            | 0.00     | 0.00        |
| <b>Provider Integration Index</b>  |                |                        |        | <b>6.22 (2.72)</b> | <b>0.50 (14% of facilities)</b>                    | <b>6.00 (38% of facilities)</b> | <b>8.10 (48% of facilities)</b> | 0.00     | 0.00        |

**Table 5: Provider Integration Index Score Classification Differences in Facilities with High Provider Integration Index Scores Following Use of Variable Sub-Sets in the PCA**


| Provider Integration Index Score Classification | Index Score  | Base case: all variables | Omitted Variable  |  |  |  |                                  |
|---|--------------|--------------------------|---|--|--|--|----------------------------------|
|   |              |                          | Proportion providers who offer CI and at least 1 modern FP method | Proportion providers who routinely offer FP info during CI/CGM | Average FP items that a provider tells client during CHS | Proportion providers who do not request partner consent during CHS | Facility provides both CI and FP |
| Low   | 0.00 - 3.29  | 0%                       | 0%  | 0%   | 0%   | 0%   | 0%                               |
| Medium  | 3.30 - 6.59  | 0%                       | 7%  | 5%   | 0%   | 5%   | 24%                              |
| High  | 6.60 - 10.00 | 100%                     | 93%   | 95%  | 100%   | 95%  | 76%                              |

**Table 6: Facility Integration Index Score Classification Differences in Facilities with High Facility**

| Facility Integration Index Score Classification | Index Score  | Base case: all variables | Omitted Variable  |  |   |
|---|--------------|--------------------------|---|--|---|
|   |              |                          | Normal practice if FP information is wanted during CH visit | Normal practice if hormonal method of FP is wanted during CH visit | Score of days when both CI and FP are offered |
| Low   | 0.00 - 3.29  | 0%                       | 0%  | 0%   | 0%  |
| Medium  | 3.30 - 6.59  | 0%                       | 55%   | 55%  | 1%  |
| High  | 6.60 - 10.00 | 100%                     | 45%   | 45%  | 99%   |


**Integration Index Scores Following Use of Variable Sub-Sets in the PCA**

**Figures**




**Figure 1**

Scree Plot of Eigenvalues after PCA



**Figure 2**

Provider Integration Index Scores



**Figure 3**

Facility Integration Index Scores

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [200626MeasureFPCIIntegrationCoverLetter.pdf](#)