

Dispersal patterns and population structure of *Aedes albopictus* (Diptera: Culicidae) in three different climatic regions of China

Jian Gao

Beijing Institute of Microbiology and Epidemiology

Hengduan Zhang

Beijing Institute of Microbiology and Epidemiology

Xiaoxia Guo

Beijing Institute of Microbiology and Epidemiology

Dan Xing

Beijing Institute of Microbiology and Epidemiology

YanDe Dong

Beijing Institute of Microbiology and Epidemiology

CeJie Lan

Beijing Institute of Microbiology and Epidemiology

Ge Wang

Beijing Institute of Microbiology and Epidemiology

ChaoJie Li

Beijing Institute of Microbiology and Epidemiology

ChunXiao Li

Beijing Institute of Microbiology and Epidemiology

TongYan Zhao (✉ tongyanzhao@126.com)

Beijing Institute of Microbiology and Epidemiology

Research

Keywords: *Aedes albopictus*, Genetic variation, Haplotype, Dispersion pattern, Climatic regions, Microsatellite loci, Environmental factors

Posted Date: November 6th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-38761/v3>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 6th, 2021. See the published version at <https://doi.org/10.1186/s13071-020-04521-4>.

Abstract

Background:

Aedes albopictus is an indigenous primary vector for Dengue and Zika viruses in China. Compared with its insecticide resistance, biology, and vector competence, little is known about its genetic variation, which corresponds to environmental variations. Thus, the present study examines how *Ae. albopictus* varies among different climatic regions in China and deciphers its potential dispersal patterns.

Methods:

The genetic variation and population structure of 17 *Ae. albopictus* populations collected from three climatic regions of China were investigated with 11 microsatellite loci and the mitochondrial *coxI* gene.

Results:

Of 44 isolated microsatellite markers, 11 pairs were chosen for genotyping analysis and had an average PIC value of 0.713, representing high polymorphism. The number of alleles was high in each population, with the n_e value increasing from the temperate region (3.876) to the tropical region (4.144). Twenty-five *coxI* haplotypes were detected, and the highest diversity was observed in the tropical region. The mean H_o value (ca. 0.557) of all the regions was significantly lower than the mean H_e value (ca. 0.684), with nearly all populations significantly departing from HWE and displaying significant population expansion (p -value < 0.05). Two genetically isolated groups and three haplotype clades were evaluated via STRUCTURE and haplotype phylogenetic analyses, and the tropical populations were significantly isolated from those in the other regions. Most genetic variation in *Ae. albopictus* was detected within populations and individuals at 31.40% and 63.04%, respectively, via the AMOVA test, and a relatively significant positive correlation was observed among only the temperate populations via IBD analysis ($R^2 = 0.6614$, $p = 0.048$). Recent dispersions were observed among different *Ae. albopictus* populations, and four major migration trends with high gene flow ($Nm > 0.4$) were reconstructed between the tropical region and the other two regions. Environmental factors, especially temperature and rainfall, may be the leading causes of genetic diversity in different climatic regions.

Conclusions:

Continuous dispersion contributes to the genetic communication of *Ae. albopictus* populations across different climatic regions, and environmental factors, especially temperature and rainfall, may be the leading causes of genetic variation.

Background

Aedes (Stegomyia) albopictus, also known as the Asian tiger mosquito, is an epidemiologically critical vector for its transmission of several arboviruses, including Dengue virus (DENV), Zika virus (ZIKV), Chikungunya virus and filarial nematodes, such as *Dirofilaria immitis* [1-4]. Compared with *Aedes aegypti*,

the primary vector for DENV and ZIKV, *Ae. albopictus* is generally considered to be a less efficient vector due to its less anthropophilic behavior and inferior adaptation to urban domestic environments [5]. However, its broad temperature adaptability and high competence for several arboviruses make it approximately as important a vector as *Ae. aegypti*. Meanwhile, combined with its strong ecological plasticity, global trade, such as the transportation of used tires and “lucky bamboo”, has also accelerated the global spread of this species from its original Southeastern Asian distribution to every continent except Antarctica [6]. Currently, it is among the top 100 worst invasive species and a threat to human health worldwide [7].

Natural environmental variation is responsible for fluctuations in insect population dynamics, distribution, and biology, including population number, intensity, and feeding behavior [8]. An important and rapidly adapting insect with high fecundity and a short life cycle, as *Ae. albopictus* has population dynamics and vector competence that are also greatly influenced by environmental conditions [9, 10]. Numerous previous studies have discussed the effects of environmental variations on *Ae. albopictus* populations, mainly based on their influence on mosquito abundance, survival, size, fecundity, and competence for certain arboviruses [11-15]. However, less research has focused on changes in the genetic diversity and population structure of *Ae. albopictus* populations associated with various environmental conditions [79,82,83], although this information is essential for reconstructing the dispersion patterns of *Ae. albopictus* in certain regions and providing necessary information for subsequent mosquito control.

With the advent of urbanization in China, the natural environment was altered tremendously; more people have moved from rural areas to cities, and trade has increased between cities and rural areas. All these processes create suitable habitats (e.g. waste tires, vases, plastic pails, and concrete tanks, etc.) for *Ae. albopictus* and facilitate its breeding and dispersal throughout the country [16-19]. Meanwhile, the dispersion of *Ae. albopictus* from its original environment to a new environment facilitates the manifestation of new phenotypes, or that the increasing of mutated alleles in frequency, and thriving international travel and trade provide essential routes for the continuous introduction of arboviruses from other countries, especially those in Southeast Asia, where *Ae. albopictus* originated [20], thus putting people at risk worldwide. During the past ten years, at least 22 imported ZIKV cases were confirmed by the China CDC, and the persistent emergence of dengue in Southern China, especially in Guangdong and Yunnan provinces, is positively correlated to the widespread presence of *Ae. aegypti* and *Ae. albopictus* among these regions [21]. In contrast to *Ae. aegypti* (an invasive species distributed only in the southern area of China), *Ae. albopictus* is an indigenous Chinese mosquito, ranging from Dalian in the north to Hainan in the south [22, 23]. Moreover, it has been considered the sole agent of numerous recent dengue fever outbreaks in China [24, 25]. Its wide distribution and high competence for numerous arboviruses and nematode parasites emphasize the need to more extensively study the biology, distribution, and dispersion patterns of this species.

High genetic variation is ubiquitous in vector populations, especially in invasive vectors, such as *Ae. aegypti* and *Ae. albopictus*, where it helps them occupy diverse niches and respond quickly to evolutionary challenges [26]. Microsatellites are the preferred markers for studying genetic variation in

vectors due to their codominance, high informativity, and vast abundance throughout vector genomes [27]. To date, multiple microsatellite loci associated with *Ae. albopictus* have been successfully isolated and employed in *Ae. albopictus* population studies on a global scale, and they continue to be a popular choice of genetic marker [15, 28-30]. Simultaneously and independently from *Ae. albopictus* microsatellites, the mitochondrial *coxI* gene has frequently been used for mosquito barcoding and invasive species monitoring due to its conservation and high accuracy when distinguishing sequence variation at the interspecies level and its usefulness to investigate population dispersal routes of mosquito [31-35].

To evaluate how environmental factors affect genetic diversity and dispersal patterns of *Ae. albopictus* and decipher the potential dispersal patterns of *Ae. albopictus* among different climatic regions, a total of 17 *Ae. albopictus* populations were collected and examined via 11 microsatellite loci and the *coxI* gene. The results provide information for the future control of *Ae. albopictus* in China.

Materials And Methods

Eco-climatic characteristics of different climatic regions

All mosquitoes were sampled from the three main climatic regions of China, including tropical, subtropical and temperate regions. The annual accumulated temperature and precipitation for the tropical region is above 8000°C and 1500~2000 mm, respectively, with the lowest monthly average temperature is 15~24°C; this data for the subtropical region is between 4500~8000°C and above 1000 mm, respectively, with the lowest monthly average temperature is 0~15°C; and this data for the temperate region is between 3000~4500°C and 700 mm, respectively, with the lowest monthly average temperature is -8~0°C [36, 37].

Mosquito sampling and DNA isolation

In the present study, ca. 600 *Ae. albopictus* larvae were sampled from seventeen geographically diverse sites across three environmentally distinct regions of China from June to August 2018; all the sampling site information is described in Fig. 1 and Additional file 1: Table S1. To avoid inbreeding interference, each pool of larvae mosquitoes for a given locality was collected from at least five wild breeding places within 500 meters. For each of the samples, the larvae were reared separately and emerged at 25°C ± 1°C at 75% ± 5% relative humidity (RH) under a 14 h light/10 h dark (LD) photoperiod, and all female adult mosquitoes were identified under a microscope before DNA isolation [38, 39]. The DNA isolation was conducted with the Qiagen DNeasy Blood and Tissue Kit (No. 69504, Qiagen, Germany) according to the manufacturer's protocol, and all the DNA samples were stored at -80°C.

Microsatellite isolation, processing, and coxI gene amplification

Referring to Chambers *et al.* [31], microsatellite markers were isolated from the whole genome of *Ae. albopictus* by magnetic-bead enrichment and PCR screening, and all markers were found to be highly

polymorphic via denatured polyacrylamide gel electrophoresis (D-PAGE). A set of 11 microsatellite loci was employed for genotyping the 17 *Ae. albopictus* populations. Detailed microsatellite primer information is listed in Table 1. All PCRs were performed on a T100 Thermal Cycler (Bio-Rad, USA) in a 50 µl reaction system containing 10 ng of DNA, 0.25 U of PrimeSTAR HS DNA Polymerase (10 pmol/µl, TaKaRa, Japan), 6 µM of dNTPs (2.5 mM each, TaKaRa, Japan) and ca. 5 µM of each primer, with the program set to 35 cycles of 95°C for 30 sec, 57°C for 30 sec and 72°C for 1 min followed by a final elongation at 72°C for 10 min. All products were then checked with 2% agarose gel electrophoresis under UV light and run on a 3730XL DNA Genetic Analyzer (Applied Biosystems, USA).

As described by Bonacum *et al.* [32], *coxI* sequence polymorphism was investigated among 497 individuals (22-30 individuals per locality). Briefly, DNA amplification of a 550-bp fragment of *coxI* was performed on a T100 Thermal Cycler (Bio-Rad, USA) with the following primer set: 5'-GGAGGATTTGGAAATTGATTAGTTC-3' (F-*coxI*) and 5'-CCCGGTAAAATTAATAAACTTC-3' (R-*coxI*) in a 50 µl reaction mix containing 10 µl PCR Reaction Buffer (TaKaRa, Japan), 4 µl of dNTPs (2.5 mM each, TaKaRa, Japan), 1 µl of primers (10 pmol/µl, TaKaRa, Japan), and 0.5 µl PrimeSTAR HS DNA Polymerase (TaKaRa, Japan). The PCR amplification program was set as follows: pre-denaturation at 94°C for 3 min, followed by 35 cycles of denaturation at 94°C for 30 sec, annealing at 54°C for 45 sec, and elongation at 72°C for 1 min, with a final elongation at 72°C for 10 min. All the PCR products were detected and separated by 2% agarose gel electrophoresis. The target fragments for *coxI* were then cut from the gel under UV light and purified with the GenElute™ PCR Clean-Up Kit (NA1020, Sigma-Aldrich, USA). Each purified PCR product was then cloned into the pCR™2.1 Vector with a TA Cloning™ Kit (K202040, Invitrogen, USA) and selected by bacteria liquid PCR with the T7 promoter primers. Finally, at least 20 positive clones for each PCR product were sequenced on both strands using an ABI 3730XL automatic sequencer (Applied Biosystems, USA).

Population structure analyses, phylogenetic genotyping and migration analyses

The microsatellite data were processed with GeneMapper v.4.0 (Applied Biosystems, USA). All markers were tested for polymorphism by determining their polymorphism information content (PIC) values via PIC-Calc 0.6 [40], and the null allele frequency of each locus was assessed with Micro-Checker version 2.2.3 [41]. Allele frequency indices, including n_a and n_e , were assessed with Cervus version 3.0.7 [42]. The H_o , H_e , and FIS values of all populations were calculated with Arlequin version 3.5.2.2 [43]. Departure from Hardy-Weinberg equilibrium (HWE) and heterozygosity deficiency were assessed via Bottleneck version 1.2.02 [44]. Additionally, STRUCTURE version 2.3.4 [45] and the ΔK method of Evanno *et al.* [46] were employed to calculate the optimal K value; the parameters were set as follows: (1) K ranged from 1 to 20 with ten iterations for each K value; (2) an admixture model was chosen with a Markov chain Monte Carlo algorithm run for 100,000 iterations and 1,000,000 repetitions. The optimal K value was calculated via Structure Harvester: <http://taylor0.biology.ucla.edu/structureHarvester/> and depicted with Distribut version 1.1. The DAPC analysis was performed on the microsatellite data of each population with the R packages "Adegenet 2.1.3" [47], and the IBD analysis was performed between genetic distance, generated from microsatellite data, and geographical distance with "Genpop 4.7.5" [48], respectively; meanwhile, the

genetic variation and F_{ST} value for each population were evaluated with the AMOVA test via Arlequin version 3.5.2.2 [43].

The haplotypes of all the populations were screened by DnaSP version 6.0 [49], and the genetic relationships among all the haplotypes were displayed as a TCS network constructed by Network 10.0.0.0 [50, 51]. Moreover, BEAST version 1.8.4 [52] and the R package “pheatmap” [53] were used to build a phylogenetic tree of all the haplotypes and examine the distribution of all the haplotypes among the different climatic regions, with the best model selected by JModelTest 2.1.10 [54]. The mismatch distribution and Bayesian skyline plot analyses were conducted with Arlequin version 3.5.2.2 and BEAST version 1.8.4, respectively, to investigate the current dispersal incidences of each population.

Concurrently, the possible migration routes were rebuilt via the R package “divMigrate” with the number of bootstrap replicates set to 3, the alpha value set to 0.05, the Nm method chosen for the migration statistic, and the filter threshold value set to 0.25 [55]. Finally, the relationships between the genetic indices (i.e. n_e , Shannon index, H_d , H , and π) and environmental factors (i.e. temperature, altitude and rainfall) were described via principal component analysis (PCA) and multiple factor analysis (MFA) with the R package “FactoMineR” [56].

Results

Microsatellite marker isolation and assessment

In the present study, a total of 44 pairs of microsatellite markers were isolated from the whole genome of *Ae. albopictus*, 11 pairs of which were highly polymorphic and were therefore chosen for microsatellite genotyping analysis (Table 1). The allele number of each locus ranged from 10 to 33, with a mean of 17.545 alleles per locus. The PIC values of each locus ranged from 0.334 to 0.925, with a mean value of 0.713. According to the definition of PIC values by Allah *et al.* [57], nearly all the markers selected were highly informative (PIC value > 0.5); the exception was BW-P18, which had a value of 0.357 and was considered to be an informative marker. The Micro-Checker results suggested that null alleles were present at all loci, with frequencies ranging from 0.064 to 0.157, with an average of 0.078 (Additional file 8: Table S5); however, frequency values smaller than 0.2 are considered to have no significant effect on the accuracy of data analysis by many studies [59, 65]. The linkage disequilibrium (LD) test showed that 302 pairs of loci of a total of 1870 (16.15%) across all locations were in significant LD after Bonferroni correction, while no consistency was found among them (Additional File 5: Figure S1).

Genetic diversity and variation

According to the results of the microsatellite data analysis, the observed number of alleles (n_a) in each *Ae. albopictus* population was very high, and the mean n_a value in each climatic region ranged from 6.909 to 8.091 without a significant difference. In contrast, the effective number of alleles (n_e) ranged from 3.501 to 4.525, and the n_e value increased from the temperate region (3.876) to the tropical region (4.144). The mean value of observed heterozygosity (H_o) for all the climatic regions was ca. 0.557, which

was significantly lower than the expected heterozygosity (H_e) (ca. 0.684). The F_{IS} value of each climatic region ranged from 0.266 to 0.359, and the *Ae. albopictus* populations significantly departed from HWE, except in two locations, SHJD and KZXZ (subtropical region, Table 2). Heterozygosity tests of all 17 *Ae. albopictus* populations based on the stepwise mutation model (SMM) revealed that nearly all the populations from the temperate and subtropical regions displayed significant population expansion with p -values < 0.05 , while no significance was observed among all the tests for the *Ae. albopictus* populations from the tropical region after Bonferroni correction (Additional file 2: Table S2).

In all, a total of 25 *coxI* haplotypes were observed among 497 *Ae. albopictus* individuals (GenBank ID: MN651301-MN651325) based on an analysis of the *coxI* sequences. The haplotype indices (i.e. H_d , H , and π) changed dramatically across *Ae. albopictus* populations from different climatic regions (Additional file 3: Table S3). The haplotype diversity (H_d) ranged from 0.074 (ZGND, temperate region) to 0.750 (JYJB, tropical region), with the nucleotide diversity (π) ranging from 0.014 (BJLG, temperate region) to 0.190 (JYJB, tropical region). The average number of nucleotide differences (k) ranged from 0.067 (BJLG, temperate region) to 0.929 (JYJB, tropical region), and the number of polymorphic sites ranged from 1 to 5 across all the populations. Overall, the tropical *Ae. albopictus* populations showed the highest diversity, with mean values of H_d , π , and k reaching 0.657, 0.156, and 0.763, respectively, whereas the temperate populations showed the lowest diversity.

Population structure and differentiation based on microsatellite analysis

In the present study, according to the microsatellite analysis results, all the *Ae. albopictus* populations were adequately allocated to two clades with significant genetic differences, and the best K value, as determined via the ΔK method of Evanno *et al.*, was also equal to two (Fig. 2a). Combined with the STRUCTURE bar plot analysis, the Bayesian clustering analysis showed that all the *Ae. albopictus* populations were adequately allocated to two clades with certain locations from the subtropical and temperate regions genetically isolated from the other locations. (Fig. 2b&c). Moreover, a total of 86.4% of the variation was explained by 50 PCs in the discriminant analysis of principal components (DAPC) analysis; the results revealed two genetically isolated groups, and there was no clear relationship between the *Ae. albopictus* population structure and its distribution across climatic regions (Fig.2d).

The AMOVA results are presented in Table 3. Of the total genetic variation partitioned, 31.40% could be attributed to differences among individuals within populations and 63.04% to differences within individuals ($F_{IS}=0.33253$, $F_{IT}=0.36962$, and all $p<0.00001$). Meanwhile, the pairwise F_{ST} values between populations ranged from 0.008 (ZGND, temperate region and NNXD, subtropical region) to 0.141 (SHJD, subtropical region and JKCH, tropical region). Genetic differentiation was significant between all the sampled populations after Bonferroni correction ($p < 0.05$) except for five pairs of F_{ST} values among the subtropical populations NNXZ, NJTH, NJDX, KZXZ, and temperate populations ZGND, BHBG, QDDX, and HBSD (Additional file 4: Table S4). In contrast to the high individual variation, a slightly significant positive correlation was observed among only the temperate populations via isolation by distance (IBD)

analysis ($R^2 = 0.6614$, $p = 0.048$), while no such evidence was observed in the other two regions (Additional file 6: Figure S2).

Haplotype network and phylogenetic analysis based on *coxI* sequences

Three major haplotype clades, distributed across the three central climatic regions and closely related to each other, were reconstructed via the TCS network with 497 *coxI* sequences. Haplotype I (H1) of Clade I was the most frequent haplotype and was distributed from the tropical to the temperate region with an increasing trend. Nearly all the other haplotypes derived from H1 with one or two mutations, with Clade I mainly distributed in the tropical region and Clade II mainly distributed in the southern subtropical region (Fig. 3b).

A phylogenetic tree of all 25 haplotype sequences, combined with a heatmap analysis, demonstrated that the 25 haplotypes were divided into three major well-supported clades. As expected, Clade I was separated from Clade II and Clade III with 100% bootstrap support and included haplotypes H9, H10, H11, H12, H19, and H20, which were observed in only the tropical region. In comparison, Clade II diverged from Clade I with lower bootstrap support (70.74%) and included five haplotypes (H2~H3 and H5~H7) that were distributed in only the southern subtropical region. Clade III was an admixture group containing all the remaining 14 haplotypes, two of which were observed in the tropical region and four in the climatic region (Fig. 3a). Meanwhile, the Tajima's D and Fu's F_s values were negative, with no statistically significant p-values except in population JKCH (tropical region, Additional file 3: Table S3). Based on the *coxI* sequences, the mismatch analysis results showed that the Harpending raggedness indices for all three haplotype clades were relatively low (ranging from 0.1054 to 0.1812, $p > 0.05$), and unimodal mismatch distributions were observed among the different *Ae. albopictus* populations (Additional file 7: Figure S3).

Migration and correlation analyses between genetic indices and environmental factors

As illustrated in Fig. 4, all five genetic indices (i.e. π , Shannon index, Hd, H, and π) and two environmental factors (i.e., temperature and rainfall) contributed equivalently to the first axis of the PCA, explaining up to 95.2% of the variation; the exception was the environmental factor latitude, which contributed more to the second axis, with a proportion of 17.5%. Combining these results with hierarchical clustering via MFA, all 17 *Ae. albopictus* populations were clustered into three groups, of which cluster II and cluster III were closely related to each other. When environmental factors were regarded as the major influencing factors, the molecular diversity indices (i.e. Hd, k, and π) of the populations in the tropical region were significantly higher than those in the other regions (Additional File 3 TableS3).

Migration patterns were assessed using divMigrate networks representing all 17 *Ae. albopictus* populations. As expected, *Ae. albopictus* was observed to have migrated frequently between the tropical and temperate areas in both directions (Fig. 1). A total of four major migration trends were observed among the different climatic regions with high gene flow ($N_m > 0.4$). Two trends were found to have originated from the tropical and subtropical areas and reached Hebei and Beijing in the temperate

area, while another two arrived in Guangdong and Guangxi from the southern subtropical area. Compared with the south to north routes, the latter migration routes were substantially larger.

Discussion

Due to their high mutation rates, codominant expression, and universal distribution throughout the eukaryotic genome, microsatellite loci have been widely used to evaluate the genetic variation and population structure of mosquito, especially for those without fully annotated genomes [29, 31]. Although different sets of microsatellite loci have been identified by previous studies [28, 29, 58], the absence of PCR amplification products of mosquito samples from certain temperature zones has resulted in insufficient data. Therefore, in this study, we developed new primers for our samples to ensure the integrity of the experimental data. However, the potential impact of null alleles, such as reducing population genetic diversity and increasing genetic differentiation among populations [59-61], must be taken into consideration when microsatellite markers are used, because null alleles are more frequent in arthropods than in other species [62-64]. Mitochondrial genes are also good markers for studying population diversity and are widely used in many studies. To improve the accuracy of the findings in this study, the mitochondrial *coxI* gene was also employed, and the genetic variation of all the individuals was investigated using both *coxI* and microsatellite data.

Consistent with the studies of Zhong *et al.* [66] and Wei *et al.* [67], which were based on only mitochondrial and microsatellite diversity, respectively, the tropical *Ae. albopictus* populations showed higher diversity than the populations of the other two regions in China. An environment more conducive to survival and continuous dispersion may be the best explanation for this phenomenon. According to studies by Wei *et al.* and others [67-70], the hot and humid climate of Southern China is more suitable for the breeding and development of *Ae. albopictus* than a temperate climate, which has relatively low temperatures and dry air. Notably, during the winter, freezing temperatures bring about diapause in *Ae. albopictus* for the whole winter period, resulting in lower allele richness and population diversity in temperate *Ae. albopictus* populations [62-64]. Meanwhile, Southern China shares borders with many countries in Southeast Asia, where *Ae. albopictus* originates geographically, including Laos, the Philippines, and Myanmar [20]. Frequent border trades and resident travel among these areas result in the continuous dispersion of *Ae. albopictus*, enriching the diversity of the local population; this result was confirmed by the results of the bottleneck analysis, which showed that all the populations from temperate and subtropical regions had undergone bottleneck effects, while the populations of the tropical region had not. As suggested by Takezaki *et al.* [71], an H_e value between 0.5 and 0.8 indicates high genetic polymorphism in the population. In the present study, the H_e values of all the sampling sites ranged from 0.606~0.733, which indicates high polymorphism. Simultaneously, compared with those in the tropical region, the mean H_o values of the temperate and subtropical regions were significantly lower than the mean H_e values, and nearly all the populations significantly departed from HWE. This phenomenon may be also closely related to the less suitable climate and the need to develop diapause during wintertime among populations of these regions as mentioned before, and our present sampling method (larvae rather than adults) may be also the reason for the significant departure from HWE of all populations.

Meanwhile, the continuous dispersion of *Ae. albopictus* from Southeast Asian countries to the tropical region of China also augments the local mosquito population.

In the present study, nearly 44% (11 out of 25) of these 25 detected haplotypes were observed in only the tropical and subtropical regions. The same result for Chinese *Ae. albopictus* populations was also observed by both Guo *et al.* and Zhang *et al.* [79, 80]. Guo *et al.* identified three predominant haplotypes in *Ae. albopictus* populations: H01 (21.5%) from Guangdong Province (subtropical region), H19 (22.1%) from Yunnan (tropical region) and Hainan provinces (tropical region), and H30 (10%) from Hainan Province (tropical region); likewise, Zhang *et al.* also detected 25 haplotypes across China, and 48% (12 out of 25) of these haplotypes were detected in only tropical and subtropical regions. Both studies corroborate our findings that the haplotype diversity of the tropical and subtropical regions are higher than that of the temperate region. Continuous dispersal from neighboring Southeast Asian countries may be the reason for the higher diversity of the *Ae. albopictus* populations in the tropical and subtropical regions and that the origins of the *Ae. albopictus* populations of these two regions may be different. Furthermore, the mismatch analysis results revealed recent population dispersal among the different climatic regions, which is the main reason for the universal distribution of the remaining haplotypes. As an important and rapidly adapting insect, *Ae. albopictus* has biological characteristics that are strongly influenced by environmental conditions [9, 10]. In tropical and subtropical regions, the climate is hot and wet, making it more suitable for the survival of *Ae. albopictus*; when this mosquito migrates from a tropical region to a temperate region, it needs time to adapt to the new environment, especially to survive during the winter period, which causes diapause in *Ae. albopictus* [4]. According to a study by Roiz *et al.* about the effects of environmental factors on *Ae. albopictus*, the activity of host-seeking females was positively affected by temperature and rainfall [81]. The PCA in the present study examined genetic indices and environmental factors (such as temperature, rainfall and altitude) and revealed that temperature and rainfall may be the leading causes of differences in genetic diversity among *Ae. albopictus* populations from different climatic regions.

Corresponding to the results of Kotsakiozi and Manni *et al.* [29, 77], which showed no genetic differentiation within the native Asian range, our population differentiation analysis results showed that even though nearly all the pairwise F_{ST} values between populations were significant, the values were not high, which indicates potential anthropogenic exchange between the populations. A study by Schmidt *et al.* [78] of the genetic structure of *Ae. albopictus* collected from 12 sample sites in Guangzhou also found that human transportation networks, particularly shipping terminals, influenced the genetic structure of *Ae. albopictus* populations. With the development of China, the rapid expansion of high-speed rail, aircraft routes, and highways between different temperature zones has not only facilitated trade among different cities but also accelerated the spread of mosquitoes. Accordingly, only two genetically separated clades were observed among these populations, and molecular variations within populations and individuals were found to contribute to the genetic differentiation between these populations. As a global invasive species, *Ae. albopictus* has developed several strategies to cope with a broad range of temperatures and adapt to local thermal conditions [9, 68], which helps this mosquito disperse and

colonize different locations successfully. Here, the migration analysis results displayed four major migration trends with high gene flow ($Nm > 0.4$) among the different climatic regions, corresponding with China's main trade routes (e.g. pan-Pearl River Delta and Bohai Rim Business District, etc.) and tourism routes (e.g. Beijing-Guangzhou railway, Beijing-Shanghai railway, and Beijing-Kowloon railway, etc.). All these results revealed that human-aided dispersion might be the main reason for the similarity of populations across the different regions [22, 67-78]. This hypothesis was also verified via IBD analysis, which indicated that long distances were not significantly associated with individual genetic variation, except among individuals from the tropical region with a significant p -value of the Mantel test.

Overall, strong dispersal patterns were observed in Southern China, and frequent dispersals have occurred from the tropical and subtropical regions to the temperate region. Monitoring the population structure and potential migration routes of *Ae. albopictus* in different climatic regions will help in the selection of appropriate mosquito control strategies, and since dengue fever continues to break out in Southern China every year, monitoring the potential migration routes of *Ae. albopictus* can also predict the potential transmission routes of this virus and help prevent its further spread.

Conclusion

The present study systematically evaluated the genetic variation, population structure, and haplotype-based phylogenetic relationships of *Ae. albopictus* populations across different climatic regions of China. All 17 *Ae. albopictus* populations were genetically structured into two isolated clades, in accordance with their locality or region of origin, and three major haplotype clusters were observed via phylogenetic analysis of *cox1*. These results suggest different evolutionary histories in changing environments, especially differences in temperature and rainfall. Meanwhile, four major migration trends were observed among the different climatic regions with high gene flow; these trends contribute to the similarity of the *Ae. albopictus* populations.

Abbreviations

cox1: Mitochondrial cytochrome c oxidase subunit 1; PIC: Polymorphism Information Content; H_o : Observed heterozygosity; H_e : Expected heterozygosity; HWE test: Hardy-Weinberg equilibrium test; AMOVA test: Analysis of molecular variation test; IBD: Isolation by distance; WHO: World Health Organization; CDC: Centers for Disease Control; n_a : Observed number of alleles; n_e : Effective number of alleles; F_{IS} : Inbreeding coefficient; F_{ST} : Genetic differences among populations; F_{IT} : Genetic differences within individuals; PCA: Principal Component Analysis; MFA: Multiple Factor Analysis; LD: Linkage disequilibrium; Hd: Haplotype diversity; DAPC: Discriminant analysis of principal components; Nm: Number of migrants.

Declarations

Acknowledgements

Not applicable.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and material

Data are available on request to the corresponding author.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was funded by grants from the Infective Diseases Prevention and Cure Project of China (No.2017ZX1000303404)

Authors' contributions

JG, HDZ, CXL and TYZ jointly designed and coordinated the study, with contributions from XXG, DX and YDD. JG drafted the article with contributions from HDZ. HDZ, JG, CJL, and GW collected samples from different climatic regions of China. JG, HDZ, GW, and CJL carried out the laboratory work and performed the statistical analysis. All authors read and approved the final manuscript.

References

1. Cancrini G, Frangipane di Regalbono A, Ricci I, Tessarin C, Gabrielli S, Pietrobelli M. *Aedes albopictus* is a natural vector of *Dirofilaria immitis* in Italy. *Vet Parasitol.* 2003;118(3-4):195-202.
2. Wong PS, Li MZ, Chong CS, Ng LC, Tan CH. *Aedes (Stegomyia) albopictus* (Skuse): a potential vector of Zika virus in Singapore. *PLoS Negl Trop Dis.* 2013;7(8):e2348.
3. Grard G, Caron M, Mombo IM, Nkoghe D, Mboui Ondo S, Jiolle D, et al. Zika virus in Gabon (Central Africa)-2007: a new threat from *Aedes albopictus*? *PLoS Negl Trop Dis.* 2014;8(2):e2681.
4. Kress A, Kuch U, Oehlmann J, Muller R. Effects of diapause and cold acclimation on egg ultrastructure: new insights into the cold hardiness mechanisms of the Asian tiger mosquito *Aedes (Stegomyia) albopictus*. *J Vector Ecol.* 2016;41(1):142-50.
5. Rezza G. *Aedes albopictus* and the reemergence of Dengue. *BMC Public Health.* 2012;12:72.

6. Kraemer MU, Sinka ME, Duda KA, Mylne AQ, Shearer FM, Barker CM, et al. The global distribution of the arbovirus vectors *Aedes aegypti* and *Ae. albopictus*. *Elife*. 2015;4:e08347.
7. Bonizzoni M, Gasperi G, Chen X, James AA. The invasive mosquito species *Aedes albopictus*: current knowledge and future perspectives. *Trends Parasitol*. 2013;29 (9):460-8.
8. Ayres JS, Schneider DS. The role of anorexia in resistance and tolerance to infections in *Drosophila*. *PLoS Biol*. 2009;7(7):e1000150.
9. Culbert NJ, Gilles JRL, Bouyer J. Investigating the impact of chilling temperature on male *Aedes aegypti* and *Aedes albopictus* survival. *PLoS One*. 2019;14(8):e0221822.
10. J. Kweka E, Baraka V, Mathias L, Mwang'onde B, Baraka G, Lyaruu L, et al. Ecology of *Aedes* Mosquitoes, the Major Vectors of Arboviruses in Human Population. In: Jorge Abelardo Falcón-Lezama, Miguel Betancourt-Cravioto, and Roberto Tapia-Conyer, editors. *Dengue Fever-a Resilient Threat in the Face of Innovation*. London: IntechOpen; 2019. p. 40-50.
11. Catherine J. Westbrook MHR, Kendra N. Pesko, Krystle E. Greene, and L. Philip Lounibos. Larval Environmental Temperature and the Susceptibility of *Aedes albopictus* Skuse (Diptera: Culicidae) to Chikungunya Virus. *VECTOR-BORNE AND ZOOLOGIC DISEASES*. 2010;10:241-7.
12. Ephantus J. Muturi MJB, and Allison Montgomery. Temperature and density-dependent effects of larval environment on *Aedes aegypti* competence for an alphavirus. *Journal of Vector Ecology*. 2012;37(1):154-61.
13. Murdock CC, Evans MV, McClanahan TD, Miazgowicz KL, Tesla B. Fine-scale variation in microclimate across an urban landscape shapes variation in mosquito population dynamics and the potential of *Aedes albopictus* to transmit arboviral disease. *PLoS Negl Trop Dis*. 2017;11(5):e0005640.
14. Costanzo KS, Westby KM, Medley KA. Genetic and environmental influences on the size-fecundity relationship in *Aedes albopictus* (Diptera: Culicidae): Impacts on population growth estimates? *PLoS One*. 2018;13(8):e0201465.
15. Lubinda J, Trevino CJ, Walsh MR, Moore AJ, Hanafi-Bojd AA, Akgun S, et al. Environmental suitability for *Aedes aegypti* and *Aedes albopictus* and the spatial distribution of major arboviral infections in Mexico. *Parasite Epidemiol Control*. 2019;6:e00116.
16. Chadee DD, Martinez R. *Aedes aegypti* (L.) in Latin American and Caribbean region: with growing evidence for vector adaptation to climate change? *Acta Trop*. 2016;156:137-43.
17. Reba M, Reitsma F, Seto KC. Spatializing 6,000 years of global urbanization from 3700 BC to AD 2000. *Sci Data*. 2016;3:160034.
18. Jia P, Chen X, Chen J, Lu L, Liu Q, Tan X. How does the dengue vector mosquito *Aedes albopictus* respond to global warming? *Parasit Vectors*. 2017;10(1):140.
19. Johnson MTJ, Munshi-South J. Evolution of life in urban environments. *Science*. 2017;358:6363.
20. Lai S, Johansson MA, Yin W, Wardrop NA, van Panhuis WG, Wesolowski A, et al. Seasonal and interannual risks of dengue introduction from South-East Asia into China, 2005-2015. *PLoS Negl Trop Dis*. 2018;12(11):e0006743.

21. Xiang B, Gao P, Kang Y, Ren T. Importation of Zika Virus in China: A significant risk in southern China. *J Infect.* 2017;74(3):328-30.
22. Wu F, Liu Q, Lu L, Wang J, Song X, Ren D. Distribution of *Aedes albopictus* (Diptera: Culicidae) in northwestern China. *Vector Borne Zoonotic Dis.* 2011;11(8):1181-6.
23. Zheng X, Zhong D, He Y, Zhou G. Seasonality modeling of the distribution of *Aedes albopictus* in China based on climatic and environmental suitability. *Infect Dis Poverty.* 2019;8(1):98.
24. Wu JY, Lun ZR, James AA, Chen XG. Dengue Fever in mainland China. *Am J Trop Med Hyg.* 2010;83(3):664-71.
25. Lei Luo L-YJ, and Zhi-Cong Yang. The dengue preface to endemic in mainland China: the historical largest outbreak by *Aedes albopictus* in Guangzhou, 2014. *Infectious Diseases of Poverty.* 2017;6:148.
26. Powell JR. Genetic Variation in Insect Vectors: Death of Typology? *Insects.* 2018;9(4):139.
27. Vieira ML, Santini L, Diniz AL, Munhoz Cde F. Microsatellite markers: what they mean and why they are so useful. *Genet Mol Biol.* 2016;39(3):312-28.
28. Porretta D, Gargani M, Bellini R, Calvitti M, Urbanelli S. Isolation of microsatellite markers in the tiger mosquito *Aedes albopictus* (Skuse). *Molecular Ecology Notes.* 2006;6(3):880-1.
29. Manni M, Gomulski LM, Aketarawong N, Tait G, Scolari F, Somboon P, et al. Molecular markers for analyses of intraspecific genetic diversity in the Asian Tiger mosquito, *Aedes albopictus*. *Parasit Vectors.* 2015;8:188.
30. Multini LC, Marrelli MT, Wilke AB. Microsatellite loci cross-species transferability in *Aedes fluviatilis* (Diptera:Culicidae): a cost-effective approach for population genetics studies. *Parasit Vectors.* 2015;8:635.
31. Chambers EW, Meece JK, McGowan JA, Lovin DD, Hemme RR, Chadee DD, et al. Microsatellite isolation and linkage group identification in the yellow fever mosquito *Aedes aegypti*. *J Hered.* 2007;98(3):202-10.
32. Bonacum, J., R. DeSalle, P. O'Grady, D. Olivera, J. Wintermute, and M. Zilversmit. New nuclear and mitochondrial primers for systematics and comparative genomics in Drosophilidae. *Dros. Inf. Serv.* 2001;84:201-4.
33. Park DS, Suh SJ, Hebert PD, Oh HW, Hong KJ. DNA barcodes for two scale insect families, mealybugs (Hemiptera: Pseudococcidae) and armored scales (Hemiptera: Diaspididae). *Bull Entomol Res.* 2011;101(4):429-34.
34. Shaikevich E, Talbalaghi A. Molecular Characterization of the Asian Tiger Mosquito *Aedes albopictus*(Skuse) (Diptera: Culicidae) in Northern Italy. *ISRN Entomology.* 2013;2013:1-6.
35. L MH-T, V AB, N IN, Ignacio R-A, Barrero E, Thorne L, et al. DNA barcoding of British mosquitoes (Diptera, Culicidae) to support species identification, discovery of cryptic genetic diversity and monitoring invasive species. *Zookeys.* 2019;832:57-76.

36. Sun F. Achieving suitable thermal performance in residential buildings in different climatic regions of China. *Energy and Buildings*. 2013;67:11-21.
37. Kang L, Zhang H. A Comprehensive Study of Agricultural Drought Resistance and Background Drought Levels in Five Main Grain-Producing Regions of China. *Sustainability*. 2016;8(4):1-12.
38. Baolin Lu and Houyong Wu. Classification and identification of important medical insects of China. 1st ed. Henan Science and Technology Publishing House; 2003.
39. Norbert Becker aDP. Mosquitoes: Identification, Ecology and Control. 3rd ed. Springer International Publishing; 2019.
40. Nagy S, Poczai P, Cernak I, Gorji AM, Hegedus G, Taller J. PICcalc: an online program to calculate polymorphic information content for molecular genetic studies. *Biochem Genet*. 2012;50(9-10):670-2.
41. Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P. micro-checker: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*. 2004;4(3):535-8.
42. Zhang H-R, Niu S-F, Wu R-X, Zhai Y, Tian L-T. Development and characterization of 26 polymorphic microsatellite markers in *Lateolabrax maculatus* and cross-species amplification for the phylogenetically related taxa. *Biochemical Systematics and Ecology*. 2016;66:326-30.
43. Excoffier L LH. An Integrated Software Package for Population Genetics Data Analysis. *Evol Bioinform Online [Internet]*. 2015;1(1):47-50.
44. Cristescu R, Sherwin WB, Handasyde K, Cahill V, Cooper DW. Detecting bottlenecks using BOTTLENECK 1.2.02 in wild populations: the importance of the microsatellite structure. *Conservation Genetics*. 2009;11(3):1043-9.
45. Earl DA, vonHoldt BM. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conservation Genetics Resources*. 2011;4(2):359-61.
46. Evanno G, Regnaut S, Goudet J. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol Ecol*. 2005;14(8):2611-20.
47. Jombart T. adegenet: a R package for the multivariate analysis of genetic markers. *Bioinformatics*. 2008;24(11):1403-5.
48. Mahesh BA, Kannan E, Davis GDJ, Venkatesan P, Ragunath PK. GenPop-An Online Tool to Analyze Human Population Genetic Data. *Bioinformation*. 2020;16(2):149-52.
49. Rozas J, Sanchez-DelBarrio JC, Messeguer X, Rozas R. DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics*. 2003;19(18):2496-7.
50. Frohlich F, Schmidt SL. Rational design of transcranial current stimulation (TCS) through mechanistic insights into cortical network dynamics. *Front Hum Neurosci*. 2013;7:804.
51. Murias dos Santos A, Cabezas MP, Tavares AI, Xavier R, Branco M. tcsBU: a tool to extend TCS network layout and visualization. *Bioinformatics*. 2016;32(4):627-8.
52. Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Mol Biol Evol*. 2012;29(8):1969-73.

53. Zhao, S., Guo, Y., Sheng, Q. *et al.*. Heatmap3: an improved heatmap package with more powerful and convenient features. *BMC Bioinformatics* 2014;15:16.
54. Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol.* 2003;52(5):696-704.
55. Kharzinova VR, Dotsev AV, Kramarenko AS, Layshev KA, Romanenko TM, Solov'eva AD, et al. Study of the allele pool and the degree of genetic introgression of semi-domesticated and wild populations of reindeer (*Rangifer tarandus* L., 1758) using microsatellites. *Sel'skokhozyaistvennaya Biologiya.* 2016;51(6):811-23.
56. Sébastien L^êe JJ, François Husson. FactoMineR: An R Package for Multivariate Analysis. *Journal of Statistical Software.* 2008;25(1):1-18.
57. Ditta A, Zhou Z, Cai X, Shehzad M, Wang X, Okubazghi K, et al. Genome-Wide Mining and Characterization of SSR Markers for Gene Mapping and Gene Diversity in *Gossypium barbadense* L. and *Gossypium darwinii* G. Watt Accessions. *Agronomy.* 2018;8(9):1-12.
58. Delatte H, Toty C, Boyer S, Bouetard A, Bastien F, Fontenille D. Evidence of habitat structuring *Aedes albopictus* populations in Reunion Island. *PLoS Negl Trop Dis.* 2013;7(3):e211.
59. Chapuis MP, Estoup A. Microsatellite null alleles and estimation of population differentiation. *Mol Biol Evol.* 2007;24(3):621-31.
60. Girard P. A robust statistical method to detect null alleles in microsatellite and SNP datasets in both panmictic and inbred populations. *Stat Appl Genet Mol Biol.* 2011;10:article9.
61. Rico C, Cuesta JA, Drake P, Macpherson E, Bernatchez L, Marie AD. Null alleles are ubiquitous at microsatellite loci in the Wedge Clam (*Donax trunculus*). *PeerJ.* 2017;5:e3188.
62. T. Lehmann, N. J. Besansky, W. A. Hawley, T. G. Fahey, L. Kamau, and F. H. Collins. Microgeographic structure of *Anopheles gambiae* in western Kenya based on mtDNA and microsatellite loci. *Molecular Ecology.* 1997;6:243-53.
63. Meglecz E, Petenian F, Danchin E, D'Acier AC, Rasplus JY, Faure E. High similarity between flanking regions of different microsatellites detected within each of two species of Lepidoptera: *Parnassius apollo* and *Euphydryas aurinia*. *Mol Ecol.* 2004;13 (6):1693-1700.
64. Chapuis M-P, Loiseau A, Michalakis Y, Lecoq M, Estoup A. Characterization and PCR multiplexing of polymorphic microsatellite loci for the locust *Locusta migratoria*. *Molecular Ecology Notes.* 2005;5(3):554-7.
65. Dakin EE, Avise JC. Microsatellite null alleles in parentage analysis. *Heredity (Edinb).* 2004;93(5):504-9.
66. Zhong D, Lo E, Hu R, Metzger ME, Cummings R, Bonizzoni M, et al. Genetic analysis of invasive *Aedes albopictus* populations in Los Angeles County, California and its potential public health impact. *PLoS One.* 2013;8(7):e68586.
67. Wei Y, Wang J, Song Z, He Y, Zheng Z, Fan P, et al. Patterns of spatial genetic structures in *Aedes albopictus* (Diptera: Culicidae) populations in China. *Parasit Vectors.* 2019;12 (1):552.

68. Rafael A. Marinho EBB, Maria A. Bezerra-Gusmão, Valbia de S. Porto, Ricardo A. Olinda, and Carlos A. C. dos Santos. Effects of temperature on the life cycle, expansion, and dispersion of *Aedes aegypti* (Diptera: Culicidae) in three cities in Paraíba, Brazil. *Journal of Vector Ecology*. 2016;41(1):10.
69. Mogi M, Armbruster PA, Tuno N, Aranda C, Yong HS. The Climate Range Expansion of *Aedes albopictus* (Diptera: Culicidae) in Asia Inferred From the Distribution of *Albopictus* Subgroup Species of *Aedes* (Stegomyia). *J Med Entomol*. 2017;54(6):1615-25.
70. Jia P, Liang L, Tan X, Chen J, Chen X. Potential effects of heat waves on the population dynamics of the dengue mosquito *Aedes albopictus*. *PLoS Negl Trop Dis*. 2019;13(7): e0007528.
71. Naoko Takezaki aMN. Genetic Distances and Reconstruction of Phylogenetic Trees From Microsatellite DNA. *GENETICS*. 1996;144(1):389-99.
72. Wondji C, Simard F, Lehmann T, Fondjo E, Same-Ekobo A, Fontenille D. Impact of insecticide-treated bed nets implementation on the genetic structure of *Anopheles arabiensis* in an area of irrigated rice fields in the Sahelian region of Cameroon. *Mol Ecol*. 2005;14(12):3683-9.
73. Chen H, Li K, Wang X, Yang X, Lin Y, Cai F, et al. First identification of *kdr* allele F1534S in VGSC gene and its association with resistance to pyrethroid insecticides in *Aedes albopictus* populations from Haikou City, Hainan Island, China. *Infect Dis Poverty*. 2016;5:31.
74. Li Y, Xu J, Zhong D, Zhang H, Yang W, Zhou G, et al. Evidence for multiple-insecticide resistance in urban *Aedes albopictus* populations in southern China. *Parasit Vectors*. 2018;11(1):4.
75. Su X, Guo Y, Deng J, Xu J, Zhou G, Zhou T, et al. Fast emerging insecticide resistance in *Aedes albopictus* in Guangzhou, China: Alarm to the dengue epidemic. *PLoS Negl Trop Dis*. 2019;13(9):e0007665.
76. Liu H, Liu L, Cheng P, Yang L, Chen J, Lu Y, et al. Bionomics and insecticide resistance of *Aedes albopictus* in Shandong, a high latitude and high-risk dengue transmission area in China. *Parasit Vectors*. 2020;13(1):11.
77. Kotsakiozi P, Richardson JB, Pichler V, Favia G, Martins AJ, Urbanelli S, et al. Population genomics of the Asian tiger mosquito, *Aedes albopictus*: insights into the recent worldwide invasion. *Ecol Evol*. 2017;7(23):10143-57.
78. Schmidt TL RiG, Zhang D, Zheng X, Xi Z, Hoffmann AA. Genome-wide SNPs reveal the drivers of gene flow in an urban population of the Asian Tiger Mosquito, *Aedes albopictus*. *PLOS Neglected Tropical Diseases*. 2017;11(10):e0006009.
79. Guo Y, Song Z, Luo L, Wang Q, Zhou G, Yang D, et al. Molecular evidence for new sympatric cryptic species of *Aedes albopictus* (Diptera: Culicidae) in China: A new threat from *Aedes albopictus* subgroup? *Parasit Vectors*. 2018;11(1):228.
80. Ruiling Z, Peien L, Xuejun W, Zhong Z. Molecular analysis and genetic diversity of *Aedes albopictus* (Diptera, Culicidae) from China. *Mitochondrial DNA A DNA Mapp Seq Anal*. 2018;29(4):594-9.
81. David Roiz RR, Daniele Arnoldi, and Annapaola Rizzoli. Effects of Temperature and Rainfall on the Activity and Dynamics of Host-Seeking *Aedes albopictus* Females in Northern Italy. *VECTOR-BORNE*

AND ZOONOTIC DISEASES. 2010;10(8):811-6.

82. Latreille A.C., Milesi P, Magalon H. et al. High genetic diversity but no geographical structure of *Aedes albopictus* populations in Réunion Island. *Parasites Vectors*.2019;12: 1-12.
83. EunJung Lee, Seong-Chan Yang, Tae-Kyu Kim et al. Geographical Genetic Variation and Sources of Korean *Aedes albopictus* (Diptera: Culicidae) Populations. *Journal of Medical Entomology*.2020; 57(4) :1057-68.

Additional Files

Additional file 1: Table S1. Sampling information of 17 *Ae. albopictus* populations collected from three different climatic zones of China.

Additional file 2: Table S2. Heterozygosity tests of all 17 *Ae. albopictus* populations based on SMM model.

Additional file 3: Table S3. Haplotype diversity of 17 *Ae. albopictus* populations based on *coxI* collected from three different climatic regions of China.

Additional file 4: Table S4. Population differentiation estimation of the F_{ST} value (below the diagonal) and Geographic distance (above the diagonal) between all 17 *Ae. albopictus* populations.

Additional file 5: Figure S1. Linkage disequilibrium analysis at each pair of Loci across all 17 *Ae. albopictus* populations.

Additional file 6: Figure S2. Isolation by distance analysis of all 17 *Ae. albopictus* populations.

Additional file7: Figure S3. Historical demography analysis of *Ae. albopictus* inferred from mtDNA *coxI* sequences.

Additional file 8: Table S5. Genetic diversity of 11 microsatellite loci developed for *Ae. albopictus* based on samples(n=502) collected from three different climatic regions of China.

Additional file9: Figure S4. Migration analysis of all 17 *Ae. albopictus* populations inferred from microsatellite data.

Tables

Table 1 Information of microsatellite primers used for genotyping all the 17 *Ae. albopictus* populations.

No.	Primer name	Forward sequence (5'-3')	Reverse sequence (5'-3')	Repeat Motifs	Size range (bp)	Dye	GenBank No.
1	BW-P1	TTAGCATCCATCTATTCTGGC	AAACATTCCTACGCATTTCCAC	(GT)6	230-260	5'- HEX	MT642446
2	BW-P3	GAAAATATGGTCTATCAAATG	AAGTCAGTAAAACAGGAGTCT	(GT)3GC(GT)3TTT	132-178	5'- FAM	MT642447
3	BW-P6	GAATTGGGAGCTTGGTAAAAC	CGCCTACTTGAGAAACTGA	(TG)5	124-200	5'- HEX	MT642448
4	BW-P18	CACTGGTTCTCTATCGAATGC	GTGTTATCAGTTGGAAGCGTT	(GT)24	158-203	5'- HEX	MT642449
5	BW-P22	GGCGTCCCCCAACATACATC	CAGCTCCGTCCTCTCTTCCC	(GT)5(GCGT)2(GT)3	187-244	5'- HEX	MT642450
6	BW-P23	GGATAAGAATGACACAGGCAC	CAAAGAGGAACACCATAGGAA	(GAC)7	133-177	5'- FAM	MT642451
7	BW-P24	ACGAAACATACTTACAATTGCA	AACCTAGAGTCCGAGAGAGAAC	(AC)8	145-239	5'- FAM	MT642452
8	BW-P26	CGTGGTGGTTAGGTCCATGTT	TCGCTTTTCGGCTCTAGTCAAT	(GT)5	107-233	5'- HEX	MT642453
9	BW-P27	TTATACAAAAAGCGAACATCC	CACACACATAGAAAAAAGCAA	(ACG)6	249-281	5'- FAM	MT642454
10	BW-P35	TATTTGCACATCCATTTCTGCT	TTCAAAACCTGATTTCCGACTG	(CA)7TTT	83-120	5'- HEX	MT642455
11	BW-P36	GTCATGTAGTCTCACAGTCAC	ATATGGATCATAGATGATGGAG	(AC)6CCC	159-179	5'- FAM	MT642456

Table 2 Genetic diversity indices for 17 *Ae. albopictus* populations investigated by 11 microsatellite loci.

Regions	SC	SS	n _a	n _e	F _{IS}	Ho	He	HWE
Tropical region	HKWN	30	8.455	4.525	0.388	0.561	0.733	0.442***
	JYJB	30	7.636	4.355	0.385	0.584	0.699	0.535***
	JKCH	25	7.000	3.553	0.292	0.532	0.671	0.246***
	Mean	29	7.697	4.144	0.355	0.559	0.701	0.408***
South-subtropical region	NNXD	30	7.909	4.000	0.336	0.533	0.700	0.347**
	NNXZ	30	7.909	4.190	0.473	0.627	0.725	0.480***
	GZTH	30	8.455	4.292	0.172	0.467	0.714	0.192*
	Mean	30	8.091	4.161	0.327	0.542	0.713	0.340**
North-subtropical region	NJTH	30	7.455	3.836	0.392	0.590	0.682	0.473***
	NJDX	30	8.455	4.119	0.465	0.637	0.706	0.551***
	KZXZ	30	6.182	3.681	0.189	0.474	0.677	0.331
	SHJD	30	5.455	3.106	0.035	0.455	0.606	0.123
Temperate region	Mean	30	6.909	3.677	0.266	0.542	0.660	0.369*
	HNDX	30	7.000	3.642	0.247	0.553	0.627	0.365**
	QDDX	30	7.909	3.921	0.361	0.570	0.664	0.380**
	BHBG	22	8.091	3.838	0.433	0.641	0.637	0.447***
	BJLG	30	8.636	4.341	0.408	0.574	0.728	0.456***
	ZGND	30	6.909	3.728	0.425	0.631	0.636	0.419***
	SXJW	30	6.455	3.501	0.270	0.572	0.622	0.421**
HBSD	30	7.546	3.926	0.256	0.508	0.687	0.341**	
Mean	29	7.591	3.876	0.359	0.583	0.662	0.411**	

SC: Sample Code; SS: Sample Size; n_a: Observed number of alleles; n_e: Effective number of alleles (Kimura and Crow,1964); HWE: Hardy-Weinberg disequilibrium; Ho: Observed Heterozygosity; He: Expected Heterozygosity (Nei's,1973)

***: $p < 0.001$; **: $p < 0.01$; *: $p < 0.05$

Table 3 Analysis of molecular variance (AMOVA) test of 17 *Ae. albopictus* populations sampled from three climatic regions.

Source of Variation	d. f.	Sum of Squares	Variance Components	Percentage of Variation	P-value	Fixation indices
Among groups	5	142.425	0.12789Va	3.62	$P < 0.00001$	$F_{CT} = 0.03620$
Among populations within groups	11	93.157	0.06845Vb	1.94	$P < 0.00001$	$F_{SC} = 0.02010$
Among individuals within populations	485	2156.367	1.10951Vc	31.40	$P < 0.00001$	$F_{IS} = 0.33253$
Within individuals	502	1118.000	2.22709Vd	63.04	$P < 0.00001$	$F_{IT} = 0.36962$
Total	1003	3509.949	3.53294			

Figures




Figure 1

Sampling map and Migration patterns for *Ae. albopictus* among three different climatic regions of China. Three climatic regions were marked with different colors, four major destinations were marked with stars of different colors, and migration routes were marked with different types of arrows. Note: The

designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.




Figure 2

The population structure analysis of all 17 *Ae. albopictus* populations based on 11 microsatellite loci. (a) K values assessed via Evanno et al.'s ΔK methods; (b) STRUCTURE bar plots ($k=2$); (c) Bayesian clustering analysis of all *Ae. albopictus* populations; (d) DAPC analysis of all *Ae. albopictus* populations, and 86.4% of variation was explained by 50 PCs.




Figure 3

The Bayesian and Haplotype network analysis of the representing 25 haplotypes among all 17 *Ae. albopictus* populations based on *cox1* sequences. (a) Bayesian tree of the representing 25 haplotypes was constructed via BEAST version 1.8.4 with the best model “HKY+G” selected by JModelTest 2.1.10; (b) TCS haplotype network for the *cox1* gene of all *Ae. albopictus* individuals (n=502) from different climatic regions of China. The sizes of circles are proportional to haplotype frequency, and each line segment represents a single mutation.




Figure 4

Multiple factor analysis (MFA) and Hierarchical clustering performed on 17 *Ae. albopictus* populations sampled from three different climatic regions. (a) Multiple factor analysis between genetic indices of all individuals and three environmental variables (temperature, rainfall and altitude) of 17 *Ae. albopictus* populations, populations of different clusters were marked with different colors; (b) Hierarchical clustering of all 17 *Ae. albopictus* populations on the factor map.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalFile1TableS1.doc](#)
- [Additionalfile9FigureS4.tif](#)
- [Graphicalabstract.tif](#)