

# Epidemiology of COVID-19 in Yemen: A Descriptive Study

Redhwan Al-Naggar (✉ [profredhwan@gmail.com](mailto:profredhwan@gmail.com))

Al-Hikmah University - Faculty of Medicine <https://orcid.org/0000-0002-6448-6930>

Lutfi Almaktari

Faculty of Medicine and Health Sciences, University of Sana'a

Salem Madram

Medical Laboratory Center, Aden

Hisham Al-shaikhli

Auckland University of Technology

---

## Research article

**Keywords:** COVID-19, Yemen, Epidemiology

**Posted Date:** July 17th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-38912/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

## **Abstract**

In Yemen, the first COVID-19 confirmed case was reported on 10 April 2020. The OBJECTIVE of this study is to identify the epidemiological characteristics of COVID-19 patients in Yemen. METHODOLOGY: From May 20 to 31, 2020, the data were obtained from the National Center of Public Health Laboratories, Ministry of Health and population, Aden, Yemen. A retrospective study used to determine the epidemiological characteristics of the first 53 confirmed cases of patients with COVID-19. This data includes the date of diagnosis, gender, age, governorates, and clinical symptoms. Data analyzed using SPSS version 22. RESULTS: A total number of 53 confirmed cases were obtained from the National Center of Public Health Laboratories, Ministry of Health and population, Aden, Yemen. The majority of the participants were male (72%), their age were less than 39 years (40%) and from Aden governorate (43%). The commonest symptoms reported among the participants were fever (%), sore throat and cough. Conclusion: Male, less than 39 years old and from Aden is the trend of COVID-19 in Yemen. The findings of this study may help provide guidance for frontline medical staff in the clinical management of the outbreak.

## **Introduction**

Coronaviruses are a family of viruses transmitted through mammals and birds, with humans being particularly susceptible to infection and transmission of the virus (1). The earlier two coronaviruses outbreaks are the Middle East Respiratory Syndrome Coronavirus (MERS) and Severe Acute Respiratory Syndrome (SARS) in 2015 and 2003; respectively and another outbreak reported in China in December 2019 which showed similarities to the novel coronavirus and is now a pandemic disease affected most countries worldwide (2). WHO declared on January 30, 2020, declared COVID-19 is a global public health concern and collaborative efforts of all countries to prevent the rapid spread of COVID-19 are needed (3).

The main known transmission method of coronavirus is from person-to-person through sneezing or coughing, touching a contaminated surface (4). The clinical features of COVID-19 are cough, fever, fatigue, pneumonia, and shortness of breath (5, 6). There is no particular antiviral treatment or vaccine so far (4). Hence, the prevention measures should be practiced firmly such as physical distancing, wearing face masks, and wear eye protection are associated with a great reduction in COVID-19 infection. In addition to hand hygiene practices to reduce the spread of coronavirus infection (5).

WHO reported that 80% of COVID-19 patients with mild symptoms recovered without any medication showed mild symptoms and. About 20% of COVID-19 infected cases showed serious symptoms such as shortness of breath, multi-organ failure and septic shock. Around 2% of cases can be fatal (3). Recent studies suggested that COVID-19 infected those with a weak immune system such as obese, older males, with chronic comorbidities, HIV patients, pregnant women, and long-term users of immunosuppressive medications (7–10). According to a recent systematic review reported that the most common comorbidities affected by COVID-19 are cardiovascular diseases, hypertension and diabetes(11).

The recent mortality rate of COVID-19 infection is predictable to be approximately 4.8% throughout the globe (12). Different mortality rates were recorded in different countries, in Germany was 1%, and Italy 11.4% (12). The possible explanation is due to the overestimated mortality rate and health authorities selected severe cases but not mild cases and no symptoms cases. Although overall fatality rate for clinical cases ranged from 2–5% globally than the previous outbreaks related to viruses such as H7N9 (39.0%), MERS (34.4%), and SARS (9.5%) pathogens

continue to emerge and spread to the population at risk, hence there is an urgent need to move from responsiveness nations to proactive management (6, 13).

The greatest way of prevention is not exposing to COVID-19 through washing hands using soap and water, wearing face masks, quarantining suspected and confirmed cases(4, 14). Prevention measures are very important and curtail especially there is no vaccine available yet (15). Some measures were taken in the whole country to minimize the exposure and transmission of COVID-19 by closing schools, universities, and activities of large crowded. In Yemen, schools, universities and wedding halls were closed before any COVID-19 confirmed cases reported and the flight to the main airport [Aden] also closed as a prevention measure. WHO endorses the main personal protective actions against the COVID-19 which washes your hands regularly, maintain physical distancing, no touching the nose, eyes, and mouth, practice respiratory hygiene and if sick stay home (14).

Yemen's health system is already greatly breakable due to the conflict for more than 5 years. In Yemen, the first confirmed case was reported on 10 April 2020 in Hadhramout, Yemen (16). The challenges that faced Yemen to reduce COVID-19 cases are many, one is a very limited number of national laboratories and the lack of PCR reagents and machines for COVID-19 detection. Second, the difficulties of implementations of international health strategies of the COVID-19 outbreak due to the civil war and conflict. The government also failed to formulate enough isolation locations at the country entry. Third, due to the long and uncontrollable coastal borders that some refugees and immigrants may enter the country without COVID-19 testing (17). WHO representative in Yemen reported that approximately 65000 deaths and 494000 will be hospitalized will occur in Yemen if there are no protection measures that take place at the right time. WHO representative added that it is extremely alarming circumstances if the Yemeni people do not make thoughtful changes in their behavior to face the COVID-19 infection (18). Therefore, the objective of this study is to identify the epidemiological characteristics of COVID-19 patients in Yemen.

## Methodology

From May 20 to 31, 2020, the data were obtained from the National Center of Public Health Laboratories, Ministry of Health and population, Aden, Yemen. A retrospective study used to determine the epidemiological characteristics of the first 53 confirmed cases of patients with COVID-19 in Aden, Yemen. Confirmed cases defined as polymerase chain reaction assay (PCR) positive. Only cases included in this study are laboratory positive cases. The ethics committees of Al-Hikma University, Sana'a-Yemen approved this study. The data were obtained from the National Center of Public Health Laboratories, Ministry of Health and population, Aden, Yemen. This data includes the date of diagnosis, gender, age, governorates, and clinical symptoms. Data were key-in into SPSS software in the principal investigator computer and protected with a PIN at Al-Hikma Research Center, Al-Hikma University, Sana'a-Yemen. Data cleaning and validation were accomplished before the data analysis. Analytical datasets were created in an anonymized manner and categorical variables into percentages using SPSS version 20.

## Results

A total number of 53 confirmed cases were obtained from the National Center of Public Health Laboratories, Ministry of Health and population, Aden, Yemen. This data collected from 20 May 2020 until 31 May 2020. The majority of the participants were male (72%), their age was less than 39 years (40%) and from the Aden governorate (43%). The commonest symptoms reported among the participants were fever and cough (90%), followed by the sore throat (81%) (Table 1).

**Table 1**  
**Socio-demographic characteristic and symptoms  
of 53 COVID-19 confirmed cases in Yemen (N =  
53)**

<b>Characteristics</b>	<b>Confirmed cases</b>	<b>N (%)</b>
<b>Age</b>	≤ 39	21(40%)
	40–49	13(25%)
	50–59	7(13%)
	60–69	6 (12%)
	≥ 70	5(10%)
	Missing	1()
<b>Sex</b>	Male	38(72%)
	Female	15(28%)
<b>Governorates</b>	Aden	23(43%)
	Abyan	8(15%)
	Lahj	16(30%)
	Adhale	4(8%)
	Missing	2(4%)
<b>Symptoms</b>	Fever	48(90%)
	Sore throat	43(81%)
	cough	48(90%)

The number of cases confirmed on the 20th of May 2020 was one case, three days later the confirmed cases jumped into 14 cases. The confirmed cases then dropped again on 24th of May 2020, with two confirmed cases, 5 days later the confirmed cases jumped again to 14 cases. These may due to the behavior of the residents and their carelessness [Figure 1].

## Discussion

Coronaviruses are large, RNA viruses, and enveloped (19). Coronaviruses considered insignificant pathogens until the early twenty-first century where we experienced two major outbreaks from MERS and SARS. As of May 28, 2020, 5·85 million people infected with COVID-19 and 359000 deaths worldwide (20). To the best of our knowledge, this is the first epidemiological study about COVID-19 in Yemen. This analysis of early confirmed COVID-19 cases in Aden, Yemen provides insight into the epidemiological characteristics and clinical features. Cases are confirmed using PCR testing which is a universal golden standard.

Our study showed that the number of men is greater than women of COVID-19 infection (72%). Similar findings reported by previous studies reported that COVID-19 infection was more likely to affect males (6, 21). A previous study indicated that 56% of infection with COVID-19 were males (2). Another study found that 50.7% were males

(22). An Italian study reported that men represent 58% of COVID-19 infected patients and 70% of COVID-19 related deaths (23). SARS has infected more men than women (24). The possible explanation is that fewer females infected with COVID-19 may due to the X chromosome protection, in addition to sex hormones which reported to have a significant role in the immunity of women (10, 25, 26). Other possible factors such as behavioral and cultural habits have also contributed such as smoking in particular has been implicated as a significant contributor to disease severity (27, 28). Overall, men are more likely to involve health-related risks (29).

The mean age of confirmed cases of COVID-19 patients in our study is **44.4**. The mean age in Yemen is lower than several studies reported in systematic review which showed that the mean age of COVID-19 patients was **52.0 years** (30). Also lower than the mean age of COVID-19 patients presenting in Chinese study was 46.5 years (24). The highest affected age group was those  $\leq$  39 years old. Our findings supported by a previous study reported that COVID-19 infection affects mostly those in the age group of 30–65 (31). Similar findings reported in previous studies during SARS outbreaks that age was a very important factor in both mortality and morbidity of SARS infection (32). United Nation reported that the growth rate of the Yemeni population is one of the highest in the world, at 3%. Approximately 50% of the Yemeni population is below 15 years of age (33).

In this study, the highest number of COVID-19 confirmed cases were recorded in Aden governorate. The possible explanation is that Aden is the temporary capital of Yemen. It's the only international entry of the country by Air. There may be some imported cases that came with the local citizens who preferred to back home during this pandemic that hit the whole world. On 11 May 2020, Aden city which is the interim capital of Yemen declared "infested city" after the flooding hit Aden which contributed to the spread of diseases like Dengue, Malaria, Cholera and the new disease COVID-19 (34). In addition to the governmental and political conflicts in Aden obstructing national and international efforts to fight the new pandemic COVID-19 infection.

Since we are only able to access to the Aden data from the National Center of Public Health Laboratories, Ministry of Health and population, Aden, Yemen. Furthermore, we only got cases from the 20th of May until the 31st of May 2020. The number of cases confirmed on the 20th of May 2020 was one case, three days later the confirmed cases jumped into 14 cases. The confirmed cases then dropped again on 24th of May 2020, with two confirmed cases, 5 days later the confirmed cases jumped again to 14 cases. These may due to the behavior of the residents and their carelessness. Furthermore, most people not able to buy the masks, not able to stay home because of the nature of their work which is a daily earned income.

The common symptoms of COVID-19 infections included fever, shortness of breath and cough, sometimes these followed by headache, sore throat, insomnia, frequent urination and nasal congestion. Some severe cases experienced GIT symptoms such as diarrhea, nausea, and vomiting. Our study also reported that 90% of the COVID-19 confirmed cases suffer from fever, cough (90%), followed by the sore throat (81%). Similar findings reported in previous studies that 90% of hospitalized patients were suffering from fever (5, 35). Similar findings reported in Chinese study where they reported that fever is the most common symptom (91.7%) then cough (75%) (22).

The limitation of this study is the small sample size with only 53 patients with confirmed COVID-19 were included. Obtaining data from other centers was not possible at this time due to political reasons. More data about the clinical outcome of the COVID-19 was not available at this moment, though, the available data in our study documented an early epidemiological assessment of COVID-19 in Yemen. Patient comorbidities were also not documented. Another limitation is that this study conducted in a short duration and limited by Aden as geographical location is due to some political and logistics reasons. Despite the limitation of our study, the study

might offer guidance for policymakers in Yemen and medical staff to manage this outbreak and specific conclusions can be drawn from this study and further studies are needed to be built in this study. More epidemiological studies in multicentre in Yemen are urgently needed which can empower epidemiologists to build the COVID-19 model for the Yemen context.

## Declarations

### **Ethics approval and consent to participate:**

The ethics committees of Al-Hikma University, Sana'a-Yemen approved this study.

### **Consent for publication:**

Not applicable

### **Availability of data and materials:**

The data were obtained from the National Center of Public Health Laboratories, Ministry of Health and population, Aden, Yemen. The datasets generated during and/or analysed during the current study are not publicly available because the National Center of Public Health Laboratories don't have online database but this data are available from the corresponding author on reasonable request.

### **Competing interests:**

The authors declare that they have no competing interests

### **Funding:**

No funding was available for this study.

### **Authors' contributions:**

RA Al-Naggar : Study design, analysis of the data and was a major contributor in writing the manuscript

L Almaktari: Collecting the data and proofreading

S Madram: Study planning and help in writing

H Alshaikhli: Help in writing, proofreading, referencing and arrange for submission

### **Acknowledgements:**

Not applicable


## References

1. Schoeman D, Fielding BC. Coronavirus envelope protein: current knowledge. *Virology journal*. 2019;16(1):69.
2. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus–infected pneumonia. *New England Journal of Medicine*. 2020.

3. WHO. WHO Director-General's opening remarks at the mission briefing on COVID-19. 2020: World Health Organization 2020 [Available from: <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-mission-briefing-on-covid-19>].
4. CDC. How COVID-19 Spreads USA: National Center for Immunization and Respiratory Diseases (NCIRD), Division of Viral Diseases; 2020 [Available from: [https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/how-covid-spreads.html?CDC\\_AA\\_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fprepare%2Ftransmission.html](https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/how-covid-spreads.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fprepare%2Ftransmission.html)].
5. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus–infected pneumonia in Wuhan, China. *Jama*. 2020;323(11):1061-9.
6. Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *The Lancet*. 2020;395(10223):507-13.
7. Wang XF, Shi GC, Wan HY, Hang SG, Chen H, Chen W, et al. Clinical features of three avian influenza H7N9 virus-infected patients in Shanghai. *The clinical respiratory journal*. 2014;8(4):410-6.
8. Badawi A, Ryoo SG. Prevalence of comorbidities in the Middle East respiratory syndrome coronavirus (MERS-CoV): a systematic review and meta-analysis. *International Journal of Infectious Diseases*. 2016;49:129-33.
9. Channappanavar R, Fett C, Mack M, Ten Eyck PP, Meyerholz DK, Perlman S. Sex-based differences in susceptibility to severe acute respiratory syndrome coronavirus infection. *The Journal of Immunology*. 2017;198(10):4046-53.
10. Jaillon S, Berthenet K, Garlanda C. Sexual Dimorphism in Innate Immunity. *Clinical Reviews in Allergy & Immunology*. 2019;56(3):308-21.
11. Yang A, Qiu Q, Kong X, Sun Y, Chen T, Zuo Y, et al. Clinical and Epidemiological Characteristics of COVID-19 Patients in Chongqing China. *Frontiers in Public Health*. 2020;8:244.
12. BNO. Tracking coronavirus: Map, data and timeline 2020 [Available from: <https://bnonews.com/index.php/2020/04/the-latest-coronavirus-cases/>].
13. Munster VJ, Koopmans M, van Doremalen N, van Riel D, de Wit E. A novel coronavirus emerging in China—key questions for impact assessment. *New England Journal of Medicine*. 2020;382(8):692-4.
14. WHO. How to prevent infection and to slow transmission of COVID-19: World Health Organization 2020a [Available from: [https://www.who.int/health-topics/coronavirus#tab=tab\\_2](https://www.who.int/health-topics/coronavirus#tab=tab_2)].
15. Qualls N, Levitt A, Kanade N, Wright-Jegede N, Dopson S, Biggerstaff M, et al. Community mitigation guidelines to prevent pandemic influenza—United States, 2017. *MMWR Recommendations and Reports*. 2017;66(1):1.
16. OCHA. The Ministry of Health in Aden confirms the first case of COVID-19 in Yemen: UN HC Yemen 2020 [Available from: <https://reliefweb.int/report/yemen/ministry-health-aden-confirms-first-case-covid-19-yemen-enar>].
17. Dureab F, Al-Awlaqi S, Jahn A. COVID-19 in Yemen: preparedness measures in a fragile state. *The Lancet Public Health*. 2020.
18. Devi S. Fears of “highly catastrophic” COVID-19 spread in Yemen. *The Lancet*. 2020;395(10238):1683.
19. de Wilde AH, Snijder EJ, Kikkert M, van Hemert MJ. Host factors in coronavirus replication. Roles of Host Gene and Non-coding RNA Expression in Virus Infection: Springer; 2017. p. 1-42.

20. Worldometer. COVID-19 coronavirus pandemic 2020 [Available from: <https://www.worldometers.info/coronavirus/>].
21. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The lancet.* 2020;395(10223):497-506.
22. Zhang J-j, Dong X, Cao Y-y, Yuan Y-d, Yang Y-b, Yan Y-q, et al. Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy.* 2020.
23. Remuzzi A, Remuzzi G. COVID-19 and Italy: what next? *The Lancet.* 2020.
24. Yang Y, Lu Q, Liu M, Wang Y, Zhang A, Jalali N, et al. Epidemiological and clinical features of the 2019 novel coronavirus outbreak in China. *MedRxiv.* 2020.
25. Sue K. The science behind “man flu”. *bmj.* 2017;359.
26. Schurz H, Salie M, Tromp G, Hoal EG, Kinnear CJ, Möller M. The X chromosome and sex-specific effects in infectious disease susceptibility. *Human genomics.* 2019;13(1):2.
27. Cai H. Sex difference and smoking predisposition in patients with COVID-19. *The Lancet Respiratory Medicine.* 2020;8(4):e20.
28. Statista. Number of individuals who smoke in Italy in 2018, by age and gender Italy: Statista Research Department; 2020 [Available from: <https://www.statista.com/statistics/501615/italy-smokers-by-age-and-gender/>].
29. Harris CR, Jenkins M. Gender differences in risk assessment: why do women take fewer risks than men? 2006.
30. Rodriguez-Morales AJ, Cardona-Ospina JA, Gutiérrez-Ocampo E, Villamizar-Peña R, Holguin-Rivera Y, Escalera-Antezana JP, et al. Clinical, laboratory and imaging features of COVID-19: A systematic review and meta-analysis. *Travel medicine and infectious disease.* 2020;101623.
31. Yao Y, Tian Y, Zhou J, Ma X, Yang M, Wang S. Epidemiological characteristics of SARS-CoV-2 infections in Shaanxi, China by 8 February 2020. *European Respiratory Journal.* 2020;55(4).
32. Cowling BJ, Muller MP, Wong IO, Ho L-M, Lo S-V, Tsang T, et al. Clinical prognostic rules for severe acute respiratory syndrome in low-and high-resource settings. *Archives of internal medicine.* 2006;166(14):1505-11.
33. UN. World Population Prospects  
The 2010 Revision New York: Department of Economic and Social Affairs Population Division; 2011 [Available from:  
[https://www.un.org/en/development/desa/population/publications/pdf/trends/WPP2010/WPP2010\\_Volume\\_I\\_Comprehensive-Tables.pdf](https://www.un.org/en/development/desa/population/publications/pdf/trends/WPP2010/WPP2010_Volume_I_Comprehensive-Tables.pdf)].
34. National T. Yemen's Aden declared 'infested city' as coronavirus spreads. *The National* 2020.
35. Wang Y, Wang Y, Chen Y, Qin Q. Unique epidemiological and clinical features of the emerging 2019 novel coronavirus pneumonia (COVID-19) implicate special control measures. *Journal of medical virology.* 2020;92(6):568-76.

## Figures



**Figure 1**

The frequency of cases per day during the study period