

Trajectories of loneliness and objective social isolation and associations between persistent loneliness and self-reported personal recovery in a cohort of secondary mental health service users in the UK

Ruimin Ma

University College London

Jingyi Wang

University College London

Brynmor Lloyd-Evans

University College London

Louise Marston

University College London

Sonia Johnson (✉ s.johnson@ucl.ac.uk)

University College London

Research Article

Keywords: Loneliness, social isolation, mental health, personal recovery, psychiatry

Posted Date: April 13th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-393505/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Loneliness is a frequent and distressing experience among people with mental health problems. However, few longitudinal studies have so far investigated the trajectories of loneliness and objective social isolation, and the extent to which both issues may impact mental health outcomes among mental health service users. Therefore, this study aims to describe the trajectories of loneliness and objective social isolation and their associations with self-rated personal recovery among people leaving crisis resolution teams (CRTs).

Methods: A total of 224 participants receiving care from CRTs (recruited for a large multi-site randomised controlled trial) were included in this longitudinal cohort study. They completed the eight-item University of California at Los Angeles Loneliness Scale (ULS-8), Lubben-Social Network Scale (LNSN-6), and the Questionnaire about the Process of Recovery (QPR) (primary outcome) at baseline, 4- and 18-month follow-up, as well as baseline sociodemographic and clinical variables.

Results: We compared groups who were persistently lonely (at all time points), intermittently lonely (at one or two time points) and never lonely. After adjusting for all potential confounders and baseline predictive variables, persistent severe loneliness was associated with worse personal recovery at 18-month follow-up compared with the never lonely (reference group) (coef. = -12.8, 95% CI -11.8, -3.8, $p < .001$), as was being intermittently lonely (coef. = -7.8, 95% CI -18.8, -6.8, $p < .001$). The persistently objectively socially isolated group (coef. = -9.8, 95% CI -15.7, -3.8, $p = .001$) also had poorer self-rated recovery at 18-month follow-up than those who were not socially isolated at any timepoint (i.e., reference category).

Conclusion: Results suggest that both persistent loneliness and objective social isolation are associated with poorer self-rated recovery following a crisis, compatible with a causal relationship. These findings suggest a potential role for interventions aimed at alleviating loneliness and objective social isolation in improving recovery outcomes for people with mental health symptoms. Increased awareness of both issues among health practitioners is also warranted.

Introduction

Social relationships are fundamental to individuals' emotional, behavioural and cognitive development [1]. Peplau and Perlman [2] define loneliness as a distressing experience that results from a perceived discrepancy between an individual's desired and actual social relationships. Loneliness is thus a subjective experience [3]. Objective social isolation describes the quantitative aspect of individuals' social relationships, such as the number of social relationships or the frequency of social contacts [4]. Wang and colleagues' [5] conceptual review put forward a comprehensive framework on loneliness and its related constructs in relation to mental health, including this distinction between loneliness and objective social isolation.

Multiple studies have shown a significant impact of loneliness and objective social isolation on our mental health, including depressive symptoms [6], onset of psychotic symptoms [7], and decline in cognitive functioning [8]. Loneliness is a more pervasive problem among people with most mental health conditions than among the general population [9]. Objective social isolation (e.g., having fewer social contacts) is also commonly experienced by people with mental disorders [10]. The underlying causes of loneliness and objective social isolation in people with mental health problems are likely to be multifactorial, including social anxiety or difficulties in initiating and maintaining social relationships with others [11, 12], and discrimination and interpersonal stigma toward people with mental health symptoms [13, 14].

Supporting personal recovery is increasingly acknowledged internationally as a key goal of mental health care and mental health policy [15, 16]. In mental health, the term 'recovery' now tends to be multidimensional; it focuses on mental health service users' personal experiences and personal goals for recovery [17]. The relational dimension of 'recovery' underscores the importance of maintaining interpersonal relationships and social contacts with family and friends [18]. While recent studies suggest a significant association between loneliness, poor social support and personal recovery among mental health service users [19], there is not as yet much evidence exploring longitudinal aspects of such associations and the likely direction of causality. Loneliness often is a transient experience; however, recent research has given grounds for particular concern about a sub-group of people who experience persistent loneliness, potentially related to factors such as maladaptive cognitive biases, social vigilance and specific attribution styles come into play [20]. Taking into consideration the association between loneliness, objective social isolation and a wide range of mental health outcomes, we should expect that being lonely or objectively socially isolated for an extended period of time may contribute to worse clinical outcomes in mental health service users. Yet, no research to date has investigated the importance of persistent loneliness or objective social isolation in mental health outcomes across a range of mental health diagnoses.

Given this knowledge gap, the aim of the current study was to advance our understanding of the persistence of loneliness and objective social isolation in a sample of people with mental health diagnoses, and whether people with persistent loneliness or objective social isolation have poorer personal recovery following recovery from a crisis. We studied a sample being discharged from crisis resolution teams (CRTs), which provide intensive home treatment as an alternative to hospital admission [21, 22]. Our study had two main aims: 1) to identify between-group differences in sociodemographic and psychiatric characteristics between groups defined by whether they were persistently (i.e. being significantly lonely across three follow-up timepoints), intermittently (i.e. being significantly lonely at one or two follow-up timepoints) or never lonely (i.e. never being lonely across three follow-up timepoints), and between groups defined by whether they were persistently (i.e. being socially isolated across three timepoints), intermittently (i.e. being socially isolated at one or two timepoints) or never (i.e. never being socially isolated across three timepoints) socially isolated and 2) to examine whether persistent loneliness and persistent social isolation were associated with poorer self-rated personal recovery over an 18-month period. We hypothesised that 1) participants with persistent severe loneliness would have the poorest self-rated personal recovery at 18-month follow-up, followed by participants who reported from intermittent severe loneliness, and then participants who were never severely lonely; and 2) Participants with persistent objective social isolation would have the poorest self-rated personal recovery at 18-month follow-up, followed by participants who suffered from intermittent objective social isolation, and then participants who were never objectively socially isolated.

Method

Participants

Participants were drawn from the Crisis Team Optimisation and Relapse Prevention (CORE) study. This large multi-site randomised controlled trial was primarily designed to test the effectiveness of a peer-provided self-management intervention for people with mental health problems, delivered following discharge from CRTs in six NHS trusts, including inner city, suburban and rural environments [23]. Trial inclusion criteria and recruitment procedures are fully described in the published trial protocol [24]. For this paper, we pool participants in both arms of the trial to form a single cohort. A detailed baseline description of this cohort and the variations associated with loneliness at baseline has been previously published [25].

Inclusion criteria for the CORE study required participants to have the capacity to provide written informed consent to participate and to have received support from one of the CRTs for a week or longer. Exclusion criteria included: 1) being clinically assessed as being too high risk to others to participate safely in the study or trial intervention; 2) lacking sufficient knowledge in English to complete the assessments; and 3) CRT discharge having occurred more than a month ago.

The CORE study was approved by the London Camden and Islington Research Ethics Committee (ref 12/LO/0988) before the commencement of data collection. Written informed consent was received from all participants prior participation. All methods were performed in accordance with the relevant guidelines and regulations.

Measures

All participants recruited for the CORE study (i.e., both experimental and control groups) were assessed at three timepoints using structured interviews: baseline, 4- and 18-month follow-up. Sociodemographic information was only collected at baseline. The measures used in this research are reported below.

Social isolation variables

Loneliness: The UCLA Loneliness Scale (ULS-8) was administered as the main measure of loneliness. The ULS-8 consists of eight self-reported items, all rated on a 4-point Likert scale. The total score of this scale ranges from 8 and 32, and a higher score indicates more severe loneliness. This scale has demonstrated good validity and reliability [26]. For the current research, a participant was considered as being significantly lonely if his/her total score on the ULS-8 was 24 or above. This cut-off point was defined as scoring an average of three on each item, which is equivalent to being lonely at least sometimes. Trajectories of loneliness were determined based on loneliness scores at baseline, 4- and 18-month follow-up: 1) persistently severely lonely, included participants who scored as severely lonely at all three timepoints; 2) intermittently severely lonely, consisting of participants who scored as severely lonely at one or two timepoints only; and 3) never severely lonely, including participants who did not score as severely lonely at any timepoint.

Objective social isolation: The Lubben Social Network Scale (LSNS-6) is the revised version of the original LSNS. The LSNS-6 was administered to measure objective social isolation, it aims to examine the number of kin relationships and non-kin relationships that participants keep in contact with, as well as the perceived quality of these relationships. The LSNS-6 has shown excellent internal consistency and convergent validity [27]. For the current study, two items (i.e., the number of relatives and of friends one sees or hears from at least once a month) relating to social network size were utilised as a measure of objective social isolation, which is conceptually distinct from loneliness. The sum of item 1 and 4 ranged from 0 to 10, with a higher score indicating lower objective social isolation.

Outcome measure (primary outcome)

The Questionnaire about the Process of Recovery (QPR): The QPR contains 22 items and yields a total score between 0 and 88. The higher the score, the better the self-rated intrapersonal and interpersonal recovery. The QPR has demonstrated stability across time, satisfactory internal consistency, construct validity and reliability [28].

Sociodemographic and psychiatric (controlled) variables

Participants' sociodemographic and clinical characteristics were recorded at baseline, including age, gender, ethnicity, marital status, employment status, educational attainment, whether they were in contact with children under 16 years old, current diagnoses, number of psychiatric inpatient hospitalisation, and number of years since first contact with mental health services. Psychiatric symptoms were measured by the Brief Psychiatric Rating Scale (BPRS), which was developed to evaluate major symptoms for psychiatric patients [29].

Statistical analyses

Case mean substitution was implemented to replace missing data on continuous variables, including loneliness, social network size and personal recovery. Given the low level of missing data on all scales at both baseline and 18-month follow-up in the current study, case mean substitution should have equivalent effectiveness as other missing data techniques [30]. Following established guidance [31, 32], cases with over 25% of data missing on a single scale were excluded from the final analysis. Pearson's correlation was used to check for potential collinearity between all baseline independent variables.

T-test and chi-square tests were used to check predictors of missingness for the outcomes. When baseline variables were found to be different between the participants who completed the ULS-8 and LSNS-6 at all three timepoints (i.e., completers) and those who did not (i.e., non-completers), these variables were controlled for in the final analysis. The primary outcome, that is the association between the three loneliness groups and personal recovery at 18-month follow-up was tested by multivariable linear regression analyses. Firstly, a univariate linear regression model was carried out. Then four blocks of baseline variables were added into the multivariable regression models, in the following steps: 1) baseline social network size; 2) baseline sociodemographic variables (i.e. age, gender, ethnicity, educational attainment, and whether employed or in education or not) and baseline social network size; 3) baseline sociodemographic variables, baseline psychiatric variables (i.e. number of years since first contact with mental health services, the Brief Psychiatric Rating

Scale (BPRS) total score, number of psychiatric inpatient hospitalisation), and baseline social network size; and 4) baseline sociodemographic and clinical variables, baseline social network size, and baseline QPR score. Comparable five steps were also carried out to examine the association between the three objective social isolation groups and personal recovery at 18-month follow-up.

All analyses were performed in STATA 12.1. Associations were considered as significant if a p-value of less than 0.05 was calculated.

Results

Baseline descriptive results

In total, 399 participants were included in the baseline analysis. Baseline characteristics of our sample are described in Appendix 1. A detailed description of participants at baseline has been previously published [25]. Of these 399 participants, only 224 participants completed measures at all three timepoints and could be included in the final analysis for research question 2. Please see Table 1 for the characteristics of participants at 18-month follow-up. The mean age of these participants was 40.0 (SD = 12.5), and about 40% were male. Approximately 30% of this sample was diagnosed with schizophrenia or other psychotic disorders. The mean of the QPR reported by these participants was 43.3 (SD = 10.7). Reported loneliness at 18-month follow-up had a mean score of 22.1 (SD = 4.5), and the mean of 18-month social network size was 5.0 (SD = 2.3).

Educational attainment and employment status were the only two variables identified as predictors of missingness at a $p = 0.05$ level of significance, so they were included in multivariable analyses for research question 2. Pearson's correlation revealed that the correlation between baseline variables ranged from -0.27 to 0.39. Based on the cut-off proposed by existing literature [33], collinearity was unlikely to be a problem.

Loneliness and objective social isolation groups

The proportion of participants in each loneliness group is shown in Fig. 1. Of the 224 participants, 36 (16%) participants were in the persistently severely lonely group, 113 (50%) participants were never severely lonely, and the rest of the sample ($n = 75$, 34%) was intermittently severely lonely. Of these 75 participants in the intermittently severely lonely group, 32 participants were lonely at an earlier timepoint but not subsequently (Groups 3 and 4); 19 were not lonely at an earlier timepoint but reported being so later (Groups 6 and 7), and another 24 participants experienced fluctuating loneliness (Groups 5 and 8).

Of the same 224 participants, 28 participants (13%) met our criteria for persistent objective social isolation, 124 (55%) participants were not objectively socially isolated at any timepoint, and the rest of the sample ($n = 72$, 32%) met criteria for the intermittent social isolation group. Of these 72 participants, a pathway out of objective social isolation was observed for 28 participants (Groups 3 and 4); a pathway into objective social isolation was observed for 19 participants (Groups 6 and 7), and another 25 participants experienced fluctuating objective social isolation (Groups 5 and 6).

The characteristics of people in different loneliness and objective social isolation groups at baseline

The characteristics of participants in different loneliness groups are shown in Table 2. Not only were the participants in the persistent severe loneliness group more likely to be single, separated, divorced or widowed ($p < 0.001$), they were also less likely to be employed, or in education, or any full-time caring role ($p = 0.002$) than those who were never severely lonely. The persistent severe loneliness group had the smallest social network, followed by the intermittent severe loneliness group. There were significant differences in the QPR score at baseline between the three loneliness groups: persistent severe loneliness group scored the lowest on the QPR. This group was followed by the intermittent severe loneliness group. The group who was never severely lonely scored the highest on the QPR.

The characteristics of participants in different social isolation groups are described in Table 3. The comparisons between the three objective social isolation groups indicated that the participants in the persistent objective social isolation group were less likely to be born in the UK, were less likely to be employed, in education or any full-time caring role ($p = 0.01$), and they were less likely to have been admitted as a psychiatric inpatient previously ($p = 0.001$), compared to the participants who were never objectively socially isolated during the study period. They were also more likely to have a diagnosis of depression, or anxiety disorder or post-traumatic stress disorder ($p = 0.03$), rather than having a diagnosis of psychosis, bipolar disorders, personality disorders or other mental health diagnoses. Statistically significant differences were also reported for their baseline loneliness score: the persistent objective social isolation group scored the highest on the ULS-8, followed by the intermittent objective social isolation group. The persistent objective social isolation group also scored lowest on the QPR. The group who was not socially isolated at any timepoint had the highest QPR score. The detailed results are shown in Table 3.

The association between loneliness groups and self-rated personal recovery at 18-month follow-up

Results from the multivariable linear regression analyses demonstrate a significant relationship between loneliness group and self-rated personal recovery at 18-month follow-up. As shown in model 1, with being never severely lonely as the reference category, intermittent severe loneliness was associated with a statistically significant 9.8-point lower score on the QPR (95% CI -13.59, -6.02), persistent severe loneliness was associated with a 21.75-point lower score on the QPR (95% CI -26.58, -16.93). These associations remained statistically significant in the rest of the models, after controlling for the three blocks of baseline variables (i.e., sociodemographic and psychiatric variables, and social network size) and baseline QPR. In the final model (i.e., model 5), persistent (coef. = -12.8, 95% CI -18.83, -6.83, $p < 0.001$) and intermittent severe loneliness (coef. = -7.8, 95% CI -11.80, -3.75, $p < 0.001$) were both associated with a lower QPR score at 18-month follow-up than for the group who did not report severe loneliness at any timepoint. One clinical variable '2–10 years since first contact with mental health services' (i.e., compared with 'less than 3 months since first contact with mental health services' as a reference category) was also negatively associated with the QPR at 18-month follow-up in model 5. The detailed results are shown in Table 4.

The association between objective social isolation groups and self-rated personal recovery at 18-month follow-up

Multivariable linear regression analyses demonstrate a significant relationship between the objective social isolation groups and self-rated personal recovery at 18-month follow-up. In model 1, persistent objective social isolation was the only group significantly associated with a lower QPR at 18-month follow-up (with being never objectively socially isolated as a reference category). This relationship remained statistically significant even after controlling for the three blocks of baseline variables and baseline QPR score (coef. =-9.8, 95% CI -15.71, -3.79, $p = 0.001$). In the final model (i.e., model 5), the QPR at 18-month follow-up was also negatively associated with '2–10 years and over 10 years since first contact mental health services' (i.e., with 'less than 3 months since first contact mental health services' as a reference category). The detailed results are presented in Table 5.

Discussion

Sample characteristics

Our study included a diagnostically diverse sample, and the median baseline loneliness was equivalent to a moderate level of loneliness. Baseline loneliness score for our cohort is comparable to people with diagnoses across the entire spectrum of mental disorders [34, 35]. Our sample also reported having seen or heard from approximately five family members and friends in the previous month, which is comparable to previous studies of people with mental health problems [36, 37].

Loneliness and objective social isolation groups

Whereas much research in people with psychosis or in mixed groups of mental health service users has been cross-sectional in nature, the three timepoints in this study allowed a novel exploration of associations between personal recovery and loneliness trajectory. Being consistently lonely at three timepoints over 18 months appeared associated with considerably lower ratings for self-rated recovery. While age, sex and ethnic group did not vary by loneliness trajectory, in exploratory analyses the persistently lonely group was also more likely to be single, separated, divorced or widowed, and to be unemployed, not in education or any full-time caring role. Both marriage or cohabiting status (11% of persistently lonely vs. 38% of never lonely) and being employed at baseline (31% of persistently lonely vs. 66% of never lonely) had large associations with subsequently loneliness trajectory. The persistently lonely group also had more severe symptoms at baseline, and a large difference in self-rated recovery was already present. The group whose trajectory showed relatively severe loneliness at some but not all time points were intermediate between the persistently and never lonely groups in their initial symptom and recovery scores, and in baseline loneliness scores and social network size. Thus, at baseline, the group whose subsequent trajectory was defined as persistently lonely, were already disconnected from others in several respects, with poorer scores for both recovery and symptom severity. Prolonged loneliness has been argued to be especially significant in across age groups in the general population [20] and in our study, those who were lonely at multiple timepoints appear a group warranting attention in future research and intervention development.

People who were socially isolated at each time point, were likewise markedly more likely to be unemployed and not cohabiting or married, had more severe symptoms and rated their recovery as poorer at baseline. There was likewise a gradient between those who appeared socially isolated at each time point, those for whom this fluctuated, and those who were not isolated at any point.

The association between loneliness groups, objective social isolation groups and personal recovery at 18-month follow-up

Our primary hypothesis, that people who were lonely at multiple timepoints would show less improvement in self-rated recovery was confirmed, with a persisting effect following adjustment for clinical and social variables and baseline recovery score (with recovery already poorer at baseline). This is compatible with a causal relationship between persistent loneliness and poor improvement in self-rated recovery, although the caveats below in the limitations must be noted. Social connection is conceptualised as an aspect of personal recovery [38, 39], making interpretation of their relationship over time more complex. However, our findings at least suggest that people who are and remain severely lonely and/or social isolated, with few connections with others, are a group worthy of a specific focus as they appear to recover poorly following a crisis. The mechanisms through which persistent severe and social isolation loneliness might impact one's personal recovery remain unexplored. The interrelationships between a number of underlying factors, such as self-esteem, public and self-stigma, sociodemographic backgrounds, and social deprivation, may play a part. Despite a smaller explanatory power, a significant association was also discovered between persistent objective social isolation and poorer personal recovery at 18-month follow-up.

Strengths and limitations

The current study benefits from several main strengths: 1) we included a diagnostically varied sample which was recruited from standard secondary mental health services in the UK; 2) and they were at an illness stage that is of high clinical relevance; and 3) to the best of our knowledge, the current study is the first investigating the chronicity of loneliness and objective social isolation, and their impact on personal recovery among people with mental health problems. However, our results also subject to certain limitations: 1) our study only included mental health patients who were willing to participate in an RCT; 2) the current study had a large proportion of missing data on the diagnosis variable; 3) despite the well-established scales included in the current study, the quality of our measures is subject to limitations. For example, despite being widely used in this population [5], the ULS-8 was not originally developed for people with mental health problems, and lacks clear thresholds for severe and moderate loneliness, so that we needed to specify these pragmatically for the study. In terms of the measure for social network size, given that the LSNS-6 measures both subjective and objective aspects of one's social relationships, we had to make our own selection of items from the scale in order to focus only on objective social isolation, assessed by numbers of contacts. The measurement of loneliness at more than one timepoint is an advance over many previous studies: however, that loneliness and self-rated personal recovery were measured

over the same time period and that they are relatively closely related concepts, we cannot make confident statements regarding the causal link between the two concepts.

Research and clinical implications

Results of our study carry some important implications for future research and clinical practice. In the current study, 16% of our cohort were persistently severely lonely, and 13% suffered from persistent objective social isolation. Large-scale longitudinal cohort studies will be valuable in providing rigorous evidence concerning the extent and the impact of persistent severe loneliness and persistent objective social isolation among mental health service users. The significance of prolonged loneliness across people in different age groups in the general population has been acknowledged [20] and the current study suggests that people who are persistently lonely might also be a group of interest for further investigation and intervention development among people with longer-term mental health problems. People who are lonely or report being relatively isolated at times, but in whom improvements are observed may be a group who have more resources to remedy their own lack of connections: understanding this process better would also contribute to a greater understanding of how lack of connection may improve. Despite widespread acknowledgement of the social determinants of mental health, social interventions still tend to receive less attention in mental health care than pharmacological or psychological approaches.

Given the poor outcomes associated with persistently loneliness and social isolation, routine enquiry into loneliness and the extent of social networks as part of assessments seems desirable so that practitioners are at least aware of these potentially remediable factors. The Office for National Statistics [40] has recommended a quick screening tool, which involves three questions from the ULS-3 and a direct question as an efficient approach to identify and address loneliness in all public settings and health practices. Finally, there is a growing interest in the development of interventions targeting loneliness among mental health service users with various diagnoses [41, 42], our results suggest that targeting people who are lonely and/or socially isolated at multiple timepoints may be a fruitful approach.

Declarations

Ethics approval and consent to participate

The CORE study protocol was approved by the London Camden and Islington Research Ethics Committee (ref 12/LO/0988). Written informed consent was received from all participants.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

The study was funded by the National Institute for Health Research under its Programme Grants for Applied Research programme (reference RP-PG-0109-10078). SJ is supported by the National Institute for Health Research (NIHR) Mental Health Policy Research Unit and the NIHR CLAHRC North Thames. SJ and BLE are supported as part of the UKRI-funded Loneliness and Social Isolation in Mental Health Research Network [grant number ES/S004440/1]. These institutions were not involved in the design of the study and collection, analysis and interpretation of data, and writing of the manuscript.

Authors' Contributions

SJ, BLE and RM conceptualised the study design. RM was a major contributor in data analysis and interpretation and drafted the manuscript. JW and LM provided support for statistical analysis. RM, BLE and SJ critically revised and substantially contributed throughout the writing of the manuscript. SJ is the principal investigator of the study. SJ and BLE supervised the data collection. All authors critically reviewed and approved the final manuscript.

Acknowledgement

We would like to thank the CORE research team, CORE participants and staff from the six participating NHS Trusts.

References

1. Hawkey LC, Cacioppo JT. Loneliness and pathways to disease. *Brain Behav Immun*, 2003; 17 Suppl 1: S98-105.
2. Peplau LA, Perlman D. Perspectives on loneliness. In: Peplau, L. A., et al, editors. *Loneliness: A Sourcebook of Current Theory, Research and Therapy*; 1982. p.1. Wiley, New York
3. Andersson L. Loneliness research and interventions: a review of the literature. *Aging & Mental Health*, 1998; 2: 264–274.

4. Cornwell EY, Waite LJ. Social disconnectedness, perceived isolation, and health among older adults. *Journal of Health and Social Behavior*, 2009; 50(1): 31–48
5. Wang J-Y, Lloyd-Evans B, Giacco D, Forsyth R, Nebo C, Mann F, Johnson S. Social isolation in mental health: a conceptual and methodological review. *Soc Psychiatry Psychiatr Epidemiol*, 2017; 52(12): 1451–1461. doi: 10.1007/s00127-017-1446-1.
6. Yang J, Park E-C, Lee SA, Lee JE, Choi D-W, Chae W, Jang S-I. The association between social contacts and depressive symptoms among elderly Koreans. *Psychiatry Investig*, 2018; 15(9): 861–868
7. Lim MH, Rodebaugh TL, Zyphur MJ, Gleeson JFM. Loneliness over Time: The Crucial Role of Social Anxiety. *Journal of Abnormal Psychology*, 2016. 125(5): 620–630. doi: 10.1037/abn0000162
8. Kuiper JS, Zuidersma M, Oude Voshaar RC., Zuidema SU, van den Heuvel ER, Stolk RP, Smidt N. Social relationships and risk of dementia: a systematic review and meta-analysis of longitudinal cohort studies. *Ageing Research Reviews*, 2015; 22: 39–57
9. Lauder W, Sharkey S, Mummery K. A community survey of loneliness. *J Adv Nurs*. 2004; 46(1):88–94.
10. Giacco D, Palumbo C, Strappelli N, Catapano F, Priebe S. Social contacts and loneliness in people with psychotic and mood disorders. *Comprehensive psychiatry*, 2016; 66: 59–66
11. Kupferberg A, Bicks L, Hasler G. Social functioning in major depressive disorder. *Neuroscience and Biobehavioral Reviews*, 2016; 69: 313–332
12. Holwerda TJ, Beekman AT, Deeg DJ, Stek ML, van Tilburg TG., Visser PJ, et al. Increased risk of mortality associated with social isolation in older men: only when feeling lonely? Results from the Amsterdam Study of the Elderly (AMSTEL). *Psychol Med*, 2012; 42: 843–853.
13. Thornicroft G, Brohan E, Rose D, Sartorius N, Leese M. The INDIGO Study Group. Global pattern of experienced and anticipated discrimination against people with schizophrenia: a cross-sectional survey. *Lancet*, 2009; 373: 408–415
14. Perese EF, Wolf M. Combating loneliness among persons with severe mental illness: social network interventions' characteristics, effectiveness, and applicability, *Issues in Mental Health. Nursing*, 2005; 26(6): 591–609
15. U.S. Department of Health and Human Services. *Transforming Mental Health Care in America, Federal Action Agenda: First Steps*. 2006. Author, Rockville, MD
16. HM Government. *No health without mental health: a cross-government mental health outcomes strategy for people of all age*. 2011. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/138253/dh_124058.pdf. Accessed 22nd March 2021.
17. Slade M. *Personal recovery and mental illness. A guide for mental health professionals*. 2009. Cambridge, UK: Cambridge University Press.
18. Liberman RP, Kopelowicz AK. Recovery from schizophrenia: A criterion-based definition. In: Ralph RO, Corrigan PW, editors. *Recovery in mental illness: Broadening our understanding of wellness*, 2005; 101–130. Washington, DC: American Psychological Association
19. Roe D, Mashiach-Eizenberg M, Lysaker PH. The relation between objective and subjective domains of recovery among persons with schizophrenia-related disorders. *Schizophr Res*, 2011; 131: 133–138.
20. Qualter P, Vanhalst J, Harris R, Van Roekel E, Lodder G, Bangee M, et al. Loneliness across the life span. *Perspectives on Psychological Science*, 2015; 10(2): 250–264. DOI: 10.1177/1745691615568999
21. Wheeler C, Lloyd-Evans B, Churchard A, Fitzgerald C, Fullarton K, Mosse L, et al. Implementation of the Crisis Resolution Team model in adult mental health settings: a systematic review. *BMC Psychiatry*, 2015; 15: 74. doi: 10.1186/s12888-015-0441-x
22. Johnson, S. Crisis resolution and home treatment teams: An evolving model. *Adv Psychiatr Treat*, 2013; 19: 115–23.
23. Johnson S, Lamb D, Marston L, Osborn D, Mason O, Henderson C, et al. Peer supported self-management for people discharged from a mental health crisis team: a randomised controlled trial. *The Lancet*, 2018; 392: 409–418
24. Johnson S, Mason O, Osborn D, Milton A, Henderson C, Marston L, et al. Randomised controlled trial of the clinical and cost-effectiveness of a peer-delivered self-management intervention to prevent relapse in crisis resolution team users: study protocol. *BMJ Open*, 2017; 7(10): e015665. doi: 10.1136/bmjopen-2016-015665.
25. Wang J-Y, Lloyd-Evans B, Marston L, Mann F, Ma R, Johnson S. Loneliness as a predictor of outcomes in mental disorders among people who have experienced a mental health crisis: a 4-month prospective study. *BMC Psychiatry*, 2020; 20 (article no: 249). <https://doi.org/10.1186/s12888-020-02665-2>
26. Hays RD, DiMatteo MR. A short form measure of loneliness. *J Pers Assess*, 1987; 51(1): 69–81
27. Lubben J, Blozik E, Gillmann G, Illiffe S, von Reneln Kruse W, Beck JC, Stuck AE. Performance of an abbreviated version of the Lubben social network scale among three European community-dwelling older adult populations. *Gerontologist*, 2006; 46(4): 503–513
28. Neil S, Kilbride M, Pitt L, Nothard S, Welford M, Sellwood W, Morrison AP. The questionnaire about the process of recovery (QPR): a measurement tool developed in collaboration with service users. *Psychosis*, 2009; 1(2): 145–155
29. Overall JE, Gorham DR. The brief psychiatric rating scale. *Psychol Rep*, 1962; 10: 799–812
30. Parent MC. Handling item-level missing data: simpler is just as good. *The Counseling Psychologist*, 2013; 41(4): 568–600. DOI: 10.1177/0011000012445176
31. Eekhout I, de Vet HCW, Twisk JWR, Brand JPL, de Boer MR, Heymans MW. Missing data in a multi-item instrument were best handled by multiple imputation at the item score level. *Journal of Clinical Epidemiology*, 2014; 67(3): 335–342.
32. Donner A. The relative effectiveness of procedures commonly used in multiple regression analysis for dealing with missing values. *The American Statistician*, 1982; 36: 378–381

33. Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carre G, et al. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography*, 2012; 36(1): 27–46. <https://doi.org/10.1111/j.1600-0587.2012.07348.x>
34. Syu Y-C, Lin L-Y. Sensory overresponsivity, loneliness and anxiety in Taiwanese adults with autism spectrum disorder. *Occupational Therapy International*, eCollection, 2018; Article ID 9165978, 7 pages. <https://doi.org/10.1155/2018/9165978> Accessed on 13th December 2020
35. Jazaieri H, Goldin PR, Werner K, Ziv M, Gross JJ. A randomised trial of MBSR versus aerobic exercise for social anxiety disorder. *J Clin Psychol*, 2012; 68(7): 715–31. doi: 10.1002/jclp.21863
36. Palumbo C, Volpe U, Matanov A, Priebe S, Giacco D. Social networks of patients with psychosis: a systematic review. *BMC Research Notes*, 2015; 8: 560
37. Chrostek A, Grygiel P, Anczewska M, Wciorka J, Switaj P. The intensity and correlates of the feelings of loneliness in people with psychosis. *Compr Psychiatry*, 2016; 70: 190–199. doi: 10.1016/j.comppsy.2016.07.015.
38. Pernice-Duca F, Onaga E. Examining the contribution of social network support to the recovery process among clubhouse members. *American Journal of Psychiatric Rehabilitation*, 2019; 12: 1–30. doi: <http://dx.doi.org/10.1080/15487760802615566>
39. Lysaker PH, Ringer J, Maxwell C, McGuire A, Lecomte T. Personal narratives and recovery from schizophrenia. *Schizophr. Res*, 2010; 121(1–3): 271–276.
40. Office for National Statistics. *Measuring Loneliness: guidance for use of the national indicators on surveys*. 2018. <https://www.ons.gov.uk/peoplepopulationandcommunity/wellbeing/methodologies/measuringlonelinessguidanceforuseofthenationalindicatorsonsurvey> Accessed on 23rd March 2021
41. Ma R, Mann F, Wang J, Lloyd-Evans B, Terhune J, Al-Shihabi A, Johnson S. The effectiveness of interventions for reducing subjective and objective social isolation among people with mental health problems: a systematic review. *Social Psychiatry and Psychiatric Epidemiology*, 2019; 55(7): 839–876. doi: 10.1007/s00127-019-01800-z.
42. Lloyd-Evans B, Frerichs J, Stefanidou T, Bone J, Pinfold V, Lewis G, et al. The Community Navigator Study: Results from a feasibility randomised controlled trial of a programme to reduce loneliness for people with complex anxiety or depression. *PLoS ONE*, 2020; 15(5): e0233535. <https://doi.org/10.1371/journal.pone.0233535>

Tables

Table 1. Characteristics of participants at 18-month follow-up

Variables	Participants at 18-month follow-up (n=224)	
	Mean (SD) or %	N
Age	39.97 (12.54)	224
Gender (%)		
Male	39.73%	89
Female	60.27%	135
Ethnicity		
White British/Irish/other	61.88%	138
Black, Black British/Caribbean/African/other	21.52%	48
Asian, Asian British/Indian/Pakistani/Bangladeshi/ other, Chinese	7.62%	17
Mixed White/Black Caribbean, Mixed White/Black African, mixed White/Asian, other mixed, other ethnic groups	8.97%	20
Marital status		
Single/Separated/divorced/widowed	74.55%	167
Married/cohabiting	25.45%	57
UK born		
No	24.89%	55
Yes	75.11%	166
Housing		
Permanent/supported accommodation	96.88%	217
Unstable accommodation	3.13%	7
Contact with children under 16		
No contact	6.25%	14
Contact with dependent children	22.32%	50
Having no children	71.43%	160
Educational attainment		
No qualification	15.25%	34
Other qualification	50.67%	113
Degree	34.08%	76
Employment/education status		
Not in employment/education/full time caring role	43.89%	97
Yes	56.11%	124
Loneliness score	22.07 (4.50)	224
Social network size	4.97 (2.28)	224
Number of psychiatric inpatient hospitalisations		
Never	62.50%	140
Once	15.63%	35
2 or more	21.88%	49
Number of years since first contact mental health services		
Less than 3 months	16.07%	36
3 months - 2 years	16.07%	36
2-10 years	33.93%	76
More than 10 years	33.93%	76
Current diagnosis		
Schizophrenia or schizoaffective disorder/bipolar affective disorder/other psychosis	30.14%	66

Variables	Participants at 18-month follow-up (n=224)	
	Mean (SD) or %	N
Depression/anxiety disorder/post-traumatic stress disorder	26.94%	59
Borderline or emotionally unstable personality disorder/other personality disorder	11.42%	25
Other diagnosis	31.51%	69
BPRS total score	43.30 (10.67)	223
QPR total score	51.21 (17.61)	224

Abbreviations: N = numbers of participants; M = mean; SD = standard deviation; BPRS = the Brief Psychiatric Rating Scale; QPR = the Questionnaire about the Process of Recovery

Table 2. The comparisons of baseline variables between three loneliness groups^a

Variables	Loneliness groups						P value	95% CI
	Persistent loneliness group (Group 1)		Intermittent loneliness group (Group 2)		Never loneliness group (Group 3)			
	Mean (SD) or %	N	Mean (SD) or %	N	Mean (SD) or %	N		
Age	40.90 (11.49)	36	36.54 (12.25)	75	41.95 (12.68)	113	Group 1 vs. Group 3: 0.66 Group 2 vs. Group 3: 0.004^b Group 1 vs. Group 2: 0.08	Group 1 (37.01, 44.79) Group 2 (33.72, 39.36) Group 3 (39.58, 44.31)
Gender							0.25	
Male	27.78%	10	44.00%	33	40.71%	46		
Female	72.22%	26	5.00%	42	59.29%	67		
Ethnicity							0.77	
White British/Irish/other	62.86%	22	56.00%	42	65.49%	74		
Black, Black British/Caribbean/African/other	17.14%	6	25.33%	19	20.35%	23		
Asian, Asian British/Indian/ Pakistani/Bangladeshi/ other, Chinese	11.43%	4	9.33%	7	5.31%	6		
Mixed White/Black Caribbean, mixed White/Black African, mixed White/Asian, other mixed, other ethnic groups	8.57%	3	9.33%	7	8.85%	10		
Marital status							<0.001	
Single/Separated/divorced/widowed	88.89%	32	86.87%	65	61.95%	70		
Married/Cohabiting	11.11%	4	13.33%	10	38.05%	43		
UK born							0.99	
No	25.71%	9	24.332%	18	25.00%	28		
Yes	74.29%	26	75.68%	56	75.00%	84		
Housing							0.49	
Permanent/supported accommodation	100%	36	96.00%	72	96.46%	109		
Unstable accommodation	0%	0	4.00%	3	3.54%	4		
Contact with children under 16							0.78	
No contact	2.78%	1	8.00%	6	6.19%	7		
Contact with dependent children	27.78%	10	20.00%	15	22.12%	25		
Having no children	69.44%	25	72.00%	54	71.68%	81		
Educational attainment							0.11	
No qualification	2.71%	9	13.33%	10	13.27%	15		
Other qualification	37.14%	13	60.00%	45	48.67%	55		
Degree	37.14%	13	26.67%	20	38.05%	43		
Employment/education status							0.002	
Not in employment/education/full time caring role	68.57%	24	46.58%	34	34.51%	39		
Yes	31.43%	11	53.42%	39	65.49%	74		
Baseline loneliness score	27.77 (2.13)	36	23.52 (3.76)	75	19.28 (3.15)	113	Group 1 vs. Group 3: <.001 Group 2 vs. Group 3: <.001 Group 1 vs. Group 2: <.001	Group 1 (27.05, 28.49) Group 2 (22.66, 24.39) Group 3 (18.69, 19.87)
Baseline social network size	3.19 (2.10)	36	4.73 (2.37)	75	5.69 (1.93)	113	Group 1 vs. Group 3: <.001 Group 2 vs. Group 3: 0.003 Group 1 vs. Group 2: 0.001	Group 1 (2.49, 3.90) Group 2 (4.19, 5.28) Group 3 (5.33, 6.05)
Numbers of psychiatric inpatient hospitalisations							0.49	
Never	69.44%	25	64.00%	48	59.29%	67		
Once	19.44%	7	13.33%	10	15.93%	18		
More than 2 times	11.11%	4	22.67%	17	24.78%	28		
Number of years since first contact mental health services							0.08	
Less than 3 months	5.56%	2	18.67%	14	17.70%	20		
3 months - 2 years	5.56%	2	22.67%	17	15.04%	17		
2-10 years	41.67%	15	32.00%	24	32.74%	37		
More than 10 years	47.22%	17	26.67%	20	34.51%	39		
Current diagnosis							0.09	
Psychosis or bipolar disorders ^c	19.44%	7	21.92%	16	39.09%	43		
Depression or anxiety disorders ^d	36.11%	13	30.14%	22	21.82%	24		
Personality disorders ^e	16.67%	6	13.70%	10	8.18%	9		

Variables	Loneliness groups							
	Persistent loneliness group (Group 1)		Intermittent loneliness group (Group 2)		Never loneliness group (Group 3)			
	Mean (SD) or %	N	Mean (SD) or %	N	Mean (SD) or %	N	P value	95% CI
Other diagnosis	27.78%	10	34.25%	25	30.91%	34		
Baseline BPRS total score	51.19 (12.62)	36	44.77 (9.33)	75	39.79 (9.23)	112	Group 1 vs. Group 3: <.001 Group 2 vs. Group 3: 0.0004 Group 1 vs. Group 2: 0.003	Group 1 (46.93, 55.46) Group 2 (42.63, 46.92) Group 3 (38.06, 41.51)
Baseline QPR total score	35.06 (13.92)	36	49.59 (16.83)	75	57.43 (15.65)	113	Group 1 vs. Group 3: <.001 Group 2 vs. Group 3: 0.0013 Group 1 vs. Group 2: <.001	Group 1 (30.34, 39.77) Group 2 (45.71, 53.46) Group 3 (54.52, 60.35)

Abbreviations: SD = standard deviation; CI= confidence interval; N = number of participants; BPRS = the Brief Psychiatric Rating Scale; QPR = the Questionnaire about the Process of Recovery

- a. t-test and chi-square test were conducted to examine the differences in baseline characteristics between three loneliness groups
- b. significant p-values are marked in bold.
- c. Schizophrenia or schizoaffective disorder or bipolar affective disorder or other psychosis
- d. Depression or Anxiety disorder or post-traumatic stress disorder
- e. Borderline or emotionally unstable personality disorder or other personality disorder

Table 3. The comparison of baseline variables between three objective social isolation groups^a

Variables	Objective social isolation groups						P value	95% CI
	Persistent social isolation group (Group 1)		Intermittent social isolation group (Group 2)		Never social isolation group (Group 3)			
	Mean (SD) or %	N	Mean (SD) or %	N	Mean (SD) or %	N		
Age	43.76 (10.06)	28	40.29 (11.9)	72	38.92 (13.26)	124	Group 1 vs. Group 3: 0.07 Group 2 vs. Group 3: 0.47 Group 1 vs. Group 2: 0.18	Group 1 (39.86, 47.66) Group 2 (37.48, 43.10) Group 3 (36.57, 41.28)
Gender							0.23	
Male	25%	7	43.06%	31	41.13%	51		
Female	75%	21	56.94%	41	58.87%	73		
Ethnicity							0.75	
White British/Irish/other	50%	14	59.15%	42	66.13%	82		
Black, Black British/Caribbean/African/other	28.57%	8	22.54%	16	19.35%	24		
Asian, Asian British/Indian/ Pakistani/Bangladeshi/ other, Chinese	7.14%	2	8.45%	6	7.26%	9		
Mixed White/Black Caribbean, mixed White/Black African, mixed White/Asian, other mixed, other ethnic groups	14.29%	4	9.86%	7	7.26%	9		
Marital status							0.06	
Single/Separated/divorced/widowed	78.57%	22	83.33%	60	68.55%	85		
Married/ Cohabiting	21.43%	6	16.67%	12	31.45%	39		
UK born							0.02 ^b	
No	42.86%	12	28.17%	20	18.85%	23		
Yes	57.14%	16	71.83%	51	81.25%	99		
Housing							0.59	
Permanent/supported accommodation	96.43%	27	98.61%	71	95.97%	119		
Unstable accommodation	3.57%	1	1.39%	1	4.03%	5		
Contact with children under 16							0.88	
No contact	7.14%	2	5.56%	4	6.45%	8		
Contact with dependent children	17.86%	5	26.39%	19	20.97%	26		
Having no children	75.00%	21	68.06%	49	72.58%	90		
Educational attainment							0.18	
No qualification	28.57%	8	15.49%	11	12.10%	15		
Other qualification	35.71%	10	47.89%	34	55.65%	69		
Degree	35.71%	10	36.62%	26	32.26%	40		
Employment/education status							0.01	
Not in employment/education/full time caring role	67.86%	19	47.14%	33	36.59%	45		
Yes	32.14%	9	52.86%	37	63.41%	78		
Baseline loneliness score	26.07 (3.78)	28	22.65 (4.31)	72	20.82 (4.19)	124	Group 1 vs. Group 3: <.001 Group 2 vs. Group 3: 0.004 Group 1 vs. Group 2: 0.0004	Group 1 (24.61, 27.54) Group 2 (21.64, 23.66) Group 3 (20.08, 21.57)
Baseline social network size	1.75 (0.97)	28	3.76 (1.78)	72	6.40 (1.50)	124	Group 1 vs. Group 3: <.001 Group 2 vs. Group 3: <.001 Group 1 vs. Group 2: <.001	Group 1 (1.38, 2.12) Group 2 (3.35, 4.18) Group 3 (6.13, 6.66)
Numbers of psychiatric inpatient hospitalisations							0.001	
Never	82.14%	23	45.83%	33	67.74%	84		
Once	14.29%	4	16.67%	12	15.32%	19		
More than 2 times	3.57%	1	37.50%	27	16.94%	21		
Number of years since first contact mental health services							0.08	
Less than 3 months	17.86%	5	13.89%	10	16.94%	21		
3 months - 2 years	17.86%	5	13.89%	10	16.94%	21		
2-10 years	32.14%	9	23.61%	17	40.32%	50		
More than 10 years	32.14%	9	48.61%	35	25.81%	32		
Current diagnosis							0.03	
Psychosis or bipolar disorder ^c	14.81%	4	39.44%	28	28.10%	34		
Depression or anxiety disorders ^d	37.04%	10	25.35%	18	25.62%	31		
Personality disorders ^e	25.93%	7	9.86%	7	9.09%	11		
Other diagnosis	22.22%	6	25.35%	18	37.19%	45		

Variables	Objective social isolation groups							
	Persistent social isolation group (Group 1)		Intermittent social isolation group (Group 2)		Never social isolation group (Group 3)			
	Mean (SD) or %	N	Mean (SD) or %	N	Mean (SD) or %	N	P value	95% CI
Baseline BPRS total score	49.21 (10.82)	28	44.68 (11.48)	71	41.19 (9.54)	124	Group 1 vs. Group 3: 0.0001 Group 2 vs. Group 3: 0.02 Group 1 vs. Group 2: 0.08	Group 1 (45.02, 53.41) Group 2 (41.96, 47.39) Group 3 (39.49, 42.88)
Baseline QPR total score	41.54 (15.74)	28	50.71 (17.44)	72	53.69 (17.46)	124	Group 1 vs. Group 3: 0.001 Group 2 vs. Group 3: 0.25 Group 1 vs. Group 2: 0.02	Group 1 (45.02, 53.41) Group 2 (50.58, 56.79) Group 3 (41.96, 47.39)

Abbreviations: SD = standard deviation; CI= confidence interval; N = number of participants; BPRS = the Brief Psychiatric Rating Scale; QPR = the Questionnaire about the Process of Recovery

- a. t-test and chi-square test were conducted to examine the differences in baseline characteristics between three objective social isolation groups
- b. significant p-values are marked in bold.
- c. Schizophrenia or schizoaffective disorder or bipolar affective disorder or other psychosis
- d. Depression or Anxiety disorder or post-traumatic stress disorder
- e. Borderline or emotionally unstable personality disorder or other personality disorder

Table 4. Linear regression between three loneliness groups and 18-month QPR, controlling for baseline variables^a

Variables	Model 1		Model 2		Model 3		Model 4		Model 5	
	Coef. (95% CI)	P- value	Coef. (95% CI)	P- value	Coef. (95% CI)	p- value	Coef. (95% CI)	P- value	Coef. (95% CI)	p- value
Loneliness group										
Never severely lonely group	Reference		Reference		Reference		Reference		Reference	
Intermittently severely lonely group	-9.80 (-13.59, -6.02)	<.001^b	-9.69 (-13.57, -5.81)	<.001	-9.73 (-13.88, -5.59)	<.001	-8.73 (-12.85, -4.61)	<.001	-7.78 (-11.80, -3.75)	<.001
Persistently severely lonely group	-21.75 (-26.58, -16.93)	<.001	-21.46 (-26.70, -16.22)	<.001	-19.83 (-25.55, -14.12)	<.001	-16.27 (-22.16, -10.37)	<.001	-12.83 (-18.83, -6.83)	<.001
Psychosocial variable										
Social network size			0.12 (-.69, .93)	0.77	0.05 (-.82, .91)	0.92	0.14 (-.72, 1.01)	0.75	0.05 (-.79, .89)	0.91
Sociodemographic variables										
Age (years)					-0.3 (-.18, .13)	0.75	-0.06 (-.22, .10)	0.47	-1.12 (-1.28, .038)	0.14
Gender (Reference - male)					0.03 (-3.69, 3.75)	0.99	0.81 (-2.86, 4.48)	0.67	1.24 (-2.33, 4.81)	0.49
Ethnicity										
White British/Irish/other					Reference		Reference		Reference	
Black, Black British/Caribbean/African/other					2.09 (-2.33, 6.51)	0.35	2.09 (-2.29, 6.47)	0.35	0.53 (-3.80, 4.85)	0.81
Asian, Asian British/Indian/ Pakistani/Bangladeshi/ other, Chinese					1.23 (-5.70, 8.17)	0.73	-0.76 (-7.65, 6.14)	0.83	-.65 (-7.34, 6.03)	0.85
Mixed White/Black Caribbean, mixed White/Black African, mixed White/Asian, other mixed, other ethnic groups					-3.26 (-9.57, 3.06)	0.31	-2.92 (-9.25, 3.41)	0.36	-4.28 (-10.46, 1.90)	0.17
Employment/ education status (Reference - not in employment/ education)					2.69 (-1.39, 6.76)	0.20	2.62 (-1.44, 6.69)	0.21	2.18 (-1.77, 6.13)	0.28
Educational attainment										
No qualification					Reference		Reference		Reference	
Other qualifications					1.52 (-3.94, 6.98)	0.58	1.03 (-4.40, 6.46)	0.71	.91 (-4.35, 6.18)	0.73
Degree					1.64 (-4.21, 7.49)	0.58	1.16 (-4.63, 6.96)	0.69	1.40 (-4.21, 7.02)	0.62
Psychiatric variables										
Number of psychiatric inpatient hospitalisations										
Never							Reference		Reference	
Once							1.87 (-2.97, 6.71)	0.45	2.16 (-2.53, 6.86)	0.36
2 or more							5.21 (.56, 9.86)	0.03	4.37 (-1.17, 8.90)	0.059
Number of years since first contact with mental health services										
Less than 3 months							Reference		Reference	
3 months - 2 years							-3.60 (-9.87, 2.67)	0.26	-4.48 (-10.57, 1.61)	0.15
2-10 years							-8.06 (-13.51, -2.60)	0.004	-8.20 (-13.49, -2.91)	0.003
More than 10 years							-3.79 (-9.51, 1.92)	0.19	-4.84 (-10.41, .72)	0.09
Baseline BPRS total score							-.19 (-.38, .0002)	0.05	-0.13 (-.31, .06)	0.17
Baseline QPR total score									0.22 (.10, .33)	<.001
R ² adjusted	0.273		0.270		0.244		0.286		0.329	

Abbreviation: CI= confidence interval; N = number of participants; BPRS = the Brief Psychiatric Rating Scale; QPR = the Questionnaire about the Process of Recovery; R² adjusted = adjusted- R²

a. multivariable linear regression analyses were conducted with QPR at 18-month follow-up as dependent variable and other factors as independent variables

b. significant p-values are marked in bold.

Table 5. Linear regression between objective social isolation groups and 18-month QPR, controlling for baseline variables


Variables	Model 1		Model 2		Model 3		Model 4		Model 5	
	Coef. (95% CI)	P- value	Coef. (95% CI)	P- value	Coef. (95% CI)	p- value	Coef. (95% CI)	P- value	Coef. (95% CI)	p- value
Social isolation group										
Never socially isolated group	Reference		Reference		Reference		Reference		Reference	
Intermittently socially isolated group	-3.53 (-7.68, .62)	0.10	-1.75 (-5.80, 2.30)	0.40	-0.64 (-4.77, 3.48)	0.76	-2.13 (-6.37, 2.12)	0.33	-2.23 (-6.32, 1.85)	0.28
Persistently socially isolated group	-16.35 (-22.18, -10.53)	<.001 ^b	-11.17 (-17.19, -5.16)	<.001	-10.63 (-4.77, 3.48)	0.001	-10.75 (-16.93, -4.57)	0.001	-9.75 (-15.71, -3.79)	0.001
Psychosocial variable										
Loneliness score					-8.78 (-1.33, -4.3)	<.001	-0.52 (-1.01, -.04)	0.04	-0.12 (-.63, .38)	0.63
Sociodemographic variables										
Age (years)					0.06 (-.10, .22)	0.45	0.02 (-.14, .18)	0.80	-.05 (-.22, .11)	0.51
Gender (Reference - male)					-.37 (-4.22, 3.48)	0.85	0.52 (-3.29, 4.34)	0.79	1.23 (-2.46, 4.92)	0.51
Ethnicity										
White British/Irish/other					Reference		Reference		Reference	
Black, Black British/Caribbean/African/other					2.51 (-2.08, 7.09)	0.28	2.55 (-1.99, 7.09)	0.27	0.73 (-3.73, .18)	0.75
Asian, Asian British/Indian/ Pakistani/Bangladeshi/ other, Chinese					0.89 (-6.33, 8.11)	0.81	-1.60 (-8.81, 5.61)	0.66	-1.92 (-8.85, 5.02)	0.59
Mixed White/Black Caribbean, mixed White/Black African, mixed White/Asian, other mixed, other ethnic groups					-1.62 (-8.22, 4.97)	0.63	-1.55 (-8.18, 5.08)	0.65	-3.47 (-9.91, 2.98)	0.29
Employment/ education status (Reference - not in employment/ education)					4.09 (-.08, 8.25)	0.054	3.86 (-.31, 8.04)	0.07	3.08 (-.95, 7.11)	0.13
Educational attainment										
No qualification					Reference		Reference		Reference	
Other qualifications					1.30 (-4.37, 6.97)	0.65	0.59 (-5.04, 6.22)	0.84	.30 (-5.12, 5.71)	0.91
Degree					1.89 (-4.19, 7.98)	0.54	1.26 (-4.78, 7.29)	0.68	1.36 (-4.45, 7.16)	0.65
Psychiatric variables										
Number of psychiatric inpatient hospitalisations										
Never							Reference		Reference	
Once							2.06 (-3.00, 7.12)	0.42	2.31 (-2.57, 7.18)	0.35
More than 2 times							4.92 (-1.13, 9.96)	0.056	4.50 (-3.36, 9.35)	0.07
Number of years since first contact with mental health services										
Less than 3 months							Reference		Reference	
3 months - 2 years							-3.77 (-10.29, 2.75)	0.26	-4.86 (-11.14, 1.43)	0.13
2-10 years							-9.31 (-14.97, -3.64)	0.001	-9.56 (-15.01, -4.12)	0.001
More than 10 years							-4.56 (-10.49, 1.38)	0.13	-5.90 (-11.64, -1.15)	0.04
Baseline BPRS total score							-0.20 (-.41, .0002)	0.050	-0.17 (-.36, .03)	0.10
Baseline QPR total score									0.26 (.14, .38)	<.001
R ² adjusted	0.115		0.188		0.182		0.227		0.285	

Abbreviations: CI= confidence interval; N = number of participants; BPRS = the Brief Psychiatric Rating Scale; QPR = the Questionnaire about the Process of Recovery; R² adjusted = adjusted- R²

a. multivariable linear regression analyses were conducted with QPR at 18-month follow-up as dependent variable and other factors as independent variables

b. significant p-values are marked in b

Figures



Group 1: persistently severely lonely (N=36)

Group 2: never severely lonely (N=113)

Group 3: severely lonely – not severely lonely – not severely lonely (N=21)

Group 4: severely lonely – severely lonely – not severely lonely (N=11)

Group 5: not severely lonely- severely lonely – not severely lonely (N=14)

Group 6: not severely lonely – severely lonely – severely lonely (N=6)


Group 7: not severely lonely – not severely lonely – severely lonely (N=13)

Group 8: severely lonely – not severely lonely – severely lonely (N=10)

Abbreviation: N = numbers of participants

Figure 1

Percentage of participants in severely lonely groups



Group 1: persistently socially isolated (N=28)

Group 2: never socially isolated (N=124)

Group 3: socially isolated – not socially isolated – not socially isolated (N=19)

Group 4: socially isolated – socially isolated – not socially isolated (N=9)

Group 5: not socially isolated – socially isolated – not socially isolated (N=18)

Group 6: not socially isolated – socially isolated – socially isolated (N=9)

Group 7: not socially isolated – not socially isolated – socially isolated (N=10)

Group 8: socially isolated – not socially isolated – socially isolated (N=7)

Abbreviation: N = numbers of participants

Figure 2

Percentage of participants in objectively socially isolated groups

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarymaterial.docx](#)