

CD8⁺ T Cells in the Central Nervous System of Mice With Herpes Simplex Infection are Highly Activated and Express High Levels of CCR5 and CXCR3

Liza Lind

Goteborgs universitet Sahlgrenska Akademin

Alexandra Svensson (✉ Alexandra.Svensson@gu.se)

Goteborgs universitet Sahlgrenska Akademin <https://orcid.org/0000-0002-0832-4187>

Karolina Thörn

Goteborgs universitet Sahlgrenska Akademin

Malgorzata Krzyzowska

Goteborgs universitet Sahlgrenska Akademin

Kristina Eriksson

Goteborgs universitet Sahlgrenska Akademin

Research

Keywords: Central nervous system inflammation, Herpes Simplex Virus, CD8⁺ T cells, Chemokines

Posted Date: July 6th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-39368/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background:

Herpes simplex virus type 2 (HSV-2) is a neurotropic virus that can cause meningitis, an inflammation of the meninges in the central nervous system. T cells are key players in viral clearance, and these cells migrate from peripheral blood into the central nervous system upon infection. Several factors contribute to T cell migration, including the expression of chemokines in the inflamed tissue that attract T cells through their expression of chemokine receptors. Here we investigated CD8+ T cell profile in the spinal cord in a mouse model of herpes simplex virus type 2 neuroinflammation.

Method:

Mice were infected with HSV-2, and sacrificed when showing signs of neuroinflammation. Cells and/or tissue from spinal cord, spleen and blood were analyzed for viability, expression of activation markers, chemokine receptors and chemokines. Statistics were calculated using student's T test and one-way ANOVA.

Results:

High numbers of CD8+ T cells were present in the spinal cord following genital HSV-2 infection. CD8+ T cells were highly activated and HSV-2 glycoprotein B-specific effector cells, some of which showed signs of recent degranulation. They also expressed high levels of many chemokine receptors, in particular CCR2, CCR4, CCR5 and CXCR3. Investigating corresponding receptor ligands in spinal cord tissue revealed markedly increased expression of the cognate ligands CCL2, CCL5, CCL8, CCL12 and CXCL10.

Conclusion:

This study shows that during herpesvirus neuroinflammation anti-viral CD8+ T cells accumulate in the CNS. CD8+ T cells in the CNS also express chemotactic receptors cognate to the chemotactic gradients in the spinal cord. This indicates that anti-viral CD8+ T cells may migrate to infected areas in the spinal cord during herpesvirus neuroinflammation in response to chemotactic gradients.

Introduction

Herpes simplex virus type 2 (HSV-2) is a double-stranded DNA virus that infects through mucosal tissue to establish a latent infection in the central nervous system (CNS) (1, 2). HSV-2 commonly manifests as genital blisters or remains asymptomatic (3). In rare cases HSV-2 can cause meningitis (incidence of 2–4/100 000), an inflammation of the CNS with symptoms such as fever, headaches and photosensitivity (4–6). Although usually benign, HSV-2 meningitis often presents with recurrent episodes of disease (4, 5).

T cells are key players in cell-mediated immunity and are reported to be essential in HSV-2 clearance in mice (7, 8). A distinct T cell subset are the CD8+ cytotoxic T cells, which are involved in direct killing of

virus-infected cells by secreting granzymes, inducing target cell apoptosis (9). Accumulated evidence implies that CD8 + T cells are important not only for the control of acute CNS infection but are also necessary for maintenance of HSV latency in infected ganglia (10).

In neuroinflammation T cells migrate from peripheral tissue, crossing the blood-brain barrier to enter CNS tissue and fight the underlying cause. The mechanism of T cell recruitment to the CNS is not fully understood, but it is clear that chemokines, chemotactic cytokines, expressed by cells in the inflamed tissue, are involved (11). Chemokines bind chemokine receptors, which are G protein-linked transmembrane receptors expressed on all immune cells, but expression differs between cell types, activation states and tissues (12, 13). CXCR3 and its ligand CXCL10 is known to have protective roles in HSV-2 infection as well as to facilitate migration into the CNS (14, 15). Chemokines CCL3, CCL5 and CXCL2 have been implicated in enhanced viral clearance and/or survivability (16–18), and in humans chemokines have been characterized in the acute phase of HSV-2 meningitis (19). However, the chemokine-receptor profile of CD8 + T cells transmigrating to the CNS in HSV-2 meningitis is not known.

The present study was initiated to elucidate chemokine and chemokine receptor expression affecting CD8 + T cell migration into the CNS of HSV-2-infected mice. Using a mouse model of HSV-2-induced neuroinflammation, we show that effector CD8 + T cells express a chemokine repertoire that suggests that they have migrated to the CNS. These cells are highly activated, granzyme B-expressing cells with high specificity for HSV-2 glycoprotein B. In the CNS, the CD8 + T cells express a wide range of chemokine receptors; CCR2, CCR4, CCR5, CXCR3 and also CXCR4. Several ligands of these receptors, especially CCL2, CCL5, CCL8 and CXCL10, were highly upregulated in the CNS of mice with HSV-2-induced neuroinflammation. Our data indicate that CD8 + T cells migrate to and accumulate in the CNS, where they exert their function to battle HSV-2.

Materials And Methods

Infection

12-20-week old female C57bl/6 mice (Charles River) were infected with HSV-2 strain 333 as previously described (20). In short, mice were subcutaneously injected with 2 mg medroxyprogesterone acetate (Depo-Provera, Pfizer). 6 days post treatment mice were infected with 40 000 PFU HSV-2 strain 333 intravaginal and sacrificed 2, 4, 6 or 8 days post infection. Control mice were left uninfected. Mice were monitored and scored according to the following: 0, no signs of infection/inflammation; 1, mild genital redness; 2, moderate genital inflammation; 3, obvious damage of the genital mucosa, impaired general condition; 4, hind leg paralysis (neuroinflammation). The numbers of mice used for each experiment were as follows: flow cytometry for activation markers, n = 5 mice/group; flow cytometry for chemokine receptor expression, n = 4 mice/group and n = 15 mice/group at day 0 and day 8, respectively; chemokine and cytokine expression in spinal cord, n = 10 and n = 11 at day 0 and 8, respectively. Mice were kept under pathogen-free conditions at the Department of Rheumatology and Inflammation Research,

University of Gothenburg (Gothenburg, Sweden). This study was approved by the ethical committee for animal experiments (Gothenburg, Sweden).

Flow cytometry

Spleens and spinal cords were collected and mashed through a 70 µM cell strainer. Spleen red blood cells were lysed using ammonium chloride. Cells were counted using an automated hematology analyzer (KX-21N, Sysmex), and 10⁶ million cells/sample were subsequently used for flow cytometric staining. Spinal cords were resuspended in 2 ml PBS, and 200 µL were used for staining one sample. After blocking of unspecific binding for 10 minutes using 5 µg/mL rat anti-mouse CD16/CD32 (cat no 553142), cells were stained for 20 minutes using the following mAbs, CD44 PE (cat no 561860), CD8 BV421 (cat no 563898), CCR7 PerCp-Cy5.5 (cat no 560812) (BD Biosciences), CCR2 PE (cat no FAB5538P), CXCR3 PE (cat no FAB1685P), CXCR4 FITC (cat no FAB21651F) (R&D Systems), CCR4 APC (cat no 131211), CCR4 PE (cat no 131203), CCR5 APC (cat no 107011), CCR6 PerCp-Cy5.5 (cat no 129810), CXCR3 FITC (cat no 126535), CXCR4 APC (cat no 146507), CD107a PerCP-Cy5.5 (cat no 121625), CD3 FITC (cat no 100306), (Biolegend), CD62L APC (cat no 17-0621-82, eBioscience) and SSIEFARL APC (ProImmune). For intracellular staining, BD cytofix/cytoperm fixation/permeabilization kit (BD) was used according to manufacturer's instructions. Intracellular staining was done using FITC rat anti-granzyme B (cat no 15208379, eBioscience). Stained cells were acquired using BD FACSVerse (BD Biosciences), and analyzed using FlowJo software (Tree Star, Ashland, OR, USA). Fluorescence minus one was used as control for cell-surface staining and isotype controls were used for intracellular staining.

Chemokine and cytokine analysis

Whole blood was collected and processed into serum by allowing clotting in room temperature, followed by separation of the serum phase by centrifugation. Extractions of tissue was achieved by collecting spinal cords in PBS with 0.1 mg/ml soybean trypsin inhibitor (Sigma), 1.5 mM Pefabloc (Coatech AB), 50 mM EDTA and 0.1% bovine serum albumin. Samples were weighed and frozen. Upon thawing, samples were permeabilized using 2% saponin over night at 4 °C, and the supernatant was subsequently collected by centrifugation. Chemokine and cytokine analysis was performed using a mouse magnetic Luminex assay (CCL2, CCL3, CCL4, CCL5, CCL7, CCL8, CCL11, CCL12, CCL20, CCL22, CXCL1, CXCL2, CXCL5, CXCL10, CXCL12, GM-CSF, IFN-gamma IL-1alpha, IL-1beta, IL-2, IL-4, IL-6, IL-10, TNF-alpha) from R&D Systems according to manufacturer's instructions and analyzed using Bio-Plex 200 system (Bio-Rad).

Statistical analysis

All data are expressed as mean +/- SEM. Unpaired T test was used when two groups were compared, and one-way ANOVA with Dunnett's or Tukey's multiple comparisons test was used when three or more groups were compared. Statistical analysis was performed using GraphPad Prism version 7 (GraphPad Software).

Results

Increase of CD8 + T cells in the spinal cord upon HSV-2 infection

To investigate the CD8 + T cell profile in the CNS, mice were infected with HSV-2. Neurological symptoms were observed 7–9 days post infection (i.e. hind leg paralysis). Animals were then euthanized and cells from spinal cord (CNS tissue) and spleen (peripheral tissue) were used for flow cytometric analysis both before and after infection. Low numbers of CD8 + T cells were detected in spinal cord from healthy animals. Following HSV-2 infection the numbers increased almost 100-fold, compared to healthy animals (Fig. 1A). This was associated with elevated levels of IFN-gamma in spinal cord extracts (Fig. 1B). General gating strategy for flow cytometry of CD8 + T cells derived from spinal cord and spleen is shown in Fig. 1C. Thus, there is a vast increase of CD8 + T cells in the spinal cord during herpes simplex infection, indicating that CD8 + T cells traffic into the CNS during viral neuroinflammation.

CD8 + T cells in the spinal cord of HSV-2-infected mice have a highly activated HSV-2 specific phenotype

To prove the hypothesis that CD8 + T cells found in the CNS are virus-specific effector cells we characterized the viable CD8 + T cells by flow cytometry. In spinal cord almost all CD8 + T cells were highly activated, defined as CD44 + CD62L⁻, while CD8 + T cells in the spleen showed a significantly lower activation frequency (Fig. 2A). Staining cells with the HSV-2 specific MHC class 1-pentamer SSIEFARL showed that a 15-fold higher proportion of CD8 + T cells were specific for SSIEFARL in spinal cord compared to spleen (Fig. 2B). Most CD8 + T cells in the spinal cord expressed the cytotoxic enzyme granzyme B (Fig. 2C), and approximately 4% of the cells showed signs of recent degranulation as indicated by the marker CD107a (Fig. 2D). In summary, CD8 + T cells present in the spinal cord during herpes simplex induced neuroinflammation are highly activated virus-specific effector T cells.

Chemokine receptor expression in CD8 + T cells in the spinal cord of herpes simplex induced neuroinflammation

To assess contributing factors of T cell presence in the CNS upon HSV-2 infection, chemokine receptors on the CD8 + T cells from mice 8 days post infection were analyzed by flow cytometry and compared to the levels in uninfected mice and to the levels on spleen CD8 + T cells.

CD8 + T cells in spinal cord from uninfected mice had a restricted chemokine receptor repertoire with 20% of the cells expressing CCR5 and 40% expressing CXCR3 (Fig. 3, black bars). The CD8 + T cells that appeared in the spinal cord after infection had a more diverse chemokine receptor expression pattern compared to both CD8 + T cells in the spinal cord of uninfected mice (Fig. 3, green bars) and to CD8 + T cells in the spleen of HSV-2 infected animals (Fig. 3, grey bars), and there were also higher frequencies of chemokine-receptor-positive CD8 + T cells (Fig. 3). The majority of CD8 + T cells in the spinal cord of HSV-2 infected mice expressed CCR5 and CXCR3 (Fig. 3, green bars). The frequencies of CD8 + T cells expressing CCR2, CCR4 or CCR7 were also higher in the spinal cord in HSV-2-infected mice compared to spinal cord CD8 + T cells in uninfected mice and also compared to CD8 + T cells in spleen from infected

animals, with frequencies in the range of 20–30% of CD8 + T cells. CXCR4 on the other hand was expressed by a majority of spleen CD8 + T cells, less so by CD8 + T cells in spinal cord from infected mice, and hardly at all by spinal cord CD8 + T cells from uninfected mice (Fig. 3). Taken together, these data show that CD8 + T cells present in the CNS express CCR5 and CXCR3 during HSV-2 induced neuroinflammation.

Chemokine expression pattern in the spinal cord of mice with HSV-2-induced neuroinflammation

We used multiplex analysis of spinal cord tissue extractions to evaluate if cognate ligands to CCR2, CCR4, CCR5, CXCR3 and CXCR4 are induced in the spinal cord of mice upon HSV-2 infection, and also compared this to the levels induced in serum upon HSV-2 infection. There was a strong induction of CCL5 and CCL8 (which are ligands to CCR5) and CXCL10 (which is a CXCR3 ligand) in the spinal cord of HSV-2-infected mice (Fig. 4A). Also CCL2 (ligand for CCR2 and CCR4), CCL3 (ligand for CCR4 and CCR5), CCL7 and CCL12 (ligands for CCR2) showed a moderate but significant increase in spinal cord at day 8 compared to uninfected mice (day 0), while CCL4, CCL20, CCL22, CXCL1, CXCL2, CXCL5 and CXCL12 remained low.

In blood, CCL8 and CXCL5 dominated in healthy animals, but CXCL12 was also detected (Fig. 4B). Upon HSV-2 infection, the serum levels of CCL8 and CXCL12 decreased slightly while CXCL1 was increased (Fig. 4B).

Discussion

In this study we show that highly activated virus-specific CD8 + T cells expressing CCR5 and CXCR3 are found in the spinal cord of HSV-2-infected mice. This was associated with elevated levels of cognate chemokines in the spinal cord during HSV-2 infection.

We confirm previous studies showing that there are few CD8 + T cells in the spinal cord of naïve mice (21). For the first time though, we can show that the majority of these CD8 + T cells have a limited expression of chemokine receptors. 20–30% of the cells expressed CCR5 and/or CXCR3 and few/none expressed any of the other chemokine-receptors investigated; CCR2, CCR4, CCR6, CCR7, CXCR2 or CXCR4. In accordance, also the chemokine levels in the spinal cord were low in the absence of infection, as previously documented both in humans and in mice (14, 19). Thus, a limited number of CD8 + T cells may patrol the healthy spinal cord in what appear to be a chemokine-independent fashion.

Neuroinflammation caused by HSV-2 infection led to a pronounced increase of CD8 + T cells in the spinal cord, as shown by us here and by others (14). The viable fraction of the spinal cord CD8 + T cells in HSV-2 infected mice were CD44+ (and CD62L-) effector/memory cells, similar to what has been shown in other inflammatory diseases of the CNS (22) and in cell adhesion and migration to the CNS in general (23). The majority of the cells expressed granzyme B, and a few percent of them had recently degranulated. As

previously shown (14), CD8 + T cells that were specific for the HSV-2 peptide SSIEFARL predominated in the spinal cord of HSV-2-infected mice, in accordance with the notion that only antigen-specific effector cells translocate to the CNS. We conclude that HSV-2 induced neuroinflammation in mice is associated with the presence of highly activated virus-specific CD8 + T cells in the spinal cord.

The majority of the spinal cord CD8 + T cells expressed CCR5 and several CCR5 ligands were expressed in the HSV-2-infected spinal cord. Studies in gene-deficient mice show that albeit CCR5 deficiency is associated with exacerbated CNS disease, it does not affect CD8 + T cell numbers in the CNS, neither during HSV-2 infection nor during intracranial infection with coronavirus (24, 25). This is in contrast to studies on non-viral CNS inflammation where CCR5 deficiency leads to an attenuated influx of CD8 + T cells and other immune cells to the spinal cord (26, 27). Many ligands for CCR5 are however expressed in the HSV-2-infected spinal cord. Thapa et al have previously reported the presence of CCL3 and CCL5 (14, 15, 24), and in this study we show that also CCL4 and in particular CCL8 are upregulated. The primary binding partner of CCL8 is CCR5, which is expressed on activated T cells (28) as confirmed by us. The latter is particularly interesting as CCL8 is one of the most prominent chemokines in HSV-2 meningitis patients (19). We conclude that CCR5 is present on most CD8 + T cells in the spinal cord during viral neuroinflammation, with concurrent high levels of CCR5 ligands including in particular CCL8.

Almost 60% of the CD8 + T cells in the spinal cord of HSV-2 infected mice also expressed CXCR3, and the cognate ligand CXCL10 was markedly upregulated in the infected spinal cord. CXCR3 is an important chemokine receptor for T cell migration; however it is debated if CXCR3 has a protective or detrimental function. On one hand mice deficient in CXCR3 succumb faster to HSV-2 infection, on the other hand HSV-1-infected mice lacking CXCR3 are protected from fatal encephalitis (15, 29), indicating a mouse strain and tissue-specific role of CXCR3 (30). Lack of the CXCR3 ligands CXCL9 and CXCL10 on the other hand increase the susceptibility to HSV-2 infection in mice, and reduce T cell migration into the spinal cord despite high levels of CCL2, CCL3 and CCL5 (14), implying that CXCR3/CXCL10 represent a major pathway for spinal cord T cell recruitment. IFN-gamma, which we could show is expressed in the infected spinal cord, induces the expression of CXCL10. CXCL10 is also upregulated in other neuroinflammatory infections like Theiler's murine encephalomyelitis virus, lymphocytic choriomeningitis virus and tick-borne encephalitis (31–33). Most importantly, the enhanced levels of CXCL10 correlate with data from cerebrospinal fluid from humans with HSV-2 meningitis (19).

A moderate proportion of CD8 + T cells in the infected spinal cord also expressed CCR2 and CCR4. CCR2, usually considered a monocyte marker, is found on 2–15% of T cells in both healthy humans and mice, where it binds ligands CCL2, CCL7, CCL8 and CCL12 (28, 34, 35). These four chemokines were strongly upregulated in the spinal cord of mice with HSV-2-induced neuroinflammation, as well as in human HSV-2 meningitis patients (19). Continuing the chain of promiscuous chemokine receptor/ligand interactions CCL5, together with CCL2 and CCL3, also interacts with CCR4. This chemokine receptor is normally considered a classical T_H2 and Treg marker (36, 37). In humans it is also expressed on a subset of cytokine-producing but non-cytotoxic memory CD8 + T cells (38), which resemble a CD8 + T cells subset that are involved in the inhibition of HSV reactivation from latency (38). In summary, all measured

chemokines binding either CCR2 or CCR4 increased in the spinal cord of HSV-2-infected mice, possibly contributing to the recruitment of CCR2- and CCR4-expressing CD8 + T cells.

CXCR4 was found on a limited number of spinal cord CD8 + T cells but was abundantly expressed on T cells in the spleen, indicating an inverse relationship to CNS homing. CXCR4 interacts with CXCL12, which is a known retention signal for CD8 + T cells at the blood-brain barrier. In West Nile virus infection it has been reported that disturbing the CXCR4/CXCL12 interaction allows migration of CD8 + T cells into the brain parenchyma resulting in increased viral clearance and survival (39, 40). In HSV-2 infection, the CXCL12 levels remained low in spinal cord confirming that the CXCR4/CXCL12 pathway is not a major driving force for CD8 + T cell presence in the spinal cord.

Conclusion

We have characterized the chemokine receptor expression pattern on CD8 + T cells in spinal cord during steady state and upon HSV-2 neuroinflammation. During homeostasis the CD8 + T cells that patrol the spinal cord express few chemokine receptors. Upon HSV-2 infection there is a pronounced increase of CCR5- and CXCR3-expressing highly activated virus-specific CD8 + T cells. These data agree with the chemokine secretion pattern in the spinal cord with particularly high levels of CCL5, CCL8 and CXCL10. Importantly, the high levels of CXCL10 correspond to the levels found in cerebrospinal fluid from humans with HSV-2 meningitis. The role of chemokine receptor profile and chemokines in CD8 + T cell trafficking to the CNS should be further investigated to evaluate the contribution to disease severity and/or clearance of the infection, and as a potential target for new treatments.

Abbreviations

HSV Herpes Simplex Virus

CNS Central Nervous System

Declarations

Ethics approval and consent to participate

This study was approved by the ethical committee for animal experiments in Gothenburg, Sweden (D.nr. 12.2014 and D.nr. 5.8.18-00801/2019).

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

This study was funded by the Swedish Research Council, *the Swedish state under the agreement between the Swedish government and the country councils (the ALF-agreement)* and IngaBritt and Arne Lundbergs Forskningsstiftelse.

Authors' contributions

LL and KE designed the study. LL, AS and KE wrote the manuscript and performed the statistical analysis. LL, KT and MK performed the experiments. All authors read and approved the final manuscript.

Acknowledgements

Not applicable.

Authors' information

Liza Lind:Liza.Lind@gmail.com

Alexandra Svensson: Alexandra.Svensson@gu.se

Karolina Thörn: Karolina.Thorn@microbio.gu.se

Malgorzata Krzyzowska: Malgorzata.Krzyzowska@gu.se

Kristina Eriksson: Kristina.Eriksson@microbio.gu.se

References

1. Kramer MF, Cook WJ, Roth FP, Zhu J, Holman H, Knipe DM, et al. Latent herpes simplex virus infection of sensory neurons alters neuronal gene expression. *J Virol.* 2003;77(17):9533–41.
2. Weller SK, Coen DM. Herpes simplex viruses: mechanisms of DNA replication. *Cold Spring Harb Perspect Biol.* 2012;4(9):a013011.
3. Schiffer JT, Corey L. New concepts in understanding genital herpes. *Curr Infect Dis Rep.* 2009;11(6):457–64.
4. Bergstrom T, Vahlne A, Alestig K, Jeansson S, Forsgren M, Lycke E. Primary and recurrent herpes simplex virus type 2-induced meningitis. *J Infect Dis.* 1990;162(2):322–30.

5. Kallio-Laine K, Seppanen M, Kautiainen H, Lokki ML, Lappalainen M, Valtonen V, et al. Recurrent lymphocytic meningitis positive for herpes simplex virus type 2. *Emerg Infect Dis*. 2009;15(7):1119–22.
6. Wald A, Corey L. Persistence in the population: epidemiology, transmission. In: Arvin A, Campadelli-Fiume G, Mocarski E, Moore PS, Roizman B, Whitley R et al, editors *Human Herpesviruses: Biology, Therapy, and Immunoprophylaxis*. Cambridge 2007.
7. McDermott MR, Goldsmith CH, Rosenthal KL, Brais LJ. T lymphocytes in genital lymph nodes protect mice from intravaginal infection with herpes simplex virus type 2. *J Infect Dis*. 1989;159(3):460–6.
8. Koelle DM, Posavad CM, Barnum GR, Johnson ML, Frank JM, Corey L. Clearance of HSV-2 from recurrent genital lesions correlates with infiltration of HSV-specific cytotoxic T lymphocytes. *J Clin Invest*. 1998;101(7):1500–8.
9. Zhang N, Bevan MJ. CD8(+) T cells: foot soldiers of the immune system. *Immunity*. 2011;35(2):161–8.
10. Schiffer JT, Corey L. Rapid host immune response and viral dynamics in herpes simplex virus-2 infection. *Nat Med*. 2013;19(3):280–90.
11. Korn T, Kallies A. T cell responses in the central nervous system. *Nat Rev Immunol*. 2017;17(3):179–94.
12. Melik-Parsadaniantz S, Rostene W. Chemokines and neuromodulation. *J Neuroimmunol*. 2008;198(1–2):62–8.
13. Griffith JW, Sokol CL, Luster AD. Chemokines and chemokine receptors: positioning cells for host defense and immunity. *Annu Rev Immunol*. 2014;32:659–702.
14. Thapa M, Welner RS, Pelayo R, Carr DJ. CXCL9 and CXCL10 expression are critical for control of genital herpes simplex virus type 2 infection through mobilization of HSV-specific CTL and NK cells to the nervous system. *J Immunol*. 2008;180(2):1098–106.
15. Thapa M, Carr DJ. CXCR3 deficiency increases susceptibility to genital herpes simplex virus type 2 infection: Uncoupling of CD8 + T-cell effector function but not migration. *J Virol*. 2009;83(18):9486–501.
16. Benencia F, Gamba G, Cavalieri H, Courreges MC, Benedetti R, Villamil SM, et al. Nitric oxide and HSV vaginal infection in BALB/c mice. *Virology*. 2003;309(1):75–84.
17. Eo SK, Lee S, Chun S, Rouse BT. Modulation of immunity against herpes simplex virus infection via mucosal genetic transfer of plasmid DNA encoding chemokines. *J Virol*. 2001;75(2):569–78.
18. Harle P, Noisakran S, Carr DJ. The application of a plasmid DNA encoding IFN-alpha 1 postinfection enhances cumulative survival of herpes simplex virus type 2 vaginally infected mice. *J Immunol*. 2001;166(3):1803–12.
19. Lind L, Studahl M, Persson Berg L, Eriksson K. CXCL11 production in cerebrospinal fluid distinguishes herpes simplex meningitis from herpes simplex encephalitis. *J Neuroinflammation*. 2017;14(1):134.

20. Svensson A, Nordstrom I, Sun JB, Eriksson K. Protective immunity to genital herpes simplex [correction of simplex] virus type 2 infection is mediated by T-bet. *J Immunol.* 2005;174(10):6266–73.
21. Deb C, Howe CL. Functional characterization of mouse spinal cord infiltrating CD8 + lymphocytes. *J Neuroimmunol.* 2009;214(1–2):33–42.
22. Coque E, Salsac C, Espinosa-Carrasco G, Varga B, Degauque N, Cadoux M, et al. Cytotoxic CD8(+) T lymphocytes expressing ALS-causing SOD1 mutant selectively trigger death of spinal motoneurons. *Proc Natl Acad Sci U S A.* 2019;116(6):2312–7.
23. Brocke S, Piercy C, Steinman L, Weissman IL, Veromaa T. Antibodies to CD44 and integrin alpha4, but not L-selectin, prevent central nervous system inflammation and experimental encephalomyelitis by blocking secondary leukocyte recruitment. *Proc Natl Acad Sci U S A.* 1999;96(12):6896–901.
24. Thapa M, Kuziel WA, Carr DJ. Susceptibility of CCR5-deficient mice to genital herpes simplex virus type 2 is linked to NK cell mobilization. *J Virol.* 2007;81(8):3704–13.
25. Glass WG, Lane TE. Functional analysis of the CC chemokine receptor 5 (CCR5) on virus-specific CD8 + T cells following coronavirus infection of the central nervous system. *Virology.* 2003;312(2):407–14.
26. Gu SM, Park MH, Yun HM, Han SB, Oh KW, Son DJ, et al. CCR5 knockout suppresses experimental autoimmune encephalomyelitis in C57BL/6 mice. *Oncotarget.* 2016;7(13):15382–93.
27. Badell IR, Kitchens WH, Wagener ME, Lukacher AE, Larsen CP, Ford ML. Pathogen Stimulation History Impacts Donor-Specific CD8(+) T Cell Susceptibility to Costimulation/Integrin Blockade-Based Therapy. *Am J Transplant.* 2015;15(12):3081–94.
28. Mack M, Cihak J, Simonis C, Luckow B, Proudfoot AE, Plachy J, et al. Expression and characterization of the chemokine receptors CCR2 and CCR5 in mice. *J Immunol.* 2001;166(7):4697–704.
29. Zimmermann J, Hafezi W, Dockhorn A, Lorentzen EU, Krauthausen M, Getts DR, et al. Enhanced viral clearance and reduced leukocyte infiltration in experimental herpes encephalitis after intranasal infection of CXCR3-deficient mice. *J Neurovirol.* 2017;23(3):394–403.
30. Lundberg P, Openshaw H, Wang M, Yang HJ, Cantin E. Effects of CXCR3 signaling on development of fatal encephalitis and corneal and periocular skin disease in HSV-infected mice are mouse-strain dependent. *Invest Ophthalmol Vis Sci.* 2007;48(9):4162–70.
31. Hoffman LM, Fife BT, Begolka WS, Miller SD, Karpus WJ. Central nervous system chemokine expression during Theiler's virus-induced demyelinating disease. *J Neurovirol.* 1999;5(6):635–42.
32. Nansen A, Marker O, Bartholdy C, Thomsen AR. CCR2 + and CCR5 + CD8 + T cells increase during viral infection and migrate to sites of infection. *Eur J Immunol.* 2000;30(7):1797–806.
33. Pokorna Formanova P, Palus M, Salat J, Honig V, Stefanik M, Svoboda P, et al. Changes in cytokine and chemokine profiles in mouse serum and brain, and in human neural cells, upon tick-borne encephalitis virus infection. *J Neuroinflammation.* 2019;16(1):205.
34. Kivisakk P, Trebst C, Lee JC, Tucky BH, Rudick RA, Campbell JJ, et al. Expression of CCR2, CCR5, and CXCR3 by CD4 + T cells is stable during a 2-year longitudinal study but varies widely between

individuals. *J Neurovirol.* 2003;9(3):291–9.

35. Semple BD, Kossmann T, Morganti-Kossmann MC. Role of chemokines in CNS health and pathology: a focus on the CCL2/CCR2 and CXCL8/CXCR2 networks. *J Cereb Blood Flow Metab.* 2010;30(3):459–73.
36. Morimoto Y, Bian Y, Gao P, Yashiro-Ohtani Y, Zhou XY, Ono S, et al. Induction of surface CCR4 and its functionality in mouse Th2 cells is regulated differently during Th2 development. *J Leukoc Biol.* 2005;78(3):753–61.
37. Molinaro R, Pecli C, Guilherme RF, Alves-Filho JC, Cunha FQ, Canetti C, et al. CCR4 Controls the Suppressive Effects of Regulatory T Cells on Early and Late Events during Severe Sepsis. *PLoS One.* 2015;10(7):e0133227.
38. Kondo T, Takiguchi M. Human memory CCR4 + CD8 + T cell subset has the ability to produce multiple cytokines. *Int Immunol.* 2009;21(5):523–32.
39. Michlmayr D, Lim JK. Chemokine receptors as important regulators of pathogenesis during arboviral encephalitis. *Front Cell Neurosci.* 2014;8:264.
40. McCandless EE, Piccio L, Woerner BM, Schmidt RE, Rubin JB, Cross AH, et al. Pathological expression of CXCL12 at the blood-brain barrier correlates with severity of multiple sclerosis. *Am J Pathol.* 2008;172(3):799–808.

Figures

Figure 1




Figure 1

CD8+ T cells in spinal cord from healthy and HSV-2 infected mice. (A) Total number of CD8+ T cells and (B) IFN-gamma levels (n=10) day 0 and day 8 in the spinal cord of HSV-2 infected mice. Data are presented as mean \pm SEM using unpaired T test where * indicates $p < 0.05$. Statistical analysis was performed using GraphPad Prism version 7 (GraphPad Software). (C) Representative flow cytometry plots of spinal cord cells showing the general gating strategy for staining of spinal cord and spleen cells.

Figure 2




Figure 2

Activation marker expression on CD8+ T cells in spinal cord and spleen of HSV-2-infected mice. (A) Activation (defined as CD44+CD62L- cells), (B) HSV-2 specificity to SSIEFARL peptide, (C) granzyme B expression and (D) degranulation (defined as CD107a+ cells) of CD8+ T cell populations in spleen and spinal cord day 8 post HSV-2 infection (n=5/group). Data are presented as mean \pm SEM. Statistical analysis was performed using unpaired T test, where *** indicates $p < 0.001$; ** $p < 0.01$ and * $p < 0.05$ using GraphPad Prism version 7 (GraphPad Software).

Figure 3




Figure 3

Chemokine receptor expression on CD8+ T cells in spinal cord and spleen of HSV-2-infected mice. Data are presented as mean \pm SEM. Black bars represent spinal cord cells day 0 (n=4), green bars represent spinal cord cells day 8 (n=15), and grey bars represent spleen cells day 8 (n=15) post HSV-2 infection. Statistical analysis was performed using ANOVA with Dunnett's multiple comparisons test, where *** indicates $p < 0.001$; ** $p < 0.01$ and * $p < 0.05$ using GraphPad Prism version 7 (GraphPad Software).

Figure 4




Figure 4

Chemokine expression in spinal cord and sera of HSV-2 infected mice. Chemokine levels in (A) spinal cord extracts and (B) sera of mice infected with HSV-2 were measured at 0 (n=10) or 8 (n=11) days post infection. Data are presented as mean \pm SEM using unpaired T test where *** indicates $p < 0.001$; ** $p < 0.01$ and * $p < 0.05$ using GraphPad Prism version 7 (GraphPad Software).