Genetic Diversity Analysis of Chinese Leishmania Isolates and Development of *L. donovani* Complex Specific Markers by RAPD

Dongmei Yuan

Sichuan University West China School of Basic Medical and Forensic Medicine

Hanxiao Qin

Sichuan University West China School of Basic Medical and Forensic Medicine

Dali Chen

Sichuan University West China School of Basic Medical and Forensic Medicine

Jianping Chen (✉ jpchen007@163.com)

Research article

Keywords: Leishmania donovani, China, RAPD, SCAR

Posted Date: July 9th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-39415/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at BMC Infectious Diseases on May 21st, 2021. See the published version at <https://doi.org/10.1186/s12879-021-06163-y>.

Abstract

Background: Leishmaniasis is one of the most neglected tropical diseases in the world and is still endemic in some underdeveloped regions including western of China. The phylogeny and classification of Chinese *Leishmania* has not yet been clarified, especially within *Leishmania* (*L.*) *donovani* complex, although phylogenetic analyses based on a series of gene markers were carried out. More different analytic method and data still be needed. Random amplified polymorphic DNA (RAPD) technology could identify slight intraspecific differences sensitively, and it always be a powerful tool to seek the species-specific markers. This work aimed to identify Chinese *Leishmania* isolates from diverse geographic regions at genomic level. Meanwhile, the specific markers of *L. donovani* complex were also developed by RAPD.

Methods: The RAPD was applied on 14 Chinese *Leishmania* isolates from diverse geographic regions and 3 WHO reference strains. The polymorphic sites of amplification were transformed into data matrix, based on which genetic similarity were calculated and UPGMA dendrogram were constructed to analyze genetic diversity of these *Leishmania* isolates. Meanwhile, the specific amplification loci of *L. donovani* complex were TA-cloned, sequenced and converted into sequence characterized amplified regions (SCAR) markers, which were validated preliminarily in available 17 *Leishmania* strains in this study and analyzed by bioinformatics.

Results: The cluster analyses showed that the three *Leishmania* sp. isolates SC10H2, SD and GL clustered together and apart from others, and the strains of *L. donovani* complex clearly divided into two clades and the three isolates Cy, WenChuan and 801 formed a subclade. Three specific SCAR markers of *L. donovani* complex, i.e. 1-AD17, 2-A816 and 3-O13, were successfully obtained and validated on available 17 *Leishmania* strains in this study. Through bioinformatics analyses, Marker 1-AD17 may have more specificity on PCR detection of VL and Marker 3-O13 has the potential of encoding protein.

Conclusions: the RAPD result verified that the undescribed *Leishmania* species causing visceral Leishmaniasis (VL) in China was a unique clade distinguished from *L. donovani*, and revealed that there was genetic differentiation among Chinese *L. donovani*. The development of *L. donovani* specific markers may provide the foundation for developing new specific diagnostic markers of VL and the research of specific gene function.

Background

Leishmaniasis is a tropical disease caused by obligate intracellular protozoa genus *Leishmania* and transmits through bite of genus *Phlebotomus*, which threaten 350 million people over 98 countries, mainly in developing countries [1]. There are four main forms of Leishmaniasis according to different clinical syndromes, i.e., visceral Leishmaniasis (VL, kala-azar), post-kala-azar dermal leishmaniasis (PKDL), cutaneous Leishmaniasis (CL) and mucocutaneous Leishmaniasis (MCL), which caused by different species of *Leishmania*. There are more than 64 species of genus *Leishmania* consist of subgenera *Euleishmania*, *Paraleishmania* and *Sauroleishmania* [2], among which 20 species have been considered human-infectious [1].

Visceral leishmaniasis, which is acute and fatal if left untreated, is the main form of leishmaniasis that prevailed in China. Chinese government has taken many implements to control and eliminate VL since 1950s and it has been effectively restricted. Nevertheless, Chinese VL still presents localized and sporadic outbreaks now, and mostly occurred in Xinjiang Uygur Autonomous Region, Sichuan and Gansu province [3]. What' more, China is still one of 14 VL high-burden countries [4]. In the early stages, according to epidemiological characteristics, Chinese VL mainly divided into three different types, i.e., anthroponotic type (AVL), zoonotic mountain type (MT-ZVL) and zoonotic desert type (DT-ZVL) epidemiologically [5]. With further research, the genetic heterogeneity was found to exist in *Leishmania* isolates from different foci of China based on kDNA and nDNA analyses [6, 7]. In recent years, through a series of gene markers, such as internal transcribed spacer 1 (ITS1), cytochrome oxidase II (COX II), cytochrome *b*(*cyt b*) and HSP70 were applied to establish phylogenetic trees, *L. donovani* complex (include *L. donovani* and *L. infantum*), *Leishmania gerbilli*, *Leishmania tropica* and *Leishmania turanica* has been identified, along with an undescribed *Leishmania* species which has clustered with Lizard *Leishmania* [8–11]. However, the species classification and pathogen identification of Chinese leishmaniasis is far from illuminated, especially within *L. donovani* complex. Our previous phylogenetic analyses on HSP70 presented clear relationship within *L. donovani* complex [11]. It inferred that *L. infantum*, which is one of the causative agents of VL, is primarily distributed in western mountainous areas and plain of northwestern China, including Sichuan, Gansu, and Xinjiang provinces. While the identification of isolate MHOM/CN/80/801 from VL patients in Kashi of Xinjiang, was different from the study of ITS1 sequences [12]. Moreover, the analysis of HSP70 also concluded that *L. donovani* is the pathogen of CL in Karamay of Xinjian and *Phlebotomus major wui* is the vector, which challenged the previous determination that the *L. infantum* is the pathogen of CL in Karamay based on gene hybridization and animal inoculation [13, 14]. Thus, application of more diverse analytical methods and more classified data would be needed to add our understanding of genetic relationship and population structure of Chinese *Leishmania* isolates.

Rapid species identification is essential to early diagnosis of Leishmaniasis and conduce to accurate treatments. Multilocus enzyme electrophoresis (MLEE) is still the golden standard in identification of *Leishmania* species [15], but the costly and time-consuming procedure limits its application fields. In recent years, DNA markers have been explored and well-developed for species discrimination and phylogenetic research of *Leshmania*, including ribosomal DNA (rDNA) genes, kinetoplast DNA (kDNA) genes, nuclear genes and antigen genes [16]. The phylogenetic trees of these markers provided a lot of evidence for the taxonomy of main *Leishmania* complex, while the relatively slow evolutionary rate of these genes were insufficient to solve the species relationship within complex [17]. Furthermore, the discrimination capability is diverse among markers and which in turn makes identification of species and subspecies sometimes inconsistent [18]. The Random Amplification Polymorphic DNA (RAPD) is a technique which could make polymorphic analysis for unknown genomes on the basis of PCR [19, 20]. This technique is easy and sensitive, which always be applied to species identification [21], the correlation analysis of population differentiation with geographical origins on *Leishmania* [22]. Also, RAPD has its advantage on the taxonomy at subgeneric level [23]and species level [24]. Meanwhile, through the selection of DNA markers among differentially amplified bands, the specific genetic markers of one species can be developed to species identification or assist diagnosis by making a probe. So it proved to be an effective method to obtain genetic markers to develop relevant *Leishmania* DNA assays [25, 26].

In this study, the RAPD was applied on Chinese *Leishmania* isolates from diverse geographic regions. The polymorphic sites of amplification were transformed into data matrix, based on which genetic similarity were calculated and UPGMA dendrogram were constructed to analyze genetic diversity of these Chinese *Leishmania* isolates, which helped us more thoroughly understand the genetic difference of these isolates combining with our previous phylogenetic analyses based on HSP70 and cyt *b*. Meanwhile, the specific amplified bands of *L. donovani* complex on the gel electrophoretic maps were cut for extraction, TA-cloning and sequencing. Then the specific primers were designed on the successfully obtained gene sequences for converting of *L. donovani* complex specific sequence characterized amplified regions (SCAR) markers. Subsequently, these SCAR markers were validated preliminarily in available 17 *Leishmania* strains in this study and analyzed by bioinformatics, which may provide the foundation for the research of specific gene function and developing new specific diagnostic markers of VL.

Methods

Leishmania strains and DNA extraction

Fourteen Chinese *Leishmania* isolates and three WHO reference strains were used in this study and listed in Table 1 (see Additional file 1). The 14 Chinese *Leishmania* isolates were collected from the plain, hill, and desert foci of China. These *Leishmania* isolates were stored in the liquid nitrogen container. Before experiments, the strains were reheated and hold at 37°C to restore vitality. Then the parasites were cultured for multiplying in Medium 199 supplemented with 15% heat-inactivated fetal bovine serum, 100 U/mL penicillin (Sigma) and 100 µg/mL streptomycin (Sigma) at 26°C. The promastigotes were collected at logarithmic phase and centrifuged at 3300 × g for 10 min. The supernatants were discarded and the sediments were washed three times using PBS. Total DNA was then extracted using a commercially available DNA extraction kit (TianGen Cell DNA Kit) as per the manual. The concentrations of 17 DNA samples were detected (Thermo Scientific™ NanoDrop™ One) and adjusted to the same level before the following RAPD amplification. The integrity of the DNA samples was determined by agarose gel electrophoresis.

Table 1
Details of Chinese strains and the WHO reference strains which were used for RAPD analyses.

Strain name	Species	Origin	Host
IMJW/CN/91/KXG-918	<i>L. donovani</i>	Karamay, Xinjiang, China	<i>Phlebotomus major wui</i>
IMJW/CN/87/KXG-65	<i>L. donovani</i>	Karamay, Xinjiang, China	<i>Phlebotomus major wui</i>
MHOM/CN/93/KXG-XU	<i>L. infantum</i> *	Karamay, Xinjiang, China	<i>Homo sapiens</i>
MHOM/CN/90/9044	<i>L. donovani</i>	Shandong, China	<i>Homo sapiens</i>
MHOM/CN/94/KXG-LIU	<i>L. infantum</i> *	Karamay, Xinjiang, China	<i>Homo sapiens</i>
MHOM/CN/90/SC10H2	<i>Leishmania</i> sp.	Sichuan, China	<i>Homo sapiens</i>
MCAN/CN/08/Cy	<i>L. infantum</i>	Wudu, Gansu, China	<i>Canis lupus familiaris</i>
MCAN/CN/90/WenChuan	<i>L. infantum</i>	Wenchuan, Sichuan, China	<i>Canis lupus familiaris</i>
MRHO/CN/81/EJN1-154	<i>L. gerbilli</i>	Ejina, Inner Mongolia, China	<i>Rhombomys opimus</i>
IMJW/CN/92/KXG-927	<i>L. donovani</i>	Karamay, Xinjiang, China	<i>Phlebotomus major wui</i>
MHOM/CN/86/SC6	<i>L. donovani</i>	Nanping, Sichuan, China	<i>Homo sapiens</i>
MHOM/CN/80/801	<i>L. donovani</i> *	Kashgar, Xinjiang, China	<i>Homo sapiens</i>
SD	<i>Leishmania</i> sp.	Shandong, China	<i>Homo sapiens</i>
GL	<i>Leishmania</i> sp.	Gansu, China	<i>Homo sapiens</i>
MRHO/CN/60/GERBILLI	<i>L. gerbilli</i>	Gansu, China	<i>Rhombomys opimus</i>
MHOM/IN/80/DD8	<i>L. donovani</i>	India	
MHOM/SU/74/K27	<i>L. tropica</i>		
*: the species identification is disputed based on different technique and different gene markers applied.			

RAPD-PCR

Twenty decamer primers were selected according to previous studies [27, 28]and commercially synthesized (Invitrogen). All primers were prepared as 10 µM (10 pmol/µl) working solutions. These primers firstly were screened through three independent RAPD amplifications on the same *L. donovani* isolate DNA, then 10 primers that presenting polymorphic, reproducible and clear amplification profile were selected finally (Table 2). RAPD amplification was performed in 50 µl reactions containing 0.8 µM primer, 10 ng genomic DNA sample, 25 µl 2 × TaqMater Mix (Tsingke, China) and PCR grade distilled water. The PCR procedure were described as follows: initial denaturation at 94°C for 5 min followed by 45 cycles of 94°C for 1 min, 36°C for 1 min, 72°C for 2 min, and a final extension at 72°C for 8 min. The PCR products were separated using 1.5% agarose gel electrophoresis after GelRed (BioSharp) staining with voltage of 5V/cm, and visualized under UV light. Each PCR and electrophoresis were taken three times under the same protocol and operator to assure reproducibility. Only the primers and gel photographs that has stably same band profiles were chosen for data analysis.

Table 2
The nucleotide sequences of 10 primers and amplified results of *Leishmania* isolates.

Primers	Nucleotide Sequence 5'-3'	Total Bands
OP-AD17	GGCAAACCCCT	13
OP-AY5	TCGCTGCGTT	12
OP-AY8	AGGCTTCCCT	18
OP-O13	GTCAGAGTCC	10
OP-U3	CTATGCCGAC	11
AB1-13	TTCCCCCGCT	12
AB1-09	TGGGGGACTC	12
AB1-04	GGACTGGAGT	7
3301 ³⁵	TCGTAGCCAA	12
A816	GTGACGTAGG	14
Total		121

Phenetic analysis of RAPD results

The bands of all polymorphic RAPD gel were marked as “0” for absence and “1” for presence. The relative intensity among all bands were regardless. The 0/1 data matrix was formed in Microsoft Excel 2013 and analyzed using Numerical Taxonomy and Multivariate Analysis System (NTSYS) [29]. The similarity module was used for calculating similarity matrix. The SAHN function of the clustering module was employed for clustering analysis and the phenetic dendrogram was output under the unweighted pair-group method with arithmetic means (UPGMA).

Cloning, sequencing and verification of SCAR markers

The *L. donovani* and *L. infantum* strain-specific bands that stably reproduced in one gel electrophoresis were considered as potential SCAR markers of *L. donovani* complex to extract for DNA purifying and cloning. These gel blocks were purified using gel extraction kit (Tiangen, China) according to the user manual. Purified DNA fragments were then cloned into pGM-T vector (Tiangen, China) overnight. Recombined products were mixed carefully with DH5α competent *E. coli* and put on the ice for 30 min. Transformed *E. coli* were cultured in the Luria-Bertani (LB) medium and sieved through isopropyl-D-thiogalactopyranoside (IPTG), X-Gal and ampicillin protocol. The white colonies were picked up and identified by colony PCR, then the positive samples were proliferated in liquid LB medium and collected for DNA sequencing (Tsingke, China). For the SCAR markers sequences, specific primer pairs were designed using primer 5.0 and synthesized (Tsingke, China), then PCR was performed on now available 17 *Leishmania* strains to determine whether the SCAR markers are specific for *L. donovani* complex in this study. In the PCR conditions, annealing temperature were tested by gradient according to different primers of each SCAR marker. The PCR results were showed by 1.5% agarose gel electrophoresis and observed with UV light.

Bioinformatics analysis of SCAR markers

The obtained SCAR markers sequences were submitted to online BLAST for homologous analyses on NCBI (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>), and the base component was analyzed using Lasergene EditSeq. The open reading frames (ORFs) were predicted and located using NCBI-ORF Finder (<https://www.ncbi.nlm.nih.gov/orffinder>). To further determine whether the ORFs contained in the sequences have potential possibility to express protein, the promotor binding sites were analyzed and predicted online using Promotor Scan (<https://www.bimas.cit.nih.gov/molbio/proscan>). The ORFs that have potential protein expression were translated using Lasergene EditSeq, and the components of these presumed proteins were analyzed by Lasergene Protean. The secondary structures including α-helices and β-strands were predicted by loading Chou-Fasman method. The hydrophobic regions were calculated using Kyte-Doolittle method. The antigenic determinants were analyzed through JamesonWolf method and the surface probability was assessed using Emini method.

Results

RAPD analysis of 17 *Leishmania* strains

Fourteen Chinese *Leishmania* isolates from patients, dogs, gerbils, and sand flies in different geographic origins, with three WHO reference strains, were analyzed by RAPD. RAPD-PCR were performed using 20 decamers as random primers, among them 10 primers (Table 2) that produced reproductive gel photograph profiles under three repeated trials were chosen for next experiments. A total of 121 RAPD bands were observed, of which 120 bands were polymorphic (99.17%). An average of 12.1 bands were amplified by each primer, and segments ranged from 200 to 3000 bp. According to the gel photograph, there are differences among 17 *Leishmania* isolates. Figure 1 showed one gel photo as examples of polymorphism. While same species tend to form similar bands model. The genetic similarity of 17 *Leishmania* strains ranged from 0.4393 to 1.0000 with an average of 0.6758 (Table 3, see Additional file 2), which indicated a considerable genetic differentiation among these isolates. The UPGMA dendrogram established basing on the similarity matrix was showed as Fig. 2. The isolates SC10H2, SD and GL clustered into Clade I with a highly average similarity index of 0.9715, which had a distant genetic relationship with others. Other strains formed Clade II, which further consist of Clade A, B and C. The isolate EJN1-154 clustered with *L. gerbilli* WHO reference strain

MRHO/CN/60/GERBILLI and formed as Clade C. The strains that identified as *L. donovani* complex before did not cluster as one clade but two clades, Clade A and B, instead. The isolates Cy, WenChuan and 801 clustered with *L. donovani* reference strain DD8 as Clade A, whose average similarity index is 0.9439. The other *L. donovani* complex strains clustered as Clade B, in which there still existed genetic differences. Within Clade B, the similarity index of SC6 from Sichuan province with others is 0.9019, which significantly lower than the average index of Clade B (0.9537), so it was separated as an independent clade.

Table 3
Genetic similarity matrix among 17 *Leishmania* strains by NTSYS.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
1 DD8	-															
2 KXG-918	0.832	-														
3 KXG-927	0.850	0.944	-													
4 GERBILLI	0.495	0.477	0.477	-												
5 EJNI-154	0.514	0.495	0.495	0.981	-											
6 SD	0.486	0.505	0.467	0.505	0.505	-										
7 GL	0.477	0.495	0.458	0.495	0.495	0.991	-									
8 9044	0.869	0.944	0.981	0.477	0.495	0.467	0.458	-								
9 KXG-XU	0.850	0.963	0.981	0.477	0.495	0.467	0.458	0.981	-							
10 SC6	0.832	0.869	0.907	0.458	0.458	0.467	0.458	0.925	0.907	-						
11 KXG-65	0.860	0.972	0.972	0.486	0.505	0.477	0.467	0.972	0.991	0.897	-					
12 Cy	0.888	0.813	0.813	0.514	0.533	0.505	0.495	0.832	0.832	0.794	0.841	-				
13 WenChuan	0.888	0.813	0.813	0.514	0.533	0.505	0.495	0.832	0.832	0.794	0.841	1.000	-			
14 801	0.888	0.813	0.813	0.514	0.533	0.505	0.495	0.832	0.832	0.794	0.841	1.000	1.000	-		
15 KXG-LIU	0.850	0.963	0.981	0.477	0.495	0.467	0.458	0.981	1.000	0.907	0.991	0.832	0.832	0.832	-	
16 K27	0.617	0.636	0.598	0.598	0.617	0.533	0.523	0.617	0.617	0.617	0.626	0.617	0.617	0.617	0.617	-
17 SC10H2	0.458	0.477	0.439	0.495	0.495	0.972	0.963	0.439	0.439	0.439	0.449	0.495	0.495	0.495	0.439	0.523

Cloning, sequencing and verification of SCAR markers

Through gel purification, cloning and sequencing on the potential species-specific bands that only appeared in all *L. donovani* complex strains, finally a total of four fragments were successfully T-cloned and sequenced. The obtained fragments sequences were named partly after their primers, i.e. 1-AD17, 2-A816, 3-O13 and 4-09. For evaluating the species-specificity, an 18 base pair primer was designed for each potential marker respectively. By PCR amplification on 17 strains in this study, three of the four markers manifested strict species-specific single band at their respective corresponding loci and were tentatively converted to the potential SCAR markers of *L. donovani* complex (Fig. 3). The primers and annealing temperature of the three SCAR markers are showed in Table 4. The DNA sequences are showed in Table S1 (see Additional file 4).

Table 4
The converted SCAR markers of *L. donovani* complex in this study.

SCAR marker	Length	Primer sequences 5'-3'	Annealing temp (°C)
1-AD17	636	F: TGGCAAACCCTGTATGAGGAAAACGT R: TTGGCAAACCCTCATAGGTTGC	61
2-A816	416	F: GTGACGTAGGCATGCCAGCAAGGTGG R: GTGACGTAGGGGTGGGGATGAAGAGG	63
3-O13	1210	F: GTCAGAGTCCTCGCGGGGTATTC R: GTCAGAGTCCAGTAGATAGGATCGATGCG	65

Bioinformatics analysis of SCAR markers

All the three markers were undergone BLAST on NCBI, the query coverage was 99% and the identity percent were greater than 98% with *L. donovani/infantum* reference sequences. Besides, the E-values were close to '0'. All these indicated that the three markers sequences had a very highly homology with *L. donovani/infantum* reference sequences. According to distribution of Blast hits of SCAR markers, excepted the first two *L. donovani/infantum* reference

sequences, the marker 1-AD17 had a query coverage less than 90% and had no matching primer binding sites with other *Leishmania* species sequences (Fig. 4a). While marker 2-A816 and 3-O13 had more matching primer binding sites with other *Leishmania* species sequences excepted *L. donovani/infantum* reference sequences (Fig. 4b, c). Therefore, the primers of marker 1-AD17 has a greater specificity for amplification of *L. donovani* complex.

The results of sequence components, chromosomal assignment, ORF and promotor prediction for the three markers are listed in Table 5 (see Additional file 3). There are 4 to 7 ORFs in these markers, in which only 3-O13 has two potential promotors, located on 611–816 bp of sense strand and 1056 – 806 of antisense strand. The distance between predicted promotor sites and ORFs implied that ORF-4 has the potential of encoding proteins. The gene sequence was then translated into protein and analyzed by Lasergene. The protein sequence of ORF-4 contains 7 strong basic amino acids, 6 strong acid amino acids, 19 hydrophobic residues and 10 polar residues, and the putative isoelectric point is 8.835. The structure predication of the ORF-4 protein sequence by Lasergene Protean was showed in Fig. 5. The results showed that a clear structure of one hydrophilic β turns region (19–36 residues) was flanked by two hydrophobic α helices (1–18 and 37–50 residues), which implied that the α helices might located in interior of the protein and β turns might on the surface. It was confirmed by the surface probability analysis (Emini method). In addition, the β region has a higher antigen index. Furthermore, there is no homologous protein according to BLAST on GenBank.

Table 5
Bioinformatics analysis of three SCAR markers.

Marker	Length	Base components				chromosomal assignment	ORFs Num.	Promotor Num.	ORF with potential of translation
		A	T	C	G				
1-AD17	636	130	144	165	197	22	6	0	0
		20.44%	22.64%	25.94%	30.97%				
2-A816	416	66	112	117	121	36	4	0	0
		15.87%	26.92%	28.12%	29.09%				
3-O13	1210	238	249	327	396	36	7	2	ORF-4
		19.67%	20.58%	27.02%	32.73%				

Discussion

Generally, DNA marker is now the most widely used method in identification and classification of *Leishmania*, since it is both effective and efficient. While different evolutionary rate of diverse gene markers may lead to different classification result. Thus, to help us understand interspecific relationship more comprehensively, the usage of more dissimilar identification method would be necessary. RAPD has been widely used in genetic maps construction, breeding lines identification and gene markers screening, as has been in the realm of genetic evolution of parasites such as trypanosome, schistosome and trichinella spiralis [30–32]. As we all know, RAPD has its innate drawbacks in stability and repeatability of bands due to the highly random hybrid sites with template DNA. Therefore, to gain stable bands and repeatable results of RAPD, we followed the PCR amplified condition whose stability and repeatability has been already reported [27, 28] and had validated the results before formal experiments. Meanwhile, the usage of commercial Taq DNA Mix from the same batch replaced the addition of dNTP, Mg²⁺ and Taq DNA polymerase one by one also improved the reaction stability in this study. Theoretically, the larger the number of polymorphic sites of RAPD, the more reliable the genetic relationship can be concluded. In this study, 10 of 20 random primers has produced 121 polymorphic bands, with the proportion of polymorphic bands was more than 99%, which could reflect genetic diversity among these isolates credibly.

As is shown in Fig. 2 and Table 1, these 17 strains were clustered into two major Clades. Three isolates SC10H2, SD and GL, which identified as *L. (SauroLeishmania)* sp. based on gene marker *cyt b* and HSP70 previously [11], had a lower genetic similarity with other strains and firstly clustered as Clade I. This result from the view of genome adds to evidence that in China there exist an undescribed *Leishmania* species and it is a distinct branch which has low homology with Chinese *L. donovani* strains [9, 33, 34]. Particularly, this RAPD result also demonstrated the discrimination and differentiation in the relationship among subspecies of *L. donovani* complex. Three isolates Cy, WenChuan and 801, which isolated from Gansu, Sichuan and Xinjiang respectively, were clustered together and separated from other *L. donovani* isolates. Meanwhile, they had the closest genetic relatives with *L. donovani* reference strain DD8. This result confirmed that the genetic differentiation really existed in Chinese *L. donovani*, which was in accord with the phylogenetic analysis on HSP70 [11]. However, the dendrogram of UPGMA showed that *L. donovani* reference strain DD8 did not cluster with KXG-918, KXG-927, KXG-XU, KXG-LIU, 9044, KXG-65 and SC6 that identified as *L. donovani* previously. Instead, it had the closest genetic relatives with Cy, WenChuan and 801. This result indicated there were differences between these Chinese *L. donovani* strains and *L. donovani* reference strain from India on genomic level, which was inconsistent with the phylogenetic analysis results of gene markers [8, 34]. For this, as a gene marker only contains partial information of genome and the selective pressure is various among different genes, intraspecific genetic differentiation probably could not be reflected fully. On the other hand, great genetic variation is generally generated between species or genera in RAPD amplification, so that using an individual represents a species may cause deviation of phylogenic results. Consequently, the divergence in this study need to be further verified by enlarging the sample size or combining with other methods. In addition, the cluster dendrogram showed that the clade B divided into two small branches: one is Sichuan isolate SC6 from hill foci, another included Shandong isolate 9044 from plain foci and other five Xinjiang isolates from desert foci. It revealed that there were still differences among VL isolates from hill, plain and desert in China, which supported the previous report [35]. Although RAPD technology has been waned gradually, while it is very sensitive to identify slight intraspecific differences, which is suitable for differentiation of sibling species.

By means of RAPD, the population differentiation of *Leishmania* could be analyzed, and the random amplified segments reflected abundant polymorphic information of genome. Furthermore, the bioinformatics research of species-specific segments is benefit for further genetic information mining and the

development of specific genetic markers, which could help to species identification or disease diagnosis. In this study, we developed three *L. donovani* complex species-specific DNA markers in available 17 *Leishmania* strains. Although the markers have been verified and showed stably repeatable results in differentiating *L. donovani* complex from other *Leishmania* species, according to their distribution of Blast hits in NCBI, the primers of marker 1-AD17 has a greater specificity for amplification of *L. donovani* complex since it had no matching primer binding sites with other *Leishmania* species sequences. We considered that the differences of primer binding sites or annealing sites of amplification was the cause for generation of differential DNA fragments of diverse species in RAPD, which was speculated in previous report [25]. Thus, the SCAR marker 1-AD17 have the potentiality to be developed into a rapid diagnosis marker of kala-azar. Admittedly, the *L. donovani* complex specific DNA marker in this study still have certain limits because of diverse species of *Leishmania* and less genomic information is available. Therefore, more parasite samples and patient specimens would be needed to test the specificity.

Through bioinformatic analysis, the three markers were all located in large chromosomes instead of kinetoplast, which was similar with some other reports [28, 36, 37]. It may be related to that multicopy genes are found preferentially on disomic chromosomes [38], which would increase the probabilities of random primers binding to them. As genome sequences of different *Leishmania* species are highly conserved [38], the amplification loci of RAPD frequently located in variable regions. In this study, although there were 4 to 7 ORFs in the three markers, in which only 3-O13 has two potential promoters and its ORF-4 has the potential of encoding proteins. The following protein prediction analysis showed that the hypothetical protein had higher antigenic index and surface probability. Nevertheless, all these need further experiments to be verified.

Conclusions

Although the previous phylogenetic results based on a series of gene markers has revealed most genetic evolutionary relationship of Chinese *Leishmania*, there still existed some controversy, especially the subspecies in *L. donovani* complex. The RAPD method could identify intraspecific differences sensitively and reflect the correlation of geographic origins, which was confirmed in this study. The strains of *L. donovani* complex clearly divided into two clades and the three isolates Cy, WenChuan and 801 were formed a subclade, which reflected that the intraspecific heterogeneity existed in Chinese *L. donovani* strains. This result supported the previous phylogenetic analysis based on HSP70 which revealed that there was genetic differentiation among Chinese *L. donovani* and some strains differentiated into *L. infantum*. Meanwhile, the RAPD result also showed that the genetic diversity was found in VL isolates from hill, plain and desert in China. Nevertheless, more strains from different origins still would be needed to obtain more accurate intra-specific classification of Chinese *L. donovani*.

Moreover, three *L. donovani* complex species-specific DNA markers in available 17 *Leishmania* strains were developed, i.e., Marker 1-AD17, Marker 2-A816 and Marker 3-O13. Through Blast and bioinformatic analysis, Marker 1-AD17 has a greater specificity for amplification of *L. donovani* complex, and Marker 3-O13 has the potential of encoding protein. All these would be needed to be verified by collection of more parasite samples and patient specimens in further study.

Abbreviations

RAPD: Random amplified polymorphic DNA; SCAR: sequence characterized amplified regions; VL: visceral Leishmaniasis; cyt *b*: cytochrome *b*; HSP70: heat shock protein 70; NTSYS: Numerical Taxonomy and Multivariate Analysis System; UPGMA: unweighted pair-group method with arithmetic means.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the *National* Natural Science Foundation of China to Dongmei Yuan (81802034), and Jianping Chen (81672048) and Post-Doctor Research Project of Sichuan University to Dongmei Yuan (2019SCU12014).

The funder had in our study was the co-first author and the correspondence author.

The funding body played an important role in accomplishment of the study and writing the manuscript.

Authors' contributions

YD carried out the RAPD studies and drafted the manuscript, QH performed the statistical analysis and edited the manuscript, CD participated in discussions about the analysis, CJ conceived and coordinated the study. All authors have read and approved the final manuscript.

Acknowledgements

We thank the help of Dr. Lin Liao from Department of Parasitology, West China College of Preclinical and Forensic Medicine, Sichuan University for cultivation of the parasites in this work.

References

1. World Health Organization. Control of the leishmaniasis. World Health Organ Tech Rep Ser. 2010;(949): xii-xiii, 1-186, back cover.
2. Akhoundi M, Kuhls K, Cannet A, Votýpka J, Marty P, Delaunay P, et al. A Historical Overview of the Classification, Evolution, and Dispersion of *Leishmania* Parasites and Sandflies. PLoS Negl Trop Dis. 2016;10(3): e0004349.
3. Guan LR. Present Situation of Visceral Leishmaniasis and Prospect for its Control in China. Chin J Parasitol Parasit Dis. 2009;27(5), 394-397.
4. Leishmaniasis in high-burden countries: an epidemiological update based on data reported in 2014. Wkly Epidemiol Rec. 2016;91(22):287-296.
5. Lun ZR, Wu MS, Chen YF, Wang JY, Zhou XN, Liao LF, et al. Visceral Leishmaniasis in China: an Endemic Disease under Control. Clin Microbiol Rev. 2015;28(4):987-1004.
6. Hu X, Lu H, Luo P, Wang Z, Hu X, Yi T. Identification of *Leishmania donovani* isolates from different kala-azar foci in China by kDNA hybridization. Chin Med Sci J. 1992;7(2):63-66. (in Chinese with English abstract).
7. Lu HG, Zhong L, Guan LR, Qu JQ, Hu XS, Chai JJ, et al. Separation of Chinese *Leishmania* isolates into five genotypes by kinetoplast and chromosomal DNA heterogeneity. Am J Trop Med Hyg. 1994;50(6):763-70.
8. Yang BB, Guo XG, Hu XS, Zhang JG, Liao L, Chen DL, et al. Species discrimination and phylogenetic inference of 17 Chinese *Leishmania* isolates based on internal transcribed spacer 1 (ITS1) sequences. Parasitol Res. 2010;107(5):1049–1065.
9. Cao DP, Guo XG, Chen DL, Chen JP. Species delimitation and phylogenetic relationships of Chinese *Leishmania* isolates reexamined using kinetoplast cytochrome oxidase II gene sequences. Parasitol Res. 2011;109(1):163-173.
10. Yang BB, Chen DL, Chen JP, Liao L, Hu XS, Xu JN. Analysis of kinetoplast cytochrome *b* gene of 16 *Leishmania* isolates from different foci of China: different species of *Leishmania* in China and their phylogenetic inference. Parasit Vectors. 2013;6:32.
11. Yuan D, Qin H, Zhang J, Liao L, Chen Q, Chen D, et al. Phylogenetic analysis of HSP70 and cyt *b* gene sequences for Chinese *Leishmania* isolates and ultrastructural characteristics of Chinese *Leishmania* Parasitol Res. 2017;116(2):693-702.
12. Wang JY, Gao CH, Yang YT, Chen HT, Zhu XH, Lu S, et al. An outbreak of the desert sub-type of zoonotic visceral leishmaniasis in Jiashi, Xinjiang Uygur Autonomous Region, People's Republic of China. Parasitol Int. 2010;59(59):331–337.
13. Guan LR, Yang YQ, Xu YX, Qu JQ, Zuo XP, Wang G, et al. Leishmaniasis in Karamay XIV. Identification of promastigote isolates from naturally infected *phlebotomus major wui*. Chin J Parasitol Parasit Dis. 1994;12(4):257–261. (In Chinese with English abstract)
14. Yang YQ, Guan LR, Wu JT. Study on the parasitic character of *Leishmania infantum* in monkey in Karamay area. Endemic Diseases Bulletin. 1997;12(1):8–11. (in Chinese with English abstract)
15. Rioux JA, Lanotte G, Serres E, Pratlong F, Bastien P, Perieres J. Taxonomy of *Leishmania*. Use of isoenzymes. Suggestions for a new classification. Ann Parasitol Hum Comp. 1990;65(3):111-125.
16. Van der Auwera G, Dujardin JC. Species typing in dermal leishmaniasis. Clin Microbiol Rev. 2015;28(2):265-294.
17. Stevens JR, Noyes HA, Schofield CJ, Gibson W. The molecular evolution of *Trypanosomatidae*. Adv Parasitol. 2001;48:1-56.
18. Akhoundi M, Downing T, Votýpka J, Kuhls K, Lukeš J, Cannet A, et al. *Leishmania* infections: Molecular targets and diagnosis. Mol Aspects Med. 2017; 57:1-29.
19. Williams JG, Kubelik AR, Livak KJ, Rafalski JA, Tingey SV. DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. Nucleic Acids Res. 1990;18(22):6531-6535.
20. Babu KN, Rajesh MK, Samsudeen K, Minoo D, Suraby EJ, Anupama K, et al. Randomly amplified polymorphic DNA (RAPD) and derived techniques. Methods Mol Biol. 2014; 1115:191-209.
21. Ikram G, Dellagi K, Ismaïl R B. Random amplified polymorphic DNA technique for identification and differentiation of old world *Leishmania* Am J Trop Med Hyg. 2002; 66(2):152-156.
22. Yazidi R, Bettaieb J, Ghawar W, Jaouadi K, Châabane S, Zaatour A, et al. RAPD-PCR reveals genetic polymorphism among *Leishmania* major strains from Tunisian patients. BMC Infect Dis. 2015; 14; 15:269.
23. Adams RP, Demeke T. Systematic relationships in *Juniperus* based on random amplified polymorphic DNA. Taxon. 1993; 42:553–571.
24. Wilkie SE, Issac PG, Slater RJ. Random amplified polymorphic DNA (RAPD) markers for genetic analysis in *Allium*. Theor Appl Genet. 1993; 86:497–504.

25. Hanafi R, Barhoumi M, Ali SB, Guizani I. Molecular analyses of Old World *Leishmania* RAPD markers and development of a PCR assay selective for parasites of the *Leishmania donovani* species Complex. *Exp Parasitol*. 2001; 98: 90–99.
26. Martinez E, Alonso V, Quispe A, Thomas MC, Alonso R, Piñero JE, et al. RAPD method useful for distinguishing *Leishmania* species: design of specific primers for *braziliensis*. *Parasitology*. 2003;127(Pt 6):513-517.
27. Lu DM, Hu XS, Qiao ZD. Analysis of *Leishmania* species and strains from China by RAPD technique. *Chin J Parasitol Paras Dis*. 2001;19(5):290–293 (in Chinese with English abstract)
28. Mkada-Driss I, Lahmadi R, Chakroun AS, Talbi C, Guerbouj S, Driss M, et al. Screening and characterization of RAPD markers in viscerotropic *Leishmania* *PLoS One*. 2014;14(9):e109773.
29. Rohlf F J. NTSYS-pc - Numerical Taxonomy and Multivariate Analysis System. Applied Biostatistics Inc. New York.1988; 2.1.
30. Lun ZR, Desser SS. Analysis of isolates within species of anuran trypanosomes using random amplified polymorphic DNA. *Parasitol Res*. 1996;82(1):22-27.
31. Barral V, Morand S, Pointier JP, Théron A. Distribution of schistosome genetic diversity within naturally infected *Rattus rattus* detected by RAPD markers. *Parasitology*. 1996;113 (Pt 6):511-517.
32. Bandi C, La Rosa G, Bardin MG, Damiani G, Comincini S, Tasciotti L, et al. Random amplified polymorphic DNA fingerprints of the eight taxa of *Trichinella* and their comparison with allozyme analysis. *Parasitology*. 1995;110 (Pt 4):401-407.
33. Guan W, Cao DP, Sun K, Xu JN, Zhang JR, Chen DL, et al. Phylogenic analysis of Chinese *Leishmania* isolates based on small subunit ribosomal RNA (SSU rRNA) and 7 spliced leader RNA (7SL RNA). *Acta Parasitol*. 2012;57(2):101–113.
34. Sun K, Guan W, Zhang JG, Wang YJ, Tian Y, Liao L, et al. Prevalence of canine leishmaniasis in Beichuan County, Sichuan, China and phylogenetic evidence for an undescribed *Leishmania* in China based on 7SL RNA. *Parasite Vector*. 2012; 5:75.
35. Hu X, Bu L, Ma Y, Wang Y, Jing B, Yi T. Difference in DNA sequences in SSU rDNA variable regions among pathogens isolated from different epidemic foci of visceral leishmaniasis in China. *Chin Med J (Engl)*. 2002;115(10):1457-1459.
36. Van Eys GJ, Guizani I, Ligthart GS, Dellagi K. A nuclear DNA probe for the identification of strains within the *Leishmania donovani* *Exp Parasitol*. 1991; 72: 459–463.
37. Lewin S, Schöniat G, El Tai N, Oskam L, Bastien P, Presber W. Strain typing in *Leishmania donovani* by using sequence-confirmed amplified region analysis. *Int J Parasitol*. 2002;32(10):1267-1276.
38. Rogers MB, Hilley JD, Dickens NJ, Wilkes J, Bates PA, Depledge DP, et al. Chromosome and gene copy number variation allow major structural change between species and strains of *Leishmania*. *Genome Res*. 2011;21(12):2129-2142.

Figures

Figure 1

The RAPD profiles of 17 *Leishmania* strains with primer OP-AD17.

Figure 2

The UPGMA tree of 17 *Leishmania* strains clustering based on genetic similarity.

Figure 3

Species-specific amplification of the three SCAR markers to *L. donovani* complex. A: 1-AD17; B: 2-A816; C: 3-O13.

Figure 4

Distribution of Blast hits of SCAR markers. a. 1-AD17; b. 2-A816; c. 3-O13.

Figure 5

Protein predictive analytics of ORF-4 of Marker 3-O13 by Lasergene Protean.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile4TabS1.docx](#)
- [Additionalfile5originalfigures.docx](#)
- [Additionalfile1Tab.1.docx](#)
- [Additionalfile2Tab.3.docx](#)
- [Additionalfile3Tab.5.docx](#)