

Association Between Stroke Occurrence And Changes In Atmospheric Circulation

Jone Vencloviene (✉ jone.vencloviene@vdu.lt)

Albertus Magnus High School

Ricardas Radisauskas

Lithuanian university of Health sciences

Daina Kranciukaite-Butylkiniene

Lithuanian University of Health sciences

Abdonas Tamosiunas

Lithuanian University of Health sciences

Vidmantas Vaiciulis

Lithuanian University of health sciences

Daiva Rastenyte

Lithuanian University of Health sciences

Research article

Keywords: ischaemic stroke, haemorrhagic stroke, atmospheric circulation, Arctic oscillation, North Atlantic oscillation, East Atlantic/West Russia indices, El Niño-Southern Oscillation

Posted Date: December 28th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-39486/v3>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 6th, 2021. See the published version at <https://doi.org/10.1186/s12889-020-10052-5>.

Abstract

Background: The impact of weather on morbidity from stroke has been analysed in previous studies. As the risk of stroke was mostly associated with changing weather, the changes in the daily stroke occurrence may be associated with changes in atmospheric circulation. The aim of our study was to detect and evaluate the association between daily numbers of ischaemic strokes (ISs) and haemorrhagic strokes (HSs) and the teleconnection pattern..

Methods: The study was performed in Kaunas, Lithuania, from 2000 to 2010. The daily numbers of ISs, subarachnoid haemorrhages (SAHs), and intracerebral haemorrhages (ICHs) were obtained from the Kaunas Stroke Register. We evaluated the association between these types of stroke and the teleconnection pattern by applying Poisson regression and adjusting for the linear trend, month, and other weather variables.

Results: During the study period, we analysed 4,038 cases (2,226 men and 1,812 women) of stroke. Of these, 3,245 (80.4%) cases were ISs, 533 (13.2%) cases were ICHs, and 260 (6.4%) cases were SAHs. An increased risk of SAH was associated with a change in mean daily atmospheric pressure over 3.9 hPa (RR=1.49, 95% CI 1.14-1.96), and a stronger El Niño event had a protective effect against SAHs (RR=0.34, 95% CI 0.16-0.69). The risk of HS was positively associated with East Atlantic/West Russia indices (RR=1.13, 95% CI 1.04-1.23). The risk of IS was negatively associated with the Arctic Oscillation index on the same day and on the previous day (RR=0.97, $p<0.033$). During November-March, the risk of HS was associated with a positive North Atlantic Oscillation (NAO) (RR=1.29, 95% CI 1.03-1.62), and the risk of IS was negatively associated with the NAO index (RR=0.92, 95% CI 0.85-0.99).

Conclusions: The results of our study provide new evidence that the North Atlantic Oscillation, Arctic Oscillation, East Atlantic/West Russia, and El Niño-Southern Oscillation pattern may affect the risk of stroke. The impact of these teleconnections is not identical for various types of stroke. Emergency services should be aware that specific weather conditions are more likely to prompt calls for more severe strokes.

Background

During the past decade, many studies have focussed on the impact of weather on morbidity from stroke, and several meta-analyses or reviews have been carried out [1-5]. Air temperature (T) and atmospheric pressure (AP) are among the most closely studied weather variables; increases, decreases, and fluctuations in both have been significantly linked to numerous stroke subtypes. A statistically significant negative association has been observed between T and the incidence of ischaemic stroke (IS) [6-8], haemorrhagic stroke (HS) [9,10], intracerebral haemorrhage [11-14], and subarachnoid haemorrhage [15-17]. The results of a meta-analysis [4] have also shown a negative association between daily T and all stroke subtypes. The associations between stroke incidence and T were stronger in Europe and were not significant in North America [4]. Daily T changes have also been associated with the risk of stroke. The

results of a meta-analysis have shown that a daily increase in T was a protective factor against HS, and an increase as well as a decrease in daily T acted as a risk factor for IS [1].

Studies have shown both a negative association [1,12,18] and a positive association [14,15,19] between the risk of stroke and AP. Studies conducted predominantly in Europe have shown an increase in the incidence of stroke associated with AP changes, adjusting for T or seasonality [19-22]. According to a study conducted in Germany, rapid decreases in ambient temperature and rapid changes in relative humidity and AP increase the risk of IS under temperate climate conditions [23]. These results showed that the risk of stroke occurrence was associated with changing weather. It is possible that the risk of stroke occurrence was associated with some patterns of atmospheric variability.

The changes in atmospheric circulation are regulated through teleconnections – large-scale patterns of pressure and circulation anomalies that cover vast geographic areas. These patterns reflect large-scale changes in atmospheric waves and influence T, rainfall, storm tract, and jet streams [24]. The most important teleconnection pattern in the Atlantic-European region is the North Atlantic Oscillation (NAO), characterised as a dipole in the sea-level pressure between the Azores high and the Icelandic low [25]. Other teleconnections such as the Arctic oscillation (AO), Scandinavian pattern (SCA), and East Atlantic/West Russia pattern (EA/WR) also determine weather variability in the Northern Hemisphere [26-27]. Several studies have also suggested a consistent and statistically significant impact of the El Niño-Southern Oscillation (ENSO) over the North Atlantic and Europe [28].

The NAO is the most prominent and recurrent pattern of atmospheric variability over the middle and high latitudes of the Northern Hemisphere, especially during the cold season months (November-March) [25]. During winter, in Northern Europe, the positive NAO phase was associated with a stronger westerly wind flow, a higher T, and increased storminess and precipitation, whereas the negative NAO phase led to a weakened westerly wind, a lower temperature, and decreased storminess and precipitation [25]. Apart from the NAO, other teleconnections such as AO, SCA, EA/WR, and ENSO also determine climate variability in the Northern Hemisphere and regulate the frequency and intensity of significant weather events [29,30,31,32,33].

As the risk of stroke has been associated mostly with changing weather, the changes in the daily stroke occurrence are probably associated with changes in atmospheric circulation. The teleconnections affecting the variability in atmospheric circulation in the Baltic region are likely to determine certain weather patterns or events that are potentially associated with the risk of stroke. The aim of this study was to detect the complex association between the daily numbers of ISs and HSs in patients aged 25-64 years and teleconnection indices – daily NAO, AO, and ENSO indices and monthly indices of EA/WR and SCA, adjusting for weather variables. In our previous study [34], the impact of T, AP, relative humidity (RH), and wind speed (WS) on the risk of stroke was evaluated. As over the middle and high latitudes of the Northern Hemisphere, the most prominent and recurrent pattern of atmospheric variability is the NAO, especially during the cold season months [25], we assessed the associations between NAO indices and the risk of strokes separately during November-March and April-October.

Methods

Patients

Data on stroke patients were obtained from the Kaunas population-based Stroke Register database. The registration of stroke cases among middle-aged (25-64 years old) Kaunas residents has been carried out from 1986 to the present day. Data collected for the period of 2000-2010 were used in this study. Stroke registration was conducted according to the WHO MONICA project protocol and established quality control procedures and has been described in detail elsewhere [35, 36]. Multiple sources of information (hospital discharge records, records of outpatient departments, necropsy, medicolegal records, and death certificates) for the stroke event register were used. According to the study protocol, every stroke event must have its apparent onset within the study period and more than 28 days from any previously recorded stroke event in the same case. Multiple stroke attacks occurring within 28 days from onset were considered as a single event. All patients suspected of having died from stroke or having had a nonfatal acute stroke were registered. The codes for the specific types of stroke were confirmed by specific diagnostic examinations. For subarachnoid haemorrhage (SAH) (ICD-10 codes I60.0-I60.9), necropsy (for fatal events), brain computed tomography (CT), or cerebrospinal fluid containing blood was required to determine the diagnosis; for intracerebral haemorrhage (ICH) (ICD-10 codes I61.0-I61.9), the diagnosis had to be confirmed by CT or by necropsy. IS (ICD-10 codes I63.0-I63.9) was diagnosed when CT and/or autopsy could verify the infarction and/or exclude haemorrhage and nonvascular disease.

Environmental variables

Based on the results of other authors, the weather variables linked to cardiovascular health were used as predictors in the regression model for stroke. In this case, teleconnections associated with changes in the weather pattern of the Baltic region were chosen as predictors. The environmental variables used in our study are routinely collected in certain publicly available databases. The values of daily NAO and AO indices (NAOIs and AOIs) and the monthly EA/WR and SCA indices (EA/WRIs and SCAs) were obtained from the database of the National Oceanic and Atmospheric Administration (NOAA) (<https://www.cpc.ncep.noaa.gov/data/teledoc/telecontents.shtml>). We included the daily NINO3.4 index (Equatorial Pacific Sea Surface Temperature), which indicates the effect of ENSO. NINO3.4 indices were taken from the NOAA database (https://climexp.knmi.nl/data/inino34_weekly.dat). Data on the mean daily T (°C), AP (hPa), RH (%), cloud cover (CC) (okta), and WS (m/s) for the studied period were obtained from the Lithuanian Hydrometeorological Service Kaunas Meteorological Station located in the suburbs of Kaunas.

Statistical analysis

The independence in the daily number of SAHs, ICHs, and ISs was tested by calculating the autocorrelation function. As no significant autocorrelations were detected, the associations between teleconnection indices and the daily number of ISs, SAHs, and ICHs were evaluated by applying a multivariate Poisson regression. First, we assessed the univariate associations between the risk of all

types of stroke and weather variables, teleconnection indices, and daily changes (Δ) in T, AP, RH, and WS. As nonlinear and U-shaped associations between weather variables and the risk of acute events were found, the weather variables were used as continuous variables or were categorised. The thresholds of categorical variables were detected by using the classification and regression tree (CRT) method [37]. Apart from this, we analysed the risk of strokes in the tertiles of environmental variables. The weather variables were included one by one in the regression model with predictors: the linear trend, T, and the day of the week and the month (the day of the week and the month were used as categorical variables). In the analysis, we used the environmental variables on the day of the stroke and on the previous 1–2 days (with lags of 0, 1, and 2 days, respectively). The choice of the weather variable (categorical or continuous) and optimal lag was made using the Akaike information criterion. Second, we analysed the impact of teleconnection indices by including the corresponding variables in the multivariate regression model with the linear trend, month, day of the week, T, and other significant weather variables. In the analysis, we used teleconnection indices as continuous and categorised by the tertiles and by using the CRT method. The optimal models were created using the Akaike information criterion. We checked the autocorrelations of the residuals using partial autocorrelation functions for the created model. Third, we analysed the impact of the NAOI on the risk of strokes separately for November-March and April-October. To assess the impact of environmental variables, we presented adjusted rate ratios (RRs) in the multivariate Poisson regression model. Statistical analysis was performed using SPSS 20 software (IBM Corp. in Armonk, NY, USA).

Results

During the study period, we analysed 4,038 cases of stroke (2,226 (55.1%) men and 1,812 (44.9%) women). Of these, 3,245 (80.4%) cases were IS, 533 (13.2%) cases were ICH, and 260 (6.4%) cases were SAH. The analysis of demographic characteristics showed that all types of stroke were more prevalent in the older age group (55-64 years). Two-thirds of all IS occurred in the 55-64 year age group. Both ICH and IS had a higher prevalence among men. The distribution of strokes by sex, age, and stroke type is presented in Table 1.

Table 1. Distribution of strokes by sex, age and stroke types

Sex/age groups	Total	SAH	ICH	HS (SAH+ICH)	IS
All, n (%)	4038 (100.0)	260 (6.4)	533 (13.2)	793 (19.6)	3245 (80.4)
Men, n (%)	2226 (55.1)	118 (5.3)	306 (13.7)	424 (19.0)	1802 (81.0)
Women, n (%)	1812 (44.9)	142 (7.9)	227 (12.5)	369 (20.4)	1443 (79.6)
25-54, n (%)	1521 (37.7)	164 (10.8)	224 (14.7)	388 (25.5)	1133 (74.5)
55-64, n (%)	2517 (62.3)	96 (3.8)	309 (12.3)	405 (16.1)	2112 (83.9)

SAH - subarachnoid haemorrhage, ICH - intracerebral haemorrhage, HS - haemorrhage stroke, IS - ischemic stroke

Over the study period, the mean daily temperature was 7.5°C, and the daily change in air temperature ranged from -14.1 to 17.7°C. The mean daily atmospheric pressure was 1015 hPa, and the daily change in atmospheric pressure ranged from -36.8 to 32.9. All descriptive characteristics of the environmental variables are presented in Table 2.

Table 2. The descriptive characteristics of weather variables

Variable	Mean (SD)	Minimum	Percentiles			Maximum
			25 th	50 th	75 th	
Air temperature, °C	7.5 (8.9)	-2.52	1.0	7.8	14.8	26.6
Daily change in air temperature	0 (2.4)	-14.1	-1.4	0	1.4	17.7
Atmospheric pressure, hPa	1015 (9.6)	976	1009	1015	1021	1050
Daily change in atmospheric pressure	0 (6.0)	-36.8	-3.4	0	3.5	32.9
Relative humidity, %	80.4 (12)	37	73	82	89	100
Daily change in relative humidity	0 (7.8)	-34	-5	0	4	39
Wind speed, m/s	5.3 (2.0)	1	4	5	6	15
Cloud cover, okta	3.5 (1.6)	0.3	2.4	3.3	4.5	11.5
AO index	-0.12 (1.5)	-5.80	-0.95	-0.06	0.79	4.70
NAO index	-0.06 (0.8)	-3.25	-0.58	-0.02	0.51	2.39
NINO3.4 index	-0.07 (0.8)	-2.20	-0.60	0.00	0.50	1.90
EA/WR indices	-0.17 (0.9)	-2.05	-0.85	-0.25	0.46	1.79
SCA indices	0.06 (0.9)	-2.33	-0.53	0.00	0.68	2.11

AO - Arctic Oscillation, EA/WR - East Atlantic/West Russia, NAO - North Atlantic Oscillation, NINO3.4 - Equatorial Pacific Sea Surface Temperature, Scandinavian pattern (SCA)

In the multivariate Poisson regression model, a change in daily AP of >3.9 hPa from the previous day, RH $>96.5\%$ on the previous day, CC >3.95 , and NINO3.4 ≤ 1.14 were statistically significantly associated with an increased rate ratio of SAH (Table 3). In other words, a stronger El Niño (NINO3.4 >1.14) had a protective impact on SAH (RR=0.34, 95% CI 0.16-0.69). A dose-response association between NINO3.4 and SAH was found. A change in daily AP of >9.55 hPa from the previous day with a lag of 1 day, daily change in RH $<7.5\%$, CC >1.35 on the previous day, and EA/WRI were associated with an increased RR of ICH. The rate ratio of HS was associated with daily increases in AP and CC >3.95 and was positively associated with the EA/WR index. In addition, a dose-response association between NINO3.4 and EA/WRI and ICH was detected. The RR of IS was positively associated with an increase in daily T over 2.2 °C compared with the previous day, RH over 53.5% with a lag of 1 day, a strong positive SCAI, NINO3.4 <-1.60 , and a strongly negative EA/WR (EA/WRI <-1.81). Apart from this, the risk of IS was negatively associated with AOI (dose-response association was detected) in the model without other teleconnections but was not statistically significant in the multivariate model ($p=0.053$) (Table 3).

Table 3. Associations* of environmental variables with different types of stroke.

Variable	Lag	Univariate association			Multivariate model		
		RR	95% CI	p	RR	95% CI	p
SAH							
Δ AP (per 10 hPa)	0	1.20	0.98-1.47	0.082			
Δ AP >3.9 hPa	0	1.48	1.12-1.94	0.005	1.49	1.14-1.96	0.004
RH >96.5%	1	1.93	1.14-3.28	0.015	2.05	1.21-3.50	0.008
CC >3.95	0	1.36	1.05-1.76	0.022	1.42	1.09-1.84	0.008
NINO3.4	0	0.87	0.75-1.01	0.065			
NINO3.4>1.14	0	0.32	0.16-0.67	0.002	0.34	0.16-0.69	0.003
ICH							
Δ T >2.6 °C	0	1.27	0.98-1.63	0.071			
Δ AP >9.55 hPa	1	1.48	1.08-2.02	0.014	1.51	1.10-2.06	0.010
RH >66.5%	1	1.39	1.00-1.93	0.051			
Δ RH <7.5%	0	1.39	1.06-1.83	0.017	1.42	1.08-1.87	0.013
CC >1.35	1	2.17	1.16-4.06	0.016	2.17	1.16-4.07	0.016
EA/WRI	0	1.16	1.05-1.30	0.005	1.16	1.04-1.29	0.006
NINO3.4>0.11	0	1.23	1.04-1.46	0.017			
HS							
Δ AP >3.9 hPa	0	1.27	1.08-1.49	0.003	1.25	1.06-1.46	0.008
Δ AP >9.55 hPa	1	1.40	1.07-1.82	0.013	1.36	1.04-1.77	0.023
Δ RH <7.5%	0	1.32	1.06-1.64	0.015	1.28	1.03-1.46	0.028
CC >3.95	0	1.20	1.03-1.39	0.017	1.21	1.04-1.41	0.012
EA/WRI	0	1.13	1.04-1.23	0.006	1.13	1.04-1.23	0.006
IS							
Δ T >2.2 °C	0	1.13	1.03-1.25	0.009	1.12	1.01-1.23	0.024
RH >53.5%	1	1.46	1.12-1.90	0.005	1.41	1.09-1.84	0.010
AOI (per 1)	0	0.97	0.94-0.99	0.008	0.98	0.95-1.00	0.053
AOI (per 1)	1	0.97	0.95-1.00	0.034			
AOI (per 1)	2	0.98	0.96-1.00	0.092			
SCAI	0	1.04	1.00-1.08	0.089			
SCAI>0.25	0	1.12	1.04-1.21	0.002	1.10	1.02-1.18	0.015
NINO3.4<-1.60	0	1.25	1.09-1.44	0.002	1.24	1.08-1.43	0.003
EA/WRI<-1.81	0	1.73	1.41-2.12	<0.001	1.59	1.29-1.96	<0.001

* adjusting for the linear trend, the month, the day of the week, and air temperature.
 Δ AP = (mean daily AP on the same day) - (mean daily AP on the previous day).

The effect of EA/WR on the risk of HS was similar in men and women. However, the effects of teleconnection variables on the risk of IS were stronger in women.

No significant associations between the NAOI and the rate ratio of strokes were found throughout the period of the study. The analysis of the RRs of strokes in NAOI quartiles (adjusting for the month, air

temperature, and other environmental variables included in the multivariate model (Table 3)) during the period of November-March and April-October showed some impact of NAOI on the risk of stroke (Figure 1). During November-March, a positive NAO was associated with an increase in the RR of HS, and the RR of IS negatively correlated with the NAOI. During April-October, only NAOI <-0.5 was associated with the RR of HS (Table 4). Moreover, during the colder period, the effect of the NAOI on the risk of IS was stronger than the effect of the AOI. The ENSO variable was negatively correlated with the risk of SAH during November-March and with the risk of IS during April-October (Table 4).

Table 4. Associations* of the NAOI with different types of stroke during November-March and April-October.

Variable	Lag	RR (95% CI)		P	
		November-March	April-October	November-March	April-October
SAH NINO3.4	0	0.74 (0.61-0.90)	0.002	1.15 (0.89-1.49)	0.282
NAOI >-0.92	0	2.48 (0.98-56.28)	0.055	0.95 (0.61-1.48)	0.809
ICH EA/WRI	0	1.25 (1.05-1.48)	0.013	1.09 (0.95-1.25)	0.209
0<NAOI <0.5	0	1.43 (1.08-1.90)	0.012		
NAOI<-0.5	0			1.25 (0.98-1.59)	0.078
HS EA/WRI	0	1.14 (0.99-1.32)	0.064	1.10 (0.99-1.23)	0.089
NAOI >0	0	1.29 (1.03-1.62)	0.030		
NAOI <-0.5	0			1.25 (1.02-1.53)	0.028
IS (1) SCAI>0.25	0	1.08 (0.96-1.21)	0.189	1.12 (1.02-1.24)	0.024
NINO3.4<-1.60	0			1.54 (1.19-2.00)	0.001
EA/WRI<-1.81	0	1.92 (1.34-2.75)	<0.001	1.58 (1.22-2.04)	0.001
NAOI (per 1)	0	0.92 (0.85-0.99)	0.028	1.04 (0.98-1.11)	0.166
IS (2) SCAI>0.25	0	1.07 (0.96-1.21)	0.221	1.12 (1.01-1.24)	0.027
NINO3.4<-1.60	0			1.55 (1.19-2.01)	0.001
EA/WRI<-1.81	0	2.00 (1.40-2.87)	<0.001	1.59 (1.23-2.05)	<0.001
NAOI <0.5	0	1.25 (1.10-1.41)	0.001	0.91 (0.81-1.02)	0.097

* adjusting for the linear trend, the month, the day of the week, and T, RH, and CC variables

Discussion

In this 11-year study of 260 patients with SAH, 533 patients with ICH, and 3,245 patients with IS, we found that some patterns of RH and CC and daily changes in AP and RH were associated with the risk of

some types of stroke. In addition, we used the NAO, AO, EA/WR, SCA, and ENSO indices as predictors for the evaluation the risk of stroke. For the first time, we detected a protective effect of warmer ENSO (a stronger El Niño) on the risk of SAH, a positive association between the risk of HS and the EA/WR, a negative association between the rate ratio of IS and the AOI, and a negative impact of a strong positive SCA on the risk of IS. Apart from these results, during November-March, a higher risk of HS was related to a positive NAO, and a negative correlation between the risk of IS and the NAOI was found. In the analysis, an impact of teleconnection indices was detected, adjusting for seasonal variation, T, and other weather variables.

In our study, the risk of HS was associated with daily changes in AP above the threshold of 3.9 hPa for SAH and 9.55 hPa for ICH. These results are in line with those obtained by other authors who found a significant association between SAH and ICH and changing AP. The daily change in AP with a lag of 1 day was positively correlated with the daily number of SAHs in the English Midlands [19], and the daily change in AP >10 hPa was associated with the risk of SAH in the UK [38] and in Germany [20].

To our knowledge, the risk of SAH was associated with a higher RH and CC level, and the risk of ICH was associated with a higher CC level on the previous day and with a lower daily RH change. Some authors found a significant association between stroke and RH and daily hours of sunshine, which is the opposite variable to cloud cover. A decrease in daily sunlight hours was positively associated with the risk of SAH [17,39]. A positive association between SAH and RH was found in the humid subtropical zone [16] and in the Rhein Main area [20], while in the southern regions of France and in the areas with various climatic conditions (41 states of USA), a negative association between SAH and RH was observed [17, 39]. ICH is negatively correlated with sunshine hours [40] and positively correlated with the amount of precipitation [6], which coincides with our results.

We found a positive effect of a strong warm ENSO on human health. In our study, most of these events (NINO3.4>1.14) fell into colder months. Therefore, the protective effect of a warmer ENSO on SAH may be explained by the effect of ENSO on the weather pattern during autumn and winter. Studies have shown that the warm and cold phases of ENSO have different impacts on the pattern of weather regimes during the colder season in Europe [33]. Strong El Niño events were related to a higher sea level pressure, a lower T, and dry air in the Baltic countries [41,33,42]. According to our data, during the second half of autumn, the NINO3.4>1.14 period was characterised by a lower mean AP, more precipitation and RH, and a very significantly lower diurnal temperature range (DTR) (by 1.3 °C, p<0.001). In winter, the NINO3.4>1.14 period was characterised by a lower T, WS, and CC and a higher AP and DTR. These weather patterns may be associated with a lower risk of SAH. Some studies have shown that a higher DTR is significantly associated with higher mortality, and this effect was stronger during autumn [43,44]. Therefore, a lower DTR during autumn may have a positive effect on human health.

We found a positive association between the EA/WRI and HS and an increase in the risk of IS during a strong negative EA/WR phase. These effects were similar during both the colder and the warmer periods, and the additional inclusion of the NAOI in the model did not reduce the significance of the EA/WRI. In the

studied region, in winter, the positive EA/WR produced cold advection from the north and was characterised by a lower air temperature, a lower precipitation level, and stronger atmospheric circulation [45]. In the southeastern region of the Baltic Sea, the EA/WRI is negatively correlated with air temperature in spring [46], lake water temperature in spring-autumn [47], and precipitation amount in summer [45]. According to our data, during the positive EA/WR phase, a higher mean WS and AP in winter and a lower mean T and a higher mean AP both in spring and summer were observed. Apart from this, a higher variation was found in the daily change in AP both in winter and spring. Thus, a higher EA/WRI was related to a stronger variation in AP, colder air flow in winter, and colder air in other seasons. The complexity of these weather patterns may be associated with a higher risk of HS.

For the first time, negative correlations between IS and AO and between IS and NAO only during November-March were found. According to the results of studies by other authors, AO was associated not only with tropospheric variability but also with stratospheric variability and changes in weather patterns in Lithuania and nearby regions [48,49]. During January-March, a positive AO brings a higher surface T and a lower precipitation in middle-latitude regions [50]. In the region of the Baltic Sea, a positive correlation between T and AO was observed during January-March [51], March-May [52], July and October [46], and September-March [53]. According to our data, the AOI was positively correlated with T in all seasons, excluding summer, and negatively correlated with RH, excluding winter. It is possible that this complex of weather patterns (a higher T and a lower RH during the equinox, a lower RH in summer, and warmer winters) related to days of a higher AOI had a protective effect against the risk of IS.

The SCAI was positively correlated with T in summer and negatively correlated with T and positively correlated with AP over the region of the Baltic Sea in winter [26,31,46]; the same associations were found in our study. The positive phase of SCA indicates more likely anticyclonic conditions and a lower level of atmospheric circulation over the Baltic Sea region during autumn-spring [26], and the anticyclonic conditions over Scandinavia substantially suppress westerly zonal airflow in summer [54]. It is possible that cold outbreaks during the colder period and the atmospheric variations related to a stronger positive SCA are associated with the risk of IS.

During the colder period, a positive NAO had a protective effect against IS, but a negative NAO had a protective effect against HS. During wintertime, in the Baltic Sea region, a positive NAOI was associated with a higher T and with altered weather: a higher WS, a lower AP, and a northeastward shift in the Atlantic storm activity with enhanced activity from Newfoundland into Northern Europe [25]. As a positive NAO during the winter was associated with more changing weather, a positive NAO was risky for HS, whereas a change in T was more relevant for IS. Studies in Northern and Middle Europe have shown a higher risk of HS associated with changing weather but not with T [19,20,22,38]. A study conducted in the UK showed a significant impact of changes in T only on IS, whereas changes in AP had a significant impact only on the risk of HS [55].

According to the literature, both positive and negative NAO phases are associated with worse health outcomes. [56] found a positive association between the daily AOI with a lag of 3 days and the incidence

of and mortality from acute myocardial infarction in Northern Sweden. An inverse association between the climate index (which represents winters with a strong negative phase of the NAO) and the level of mortality from ischaemic heart disease was found in England [57]. In addition, a negative association between the NAO index and systolic and diastolic blood pressure during spring-autumn was found [58].

The pathophysiological mechanisms underlying the correlation between stroke and weather conditions have been discussed. Factors that increase the risk of stroke include high blood pressure, some diseases, and the lack of regular exercise. Blood pressure is influenced by cold, stress and physical activity [58]. Donkelaar et al. [22] hypothesised that AP changes trigger the inflammation process in the aneurysm wall. Variations in AP may influence vessel walls and endothelial function by endogenous inflammatory mechanisms [21]. Studies on thrombosis in air travel suggest that prothrombin fragments and the thrombin-antithrombin complex are activated in hypobaric conditions [59,60], which could be another clue to the underlying mechanism.

Studies on the associations between physical activity in the elderly and weather conditions in Europe showed that physical activity decreased with increasing WS, precipitation, humidity, and a shorter duration of sunshine [61,62]. These weather conditions are associated with a negative AO and NAO excluding winter months; therefore, it can be assumed that negative AOs are associated with fewer physical activity opportunities for the elderly, who are likely to be stressed. This activity can explain why a negative AO increased the risk of IS.

The present study has several **strengths** the inclusion of a large number of patients with various types and subtypes of stroke, the long study period, and standardised methods and criteria used for stroke registration. In addition, the present study analyses daily stroke incidence data by stroke subtypes, daily meteorological data, and the variation of these data with respect to the previous day and uses teleconnections such as NAO, AO, EA/WR, SCA, and ENSO. Moreover, the patients that were included coming from a small geographical area (Kaunas) contributes to the homogeneity of weather conditions. In our study, associations between atmospheric circulation patterns such as NAO, AO EA/WR, SCA, and ENSO and the risk of stroke were found for the first time.

The **limitation** is that other potential confounders such as air pollution, influenza epidemics or other respiratory infections were not directly considered in this study. In our region, infections are strongly related to the season with the highest prevalence during winter [63]. In our study, the analyses were controlled for the month and T. Residual confounding by short-term respiratory epidemics remains a possibility. Moreover, we did not consider weather-related physical activity that may have had an impact on individual exposure to outdoor T and humidity.

In our study, the influence of air pollution, which is a known trigger for cardiovascular diseases, was not examined. We did not have air pollution data for the entire study period, but the additional inclusion of the daily concentrations of PM₁₀, NO₂, or O₃ did not change the association between the risk of strokes and teleconnection indices. Based on the results published by other authors [64], we can assume that the

short-term effect of PM₁₀, NO₂, or O₃ on the risk of stroke was not significant enough to affect the results of our study. First, the level of air pollution in Kaunas is not high. Second, our stroke patients were relatively young (<65 years of age), whereas in other studies presenting a positive association between air pollution and stroke (except for those performed in subtropics or in regions with high levels of pollutants), the mean age of the patients was over 70 years [68]. Third, we did not evaluate other comorbidities such as acute myocardial infarction, ischaemic heart disease, arterial hypertension, heart failure or other risk factors such as atrial fibrillation, diabetes, dyslipidaemia, or renal or malignant diseases, which may also be associated with a higher risk of ischaemic and haemorrhagic stroke. Fourth, harmful lifestyle factors such as alcohol consumption or smoking, which increase the risk of haemorrhagic stroke, cannot be ruled out, either.

Conclusion

A protective effect of a stronger El Niño on SAH was found. The risk of HS was associated with an increase in daily AP, a higher CC, and a higher EA/WR index. The risk of IS was negatively associated with the AO index, and a negative impact of a strong positive SCA on the risk of IS was found. During the colder period, a positive NAO had a protective effect against IS, but, vice versa, a negative NAO had a protective effect against HS. The results of our study provide new evidence that the NAO, AO, SCA, EA/WR, and ENSO patterns may affect the risk of stroke. The impact of these teleconnections on various types of stroke is not identical. As U-shaped or J-shaped relationships between daily changes in weather variables and stroke events were found, we recommend using the weather variables as categorical factors in the regression models. Emergency services should be aware that specific weather conditions are more likely to prompt calls for more severe strokes.

Abbreviations

IS: ischaemic stroke; **SAHs:** subarachnoid haemorrhages; **ICHs:** intracerebral haemorrhages; **EA/WR:** East Atlantic/West Russia pattern; **HS:** haemorrhagic stroke **AO:** Arctic Oscillation; **NAO:** North Atlantic Oscillation; **T:** temperature; **AP:** atmospheric pressure; **SCA:** Scandinavian pattern; **ENSO:** El Niño-Southern Oscillation; **RH:** relative humidity; **WS:** wind speed; **WHO:** World Health Organisation; **CT:** computer tomography; **AOI:** Arctic Oscillation index; **NAOI:** North Atlantic Oscillation index; **EA/WRI:** West Russia index; **SCAI:** Scandinavian pattern index; **CC:** cloud cover; **CRT:** classification and regression tree.

Declarations

Acknowledgements

The authors are grateful to the other members of the Kaunas Stroke Registry team for their invaluable assistance in data collection and validation. We thank the organisers of the 31st Conference of the International Society for Environmental Epidemiology for the opportunity to present some of the results of this article.

Authors' contributions

JV conceived the idea for this study, designed the analysis plan, performed a statistical analysis, and interpreted the data; **RR** conceived the idea for this study, designed the analysis plan, interpreted the data, and drafted and revised the manuscript for important intellectual content; **AT** conceived the idea for this study, designed the analysis plan, and revised the manuscript for important intellectual content; **DR** conceived the idea for this study, interpreted the data, and revised the manuscript for important intellectual content; **DKB** and **VV** conducted the data analysis, interpreted the data, and drafted the manuscript. The authors had full access to the data and take responsibility for their integrity. All authors have read and agreed to the manuscript as written. The author(s) read and approved the final manuscript.

Funding

No funding has been provided by any private or public sources.

Availability of data and materials

Data that were used in this article have not open access. These data can be accessible to university scientific staff with the permission of the head of the Laboratory of Population Studies.

Ethics approval and consent to participate

All patient data were analysed anonymously and deidentified prior to analysis. All methods were performed in accordance with the relevant guidelines (MONICA project) and regulations (Lithuanian Bioethics Committee). All the patients who participated in the study signed their informed consent. The study protocol was approved by Lithuanian Biomedical Research Ethics Committee No 14-27. We also state that the study ethics comply with the Declaration of Helsinki.

Consent for publication

Not applicable.

Competing interests

The authors declared no conflict of interest.

References

1. Lian H, Ruan Y, Liang R, Liu X, Fan Z. Short-Term Effect of Ambient Temperature and the Risk of Stroke: A Systematic Review and Meta-Analysis. *Int. J. Environ. Res. Public Health*. 2015;12:9068–9088.
2. Wang X, Cao Y, Hong D, Zheng D, Richter S, Sandset EC et al. Ambient Temperature and Stroke Occurrence: A Systematic Review and Meta-Analysis. *J. Environ. Res. Public Health*. 2016;13:698.

3. Cao Y, Wang X, Zheng D, Robinson T, Hong D, Richtering S et al. Air Pressure, Humidity and Stroke Occurrence: A Systematic Review and Meta-Analysis. *Int. J. Environ. Res. Public Health*. 2016;13:675.
4. Zorrilla-Vaca A, Healy RJ, Silva-Medina M, Revealing the association between cerebrovascular accidents and ambient temperature: a meta-analysis. *Int. J. Biometeorol*. 2017;61(5):821–832.
5. Lavados PM, Olavarriza VV, Hoffmeister L. Ambient Temperature and Stroke Risk Evidence Supporting a Short-Term Effect at a Population Level From Acute Environmental Exposures. *Stroke*. 2018;49:255–261.
6. Magalhães R, Silva MC, Correia M, Bailey T. Are stroke occurrence and outcome related to weather parameters? Results from a population-based study in northern Portugal. *Dis*. 2011;32(6):542–551.
7. Mostofsky E, Penner EA, Mittleman MA. Outbursts of anger as a trigger of acute cardiovascular events: a systematic review and meta-analysis. *Heart J*. 2014;35:1404–1410.
8. Wang Q, Gao C, Wang H, Lang L, Yue T, Lin, H. Ischemic stroke hospital admission associated with ambient temperature in Jinan, China. *PLoS One*. 2013;8:1–8.
9. Goggins WB, Woo J, Ho S, Chan EY, Chau PH. Weather, season, and daily stroke admissions in Hong Kong. *Int. J. Biometeorol*. 2012;56:865–872.
10. Luo Y, Li H, Huang F, Van Halm-Lutterodt N, Xu Q, Wang A et al. The cold effect of ambient temperature on ischemic and hemorrhagic stroke hospital admissions: A large database study in Beijing, China between years 2013 and 2014-Utilizing a distributed lag non-linear analysis. *Environ Pollut*. 2018;232:90–96.
11. Morabito M, Crisci A, Vallorani R, Modesti PA, Gensini GF, Orlandini S. Innovative approaches helpful to enhance knowledge on weather-related stroke events over a wide geographical area and a large population. 2011;42(3):593–600.
12. Kim BJ, Kang HG, Kim HJ, Ahn SH, Kim NY, Warach S et al. Magnetic resonance imaging in acute ischemic stroke treatment. *J Stroke*. 2014;16(3):131–145.
13. Han MH, Yi HJ, Ko Y, Kim YS, Lee YJ. Association between haemorrhagic stroke occurrence and meteorological factors and pollutants. *BMC Neurology*. 2016;16:59.
14. Mukai T, Hosomi N, Tsunematsu M, Sueda Y, Shimoe Y, Ohshita T et al. Various meteorological conditions exhibit both immediate and delayed influences on the risk of stroke events: The HEWS-stroke study. *PLoS One*. 2017;12(6):1–13.
15. Abe T, Ohde S, Ishimatsu S, Ogata H, Hasegawa T, Nakamura T et al. Effects of meteorological factors on the onset of subarachnoid hemorrhage: a time-series analysis. *J. Clin. Neurosci*. 2008;15(9):1005–1010.
16. Gill RS, Hambridge HL, Schneider EB, Hanff T, Tamargo RJ, Nyquist P. Falling temperature and colder weather are associated with an increased risk of aneurysmal subarachnoid hemorrhage. *World Neurosurg*. 2013;79(1):136–42.
17. Lai PM, Dasenbrock H, Du R. The Association between Meteorological Parameters and Aneurysmal Subarachnoid Hemorrhage: A Nationwide Analysis. *PLoS One*. 2014;9(11): e112961.

18. Tarnoki AD, Turker A, Tarnoki DL, İyisoy MS, Szilagyi BK, Duong H et al. Relationship between weather conditions and admissions for ischemic stroke and subarachnoid hemorrhage. *Med. J.* 2017;58(1):56–62.
19. Buxton N, Liu C, Dasic D, Moody P, Hope DT. Relationship of aneurysmal subarachnoid hemorrhage to changes in atmospheric pressure: results of a prospective study. *Neurosurg.* 2001;95(3):391–392.
20. Setzer M, Beck J, Hermann E, Raabe A, Seifert V, Vatter H et al. The influence of barometric pressure changes and standard meteorological variables on the occurrence and clinical features of subarachnoid hemorrhage. *Surg. Neurol.* 2007;67:264–272.
21. Jimenez-Conde J, Ois A, Gomis M, Campello AR, Godia EC, Subirana I et al. Weather as a Trigger of Stroke. Daily meteorological factors and incidence of stroke subtypes. *Cerebrovasc. Dis.* 2008;26:348–354.
22. Van Donkelaar CE, Potgieser ARE, Groen H, Foumani M, Abdulrahman H, Sluijter R et al. Atmospheric Pressure Variation is a Delayed Trigger for Aneurysmal Subarachnoid Hemorrhage. *World Neurosurg.* 2018;112:e783–e790.
23. Rakers F, Schiffner R, Rupprecht S, Brandstädt A, Witte OW, Walther M et al. Rapid weather changes are associated with increased ischemic stroke risk: a case-crossover study. *Eur J Epidemiol.* 2016;31(2):137–146.
24. Feldstein S, Franzke C. Atmospheric Teleconnection Patterns. *Nonlinear and Stochastic Climate Dynamics.* 2017;54-104.
25. Hurrell JW, Kushnir Y, Ottersen G, Visbeck M. An overview of the North Atlantic Oscillation in The North Atlantic oscillation: climatic significance and environmental impact. *Geophys. Monogr. Ser.* 2003;134:279.
26. Bueh C, Nakamura H. Scandinavian pattern and its climatic impact. *Quarterly Journal of the Royal Meteorological Society.* 2007;133(629):2117–2131.
27. Jakobson L, Jakobson E, Post P, Jaagus J. Atmospheric teleconnections between the Arctic and the eastern Baltic Sea regions. *Earth Syst. Dynam.* 2017;8:1019–1030.
28. Rodríguez-Fonseca B, Suárez-Moreno R, Ayarzagüena B, López-Parages J, Gómara I, Villamayor J et al. Review of ENSO Influence on the North Atlantic. *Non-Stationary Signal.* 2016;7:87.
29. Thompson DWJ, Wallace JM. The Arctic oscillation signature in the wintertime geopotential height and temperature fields. *Res. Lett.* 1998;25:1297–1300.
30. Thompson DWJ, Wallace JM. Regional Climate Impacts of the Northern Hemisphere Annular Mode. *Science.* 2001; 293:85–89.
31. Trenberth KE, Caron JM. 2000. The Southern Oscillation revisited: Sea level pressures, surface temperatures, and precipitation. *Journal of Climate.* 2000;13(24): 4358-4365.
32. Moron P, Guy P. The impact of ELNÍÑO – southern oscillation upon weather regimes over Europe and the North Atlantic during borealwintervincent. *Int. J. Climatol.* 2003;23:363 – 379.

33. Liu Y, Wang L, Zhou W, Chen W. Three Eurasian teleconnection patterns: spatial structures, temporal variability, and associated winter climate anomalies. *Climate Dynamics*. 2014;42(11–12):2817–2839.
34. Tamasauskienė L, Rastenyte D, Radisauskas R, Tamosiunas A, Tamasauskas D, Vaiciulis V et al. Relationship of meteorological factors and acute stroke events in Kaunas (Lithuania) in 2000-2010. *Environ. Sci. Pollut. Res. Int.* 24, 9286–9293 (2017).
35. WHO MONICA Project, MONICA Manual. Revised ed. Geneva, Switzerland: Cardiovascular Diseases Unit, World Health Organization. 1990.
36. Radisauskas R, Malinauskienė V, Milinaviciene E, Kranciukaite-Butylkiniene D, Tamosiunas A, Bernotiene G et al. Trends in the Attack Rates, Incidence, and Mortality of Stroke during 1986-2012: Data of Kaunas (Lithuania) Stroke Registry. *PLoS One*. 2016;11(4):e0153942.
37. Breiman L, Friedman JH, Olshen RA, Stone CJ. *Classification and Regression Trees*. Wadsworth Inc, 1984.
38. Landers AT, Narotam PK, Govender ST, Van Dellen JR. The effect of changes in barometric pressure on the risk of rupture of intracranial aneurysms. *Br. J. Neurosurg.* 1997;11:191–195.
39. Lejeune JP, Vinchon M, Amouyel P, Escartin T, Escartin D, Christiaens JL et al. Association of occurrence of aneurysmal bleeding with meteorologic variations in the north of France. *Stroke*. 1994;25:338–341.
40. Tsementzis SA, Kennet RP, Hitchcock ER, Gill JS, Beevers DG. Seasonal variation of cerebrovascular diseases. *Acta Neurochir.* 1991;111(3-4):80–83.
41. Fraedrich K. An ENSO impact on Europe? *Tellus A*. 1994;46:540–552.
42. King MP, Yu E, Sillmann J. Impact of strong and extreme El Niños on European hydroclimate. *Tellus A: Dynamic Meteorology and Oceanography*. 2020;72:1-10.
43. Lim YH, Hong YC, Kim H. Effects of diurnal temperature range on cardiovascular and respiratory hospital admissions in Korea. *Sci. Total Environ.* 2012;417-418:55–60.
44. Zhang Y, Peng M, Wang L, Yu Ch. Association of diurnal temperature range with daily mortality in England and Wales: A nationwide time-series study. *Science of the Total Environment*. 2018;619–620:291–300.
45. Young-Kwon L. The East Atlantic/West Russia (EA/WR) teleconnection in the North Atlantic: climate impact and relation to Rossby wave propagation. *Climate Dynamics*. 2015; 44:11-12.
46. Rust HW, Richling A, Bissolli P, Ulbrich U. Linking teleconnection patterns to European temperature – a multiple linear regression model. *Meteorologische Zeitschrift*. 2015;24(4):411–423.
47. Graf R, Wrzesinski D. Relationship between water temperature of Polish rivers and large-scale atmospheric. *Circulation* 2019;11:1690.
48. Kodera K, Kuroda Y. Regional and hemispheric circulation patterns in the Northern Hemisphere winter, or the NAO and the AO. *Geophys. Res. Lett.* 2003;30(18):1934.

49. Christiansen B. On the physical nature of the Arctic Oscillation. *Geophys. Res. Lett.* 2002;29(16):1805.
50. Matsu K, Heki K. Anomalous precipitation signatures of the Arctic Oscillation in the time-variable gravity field by GRACE. *Geophys. J. Int.* 2012;190:1495–1506.
51. Wang D, Wang CH, Yang X, Lu J. Winter Northern Hemisphere surface air temperature variability associated with the Arctic Oscillation and North Atlantic Oscillation. *Geophys. Res. Lett.* 2005;32:L16706.
52. Creilson JK, Fishman J, Wozniak AE. Arctic Oscillation–induced variability in satellite-derived tropospheric ozone. *Geophys. Res. Lett.* 2005;32:L14822.
53. Ptak M, Arkadiusz M, Wrzesinski T, Wrzesinski D. Effect of Teleconnection Patterns on Changes in Water Temperature in Polish Lakes. *Atmosphere.* 2018;9:66.
54. Bednorz E, Pórolniczak M, Czernecki B, Arkadiusz MT. Atmospheric Forcing of Coastal Upwelling in the Southern Baltic Sea Basin. 2019;10:327.
55. Dawson J, Weir C, Wright F, Bryden C, Aslanyan S, Lees K. Associations between meteorological variables and acute stroke hospital admissions in the west of Scotland. *Acta. Neurol. Scand.* 2008;117:85–89.
56. Messner T, Lundberg V, Wikström, B. The Arctic Oscillation and incidence of acute myocardial infarction. *J. Intern. Med.* 2003;253(6):666–670.
57. McGregor GR. Winter North Atlantic Oscillation, temperature and ischaemic heart disease mortality in three English counties. *Int. J. Biometeorol.* 2005;49(3):197–204.
58. Vencloviene J, Tamosiunas A, Radisauskas R, Luksiene D, Vaiciulis V, Bernotiene G et al. The influence of the North Atlantic Oscillation index on arterial blood pressure. *J. Hypertens.* 2019;37(3):513–521.
59. Gungor L, Onar MK. Long-distance air travel: a potential new risk factor for cerebral sinus thrombosis? *Cerebrovasc. Dis.* 2007;24:312-314.
60. Schreijer AJ, Cannegieter SC, Meijers JC, Middeldorp S, Büller HR, Rosendaal FR. Activation of coagulation system during air travel: A crossover study. *Lancet.* 2006;367:832–838.
61. Klenk J, Buchele G, Rapp K, Franke S, Peter R. Walking on sunshine: effect of weather conditions on physical activity in older people. *J. Epidemiol. Community Health.* 2012;66:474–476.
62. Wu YT, Luben R, Wareham N, Griffin S, Jones AP. Weather, day length and physical activity in older adults: cross-sectional results from the European Prospective Investigation into Cancer and Nutrition (EPIC) Norfolk Cohort. *PLoS One.* 2017;12:e0177767.
63. Caplinskas S, Caplinskiene I, Davidaviciene E, Korablioviene J, Liausėdienė R, Razmuvienė D et al. Review of the incidence of infectious diseases in Lithuania 2016. *Centre for Communicable Diseases and AIDS.* 2017; ISSN 2351–6577.
64. Fu P, Guo X, Cheung FMH, Yung KKL. The association between PM2.5 exposure and neurological disorders: A systematic review and meta-analysis. *Science of the Total Environment.*

Figures




Figure 1

Rate ratios of strokes in the quartiles of the NAOI (reference category – NAOI<-0.58).