

Level of medication self-management capacity among patients on ambulatory care exiting hospital pharmacy at Muhimbili National hospital, Tanzania: a descriptive cross-sectional study

Hamu Joseph Mlyuka (✉ hmlyuka2011@gmail.com)

Muhimbili University of Health and Allied Sciences School of Pharmacy <https://orcid.org/0000-0002-5302-6218>

Hija Salehe

Muhimbili University of Health and Allied Sciences School of Pharmacy

Wigilya P Mikomangwa

Muhimbili University of Health and Allied Sciences School of Pharmacy

Manase E Kilonzi

Muhimbili University of Health and Allied Sciences School of Pharmacy

Alphonse I Marealle

Muhimbili University of Health and Allied Sciences School of Pharmacy

Ritah F Mutagonda

Muhimbili University of Health and Allied Sciences School of Pharmacy

George M. Bwire

Muhimbili University of Health and Allied Sciences School of Pharmacy

Research note

Keywords: Medication self-management, Medication management capacity, Ambulatory care

Posted Date: November 17th, 2019

DOI: <https://doi.org/10.21203/rs.2.13245/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on November 7th, 2019. See the published version at <https://doi.org/10.1186/s13104-019-4772-4>.

Abstract

Objectives: Medication management capacity of a patient on ambulatory care is directly related to adherence. To our knowledge data on medication management capacity for ambulatory care patients exiting outpatient pharmacy outlets in Tanzania are scarce. This study aimed to determine the level of medication management capacity among patients on ambulatory care exiting Muhimbili National Hospital outpatient pharmacy outlet. Results: A total of 424 patients on ambulatory care participated in the study. Three hundred eighty seven (91.3%) out of 424 interview questionnaires had complete data and qualified for data analysis. Majority (62.3%) out of 387 study participants had poor medication management capacity; 65.3% out of 387 patients were unable to correctly read the prescription and match the drugs they are carrying. More than half (57.4%) out of 387 participants were unable to correctly take the dose, 73.9% out of 387 were unable to correctly tell the dosing frequency and duration. Only 10.6% out of 155 patients with prescription containing drugs with warning or precaution or contraindication or potential side effects were aware.

Introduction

Adherence to medication is a key determinant of regimen effectiveness and safety. It is also a mediator of medical practices and patients' outcomes [1]. Almost 20-50% of patients do not take their medications as prescribed and the situation is even worse in chronic condition [2]. In USA non-adherence causes 125000 death and 10% hospital admission annually. Enormous healthcare financial burden up to US\$ 289 billion per annum are attributed by medication non adherence [3].

Adherence to medication is directly related to medication management capacity (MMC) of an individual patient [4][5]. MMC is defined as "the cognitive and functional ability to self-administer a medication regimen as it has been prescribed" [6]. Functional skills like correctly identifying the medication, opening the container, taking the right dose and timing the use of medication are components of medication self management. In an emphasis of this relationship Bailey and colleagues developed "medication self-management model" which aims to deconstruct the tasks involved in taking medication [7]. Medication management capacity tells us the accuracy of using medication while adherence based on pill count, refill or self-report provide an insight of how often one is taking the medication. The intervention focusing on boosting patients MMC have resulted into reduction in emergency department visiting and hospital admission [8].

MMC can be measured using Drug Regimen Unassisted Grading Scale (DRUGS) or Medication Management Ability Assessment (MMAA). DRUGS is used to measure MMC by considering the four steps which are identification of the right medication, accessing by opening the container, picking the correct number/amount of tablet or syrup and providing elaborative timing of using the medication [2].

So far, in Tanzania the data on level of medication management are dearth. This study aimed to assess the level of medication management capacity among clients exiting hospital pharmacy outlets at Muhimbili National Hospital, Dar es Salaam Tanzania.

Methods

Study area and design

This was a descriptive cross-sectional study conducted from February to July, 2018 at Muhimbili National Hospital (MNH) outpatient pharmacy outlets. MNH is a national referral tertiary level research center and university teaching hospital with 1,500 beds facility, attending 1,000 to 1,200 outpatients per day. The hospital is well equipped with laboratories, modern diagnostic facilities, specialized clinics as well as highly skilled manpower in all cadres of healthcare providers.

Study population and sample size

To be included in this study participant was supposed to be; 1) from outpatient clinic in MNH 2) carrying a prescription with more than one drug for oral use 3) carrying medicines he/she has received from pharmaceutical personnel at MNH outpatient pharmacy outlets 3) not blind 4) not deaf 5) sober 6) not a guardian 7) with a regimen of more than three days of use. Owing to scarcity of data on prevalence of level of medication management capacity (MMC) the 50% proportion was set as a reference population proportion to calculate the sample size. At the Z-score of 1.96 and margin of error of 5% the sample size was 385. To consider for partial filled or inconsistency 10% of initial sample was added to make 424.

Sampling technique

Consecutive sampling was employed to recruit study participants. The researcher stayed at the exit point of the outpatient pharmacy building. When the client was about to get outside the pharmacy building the researcher requested him/her for a brief discussion. If the client agreed, the purpose of the study was explained, and then he/she signed freely obtained consent form. Assessment for inclusion and exclusion criteria was done and those who fulfilled requirements were asked to continue with the interview.

Data collection

Face-to-face interview using semi-structured questionnaire was used to gather data for fulfilling this study objectives. The questionnaire consisted questions and directives to assess medication self management capacity. This was a minor modified adoption of DRUGS tool which was used by Kripalani and colleague [6] in assessing MMC. On medication identification part; patient was asked to read what has been written on the prescription, show the particular drug, tell about warning, precaution or contraindication, important possible side effects and interactions with food or other drugs on the prescription. The researcher used Medscape interaction checker online software to check for potential life threatening interaction and match with the participant's response. The patient was then asked to open the identified drug and the researcher observed the process of opening. After opening the container, the patient was asked to tell the number of tablet(s) or amount of syrup/suspension/solution which he/she is suppose to be taking per dose. Finally the patient was asked to tell the researcher about intervals of time to take the next dose and the total duration he/she will be using the medication. The steps were repeated for each of the medication on the prescription. For each question, the overall response was correct only if the participant has managed to perform a particular task correctly for all the prescribed medications on the prescription. Modified Bloom cut off point from study by Abdullah and colleagues [9] was used to categorize the MMC of the participants. The MMC was rated as poor/inadequate (score <50%), moderate (score: 50-59%), good (score: 60-69%), very good (70-79%) and excellent (80-100%).

Data analysis

Analysis was done using statistical package for social sciences (SPSS) version 23. Categorical data like gender, age groups, levels of education, level of MMC were summarized using frequency distribution and bar charts.

Results

A total of 387 responded questionnaires qualified for data analysis. Patient characteristics are summarized on Table 1. Majority of the study participants (66.4%) were women. Many participants aged between 30-49 years and more than half attained secondary level of education.

Table 1: Patients socio-demographic characteristics

Characteristics (N=387)	
Gender n (%)	
Male	130 (33.6)
Female	257(66.4)
Ages, years, n (%)	
20-29	32(8.30)
30-39	133(34.4)
40-49	145(37.5)
50-59	50(12.9)
>60	27(7)
Marital status n (%)	
Single	66(17.1)
Married	218(56.3)
Divorced	21(5.4)
Widowed	19(4.9)
Cohabit	63(16.3)
Education level n (%)	
Never	15(3.9)
Primary	88(22.7)
Secondary	208(53.7)
Tertiary	76(19.6)

Patients' medication management capacity

The overall grading showed that 241 (62.3%) of 387 study participants had poor MMC (Fig.1). Majority of these participants were not able to identify, take the right dose or timing the frequency and duration as detailed below.

Figure 1: Bar chart of prevalence of medication management capacity

Table 2 shows the response to questions and activities to measure MMC. More than half of study participants (65.3%) were not able to correctly read and interpret the instructions written on their prescriptions and match with the medications on hand. All patients demonstrated an ability to open a container and take the medication. When asked to count pills or measure the mockup amount as directed on the prescription 222(57.4%) were not able to correctly take the prescribed dose. When asked to explain the interval of hours to take the next dose and duration for each of the prescribed medications only 26.1% out of 387 participants were able to explain correctly.

Table 2: Response to attributes of measuring medication self management

Patient's responses on medication self management questions or activities (N=387)

Item question or activity	Proportion of response (%)
Can you read the prescription of your medications and match the drug? (Able if read correctly for every drug on the Rx)	
Able	34.7
Unable	65.3
Show how to open the container of your medications	
Able	100
Unable	0
Show the number of tablets/capsule or amount of suspension/solution/syrup which you are suppose to take as prescribed (Correct if every drug number or amount is correct for all drugs on Rx)	
Correct	42.6
Incorrect	57.4
How many hours are supposed to elapse before you take another dose and how long will you take each drug? (Overall correct if the frequency and duration of every drug on the Rx is correctly explained)?	
Correct	26.1
Incorrect	73.9
Are you aware of potential side effects as result of using medications?	
Aware	54
Unaware	46
What is a possible potential side effect/warning/precaution due to your medication?*1 (aware only if explained correctly for all drugs with possible side effects/warning/precaution) n=155	
Aware	10.6
Unaware	89.4

What is your key source of medication information?		
Medical personnel		40
Internet		53
Others (bulletin, brochures, banners etc)		7
Are you aware of drug interaction?		
	Aware	76
	Unaware	24
Can you tell the possible dangerous interaction with regards to your medications? ^{*2} (able if explained possible interaction by indicating the drugs on the Rx) n=60		
	Able	7
	Unable	93

^{*1} the total number of prescription with potential side effects/warning/contraindication/precaution was 155 hence denominator for the proportions of response to item.

^{*2} the total number of prescription with possible potential interaction was 60 hence the denominator for proportions of response to the item.

Discussion

This study found an alarming level of prevalence of poor medication self-management among patients on ambulatory care. Many patients had difficulties on reading and interpreting what has been written on the prescription matching with what they are carrying on the hand. The medication self-management model has six tasks i.e. fill, understands, organize, take, monitor and sustain [7]. The scope of this study was to measure the ability of patients to understand the prescription, organizing the plan to take medication as prescribed, ability to open a container and take right number or amount of the medications, patient's ability to monitor for side effects or warnings or interactions and the ability to sustain taking medication by understanding timing and duration of the regimen. Understanding the regimen on the prescription, taking the correct number or amount of dose, and timing of doses and duration of the regimen are areas which troubled

many participants in this study. It is possible that, poor understanding of what has been written on the prescription or narrated by the pharmaceutical personnel is the root cause of poor performance on many of the tasks as testified in the work of Wolf and colleagues [10]. Majority of study participants reported to be secondary level graduates. Therefore, though the study didn't focus on associated factors; it is most probable that education level of the study participants had a major influence of MMC as it was reported in study by Kripalani and colleagues [6]. This study has shown that more than half of the patients were not eager to ask their healthcare provider issues concerning their medications. Though it was not in the scope of this study, the reason may be uncomfortable platform of hospital pharmacy facilities, long waiting queue, unfriendly and brief contact time. All these have impact on ability of the patient to comprehend with written and verbal communication which play big role in understanding the regimen [11].

In this study many patients were not aware of the side effects or warning label or contraindication or precaution or drug interactions for their medications on hand. This is in line with findings from review by Bailey and colleagues [7]. This increase the risk of medication errors and adverse drug reactions which may jeopardize adherence to prescribed regimen. Most probable low level of education among study participants and brief contact time with healthcare providers may be the possible reasons [12]

Majority of the participants had poor ability to pick the correct number or amount. The key reason may still be the same a poor understanding of the prescription regimen means poor understanding of the downstream tasks.

Conclusion

The level of MMC for majority of outpatients attending national tertiary level hospital is very low. Further study need to focus on level of MMC and associated factors.

Limitations

Consecutive sampling posed the risk of selection bias. Confounders like English proficiency as a communication language barrier, differences in cognitive and health literacy levels of

the study participants were not measured. Also, sampled participants were from tertiary level hospital, with diversity in diseases and severity plus treatment durations.

Abbreviations

MMC: Medication management capacity; DRUGS: Drug Regimen Unassisted Grading Scale; MMAA: Medication Management Ability Assessment; MNH; Muhimbili National Hospital; Rx: Prescription.

Declarations

Authors' contributions

HM designed the study, data analysis and drafted the initial manuscript. HS coordinated and monitored data collection. WM, MK, AM, RM and GB participated in study design, interpretation of data and manuscript development. All authors read and approved the final manuscript.

Acknowledgements

The authors highly appreciate patients on ambulatory care exiting Muhimbili National Hospital outpatient pharmacy outlets for participating in this study.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

The dataset generated and/or analyzed during this study are available from the corresponding author upon reasonable request.

Consent for publication

Not applicable

Ethics approval and consent to participate

The study was reviewed and approved by Muhimbili University of health and allied sciences research ethics committee and permission to conduct research was obtained from MNH training and research directorate. Before interview participants willing to take part were consented. To ensure confidentiality, only numbers were used to identify participants and the questionnaires were kept under custodian of researchers only.

Funding

Not applicable

References

1. Krousel-Wood M, Muntner P, Islam T, Morisky D and Webber L. Barriers to and Determinants of Medication Adherence in Hypertension Management: Perspective of the Cohort Study of Medication Adherence among Older Adults (CoSMO) *Med Clin North Am.* 2009; 93 Suppl 3: 753-769
2. Kripalani S, Yao X and Haynes B. Interventions to Enhance Medication Adherence in Chronic Medical Conditions: A Systematic Review. *Arch Intern Med.* 2007; 167: 540-550
3. Viswanathan M, Golin C, Jones C, Ashok M, Blalock S, Wines R, Coker-Schwimmer E, Rosen D, Sista P and Lohr K. Interventions to Improve Adherence to Self-administered Medications for Chronic Diseases in the United States: A Systematic Review. *Ann Intern Med.* 2012; 157:785-795.

4. Janson S, McGrath W, Covington J, Cheng S and Boushey H. Individualized asthma self-management improves medication adherence and markers of asthma control. *J ALLERGY CLIN IMMUNOL*. 2009; 123 Suppl 4:840-846

5. Patton D, Woods S, Franklin Jr D, Cattie J, Heaton R, Collier A, Marra C, Clifford D, Gelman B, McArthur J, Morgello S, Simpson D, McCutchan A, Grant I, and CHARTER Group. Relationship of Medication Management Test-Revised (MMT-R) Performance to Neuropsychological Functioning and Antiretroviral Adherence in Adults with HIV/AIDS. *J Behav*. 2012; 16 Suppl 8: 2286-2296.

6. Kripalani S, Henderson L, Chiu E, Robertson R, Kolm P and Jacobson T. Predictors of Medication Self-management Skill in a Low-literacy Population. *Journal of GEN INTERN MED*. 2006; 21:852-856

7. Bailey S, Oramasionwu C and Wolf M. (2013) Rethinking Adherence: A Health Literacy-Informed Model of Medication Self-Management. *Journal of Health Communication: International Perspectives*. 2013; 18 Suppl 1: 20-30

8. Gadoury M, Schwartzman K, Rouleau M, Maltais F, Julien M, Beaupre´ A, Renzie P, R. Be´gin R, Nault D and Bourbeau J for the Chronic Obstructive Pulmonary Disease axis of the Respiratory Health Network. Self-management reduces both short- and long-term hospitalisation in COPD. *Eur Respir Journal*. 2005; 26: 853-857

9. Abdullahi A, Hassan A, Kadarman N, Saleh A, Baraya S and Lua P. Food safety knowledge, attitude, and practice toward compliance with abattoir laws among the

abattoir workers in Malaysia. *International Journal of General Medicine*. 2016; 9: 79-87

10. Wolf M, Davis T, Tilson H, Bass Iii P and Parker R. Misunderstanding of prescription drug warning labels among patients with low literacy. *American Society of Health-System Pharmacists*. 2006; 63: 1048-1055.

11. Koh K, Brach C, Harris M, and Parchman L. A Proposed 'Health Literate Care Model' Would Constitute A Systems Approach To Improving Patients' Engagement In Care. *HEALTH AFFAIRS*. 2013;32 Suppl 2: 357-367

12. Atak N, Gurkan T and Kose K. The effect of education on knowledge, self management behaviours and self efficacy of patients with type 2 diabetes. *AUSTRALIAN JOURNAL OF ADVANCED NURSING*. 2008; 26 Suppl 2:66-74

Figures




Figure 1

Bar chart of prevalence of medication management capacity