Citrobacter Keratitis: Predisposing Factors and Clinical Characteristics

Mohammad Soleimani

Farabi Eye Hospital

Ahmad Masoumi (✉ ahmad.masoomi1990@gmail.com)

Farabi Eye Hospital <https://orcid.org/0000-0002-3486-0821>

Seyed Ali Tabatabaei

Farabi Eye Hospital

Research Article

Keywords: Citrobacter spp., infectious keratitis, risk factor, clinical findings

Posted Date: August 11th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-396077/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Purpose: To report predisposing factors, clinical presentation, antibiotic sensitivity, and management of *Citrobacter*-caused infectious keratitis.

Methods: We retrospectively reviewed the medical records of culture-proven cases of *Citrobacter* keratitis in a tertiary referral center for 8 years (from January 2012 to September 2020).

Results: Eighteen cases of microbial keratitis due to *Citrobacter spp.* were identified. The median age of the patients was 66 years (range, 10-89). Thirteen patients were male and 5 were female. Multiple predisposing factors were identified in all eyes, including ocular surface disease (n=8), previous corneal surgery (n=6), and history of ocular trauma (n=6). Five patients were diabetic. Corrected distance visual acuity (CDVA) of patients was light perception (LP) in 8 patients, hand motion (HM) in 7, counting fingers (CF) at 1 meter in 1, and CF at 2 meters in 2 patients. Thirteen eyes exhibited hypopyon in association with corneal infiltration. An area of corneal thinning was observed in 7 eyes (38.9%). Endophthalmitis due to infectious keratitis developed in one patient. In vitro susceptibility testing confirmed reasonable sensitivity to ceftazidime and aminoglycosides. Medical management consisted primarily of topical amikacin (20 mg/ml) combined with topical cefazoline (50 mg/ml) (72.2%). Surgical tectonic procedures were carried out in 7 eyes (38.9%).

Conclusion: *Citrobacter spp.* is a rare cause of bacterial keratitis. It comprises 0.7% of all admitted keratitis cases. Previous keratoplasty and ocular surface problems are important risk factors. The prognosis is not good and surgical tectonic intervention is required in many cases to resolve the corneal infection.

Introduction

Microbial keratitis is an ocular emergency that can lead to a loss of sight and even the eye if it is not treated appropriately. Common clinical signs include conjunctival injection, epithelial defects, corneal ulceration with or without stromal infiltration, and hypopyon. Contact lens wear, extended exposure to topical steroids, dry eye disease, and anterior segment procedures such as corneal transplantation are some of the predisposing factors for the development of infectious keratitis.

Citrobacter spp. are gram-negative non-spore-forming bacilli of the family Enterobacteriaceae. They are uncommon causes for urinary and respiratory tract infections ¹. *Citrobacter spp.* are rare etiologic agents in ocular infections ², and most of the studies focused on *Citrobacter* keratitis and endophthalmitis are case reports ^{3,4}.

This study aims to investigate the epidemiology and antibiotic susceptibility pattern of *Citrobacter* keratitis that occurred over 8 years in a tertiary eye care center in Iran.

Materials And Methods

This retrospective observational case series was approved by the institutional review board of Tehran University of Medical Sciences and adhered to the tenets of the Declaration of Helsinki. The clinical and microbiological records of all culture-proven cases of *Citrobacter* keratitis, who presented to Farabi Eye Hospital from January 2012 to September 2020 were retrospectively reviewed. After slit-lamp examination, all corneal ulcers were scraped using a sterile scalpel blade for Gram stains. Fresh scalpel blades were then used to inoculate in chocolate agar and Sabouraud's dextrose agar. The culture plates were incubated at 35°C in carbon dioxide. *Citrobacter spp.* was considered as a causative agent for keratitis if there were discrete colonies of *Citrobacter* on two solid media or confluent growth of micro-organism was observed along with the site of inoculation. Antibiotic susceptibility testing was performed using the disk diffusion method. The following data were collected from patients' records: age, sex, local and systemic predisposing factors, presenting signs and symptoms including the size of corneal infiltration, presence of corneal thinning and hypopyon, antibiotic sensitivity, and mode of treatment.

The patients were admitted if any of the following criteria were present: 1. severe corneal infection according to overall clinical impression 2. presence of corneal thinning or perforation 3. The inability of the patient to instill drops intensively. The initial antibiotic instillation protocol was 1 drop/hour for 24 hours. The antibiotic regimen was then modified according to clinical response and antibiotic susceptibility. The indications for adjunctive procedures such as cyanoacrylate glue application or therapeutic penetrating keratoplasty (PKP) were determined by an experienced cornea specialist.

Results

Within the study period, eighteen microbiologically confirmed *Citrobacter* keratitis were identified. The age of the patients ranged from 10 to 89 years (median, 66 years). There were 13(72.2%) male patients and 5(28.8%) female patients. Eleven ulcers developed in the right eye and 7 in the left eye. All patients were admitted and all of them had a combination of one or more risk factors. During this time, 2522 patients with keratitis were admitted in our center and keratitis caused by *Citrobacter spp.* comprised 0.7% of admitted patients with infectious keratitis. Risk factors associated with *Citrobacter* keratitis are reported in Table 1. Preexisting ocular surface disease in the affected eye was the most common predisposing factor (8 of 18 patients, 44.4%); including bullous keratopathy in 4 patients (22.2%), history of herpes simplex virus keratitis in 2 patients (11.1%), neurotropic keratopathy in 1 patient (5.6%) and anesthetic abuse in 1 patient (5.6%). Six cases were diagnosed after previous keratoplasty (33.3%) ; 3 cases (16.7%) had received one or more PKPs and 3 (16.7%) had undergone Descemet stripping automated endothelial keratoplasty (DSAEK). Five patients (27.8%) had diabetes mellitus. Six cases (31.6%) presented with a history of ocular trauma. The ocular trauma consisted of trauma with the vegetative matter in 2 cases, bilateral thermal burn in one patient, bilateral alkali burn in one patient, corneal penetrating injury in one case, and corneal foreign body in one case.

Table 1
Predisposing factors for development of
Citrobacter keratitis (n = 18)

Ocular surface disease	8 (44.4%)
Pseudophakic bullous keratopathy	4
History of HSV keratitis	2
Neurotrophic keratopathy	1
Anesthetic abuse	1
Previous corneal surgery	6 (33.3%)
PKP	3
DSAEK	3
History of ocular trauma	6 (33.3%)
Trauma with vegetative matter	2
Thermal burn	1
Alkali burn	1
Corneal penetrating injury	1
Corneal foreign body	1
Diabetes mellitus	5 (27.8%)

Eight patients (44.4%) presented with corrected distance visual acuity (CDVA) of light perception (LP), 7 patients (38.9%) with hand motion (HM) vision, 1 patient (5.5%) with counting fingers (CF) at 1 m, and two patients (11.1%) with CF at 2 m. All patients complained of painful red eyes and decreased vision at presentation. The area of infiltration was small (< 2mm at its greatest dimension) in 2 eyes, medium (2–6 mm) in 14 eyes, and large (> 6mm) in 2 eyes. Hypopyon was present in 13 (72.2%) eyes at initial presentation. Descemet's membrane folds and endothelial plaques were detected in 7 (38.9%) and 2 (11.1%) eyes, respectively on initial slit-lamp examination. Corneal thinning in the area of infiltration was observed in 7 (38.9%) eyes and the corneal ulcer progressed to perforation in 4 (22.2%) patients. Vitreous involvement and endophthalmitis developed in 1 patient (5.6%) after admission. This patient presented with a perforated corneal ulcer in her left eye. Her vision at the presentation was LP. She was diabetic and had a history of cataract surgery and intraocular lens (IOL) exchange in the left eye.

Medical management consisted primarily of topical amikacin (20 mg/ml) combined with topical cefazoline (50 mg/ml) (72.2%) until the specific antibiogram testing was made according to which topical drops would be narrowed. Oral doxycycline and vitamin C were prescribed in 77.8% and 66.7% of patients, respectively. In total, 7 patients (38.9%) in the series ultimately required one or more surgical procedures. In the patient who progressed to endophthalmitis, pars plana vitrectomy through a temporary

keratoprosthesis associated with the subsequent graft was performed. Three patients underwent therapeutic penetrating keratoplasty. The presence of corneal melt and thinning in two patients necessitated the application of cyanoacrylate glue. Persistent epithelial defect in one patient was treated using amniotic membrane transplantation (AMT).

In vitro testing of *Citrobacter sp.* isolates in this series showed that all were sensitive to gentamicin (18/18), whereas 93.4% isolates were sensitive to amikacin (15/16), 87.5% to ciprofloxacin (14/16), 85.7% to levofloxacin (12/14), and 88.9% to ceftazidime (16/18).

Discussion

Citrobacter sp. belongs to the family Enterobacteriaceae and consists of 13 currently recognized species. *Citrobacter freundii* and *Citrobacter koseri* are mostly associated with human infections⁵. These organisms can be isolated from the intestinal tract of humans. *Citrobacter sp.* is often considered an opportunistic pathogen and it causes a broad spectrum of infections related to the urinary tract, respiratory tract, and bloodstream⁵. *Citrobacter sp.* are rare causes for ocular infections^{3,6}, and is rarely isolated in patients with bacterial keratitis^{2,7,8}.

Contact lens wear is the most common risk factor for bacterial keratitis in developed countries⁹⁻¹¹, however, none of the patients in this series were contact lens wearers. A high proportion (44.4%) of patients diagnosed with *Citrobacter* keratitis suffered from the ocular surface disease. This is comparable to an epidemiologic study of bacterial keratitis from Vancouver, Canada¹², but is much higher compared to the reports of infectious keratitis from northern California (17.7%)¹⁰, and southern Texas (17.6%)¹¹. The ocular surface disease impairs the defense mechanisms of the external eye and cornea, moreover, the decreased integrity of the corneal epithelium, persistent epithelial defects, and associated corneal inflammation can contribute to the increased risk of infectious keratitis in these patients^{13,14}.

Diabetes mellitus was the most common systemic risk factor in this series. The hyperglycemic state in the diabetic cornea impairs the immunologic defenses of the ocular surface. Moreover, delayed corneal re-epithelialization, and decreased corneal sensation that may progress to neurotrophic corneal ulcers further predisposes the eye to infectious keratitis¹⁵.

It is noteworthy that 6 of 18 patients in our series had a history of corneal transplantation. Epithelial defects, compromised immunity of the ocular surface, suture related problems and graft failure can all contribute to increased risk of graft infection¹⁶. The most common causative pathogens in infectious keratitis after PKP are gram-positive bacteria, with coagulase-negative staphylococci most commonly isolated. *Pseudomonas aeruginosa* is the most common inciting gram-negative bacterium in post-PKP infectious keratitis¹⁷. There is no report of *Citrobacter* keratitis after PKP in the literature to the best of our knowledge.

None of the eyes in this series were enucleated, however, progression to endophthalmitis necessitated pars plana vitrectomy through a temporary keratoprosthesis in one eye. The incidence of infectious keratitis associated with endophthalmitis is reported to be 0.29%¹⁸. Malihi et al. found that poor visual acuity, history of ocular surgeries, corneal perforation, topical corticosteroid use, and systemic immunocompromise increases the risk of endophthalmitis in a patient with infectious keratitis¹⁹. As stated earlier many of these predisposing factors were present in the patient who progressed to endophthalmitis.

Antibiotic resistance is an emerging problem in *Citrobacter sp.*, as in many other bacterial species, and limits the clinician's armamentarium. In vitro susceptibility testing shows that *Citrobacter*-caused keratitis may respond well to aminoglycosides. The rate of resistance to levofloxacin (14.7%) and ciprofloxacin (12.5%) was alarmingly higher compared to other reports about antibiotic sensitivity in gram-negative rods^{7, 8, 20}. Injudicious use of topical fluoroquinolones primarily for prophylaxis in cataract and refractive surgery might be responsible for this finding. Enhanced efflux of antibiotics and mutations in DNA gyrase subunit B is thought to be responsible for *C. freundii* resistance to fluoroquinolones²¹.

The requirement for surgical procedures was 38.9% in this series, which is higher than reports about *Haemophilus influenza*²² and *Moraxella* keratitis^{23, 24}. This implies the more severe presentation of infectious keratitis due to *Citrobacter spp.* and poorer visual outcomes.

As a tertiary eye care center, many of the patients seeking care in our unit are referred by clinicians who work in an outpatient setting. This may explain the high rate of patients with severe keratitis and who had a history of previous ocular surgeries. The clinicians may have a lower threshold for referral of patients who have undergone corneal surgeries.

The limitations of this study include its retrospective nature and small sample size. The absence of data about follow-up examinations further limits the results of this study. Nevertheless, to the best of our knowledge, this is the largest case series about infectious keratitis due to *Citrobacter spp.*, elucidating the clinical features and risk factors for this devastating infection.

In summary, *Citrobacter spp.* is a rare cause of bacterial keratitis. It comprises 0.7% of all admitted keratitis cases. This study shows that ocular and systemic predisposing factors play an important role in the development of *Citrobacter*-caused keratitis. Previous keratoplasty and ocular surface problems are important risk factors. Surgical tectonic intervention is required in many cases to resolve the corneal infection.

Declarations

Sources of funding: none

Conflicts of interest: none

Availability of data: data is available upon reasonable request

Code availability: Not applicable

Authors' contributions: All authors have made a significant contribution to the manuscript and accept the responsibility for the data presented

Ethics approval: The institutional review board of Tehran University of Medical Sciences approved this study.

Consent to participate: not applicable

Consent for publication: not applicable

References

1. Rogers L, Power K, Gaora P et al. *Escherichia coli* and other Enterobacteriaceae: occurrence and detection. 2016
2. Tavassoli S, Nayar G, Darcy K et al (2019) An 11-year analysis of microbial keratitis in the South West of England using brain–heart infusion broth. *Eye* 33(10):1619–1625
3. Goold LA, Warriar SK, Wittles NK et al (2010) Microbial keratitis secondary to infection with *Citrobacter koseri*. *Cornea* 29(4):479
4. Barsam A, Heatley C, Herbert L (2006) Panophthalmitis secondary to infection with *Citrobacter koseri*. *Eye* 20(12):1468–1470
5. Anderson MT, Mitchell LA, Zhao L et al (2018) *Citrobacter freundii* fitness during bloodstream infection. *Sci Rep* 8(1):1–14
6. Chen K-J, Sun M-H, Hwang Y-S et al (2008) Endophthalmitis caused by *Citrobacter* species. *Ocul Immunol Inflamm* 16(4):147–153
7. Peng MY, Cevallos V, McLeod SD et al (2018) Bacterial keratitis: isolated organisms and antibiotic resistance patterns in San Francisco. *Cornea* 37(1):84
8. Hernandez-Camarena JC, Graue-Hernandez EO et al (2015) Trends in microbiological and antibiotic sensitivity patterns in infectious keratitis: 10-year experience in Mexico City. *Cornea* 34(7):778–785
9. Bourcier T, Thomas F, Borderie V et al (2003) Bacterial keratitis: predisposing factors, clinical and microbiological review of 300 cases. *Br J Ophthalmol* 87(7):834–838
10. Jeng BH, Gritz DC, Kumar AB et al (2010) Epidemiology of ulcerative keratitis in Northern California. *Arch Ophthalmol* 128(8):1022–1028
11. Puig M, Weiss M, Salinas R et al (2020) Etiology and Risk Factors for Infectious Keratitis in South Texas. *J Ophthalmic Vis Res* 15(2):128
12. Termote K, Joe AW, Butler AL, et al (2018) Epidemiology of bacterial corneal ulcers at tertiary centres in Vancouver, BC. *Can J Ophthalmol* 53(4):330–336

13. Cheung AY, Sarnicola E, Eslani M et al (2018) Infectious keratitis after ocular surface stem cell transplantation. *Cornea* 37(11):1395–1399
14. Kang BS, Kim MK, Wee WR et al (2016) Infectious Keratitis in Limbal Stem Cell Deficiency: Stevens–Johnson Syndrome Versus Chemical Burn. *Cornea* 35(1):51–55
15. Priyadarsini S, Whelchel A, Nicholas S et al. Diabetic keratopathy: Insights and challenges. *Surv Ophthalmol*. 2020
16. Vajpayee R, Boral S, Dada T et al (2002) Risk factors for graft infection in India: a case-control study. *Br J Ophthalmol* 86(3):261–265
17. Davila JR, Mian SI (2016) Infectious keratitis after keratoplasty. *Curr Opin Ophthalmol* 27(4):358–366
18. Zapp D, Loos D, Feucht N et al (2018) Microbial keratitis-induced endophthalmitis: incidence, symptoms, therapy, visual prognosis and outcomes. *BMC Ophthalmol* 18(1):1–7
19. Malihi M, Li X, Patel S et al (2017) Infectious Keratitis–associated Endophthalmitis: A 14-year Study. *Retina* 37(4):662–666
20. Hsiao C-H, Sun C-C, Yeh L-K, et al (2016) Shifting trends in bacterial keratitis in Taiwan: a 10-year review in a tertiary-care hospital. *Cornea* 35(3):313–317
21. Tavío MdM, Vila J, Ruiz J et al (2000) In vitro selected fluoroquinolone-resistant mutants of *Citrobacter freundii*: analysis of the quinolone resistance acquisition. *J Antimicrob Chemother* 45(4):521–524
22. Yang K-S, Lin H-C, Ma DH et al (2006) Ulcerative keratitis caused by *Haemophilus influenzae*. *Cornea* 25(6):701–704
23. Das S, Constantinou M, Daniell M et al (2006) *Moraxella* keratitis: predisposing factors and clinical review of 95 cases. *Br J Ophthalmol* 90(10):1236–1238
24. Inoue H, Suzuki T, Inoue T et al (2015) Clinical characteristics and bacteriological profile of *Moraxella* keratitis. *Cornea* 34(9):1105–1109