Preconceptional and prenatal exposure to diurnal temperature variation increases the risk of childhood pneumonia

Xiangrong Zheng

Xiangya Hospital Central South University

Jian Kuang

Xiangya Hospital Central South University

Chan Lu (✉ chanlu@csu.edu.cn)

Central South University China

Qihong Deng

Central South University Xiangya School of Public Health

Haiyu Wu

Central South University Xiangya School of Medicine

Rachael Gakii Murithi

Central South University Xiangya School of Public Health

McSherry Brownel Johnson

Central South University Xiangya School of Public Health

Wang Peng

Xiangya Hospital Central South University

Maolan Wu

Xiangya Hospital Central South University

Research article

Keywords: Diurnal temperature variation, Preconceptional, Pregnancy, Trimester, Childhood pneumonia

Posted Date: November 13th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-39734/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Pediatrics on April 21st, 2021. See the published version at <https://doi.org/10.1186/s12887-021-02643-x>.

Abstract

Background: Pneumonia is the leading cause of death and hospitalization among young children worldwide, but its risk factors remain unclear.

Objective: To evaluate the effect of maternal exposure to diurnal temperature variation (DTV) during preconceptional and prenatal periods on childhood pneumonia.

Methods: A retrospective cohort study by case-control design was conducted for pneumonia (N=699) and normal (N=811) children under age of 14 who were enrolled in XiangYa Hospital, Changsha, China from May 2017 to April 2019. Demographic data including gender, age, birth season, gestational age, parity, mode of delivery, and parental atopy were collected from the electronic medical records in the hospital system. We obtained the data of daily DTV in Changsha during 2003-2019 from China Meteorological Administration. Maternal exposure to DTV during preconceptional and prenatal periods was respectively calculated by the average of daily DTV during one year and three months before conception and entire pregnancy as well as the three trimesters. The association between maternal exposure to outdoor DTV and childhood pneumonia was analyzed by multiple logistic regression model.

Results: We found that childhood pneumonia was significantly associated with exposure to an increase in DTV during one year before conception and entire pregnancy, with ORs (95% CI) = 2.53 (1.56-4.10) and 1.85 (1.24-2.76). We further identified a significant risk of pneumonia due to DTV exposure during the first and second trimester of pregnancy. Sensitivity analysis showed that boys were more susceptible to the effect of prenatal exposure to outdoor DTV during pregnancy particularly in the first two trimesters compared to girls.

Conclusions: Preconceptional and prenatal exposure to DTV plays an important role in development of childhood pneumonia, especially during the first and second trimesters of pregnancy.

Background

Pneumonia is the leading cause of death and hospitalization among children under age of 5 worldwide [1, 2]. Although pneumonia treatment has been improved significantly during past decades, the incidence of pneumonia in childhood remains high, especially in developing countries [3]. According to a recent epidemiological report, pneumonia has resulted in 808,694 deaths of children in 2017, which accounts for 15% of all deaths in children under 5 years [4]. Notably, China has witnessed a very high prevalence of childhood pneumonia over recent years [5]. Pneumonia not only influences the growth and development of children's lung, but also causes a heavy burden on family economy and public health [6]. Therefore, it is crucial to investigate the risk factors contributing to the frequent occurrence of childhood pneumonia, which would have a great implication for its effective reduction and early prevention.

Recently, there is increasing awareness that climatic variation significantly impacts childhood respiratory health. Pneumonia, as one of the most adverse infectious disease among children, has been suggested

to be closely related to a continuous climate change [7, 8]. Mounting evidences indicate that children suffer most from climate change due to their susceptibility for the extreme temperature and the dramatic weather change [9, 10]. Some recent epidemiological studies from different regions have found an association between outdoor temperature and childhood pneumonia [11, 12]. Diurnal temperature variation (DTV), reflecting a degree in the daily temperature changes, has been found to be correlated with both mortality and morbidity of respiratory diseases [13-18]. Particularly, one of our recent cohort studies suggests that outdoor DTV is an important risk factor for pneumonia development in young children [19].

Early life exposure to harmful environmental factors could not only have an adverse effect on organ development and childhood growth, but also plays a critical role in the onset and progression of diseases in adulthood and has a long-term impact in the later life [20, 21]. Some studies have suggested that early-life exposure to temperature and DTV contributed to an increased risk of childhood respiratory diseases [19, 22-24]. Although a few studies have linked maternal exposure to traffic-related air pollution (TRAP) during preconceptional period with increased risk of allergic diseases in preschool children [25, 26]. Thus, maternal exposure to environmental factors before pregnancy could play an important role in the development of health outcomes during postnatal period. However, the effect of preconceptional exposure to outdoor DTV on respiratory outcomes, such as childhood pneumonia, have scarcely been investigated.

Due to the lack of studies on early life exposure to climatic change particularly before birth and children's health, we hypothesized that exposure to outdoor DTV during preconceptional and prenatal periods may be associated with later development of childhood pneumonia. To test this hypothesis and identify the critical timing window(s), we performed a large retrospective cohort study via case-control design in Changsha, China.

Methods

1. Study population

A retrospective cohort study by case-control design was conducted in the Department of Pediatrics, XiangYa hospital, Central South University in Changsha, China from May 2017 to April 2019. We recruited 699 children under 14 years old who were diagnosed with pneumonia and set them as the case group, and 811 healthy children who registered for physical examination as the control group. Children were excluded if they had abnormal congenital development in the airway and lung parenchyma, chronic lung disease, obstruction in the airway or compression outside the tube, congenital heart disease, tuberculosis, heart, liver and kidney diseases, or a known diagnosis of immunodeficiency.

The case group consisted of children with a diagnosis of pneumonia made in accordance with the following WHO diagnosis criteria [27]. Cough or difficult breathing plus at least one of the following signs: (1) fast breathing (rate >60 breaths per minute if aged <2 months, >50 breaths per minute if aged 2-11 months, and >40 breaths per minute if aged 12-59 months); or (2) lower chest wall indrawing. In addition, either crackles or pleural rub may be present on chest auscultation. The control group included healthy

children who didn't have a history of pneumonia diagnosis based on WHO criteria at the time of recruitment.

2. Demographic data

We collected the demographic data from the electronic medical records system in the hospital, including children's gender, age, birth season, gestational weeks, parity, mode of delivery, and parental atopy. Parental atopy was defined by the presence of a history of any allergic diseases among child's mother and/or father.

3. Exposure assessment

3.1. Exposure timing windows

Exposure timing-windows included preconceptional and prenatal periods in this study. The preconceptional exposure involved in two timing windows: one year before conception and three months before conception. The prenatal period was defined from the first day of the mother's last menstrual period to the delivery day. The prenatal exposure was further divided into three trimesters: the first trimester (from the 1st to 12th weeks of gestation), second trimester (the 13th to 27th weeks of gestation), and third trimesters (from the 28th gestational week to the birthday of the child).

3.2. Personal exposure to diurnal temperature variation (DTV)

We obtained the data of daily temperature (including the mean, maximum, and minimum temperatures) at ten different monitoring stations in Changsha during 2003-2019 from China Meteorological Administration: Kaifu district, Yuhua district, Furong district, Liuyang county, Mapoling, Ningxiang county, Tianxin district, Wangcheng district, Yuelu district, and Changsha county. The diurnal temperature variation (DTV) was calculated as the difference between the daily maximum and minimum temperature. Children's exposure was estimated by DTV at the station where their residence was located. Then, the individual exposure to outdoor DTV for each child was calculated as follows: (1) Preconceptional exposure to DTV was calculated as the average of daily DTV during one year and three months before conception; (2) Prenatal exposure to DTV was calculated by the average of daily DTV during the period from the mother's last menstrual period to the delivery day; (3) The exposure to DTV in the first, second, and third trimester was respectively calculated as the average of daily DTV during the periods of 1st–12th gestational weeks, 13th–27th gestational weeks, and from the 28th gestational week till the date of birth.

3.3 Exposure to outdoor air pollution

Exposure to outdoor air pollution was considered as an important confounding variable for pneumonia risk in this study. We obtained daily 24h-averaged concentrations of three main air pollutants, including nitrogen dioxide (NO₂), sulfur dioxide (SO₂), and particulate matter $\leq 10 \mu\text{m}$ in diameter (PM₁₀), from 7 municipal air quality monitoring stations in Changsha from 2003 to 2019. Individual exposure to ambient

air pollution during different timing windows was estimated by the inverse distance weight (IDW) method described in our previous work [28].

4. Statistical analysis

Statistical analyses were performed by SPSS software (version 23.0, SPSS Inc., Chicago, USA). The relationship between exposure to outdoor DTV during preconceptional and prenatal periods and childhood pneumonia was assessed by using multiple logistic regression model, with adjusting potential variables in Table 1 and the three air pollutants (NO₂, SO₂, and PM₁₀). The associations in regression analysis was calculated by odds ratio (OR) of 95% confidence interval (95% CI). In our study, OR (95% CI) was estimated by per 1 °C increase in exposure to outdoor DTV. P-value ≤ 0.05 was considered as statistically significant.

Results

Our study included 699 children with pneumonia and 811 healthy children without pneumonia or other respiratory diseases. Table 1 presents the demographics and prevalence of pneumonia stratified by the covariates. A single factor analysis of variance was used to examine the association between potential covariates and childhood pneumonia. We found that the prevalence of childhood pneumonia was significantly higher in children with male sex, under 1 year old, preterm birth, caesarean birth, and parental atopy than those with female sex, equal to and older than 1 year old, non-preterm birth, non-caesarean birth, and without parental atopy, with the p-value reached the statistically significant level (p < 0.05). Our analysis showed that all these covariates were closely related to the risk of childhood pneumonia.

DTV exposure levels during preconceptional and prenatal periods between the case and control group were compared in Table 2. The average of individual exposure to DTV during one year before conception was respectively 7.3°C and 7.2°C in pneumonia and healthy children, with a relatively small difference. Comparatively, the averaged individual exposure to DTV during entire pregnancy was relatively greater in children with pneumonia (7.3°C) than that in healthy children (7.0°C). Therefore, the average individual exposure level of DTV during both preconceptional and prenatal periods was significantly higher among children with pneumonia compared to healthy children (p < 0.001).

The relationship between children's exposure to DTV during different timing window and pneumonia risk was further analyzed in Table 3. We observed a significant association between pneumonia and preconceptional exposure to increased DTV, particularly for a long-term exposure, with adjusted OR (95% CI) = 2.53 (1.56-4.10) for per 1°C increase in DTV exposure during one year before conception. Childhood pneumonia was also significantly associated with prenatal exposure to outdoor DTV during entire pregnancy especially in the first and second trimester, respectively with adjusted ORs (95% CI) = 1.85 (1.24-2.76), 1.63 (1.32-2.00) and 1.43 (1.12-1.81).

The association between maternal exposure to outdoor DTV during prenatal period and childhood pneumonia stratified by children's gender was showed in Figure 1. We observed that boys were more susceptible to the effect of exposure to outdoor DTV during pregnancy and the first two trimesters on pneumonia risk compared to girls.

Discussion

Our retrospective cohort study by case-control design, to the best acknowledge, is the first to find an association between maternal exposure to outdoor DTV during one year before conception and childhood pneumonia. The association was also observed in prenatal exposure, particularly in the first and second trimester of pregnancy. Our sensitivity analysis indicated that boys were more vulnerable to the pneumonia risk of prenatal exposure to DTV than girls.

Previous studies have suggested that DTV exposure may play a significant role in the increased risk of respiratory diseases [29-31], including asthma, allergy and respiratory tract infection (RTI) among young children [32-34]. However, the evidence in the association between DTV and childhood pneumonia is still controversial [19, 35]. Our study newly indicates that early-life exposure to a large DTV could be a risk factor contributing to the development of childhood pneumonia.

The present study, to our acknowledge, is the first to found that an increase in DTV exposure during the year prior to conception showed an adverse effect on the later development of childhood pneumonia. Previous studies mainly associated preconceptional exposure to air pollution with childhood health problems. A cohort study in China suggested that preconceptional exposure to traffic- and industry-related air pollutants was significantly related with childhood asthma [26]. NO₂ exposure during three months before conception was reported to be linked with eczema risk in children [25]. Another recent retrospective study in US proved a significant relationship between preconceptional exposure to NO_x and SO₂ and an increased risk of gestational diabetes mellitus (GDM) [36]. Furthermore, a multi-site study found that preconceptional exposure to PM₁₀, SO₂ and CO was linked with increased risk of oral cleft [37]. Therefore, it is reasonable to believe that preconceptional exposure to DTV significantly increased the risk of childhood pneumonia, which may indicate a hypothesis of "pre-fetal origin of childhood infection".

Professor David Barker, a British epidemiologist, first proposed the theory of the development origins of health and diseases (DOHaD): early life (prenatal and early postnatal periods) exposure to environmental factors affects the plasticity of development, leading to the development and/or deterioration of various complex diseases in adulthood. Therefore, early life was considered as a critical timing window of environmental exposure which is related to human health. In this study, we observed that prenatal exposure to a large DTV was a significant risk factor for pneumonia development in children, which is in line with a recent cohort study from China [19]. We further identified that the first and second trimester of pregnancy was the critical time windows for DTV exposure contributing to the development of childhood pneumonia, which was scarcely observed in previous literatures. Exposure to a large DTV in the first trimester of pregnancy was found to be associated with frequent common colds among preschool

children in China [22]. One of our recent studies have indicated a significant association between exposure to industrial air pollutant SO₂ during the first trimester of pregnancy and increased risk of childhood ear infection [38]. The mechanisms underlying the association between childhood pneumonia and prenatal exposure to DTV during the first and second trimesters of pregnancy remain unclear. It was suggested that lung development starts at about the 4th gestational week, and the airway branching morphogenesis occurs in the first two trimesters of gestation [39]. Hence, long-term exposure to environmental factors in utero may lead to an improper development of lung at various degree, which could affect postnatal development of lung function and induce a risk of respiratory diseases later in life. However, the pathogenic mechanisms underlying childhood pneumonia risk of intrauterine exposure to DTV warrants further investigations.

A recent study [13] and one of our recent work [19] indicated that boys were more susceptible to the effect of prenatal exposure to DTV in pregnancy on pneumonia risk compared to girls. According to the reports from Global Burden of Disease Study 2017 (GBD 2017), the prevalence of respiratory tract infections in males was significantly higher than females [40]. Some evidence suggested that the gender difference of environmental susceptibility was mainly related to immune response [41]. In general, females are more responsive to pathogenic stimuli and vaccines than males, and thus females are more resistant to infectious diseases [42]. Whereas, the mechanisms in the sex-related susceptibility in the effect of DTV exposure on childhood infections including pneumonia remain unclear, which is profound to further study.

Our study had several limitations for the data collection and analysis. Firstly, the data of personal exposure to outdoor air temperature was only obtained from one meteorological station, this may result in exposure misclassification. In spite of the fact that environmental temperature changes in different areas of one city (Changsha) are small, this may lead to exposure classification errors, and using temperature exposure with high spatial resolution to capture the temperature difference in the areas of inner-city may perform a better exposure assessment [43]. Secondly, indoor air temperature was not considered in the present study. However, as people especially pregnant women spend most of their time indoors, the long-term exposure to indoor air temperature may impact the health condition among pregnant mothers and young childhood. Thirdly, other meteorological parameters including relative humidity, wind speed, and precipitation were not included in our study. These climatic factors may affect air temperature or temperature change, and thus their potential influence should not be ignored. At last, we didn't consider the possibility of the change of residence of the mother during the preconception or prenatal period, however, most of them lived in Changsha during pregnancy and one year before conception.

Conclusions

This retrospective cohort study via case-control design reveals that preconceptional (one year before conception) and prenatal exposure to DTV was significantly associated with childhood pneumonia. We identified the first two trimesters of pregnancy as the critical time windows. Our results have a clinical

significance in assessing the onset and development of pneumonia. Our study also provide an novel strategy for effective reduction and early prevention of childhood pneumonia, i.e., reducing outdoor activities among women preparing for pregnancy when the temperature changes sharply, and avoiding extra exposure to a large variation in daily temperature during pregnancy especially in the first and second trimesters.

Abbreviations

DTV: diurnal temperature variation; IDW: inverse distance weight

Declarations

Acknowledgements

Not applicable.

Authors' contributions

XR conducted the study, analyzed the data, and drafted the initial manuscript. JK collected and analyzed the data, and drafted part of the manuscript. CL conceptualized, designed and performed the study, supervised the data analysis, and drafted and revised the manuscript. QD guided and revised the manuscripts, as well as critically reviewed the manuscript for important intellectual content. HW, RGM, MBJ, WP, MW conducted the study, analyzed the data, and reviewed the revised manuscript. All authors have read and approved the manuscript.

Funding

This study was supported by the National Natural Science Foundation of China (No. 81873851 and 42007391), the Natural science foundation of Hunan Province (NO.2018JJ6141), and Health and Family Planning Commission of Hunan Province (No. B20180881). The funders had no role in study design, data collection and analysis, decision for publish, or preparation of the manuscript.

Availability of data and materials

The data sets used and/or analyzed during the current present study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Ethics approval and consent to participate

This study was approved by the academic ethics committee of XiangYa Hospital (201904107).

Informed consent was obtained from a parent or guardian for all individual participants included in the study.

Consent for publication

Not applicable.

References

1. Walker CF, Rudan I, Liu L, Nair H, Theodoratou E, Bhutta ZA, et al. Global burden of childhood pneumonia and diarrhoea. *Lancet*. 2013; 381(9875):1405-1416.
2. Liu L, Oza S, Hogan D, Perin J, Rudan I, Lawn JE, et al. Global, regional, and national causes of child mortality in 2000–13, with projections to inform post-2015 priorities: an updated systematic analysis. *Lancet*. 2015; 385(9966):430-440.
3. Rudan I, Boschi-Pinto C, Biloglav Z, Mulholland K, Campbell H. Epidemiology and etiology of childhood pneumonia. *Bulletin of the world health organization*. 2008; 86:408-416.
4. World Health Organization. pneumonia. 2019. <https://www.who.int/news-room/fact-sheets/detail/pneumonia>. Accessed 2 August 2019.
5. Jiang W, Lu C, Miao Y, Xiang Y, Chen L, Deng Q. Outdoor particulate air pollution and indoor renovation associated with childhood pneumonia in China. *Atmospheric Environment*. 2018; 174:76-81.
6. McMichael AJ. Globalization, climate change, and human health. *N Engl J Med*. 2013; 368(14):1335-1343.
7. Kim J, Kim JH, Cheong HK, Kim H, Honda Y, Ha M, et al. Effect of Climate Factors on the Childhood Pneumonia in Papua New Guinea: A Time-Series Analysis. *Int J Environ Res Public Health*. 2016; 13(2):213.
8. Paynter S, Ware R, Weinstein P, Williams G, Sly PD. Childhood pneumonia: a neglected, climate-sensitive disease? *Lancet*. 2010; 376(9755):1804-1805.
9. McKie R: Children will suffer most as climate change increases in coming decades, say scientists. *BMJ*. 2013; 347:f5799.
10. Xu Z, Etzel RA, Su H, Huang C, Guo Y, Tong S. Impact of ambient temperature on children's health: a systematic review. *Environ Res*. 2012; 117:120-131.
11. Lam HCY, Chan EYY, Goggins WB: Short-term Association Between Meteorological Factors and Childhood Pneumonia Hospitalization in Hong Kong: A Time-series Study. *Epidemiology*. 2019; 30 Suppl 1:S107-S114.
12. Xu Z, Liu Y, Ma Z, Li S, Hu W, Tong S. Impact of temperature on childhood pneumonia estimated from satellite remote sensing. *Environ Res*. 2014; 132:334-341.
13. Ding Z, Guo P, Xie F, Chu H, Li K, Pu J, et al. Impact of diurnal temperature range on mortality in a high plateau area in southwest China: A time series analysis. *Sci Total Environ*. 2015; 526:358-365.

14. Zhou X, Zhao A, Meng X, Chen R, Kuang X, Duan X, et al. Acute effects of diurnal temperature range on mortality in 8 Chinese cities. *Sci Total Environ*. 2014; 493:92-97.
15. Yang J, Liu HZ, Ou CQ, Lin GZ, Zhou Q, Shen GC, et al. Global climate change: impact of diurnal temperature range on mortality in Guangzhou, China. *Environ Pollut*. 2013; 175:131-136.
16. Chen C, ZHOU W, GUI Y, Kan H. Diurnal temperature range as a novel risk factor for sudden infant death. *Biomed Environ Sci*. 2011; 24(5):518-522.
17. Sun S, Laden F, Hart JE, Qiu H, Wang Y, Wong CM, et al. Seasonal temperature variability and emergency hospital admissions for respiratory diseases: a population-based cohort study. *Thorax*. 2018; 73(10):951-958.
18. Cheng J, Xu Z, Zhu R, Wang X, Jin L, Song J, Su H. Impact of diurnal temperature range on human health: a systematic review. *Int J Biometeorol*. 2014; 58(9):2011-2024.
19. Zeng J, Lu C, Deng Q. Prenatal exposure to diurnal temperature variation and early childhood pneumonia. *J Therm Biol*. 2017; 65:105-112.
20. Heindel JJ, Vandenberg LN. Developmental origins of health and disease: a paradigm for understanding disease cause and prevention. *Curr Opin Pediatr*. 2015; 27(2):248-253.
21. Barker DJ, C O. Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales. *Lancet*. 1986; 327:1077-1081.
22. Lu C, Miao Y, Zeng J, Jiang W, Shen YM, Deng Q. Prenatal exposure to ambient temperature variation increases the risk of common cold in children. *Ecotoxicol Environ Saf*. 2018; 154:221-227.
23. Miao Y, Shen YM, Lu C, Zeng J, Deng Q. Maternal exposure to ambient air temperature during pregnancy and early childhood pneumonia. *J Therm Biol*. 2017; 69:288-293.
24. Gluckman PD, Hanson MA, Cooper C, Thornburg KL. Effect of in utero and early-life conditions on adult health and disease. *N Engl J Med*. 2008; 359(1):61-73.
25. Lu C, Deng L, Ou C, Yuan H, Chen X, Deng Q. Preconceptional and perinatal exposure to traffic-related air pollution and eczema in preschool children. *J Dermatol Sci*. 2017; 85(2):85-95.
26. Deng Q, Lu C, Ou C, Chen L, Yuan H. Preconceptional, prenatal and postnatal exposure to outdoor and indoor environmental factors on allergic diseases/symptoms in preschool children. *Chemosphere*. 2016; 152:459-467.
27. World Health Organization. Pocket book of hospital care for children: guidelines for the management of common illnesses with limited resources. 2nd ed. Geneva: World Health Organization; 2013.
28. Deng Q, Lu C, Norback D, Bornehag CG, Zhang Y, Liu W, Yuan H, Sundell J. Early life exposure to ambient air pollution and childhood asthma in China. *Environ Res*. 2015; 143(Pt A):83-92.
29. Wang MZ, Zheng S, He SL, Li B, Teng HJ, Wang SG, et al. The association between diurnal temperature range and emergency room admissions for cardiovascular, respiratory, digestive and genitourinary disease among the elderly: a time series study. *Sci Total Environ*. 2013; 456-457:370-375.

30. Lim YH, Hong YC, Kim H. Effects of diurnal temperature range on cardiovascular and respiratory hospital admissions in Korea. *Sci Total Environ.* 2012; 417:55-60.
31. Liang WM, Liu WP, Kuo HW. Diurnal temperature range and emergency room admissions for chronic obstructive pulmonary disease in Taiwan. *Int J Biometeorol.* 2009; 53(1):17-23.
32. Li T, Yang Z, Wang M. Diurnal temperature range related to respiratory tract infection in childhood. *Pediatr Infect Dis J.* 2014; 33(12):1309-1310.
33. Ge WZ, Xu F, Zhao ZH, Zhao JZ, Kan HD. Association between diurnal temperature range and respiratory tract infections. *Biomed Environ Sci.* 2013; 26(3):222-225.
34. Xu Z, Huang C, Su H, Turner LR, Qiao Z, Tong S. Diurnal temperature range and childhood asthma: a time-series study. *Environ Health.* 2013; 12(1):12.
35. Xu Z, Hu W, Tong S. Temperature variability and childhood pneumonia: an ecological study. *Environ Health.* 2014; 13(1):51.
36. Robledo CA, Mendola P, Yeung E, Männistö T, Sundaram R, Liu D, et al. Preconception and early pregnancy air pollution exposures and risk of gestational diabetes mellitus. *Environ Res.* 2015; 137:316-322.
37. Zhu Y, Zhang C, Liu D, Grantz KL, Wallace M, Mendola P. Maternal ambient air pollution exposure preconception and during early gestation and offspring congenital orofacial defects. *Environ Res.* 2015; 140:714-720.
38. Deng Q, Lu C, Li Y, Chen L, He Y, Sundell J, et al. Association between prenatal exposure to industrial air pollution and onset of early childhood ear infection in China. *Atmospheric Environment.* 2017; 157:18-26.
39. Schittny JC. Development of the lung. *Cell Tissue Res.* 2017; 367(3):427-444.
40. Spencer L James, Degu Abate, Kalkidan Hassen Abate, Solomon M Abay, Cristiana Abbafati, Nooshin Abbasi, Hedayat Abbastabar, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet.* 2018; 392:1789-1858.
41. Muenchhoff M, Goulder PJJTJoid. Sex differences in pediatric infectious diseases. *JID.* 2014; 209(suppl_3):S120-S126.
42. Edwards M, Dai R, Ahmed SA. Our Environment Shapes Us: The Importance of Environment and Sex Differences in Regulation of Autoantibody Production. *Front Immunol.* 2018; 9:478.
43. Zhong Q, Lu C, Zhang W, Zheng X, Deng Q. Preterm birth and ambient temperature: strong association during night-time and warm seasons. *J Therm Biol.* 2018; 78:381-390.

Tables

Table 1 Demographic information of covariates among children with (case) and without (control) doctor-diagnosed pneumonia (n=1,510).

	Total	Case	Control	P-value
	N (%)	N (%)	N (%)	
Total	1,510 (100.0)	699 (46.3)	811 (53.7)	—
Sex				<0.001
Boys	865 (57.3)	436 (50.4)	429 (49.6)	
Girls	645 (42.7)	263 (40.8)	382 (59.2)	
Age (years)				<0.001
<1	402 (26.6)	334 (47.5)	68 (16.9)	
=1	323 (21.4)	130 (19.0)	193 (59.8)	
>1	785 (52.0)	235 (33.5)	550 (70.1)	
Birth season				0.048
Spring	342 (22.6)	138 (19.7)	204 (59.6)	
Summer	392 (26.0)	184 (26.3)	208 (53.1)	
Autumn	422 (27.9)	197 (28.2)	225 (53.3)	
Winter	354 (23.4)	180 (25.8)	174 (49.2)	
Parity				0.046
1 st	1,025 (67.9)	454 (64.9)	571 (55.7)	
2 nd – 5 th	478 (31.7)	238 (34.0)	240 (50.2)	
Gestational age (weeks)				<0.001
<37	92 (6.1)	62 (8.9)	30 (32.6)	
≥37	1,418 (93.9)	637 (91.1)	781 (55.1)	
Mode of delivery				<0.001
Natural labour	863 (57.2)	360 (51.5)	503 (58.3)	
Caesarean birth	641 (42.5)	333 (47.6)	308 (48.0)	
Parental atopy				<0.001
No	1,432 (94.8)	646 (92.4)	786 (54.9)	
Yes	78 (5.2)	53 (7.6)	25 (32.1)	

Sum of the number is not 1,510 due to missing data. The p-values < 0.05 were in bold.

Table 2 Descriptive statistics for outdoor temperature and DTV during different time windows attributed to the children (n=1,510).

	Total	Case	Control	P-value
	Mean (SD)	Mean (SD)	Mean (SD)	
Preconceptional				
1 year before conception	7.2 (0.5)	7.3 (0.4)	7.2 (0.5)	<0.001
3 months before conception	7.2 (0.8)	7.3 (0.7)	7.1 (0.8)	<0.001
Prenatal				
1 st trimester	7.2 (0.8)	7.3 (0.7)	7.0 (0.8)	<0.001
2 nd trimester	7.1 (0.7)	7.3 (0.7)	6.9 (0.7)	<0.001
3 rd trimester	7.1 (0.7)	7.3 (0.7)	6.9 (0.7)	<0.001
Entire pregnancy	7.1 (0.5)	7.3 (0.5)	7.0 (0.5)	<0.001

DTV (°C) = Tmax - Tmin. The p-values < 0.05 were in bold.

Table 3 Odds ratio (95%CI) of childhood pneumonia for exposure to outdoor DTV during different timing-windows (n=1,510).

	Crude OR	Adjusted model#	Adjusted model†
Preconceptional			
3 months before conception	1.40 (1.22, 1.61)***	1.98 (1.67, 2.34)***	1.15 (0.91, 1.46)
1 year before conception	1.86 (1.48, 2.32)***	10.21 (7.08, 14.71)***	2.53 (1.56, 4.10)***
Prenatal			
1 st trimester	1.72 (1.50, 1.97)***	2.30 (1.94,2.72)***	1.63 (1.32, 2.00)***
2 nd trimester	1.99 (1.71, 2.33)***	2.55 (2.12,3.07)***	1.43 (1.12, 1.81)**
3 rd trimester	1.88 (1.62, 2.18)***	2.29 (1.91,2.73)***	1.20 (0.96, 1.51)
Entire pregnancy	4.45 (3.52, 5.6)***	8.36 (6.21,11.27)***	1.85 (1.24, 2.76)**

OR (95% CI) was estimated by 1°C increase in outdoor DTV during each time window.

Model was adjusted for the covariates in Table 1.

† Model was further adjusted for outdoor air pollution exposure during each time window based on model.

* p≤0.05.

** p≤0.01.

*** p≤0.001.

Figures

Figure 1

Odds ratio (95%CI) of childhood pneumonia for exposure to diurnal temperature variation (DTV) stratified by gender during pregnancy (n=1,510). ORs was adjusted for all covariates in Table 1 and outdoor air pollutants (PM10, SO2 and NO2).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [GraphicalAbstract.docx](#)