Performance and Host Association of Spotted Lanternfly (*Lycorma Delicatula*) among Common Woody Ornamentals

Osariyekemwen Uyi

University of Benin

Keller Joseph A.

Pennsylvania State University

Kelli Hoover (✉ kxh25@psu.edu)

Pennsylvania State University

Research Article

Keywords: performance, preference, offspring, fitness, *Ailanthus altissima*, insect-plant 17 interactions.

Posted Date: April 14th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-397670/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on August 4th, 2021. See the published version at <https://doi.org/10.1038/s41598-021-95376-x>.

Performance and host association of spotted lanternfly (*Lycorma delicatula*) among common woody ornamentals

Osariyekemwen Uyi^{1,2}, Joseph A. Keller¹ & Kelli Hoover^{1*}

¹Department of Entomology, Pennsylvania State University, University Park, PA 16802, United States of America

²Department of Animal and Environmental Biology, University of Benin, P.M.B. 1154, Benin City, Nigeria

***Correspondence:** Kelli Hoover, Department of Entomology, 501 ASI Building, Pennsylvania State University, University Park, PA 16802, United States of America E-mail: kxh25@psu.edu

Running title: Performance of spotted lanternfly among woody ornamentals

Keywords: performance; preference; offspring; fitness; *Ailanthus altissima*; insect-plant interactions

Abstract

Despite the broad host range of *Lycorma delicatula*, the performance of this invasive pest on non-*Ailanthus* host plant species is difficult to document realistically without using field conditions given this pest's requirements for vast amounts of phloem. In spring 2020, we examined the performance of *L. delicatula* with and without access to *Ailanthus altissima* by tracking development, survival, host tree association and oviposition in large enclosures planted with one each of *Juglans nigra*, *Salix babylonica* and *Acer saccharinum* along with either one *A. altissima* or one *Betula nigra*. We reared nymphs with and without access to *A. altissima*, released them into the corresponding large enclosures as third instars, and monitored them from early July 2020 through November 2020. Insect survival was higher and development faster with access to *A. altissima*. Third and fourth instar nymphs were most frequently observed on *A. altissima* when it was present, while adults were equally associated with *A. saccharinum* and *A. altissima*. In the absence of *A. altissima*, nymphs were most frequently found on *S. babylonica*, while adults were most often on *A. saccharinum*. Females with access to *A. altissima* deposited nearly 7-fold more egg masses than those without access to *A. altissima*. In another experiment, the offspring of parents that had been reared without access to *A. altissima* showed similar survival and development time from egg to adult as offspring from parents that never had access to *A. altissima*. These findings suggest that managers need to be aware that even in the absence of *A. altissima* in the landscape, several hardwood host trees can be utilized by *L. delicatula* to develop and reproduce.

Keywords: performance, preference, offspring, *Lycorma delicatula*, *Ailanthus altissima*, insect-plant interaction

51

52 **Introduction**

53 Since 2014, the spotted lanternfly (SLF), *Lycorma delicatula* (White) (Hemiptera: Fulgoridae),
54 an exotic planthopper, has invaded nine states in the Northeast, mid-West, and mid-Atlantic
55 regions of the U.S. Native to China, Taiwan and Vietnam, *L. delicatula* has also expanded its
56 range to include South Korea and Japan [1-4]. Following its initial detection in Berks County,
57 Pennsylvania (Barringer et al. 2015), *L. delicatula* has spread to New Jersey, New York,
58 Virginia, Maryland, Delaware, West Virginia, Connecticut, and Ohio [4-5]. The invasion success
59 of this pest may be partly due to its apparent capacity for dispersal [6-7] and its potential to
60 occupy a wide range of climatic conditions and ecosystems, especially disturbed habitats where
61 the preferred host *Ailanthus altissima* (Mill.) Swingle (Simaroubaceae) (tree-of-heaven) is
62 abundant [3, 8, 9].

63

64 As a generalist phloem feeder, *L. delicatula* has proven to be a prominent pest in forest and
65 agricultural ecosystems and a nuisance pest in suburban landscapes, causing significant
66 economic losses to vineyards, nurseries, and sawmills in the invaded regions [4, 10]. For the
67 ornamentals and forest products industries, the major economic impacts are primarily due to the
68 cost of best management practices to comply with quarantine restrictions, and in some cases,
69 reduced sales [10-11]. *Lycorma delicatula* is projected to cause \$42.6 million in damages
70 annually if it spreads across Pennsylvania [11]. Although studies on the impacts of feeding by *L.*
71 *delicatula* on host plant health are in their infancy, phloem feeding by nymphs and adults is
72 known to cause physiological stress in young stems that may in turn cause infested branches to
73 wilt, lose vigor or die following heavy infestations [10, 12]. Feeding by *L. delicatula* results in

the production of copious amounts of honeydew, which promotes the growth of sooty mold on plants below feeding sites, impeding photosynthesis of affected plants [13-16]. Damage caused by *L. delicatula* decreases grapevine growth, reducing crop yield by up to 90% and sometimes killing the vines (*Vitis* spp. L. [Vitales: Vitaceae]) [10, 12]. Additionally, *L. delicatula* poses a threat to the growth of several hardwood trees, including silver maple (*Acer saccharinum* L. [Sapindaceae]), red maple (*A. rubrum* L. [Sapindaceae]), black walnut (*Juglans nigra* L. [Juglandaceae]), weeping willow (*Salix babylonica* L. [Salicaceae]), river birch (*Betula nigra* L. [Betulaceae]) and tulip tree (*Liriodendron tulipifera* L. [Magnoliaceae]) [1, 2, 17-19].

Lycorma delicatula prefers *A. altissima* in both its native and invasive ranges, though it can utilize over 103 plant species across 33 families [4, 17, 18]. In North America, feeding by *L. delicatula* has been observed on 56 plant species, which include native, cultivated, and nonnative species [18]. The rapid spread of this pest is likely facilitated by the prevalence of *A. altissima*, in addition to other suitable host plant species [18]. Despite the broad host range of *L. delicatula*, basic information is lacking on its host preferences and relative performance when feeding on common woody ornamentals (but see [19-20]). Our previous study in 2019 [19] found that it can complete development and reproduce on young weeping willow (*S. babylonica*), silver maple (*A. saccharinum* L.) and river birch (*B. nigra*) without access to *A. altissima*. Development to adult was delayed by one week and the number of egg masses laid by females were significantly fewer in enclosures with no access to *A. altissima*. It is, however, unclear whether *L. delicatula* without access to *A. altissima* can perform as well as those with *A. altissima* if provided larger trees to provide a greater quantity of phloem. Thus, in 2020, we investigated the performance and host association of *L. delicatula* among planted common woody ornamentals that were larger in 2020

than in 2019. A further objective of this study was to determine whether offspring (F_1 generation) from parents that were reared from egg to adult without access to *A. altissima* suffer any lingering fitness effects as they develop into adults.

Materials and methods

Origin and maintenance of trees and insect cultures.

In Macungie, Berks County, PA, we set up ten 5.76 m² plots in mid-September 2018; five plots each were randomly allocated to two treatments: ‘with *Ailanthus*’ and ‘without *Ailanthus*’ [19]. All plots contained one *S. babylonica*, one *A. saccharinum*, and one *J. nigra* spaced 1 m apart; in addition, the ‘with *Ailanthus*’ plots were also planted with one *A. altissima*, while the ‘without *Ailanthus*’ plots contained one river birch (*B. nigra*) in place of *A. altissima*. We purchased the *Salix babylonica* and *B. nigra* as 15-gallon potted trees at New Hanover Gardens (Perkiomenville, PA), and *A. saccharinum* as 7-gallon and *J. nigra* as 2-gallon trees from Octoraro Native Plant Nursery (Kirk, PA). Since *A. altissima* is a ubiquitous invasive plant, small trees were dug up and transplanted from a local property with the owner’s permission. We selected these species because they common in this region of the U.S. and are often used as hosts trees by *L. delicatula* in Pennsylvania. *L. delicatula* were confined to each plot using a rectangular screen enclosure made of PAK25 Anti-Insect Mesh (Hummert International, MO) attached to a frame (3 × 2.4 × 2.4 m) made of galvanized steel tubes (38 mm diameter) at each corner to prevent collapse during windy thunderstorm. To obtain access to each enclosure we instaleed a 1.8 m zipper on one side. To prevent *L. delicatula* nymphs or adults from escaping from the enclosures, sandbags (7.6 × 99 cm) were sewn into the bottom edge of the enclosure to

weigh down the sides. We installed drip irrigation connected to the well on the property. In the early spring, trees were pruned to minimize crowding.

In January 2020, egg masses of *L. delicatula* were field collected in Allentown, PA and kept in two pop-up cages (90 × 60 × 60 cm) under ambient environmental conditions in an unheated greenhouse in Macungie, PA. In early May, egg masses were transferred to 20 pop-up cages (90 × 60 × 60 cm) containing potted host plants and housed under a canopy at the experimental site. In this way, all plants received direct sunlight for part, but not all, of the day. These pop-up cages were divided between two treatments: ‘with *Ailanthus*’ (10 cages) and ‘without *Ailanthus*’ (10 cages). Pop-up cages ‘with *Ailanthus*’ contained one potted ~60 cm tall *A. altissima* (planted from field-collected seeds), one potted ~45 cm tall *V. vinifera* (cv. Cabernet-Franc; Hermann J. Wiemer Vineyard, Dundee, NY), one potted ~45 cm tall strawberry and one potted ~45 cm tall sunflower, while pop-up cages ‘without *Ailanthus*’ contained one potted ~45 cm tall *V. vinifera*, strawberry and sunflower. Neonates in both treatments were reared to third instar until release into each multi-tree enclosure.

Experiment I: *Lycorma delicatula* survival, development and host plant associations.

Between July 7 and July 10, 2020, 120 third instars were released into each multi-tree enclosure. Enclosures ‘with *Ailanthus*’ received nymphs with prior access to *A. altissima* from the pop-up cages with potted *A. altissima* (see details above) and vice versa [19]. Each week we monitored plots for survival, life stage, and which host tree the insects were found on by counting the numbers of nymphs (from July 15, 2020 to September 2, 2020) and adults (August 12, 2020 to November 4, 2020) present on each host tree species and on other surfaces in each enclosure.

After reaching adulthood, we began monitoring and recording for oviposition weekly from through November 4, 2020 when a hard freeze killed all remaining adults.

To determine how *L. delicatula* feeding affected tree growth, we measured trunk diameters at 10 cm above the soil line using a caliper on July 8, 2020 before releasing *L. delicatula* into the enclosures and on November 3, 2020 at the end of the experiment. To document growth between 2019 and 2020, we compared trunk diameter of trees by species between 2019 and 2020 on June 4 and October 31. Trunk diameter measurements for *J. nigra* were not available because most of these trees had died during the 2019 studies and were replaced in spring of 2020.

Experiment II: Performance of *L. delicatula* offspring.

To compare fitness of offspring from parents that did or did not have access to *A. altissima* trees during development from egg to adult, we collected egg masses from the prior experiment (as above) conducted in 2019 [19]. In early-June of 2020 these egg masses were carefully collected from the trees in each multi-tree plot, placed in Petri dishes and held in pop-up cages (90 × 60 × 60 cm) containing two potted *A. altissima* trees, with 4 replicates per treatment. The pop-up cages were placed in open field conditions in the same location but were sheltered from direct rainfall. Trees were watered thrice per week and replaced once per month with fresh potted *A. altissima* trees. Weekly, we recorded survival and development of *L. delicatula* by counting the numbers of nymphs and adults present in each cage.

Statistical analysis.

To evaluate the effect of the presence or absence of *A. altissima* on the survival of *L. delicatula* in Experiments I and II, we fit a generalized linear mixed model with a binomial error

distribution and logit link function using the *glmmTMB* function [19] in R [21-22]. Our model included treatment, date of observation, and the interaction between these two factors as predictors, and cage as a random effect to account for repeated observations. Similarly, to assess the effect of *A. altissima* presence on the timing at which individuals reached the fourth instar and adulthood in Experiments I and II, we fit a GLM mixed effects model with a binomial error distribution and a logit link function. We predicted the proportion of individuals in each cage that had reached the fourth instar and adult stages based on the date, treatment (with or without *A. altissima*) and interaction of these two factors. We again included cage as a random effect in the model to account for repeated observations. We assessed the significance of the treatment by conducting a likelihood ratio test to compare the full model against a reduced model, excluding treatment as a predictor. Repeated measures GLM ANOVA was used to compare the host plant association (i.e., the proportion of nymphs and adults of *L. delicatula* on a given tree species) of *L. delicatula* in the presence or absence of *A. altissima* using SPSS version 20.0 (IBM, SPSS Inc. Chicago, IL). If the overall model was significant, differences between trees was determined using the Bonferroni *post hoc* test. The Mann Whitney U test was used to compare the number of eggs masses between treatments using SPSS version 20.0. The impact of *L. delicatula* feeding on trunk diameter and trunk diameter growth of tree species between 2019 and 2020 was evaluated with a GLM ANOVA using SPSS version 20.0. If the overall model was significant, differences between trees was determined using the Bonferroni *post hoc* test. *Juglans nigra* trees were not included in the analysis because most died during the 2019 studies and had to be replaced in spring of 2020.

Results

Survival, development time, host association and egg mass count.

Following release into multi-tree enclosures, the proportion of individuals of *L. delicatula* (from third instar nymph to adult) that survived was approximately 10% higher in enclosures with *A. altissima* compared to the non-*Ailanthus* enclosures throughout the season ($\chi^2 = 16.29$, $df = 1$, $P = 0.001$; Fig. 1). Third instar nymphs in *A. altissima* enclosures developed slightly faster to fourth instar ($\chi^2 = 56.26$, $df = 1$, $P = 0.001$; Fig. 2); fitted logistic regression curves showed that 50% of third instars with *Ailanthus* reached the fourth instar 2.1 days earlier than those without *Ailanthus*. In cages containing *A. altissima* 50% of fourth instars reached adulthood 8.4 days earlier than those in enclosures without *A. altissima* ($\chi^2 = 9.67$; $P = 0.001$; Fig. 3).

Host plant association of third and fourth instar nymphs as well as adults in *A. altissima* enclosures were influenced by the available tree species but did not differ over time (Table 1; Fig. 4a-c). In enclosures with *A. altissima*, third instars were most frequently observed on *A. altissima* (Fig. 4a), while fourth instars were found with equal frequency on *A. altissima*, *S. babylonica* and *A. saccharinum* (Fig. 4b). Adults were mostly observed on *A. saccharinum* despite the presence of *A. altissima* (Fig. 4c). In enclosures without *A. altissima*, host plant association of nymphs and adults differed as a function of tree species but did not differ over time (Table 2; Fig. 5a-c). In enclosures without *A. altissima*, third instars were most frequently observed on *S. babylonica* (Fig. 5a), while fourth instars were found with equal frequency on *S. babylonica* and *B. nigra* (Fig. 5b). Again, adults were most commonly associated with *A. saccharinum* (Fig. 5c).

Table 1. Repeated measures binomial GLM ANOVA for data on the effect of tree species and time on the proportion of *Lycorma delicatula* nymphs and adults present on individual trees in cages where they had access to *Ailanthus altissima*.

Variable	Source of variation	df	MS	F-value	P-value
Third instars	Tree species	3	3495	11.20	0.001
	Time	16	0.023	0.001	0.999
	Tree species $\times$ Time	9	388.3	1.24	0.301
	Error	59			
Fourth instars	Tree species	4	5218	20.68	0.001
	Time	5	0.067	0.011	0.967
	Tree species $\times$ Time	20	414.2	1.84	0.061
	Error	100			
Adults	Tree species	3	14347	13.45	0.001
	Time	11	1.03	0.009	0.991
	Tree species $\times$ Time	33	358.9	2.98	0.063
	Error	176			

df: degrees of freedom; MS: mean squares.

Statistically significant values are indicated in bold.

Table 2. Repeated measures binomial GLM ANOVA for analysis of data on the effect of tree species and time on the proportion of *Lycorma delicatula* nymphs and adults present on individual trees in cages without *Ailanthus altissima*.

Variable	Source of variation	df	MS	F-value	P-value
Third instars	Tree species	3	1241	10.42	0.001
	Time	3	0.09	0.05	0.970
	Tree species $\times$ Time	9	292.3	2.45	0.057

	Error	59			
Fourth instars	Tree species	4	4071	8.22	0.001
	Time	5	0.021	0.001	0.999
	Tree species $\times$ Time	20	342.5	1.42	0.129
	Error	100			
Adults	Tree species	3	14352	13.05	0.001
	Time	11	0.32	0.003	0.993
	Tree species $\times$ Time	33	3231	2.11	0.071
	Error	176			

DF: degrees of freedom; MS: mean squares.

Statistically significant values are indicated in bold.

The number of egg masses laid by females was 6.8-fold higher in enclosures with *A. altissima* ('with *Ailanthus*': 19.20 ± 2.13 (mean $\pm$ SE) than 'without *Ailanthus*' 2.80 ± 0.42 ; $U = 20.0$, $P = 0.040$). Egg masses were first observed on September 30, 2020 in the presence of *A. altissima* and oviposition continued until November 4 when field temperature dropped below 0 °C, killing the remaining adults in both treatments. Of these egg masses, 23 were laid on *A. altissima*, 21 on *A. saccharinum*, 30 on *S. babylonica*, 1 on *J. nigra*, and 14 and 7 on wooden planks and red maple logs placed in the enclosures, respectively. Oviposition occurred in all five of these enclosures for a total of 96 egg masses. In enclosures without *A. altissima*, 14 egg masses were recorded, with at least one egg mass in each of the 5 enclosures. Oviposition in these cages occurred between October 14 and November 4, 2020. Of these 14 egg masses, one was on *B. nigra*, 2 on *S. babylonica*, 6 on *A. saccharinum*, 2 on planks and 3 on red maple logs.

Impact of *Lycorma delicatula* feeding on tree species.

For both treatments, trunk diameter differed significantly by year ('with *Ailanthus*': $F_{1,29}=79.27$; $P = 0.0001$; 'without *Ailanthus*': $F_{1,29}=40.84$; $P = 0.0001$) and tree species ('with *Ailanthus*': $F_{2,29}=45.47$; $P = 0.0001$; 'without *Ailanthus*': $F_{2,29}=101.87$; $P = 0.0001$). There were no

significant tree species $\times$ year interactions ('with *Ailanthus*': $F_{2,29}=1.15$; $P = 0.334$; 'without *Ailanthus*': $F_{2,29}=2.25$; $P = 0.070$). Trees used in 2020 were on average $26.0 \pm 2.3\%$ larger than the same trees were in 2019 (Fig. 6a and b). Trunk diameter increase differed significantly by year ('with *Ailanthus*': $F_{1,29}=14.47$; $P = 0.0001$; 'without *Ailanthus*': $F_{1,29}=13.25$; $P = 0.0001$) but not by tree species ('with *Ailanthus*': $F_{2,29}=1.51$; $P = 0.252$; 'without *Ailanthus*': $F_{2,29}=0.10$; $P = 0.903$). There were no significant tree species $\times$ year interactions ('with *Ailanthus*': $F_{2,29}=0.11$; $P = 0.899$; 'without *Ailanthus*': $F_{2,29}=3.10$; $P = 0.057$). Trees in both treatments and of every species except river birch grew ≥ 2 -times more in diameter in 2019 on average than the same trees grew in 2020; river birch grew 1.2-times more in 2019 than in 2020 (Figs. 7a and b).

Offspring performance.

After being released in popup cages with access to healthy *A. altissima*, the proportion of hatched offspring from eggs laid the previous fall did not differ as a function of parental diet ($\chi^2 = 0.70$, $df = 1$, $P = 0.401$). Development time (hatch to adults) also did not differ between offspring from parents with or without access to *A. altissima* ($\chi^2 = 0.13$, $df = 1$, $P = 0.722$).

Discussion

Lycorma delicatula can complete development and reproduce without access to *A. altissima*, confirming our findings from the previous year [19], and providing additional insight into host association and the impact of *L. delicatula* on diameter growth of common woody ornamental trees in response to feeding pressure from *L. delicatula*. Fitness of the insect was greater in the presence of *A. altissima*. Survival was higher, development time was faster, and the number of egg masses was 6.8-fold higher than for insects that lacked access to *A. altissima*. This study

also showed that offspring from parents that had been reared without access to *A. altissima* did not suffer fitness effects into the next generation. We hypothesize that reduced egg mass production in cages without *Ailanthus* was caused by slower development resulting in a shorter time span between adult emergence and freezing-induced mortality during which adults can mate and lay eggs. However, in southern climates where freezing temperatures appear several months later in the year or not at all, lack of access to *A. altissima* may have less impact on fitness since there could be time for slower developing adults to continue oviposition into the early winter.

In this study, survival to adult was high for both treatments (with and without *Ailanthus*), although survival was slightly higher (10% difference) in the presence of *A. altissima*. By early September 2020, approximately 63% of the individuals released in *A. altissima* enclosures and non-*Ailanthus* enclosures were still alive. This is in contrast to our 2019 study where survival on younger, smaller trees was less than 20% for both treatments (with and without *Ailanthus*) by September 2 when most lanternflies were adults [19]; however, the previous study started with the release of newly hatched first instars. Lower survival in 2019 could have been due to early mortality of first and second instar nymphs and/or having access to smaller trees in 2019 than in 2020 in that larger and vigorous trees may provide a greater volume of phloem and nutrients over time.

The faster development time of *L. delicatula* in enclosures with *A. altissima* suggests that adults could reach sexual maturity faster in areas where *A. altissima* is abundant. Development time remains one of the most crucial fitness indicators of host plant nutritional quality and can influence insect survival, behavior and physiology [23]. For example, several studies have found

that slow development in phytophagous insects can cause high mortality of immature stages in the field by exposing them to unfavorable environmental conditions for a longer time period [24-25]. A key fitness cost caused by prolonged development in *L. delicatula* was narrowing of the reproductive window for adults, which was reflected in markedly fewer eggs being laid before freezing temperatures killed the remaining adults [19].

Third and fourth instar nymphs were mostly associated with *A. altissima* when it was present, however, more adults were found on *A. saccharinum* than on *A. altissima* when both tree species were present. This may occur when the pest has exhausted the ability of *A. altissima* to provide sufficient sap flow, which also coincides with the onset of senescence of this tree species in mid-September. In the absence of *A. altissima*, *B. nigra*, *A. saccharinum*, *J. nigra* and *S. babylonica* together appeared to provide sufficient nutrition for growth and reproduction, but sap flow may be lower in these species, which could explain the delayed development time in the absence of *A. altissima*. However, we cannot rule out that if given access to mature and vigorous suitable hosts such as *A. saccharinum* or *A. rubrum*, sexual maturity could have occurred at the same rate as those with access to *A. altissima*. These two maple species are heavily utilized by *L. delicatula* adults in the field in the fall (D.D. Calvin et al. unpublished data). Although the trees in our study were significantly larger than they were the previous year, they were still smaller than trees that are selected by wild populations of *L. delicatula* adults in the field where profuse feeding is associated with reproductive maturation (pers. observations).

Several factors ranging from sap flux to host-tree bark characteristics and the presence or absence of defensive chemicals may influence feeding preference in *L. delicatula* [26-27]. For

example, *L. delicatula* has been reported to survive longer on, and prefer, host trees with a high concentration of sugars similar to those produced by *A. altissima* and *V. vinifera* [28]. In prior work by Lee and colleagues [19], the authors reasoned that the improved performance on and preference for *A. altissima* and *V. vinifera* is related to high concentrations of sucrose and glucose in *A. altissima* phloem, and high proportions of sucrose and fructose in *V. vinifera* phloem. Several authors have also noted that *L. delicatula* seems to prefer hosts with trunks and branches that do not have thick bark but have high sap turgor pressure, allowing for ingestion of phloem at a greater rate [17, 18, 20]. The trees used in our study were chosen because they are frequently infested in the field and do not have the thick bark that occurs on oak trees, for example, which are infrequent hosts [29].

Although egg masses were laid on all host trees as well as planks and logs, *S. babylonica*, *A. saccharinum*, and *A. altissima* were generally preferred for oviposition. Oviposition substrate selection can be vital to the reproductive success of insect herbivores. The preference-performance hypothesis (a.k.a. mother knows best hypothesis) predicts correlation between oviposition preference by the female parent and host suitability for offspring development [30-31]. However, it does not translate well to species like *L. delicatula* with immature stages that are highly mobile or instances where the adults and immature life stages feed on different hosts [32]. Although tree species and branch structure may play a role in the selection of sites for oviposition [17], why *L. delicatula* oviposits on non-living materials remains to be explained. The basis for oviposition preferences is not always easy to empirically verify [31]. Some *L. delicatula* researchers think that females lay most of their eggs in the vicinity of where they fed to fatten up and become reproductively mature because it's harder for them to fly once they've

gained so much weight [33]. Alternatively, as is evident in some insects and mite species [31, 34], *L. delicatula* may be following a strategy where females do not necessarily oviposit on suitable hosts in order to make it harder for predators to find their eggs and early instars.

The ability of *L. delicatula* to survive, develop and produce egg masses with or without *A. altissima* may be due to the presence of multiple, suitable host plant species in the enclosures. It is not uncommon for generalist feeders to require diet mixing to acquire the necessary nutrients for development and reproduction [35-36], and *L. delicatula* appears to demonstrate better fitness when provided multiple suitable host plants. While they can be reared from nymphs to adults on a single species such as *V. vinifera* or *A. altissima*, without diet mixing, adults may not produce eggs. For example, *L. delicatula* that were reared exclusively on *A. altissima* or *V. vinifera* failed to produce egg masses, whereas their counterparts that received a combination of both did (Tracy Leskey, USDA/ARS, personal communication).

The fact that trunk diameter growth of trees was significantly greater (by 3 mm) in 2019 compared to 2020 suggests that exposure of host trees to *L. delicatula* feeding pressure for two seasons in a row may have limited the growth of these trees, although abiotic factors such as climate could play a role. Our results also suggest that economically important ornamentals trees and other indigenous host tree species may be significantly impacted when exposed to feeding pressure for more than one season. Feeding by nymphs and adults of *L. delicatula* can reduce photosynthesis and carbon storage (Lavelly et al. unpublished data; Centinari et al. unpublished data), potentially causing stunted growth and/or mortality of host plants [37-38]. Although feeding by *L. delicatula* can result in oozing trunk wounds, wilting, branch death, and sooty

mold [15], the extent of damage and effects of herbivory on woody ornamental trees in North America are only beginning to be investigated. Our results on the impact of *L. delicatula* feeding should be considered preliminary and need to be interpreted with caution as our study lacked the adequate control (trunk diameter growth) data to make a definitive statement on the impact of *L. delicatula* feeding on host tree species.

Although *L. delicatula* developed more slowly without access to *A. altissima*, the offspring from eggs produced by these adults did not suffer ongoing fitness effects, which could have implications for the spread of this pest in the United States. While range expansion of *L. delicatula* is likely facilitated by the prevalence of *A. altissima*, it is clear that other hardwood tree species can serve as suitable hosts to support development and reproduction. It's possible that in a warmer southern climate, adults that develop more slowly in a region without *A. altissima* may have time to catch up and lay as many egg masses as they would have if there was access to *A. altissima*. Overall, our findings suggest that managers need to be aware that even in the absence of *A. altissima* in the landscape, several hardwood host trees can be utilized by *L. delicatula* to develop and reproduce.

Acknowledgements

The authors would like to thank David Long, Anne Johnson, Elizabeth Wagner, Emelie Swackhamer and Scott Alan Swackhamer for assistance during the field study. Funding was provided by Cooperative Agreements with USDA Forest Service Forest Health Protection (Grant 18-CA-11420004-198) and USDA APHIS (Award No: AP19PPQS&T00C080) and grants from the USDA NIFA and McIntire-Stennis Appropriations under Project #PEN04684 and Accession

#1018126, the Pennsylvania Department of Agriculture (Grant No. 44187340), and the Horticultural Research Institute. This work was also supported in part by the USDA National Institute of Food and Agriculture and Hatch Appropriations under Project #PEN04728 and Accession #1021211.

Competing interests

The author declares no competing interests.

Data Availability Statement

All relevant data are within the paper

Author contributions

KH and OU conceptualized and designed the study; OU performed the field experiments; OU and JK performed the statistical analysis; OU wrote the manuscript; KH and JK critically reviewed and amended the manuscript.

IUCN Policy Statement

Collection of plant material was done in compliance with relevant institutional, national, and international guidelines and legislation.

References

1. Barringer, L.E., Donovall, L.R. Spichiger, S.E., Lynch, D & Henry, D. The first New World record of *Lycorma delicatula* (Insecta: Hemiptera: Fulgoridae). *Entomological News* **125**, 20–23 (2015).
2. Dara, S.K., Barringer, L. & Arthurs, S.P. *Lycorma delicatula* (Hemiptera: Fulgoridae): a

new invasive pest in the United States. *Journal of Integrated Pest Management* **6**, 20 (2015).

3. Jung, J.M., Jung, S., Byeon, D. & Lee, W. Model-based prediction of potential distribution of the invasive insect pest, spotted lanternfly *Lycorma delicatula* (Hemiptera: Fulgoridae), by using CLIMEX. *Journal of Asia- Pacific Biodiversity* **10**, 532–538 (2017).
4. Lee, D.-H., Park, Y.-L. & Leskey, T.C. A review of biology and management of *Lycorma delicatula* (Hemiptera: Fulgoridae), an emerging global invasive species. *Journal Asia-Pacific Entomology* **22**, 589–596 (2019).
5. (NYSIPM) New York State Integrated Pest Management. 2020. Spotted lanternfly. <https://nysipm.cornell.edu/environment/invasive-species-exotic-pests/spotted-lanternfly/> (accessed 18 January 21).
6. Park, M., Kim, S.M. & Lee, J.H. Genetic structure of *Lycorma delicatula* (Hemiptera: Fulgoridae) populations in Korea: implication for invasion processes in heterogeneous landscapes. *Bulletin of Entomological Research* **103**, 414–424 (2013).
7. Keller, J.A., Johnson, A.E., Uyi, O., Wurzbacher, S., Long, D., & Hoover, K. Dispersal of *Lycorma delicatula* (Hemiptera: Fulgoridae) nymphs through contiguous, deciduous forest. *Environmental Entomology* **49**, 1012–1018 (2020).
8. Smyers, E.C., Urban, J.M., Dechaine, J.D., Pfeiffer, D.G., Crawford, S.R., Calvin, D.D. (2020). Spatio-temporal model for predicting spring hatch of the spotted lanternfly (Hemiptera: Fulgoridae). *Environmental Entomology* **50**, 126-137.
9. Wakie, T.T., Neven, L.G., Yee, W.L. & Lu, Z. The establishment risk of *Lycorma delicatula* (Hemiptera: Fulgoridae) in the United States and globally. *Journal of*

- Economic Entomology* 113, 306–314 (2019).
10. Urban, J.M. 2020. Perspective: shedding light on spotted lanternfly impacts in the USA. *Pest Management Science* **76**, 10-17.
11. Harper, J.K., Stone, W., Kelsey, T.W. & Kime, L.F. Potential Economic Impact of the spotted lanternfly on agriculture and forestry in Pennsylvania. The Center for Rural Pennsylvania, Harrisburg, PA. 84p (2019).
12. Song, M.K. Damage by *Lycorma delicatula* and chemical control in vineyards [M.S. thesis]. Korea: Chunbuk National University. 39p (2010).
13. Tedders, W.L., & Smith J.S. Shading effect on pecan by sooty mold growth. *Journal Economic Entomology* **69**, 551-553 (1976).
14. Lemos-Filho, J.P.D. & Paiva, É.A.S. The effects of sooty mold on photosynthesis and mesophyll structure of mahogany (*Swietenia macrophylla* King., Meliaceae). *Bragantia* **65**, 11-17 (2006).
15. Han, J.M., Kim, H.J., Lim, E.J. Lee, S.H., Kwon, Y.J. & Cho, S.W. *Lycorma delicatula* (Hemiptera: Auchenorrhyncha: Fulgoridae: Aphaeninae) finally, but suddenly arrived in *Korean Entomological Research* **38**, 281–286 (2008).
16. Park, J.D., Kim, M.Y., Lee, S.G., Shin, S.C., Kim, J.H. and Park, I.K. Biological characteristics of *Lycorma delicatula* and the control effects of some insecticides. *Korean Journal Applied Entomology* **48**, 53–57 (2009).
17. Liu, H. 2019. Oviposition substrate selection, egg mass characteristics, host preference, and life history of the spotted lanternfly (Hemiptera: Fulgoridae) in North America. *Environmental Entomology* **48**, 1452-1468.
18. Barringer, L.E. & Ciafré, C.M. Worldwide feeding host plants of spotted lanternfly, with

- significant additions from North America. *Environmental Entomology* **49**, 999-1011 (2020).
19. Uyi, O., Keller, J.A., Johnson, A., Long, D., Walsh, B. & Hoover, K. Spotted lanternfly (Hemiptera: Fulgoridae) can complete development and reproduce without access to the preferred host, *Ailanthus altissima*. *Environmental Entomology* **49**, 1185-1190 (2020).
20. Murman, K. Distribution, survival, and development of spotted lanternfly on host plants found in north America. *Environmental Entomology* **49**, 1270-1281 (2020)
21. Magnusson, A. glmmTMB: Generalized linear mixed models using template model builder. R package v. 0.1.3. [https://github.com/ glmmTMB](https://github.com/glmmTMB). (2017).
22. R Development Core Team. *R: a language and environment for statistical computing computer program, version 3.6.3*. R Development Core Team, Vienna, Austria (2020).
23. Kariyat, R.R. & Portman, S.L. Plant–herbivore interactions: Thinking beyond larval growth and mortality. *American Journal of Botany* **103**,1–3 (2016).
24. Fordyce, J.A., & Shapiro, A.M. Another perspective on the slow-growth/high-mortality hypothesis: chilling effects on swallowtail larvae. *Ecology* **84**, 263–268 (2003).
25. Uesugi, A. 2015. The slow-growth high-mortality hypothesis: direct experimental support in a leaf mining fly. *Ecological Entomology* **40**, 221-228.
26. Song, S., Kim, S., Kwon, S.W., Lee, S-I. & Jablonski, P.G. 2018. Defense sequestration associated with narrowing of diet and ontogenetic change to aposematic colours in the spotted lanternfly. *Scientific Reports* **8**, 16831 (2018).
27. Domingue, M.J., & Baker, T.C. Orientation of flight for physically disturbed spotted lanternflies, *Lycorma delicatula*, (Hemiptera, Fulgoridae). *Journal of Asia Pacific Entomology* **22**, 117–120 (2019).

28. Lee, J.E., Moon, S.R., Ahn, H.G., Cho, S.R. Yang, J.O., Yoon, C. & Kim, J.H. Feeding behavior of *Lycorma delicatula* (Hemiptera: Fulgoridae) and response on feeding stimulants of some plants. *Korean Journal of Applied Entomology* **48**, 467–477 (2009).
29. Liu, H. Seasonal development, cumulative growing degree-days, and population density of spotted lanternfly (Hemiptera: Fulgoridae) on selected hosts and substrates. *Environmental Entomology* **49**, 1171–1184 (2020).
30. Jaenike, J. On optimal oviposition behavior in phytophagous insects. *Theoretical Population Biology* **14**, 350–356 (1978).
31. Gripenberg, S., Mayhew, P.J., Parnell, M. & Roslin, T. A meta-analysis of preference–performance relationships in phytophagous insects. *Ecology Letters* **13**, 383–393 (2010).
32. Fujiyama, N., Torii, C., Akabane, M. & Katakura, H. Oviposition site selection by herbivorous beetles: a comparison of two thistle feeders: *Cassida rubiginosa* and *Henosepilachna niponica*. *Entomologia Experimentalis et Applicata* **128**, 41–48 (2008).
33. Wolfin, M.S., Myrick, A.J. & Baker, T.C. Flight duration capabilities of dispersing adult spotted lanternflies, *Lycorma delicatula*. *Journal of Insect Behavior* **33**, 125–137 (2020).
34. Faraji, F., Janssen, A., Sabelis, M.W. Oviposition patterns in a predatory mite reduce the risk of egg predation caused by prey. *Ecological Entomology* **27**, 660–664 (2002).
35. Behmer, S.T. & Joern, A. Coexisting generalist herbivores occupy unique nutritional feeding niches. *Proceedings of the National Academy of Sciences, USA*. **105**, 1977–1982 (2008).
36. Behmer, S.T. Insect herbivore nutrient regulation. *Annual Review of Entomology* **54**, 165–187 (2009).
37. Ding, J, Wu, Y., Zheng, H., Fu, W., Reardon, R. & Liu, M. Assessing potential biological

control of the invasive plant, tree-of-heaven, *Ailanthus altissima*. *Biocontrol Science and Technology* **16**, 547-566 (2006).

38. Anderson, H. et al. Pest risk analysis for *Lycorma delicatula* (Hemiptera: Fulgoridae). European and Mediterranean Plant Protection Organization 16-21777 (2015).

Figure legends

Figure 1. Proportion of *Lycorma delicatula* (nymphs and adults combined) that survived in enclosures with planted *Ailanthus altissima*, *Acer saccharinum*, *Juglans nigra* and *Salix babylonica* ('*Ailanthus*,' $n = 5$ enclosures) or enclosures with the same tree species except for the presence of *Betula nigra* in place of *A. altissima* ('without *Ailanthus*,' $n = 5$ enclosures) from July 15 to November 4, 2020. Fitted binomial regression lines are shown, with shaded areas indicating 95% confidence intervals for predicted regression means. In total, 120 third-instar nymphs were released in each enclosure and their numbers and host tree associations were recorded weekly.

Figure 2. Proportion of *L. delicatula* that developed from the third to the fourth instar over time, with fitted logistic regression lines, in enclosures with and without *Ailanthus altissima*.

Figure 3. Proportion of *L. delicatula* that emerged as adults over time, with fitted logistic regression lines, for enclosures with and without *Ailanthus altissima*.

Figure 4. Mean ($\pm$ SE) proportion of *Lycorma delicatula* third instars (A) fourth instars (B) and adults (C) present on each tree species in enclosures with *Ailanthus* (*Ailanthus altissima*), silver maple (*Acer saccharinum*), black walnut (*Juglans nigra*) and willow (*Salix babylonica*) ($n = 5$ enclosures) from July 15 to November 4, 2020. Nymphs and adults found on annual plants (2%) within the enclosure or on enclosure walls (98%) are included as ‘Others.’ Means within a column followed by different letters are significantly different (Bonferroni post hoc test: $P < 0.05$). No adults were found on black walnut and no third instar nymphs were found on other surfaces.

Figure 5. Mean ($\pm$ SE) proportion of *Lycorma delicatula* third instars (A), fourth instars (B) and adults (C) present on each tree species in enclosures with silver maple (*Acer saccharinum*), black walnut (*Juglans nigra*), willow (*Salix babylonica*) and river birch (*Betula nigra*) (without *Ailanthus altissima*, $n = 5$ enclosures) from July 15 to November 4, 2020. Nymphs and adults found on other plants (2%) within the enclosure or on enclosure walls (98%) are included as ‘Others.’ Means within a column followed by different letters are significantly different (Bonferroni post hoc test: $P < 0.05$). No adults were found on black walnut and no third instar nymphs were found on other surfaces.

Figure 6. Mean ($\pm$ SE) trunk diameter (mm) of tree species in enclosures with *Ailanthus* (A) and without *Ailanthus* (B) ($n = 5$ enclosures) in 2019 and 2020. Measurements were made on October 31, 2019 and November 3, 2020. Means within a column followed by different letters are significantly different (Bonferroni post hoc test: $P < 0.05$). Black walnut trees were not included in the analysis because most died during the 2019 studies and had to be replaced in spring of 2020.

587 **Figure 7.** Mean ($\pm$ SE) increase in trunk diameter (mm) of tree species in enclosures with
588 *Ailanthus* (A) and without *Ailanthus* (B) ($n = 5$ enclosures) in 2019 and 2020. Measurements
589 were made on June 4 and October 31, 2019, and July 8 and November 3, 2020. Increase in trunk
590 diameter was obtained by subtracting trunk diameter data collected at the start of the experiment
591 from those collected at the end of the experiment. Diameter increase did not differ ($P > 0.05$)
592 between species in 2019 and 2020 but differed ($P < 0.05$) between 2019 and 2020 in both sets of
593 cages (in ‘with *Ailanthus* cages’: $F_{1,29}=14.47$; $P = 0.0001$; in ‘without *Ailanthus*’ cages:
594 $F_{1,29}=13.25$; $P = 0.0001$). Means within a column followed by asterisks are significantly different
595 (Bonferroni post hoc test: $P < 0.05$). Black walnut trees were not included in the analysis because
596 most died during the 2019 studies and had to be replaced in spring of 2020.

Figures

Figure 1

Proportion of *Lycorma delicatula* (nymphs and adults combined) that survived in enclosures with planted *Ailanthus altissima*, *Acer saccharinum*, *Juglans nigra* and *Salix babylonica* ('Ailanthus,' n = 5 enclosures) or enclosures with the same tree species except for the presence of *Betula nigra* in place of *A. altissima* ('without Ailanthus,' n = 5 enclosures) from July 15 to November 4, 2020. Fitted binomial regression lines are shown, with shaded areas indicating 95% confidence intervals for predicted regression means. In total, 120 third-insta nymphs were released in each enclosure and their numbers and host tree associations were recorded weekly.

Figure 2

Proportion of *L. delicatula* that developed from the third to the fourth instar over time, with fitted logistic regression lines, in enclosures with and without *Ailanthus altissima*.

Figure 3

Proportion of *L. delicatula* that emerged as adults over time, with fitted logistic regression lines, for enclosures with and without *Ailanthus altissima*.

Figure 4

Mean ($\pm$ SE) proportion of *Lycorma delicatula* third instars (A) fourth instars (B) and adults (C) present on each tree species in enclosures with *Ailanthus* (*Ailanthus altissima*), silver maple (*Acer saccharinum*), black walnut (*Juglans nigra*) and willow (*Salix babylonica*) (n = 5 enclosures) from July 15 to November 4, 2020. Nymphs and adults found on annual plants (2%) within the enclosure or on enclosure walls (98%) are included as 'Others.' Means within a column followed by different letters are significantly different (Bonferroni post hoc test: P < 0.05). No adults were found on black walnut and no third instar nymphs were found on other surfaces.

Figure 5

Mean ($\pm$ SE) proportion of *Lycorma delicatula* third instars (A), fourth instars (B) and adults (C) present on each tree species in enclosures with silver maple (*Acer saccharinum*), black walnut (*Juglans nigra*), willow (*Salix babylonica*) and river birch (*Betula nigra*) (without *Ailanthus altissima*, $n = 5$ enclosures) from July 15 to November 4, 2020. Nymphs and adults found on other plants (2%) within the enclosure or on enclosure walls (98%) are included as 'Others.' Means within a column followed by different letters are significantly different (Bonferroni post hoc test: $P < 0.05$). No adults were found on black walnut and no third instar nymphs were found on other surfaces.

Figure 6

Mean ($\pm$ SE) trunk diameter (mm) of tree species in enclosures with *Ailanthus* (A) and without *Ailanthus* (B) ($n = 5$ enclosures) in 2019 and 2020. Measurements were made on October 31, 2019 and November 3, 2020. Means within a column followed by different letters are significantly different (Bonferroni post hoc test: $P < 0.05$). Black walnut trees were not included in the analysis because most died during the 2019 studies and had to be replaced in spring of 2020.

Figure 7

Mean ($\pm$ SE) increase in trunk diameter (mm) of tree species in enclosures with Ailanthus (A) and without Ailanthus (B) ($n = 5$ enclosures) in 2019 and 2020. Measurements were made on June 4 and October 31, 2019, and July 8 and November 3, 2020. Increase in trunk diameter was obtained by subtracting trunk diameter data collected at the start of the experiment from those collected at the end of the experiment. Diameter increase did not differ ($P > 0.05$) between species in 2019 and 2020 but differed ($P < 0.05$) between 2019 and 2020 in both sets of cages (in 'with Ailanthus cages': $F_{1,29}=14.47$; $P = 0.0001$; in 'without Ailanthus' cages: $F_{1,29}=13.25$; $P = 0.0001$). Means within a column followed by asterisks are significantly different (Bonferroni post hoc test: $P < 0.05$). Black walnut trees were not included in the analysis because most died during the 2019 studies and had to be replaced in spring of 2020.