Aphid Species Specializing on Milkweed Have Distinct Bacterial Symbiont Communities

Laramy Enders (✉ lenders@purdue.edu)

Purdue University <https://orcid.org/0000-0002-7569-1171>

Thorsten Hansen

Purdue University

Kirsten Brichler

Virginia Polytechnic Institute and State University

John Couture

Purdue University

Elizabeth French

Purdue University

Research Article

Keywords: Milkweed, aphid, symbiont, microbiome, host plant specialization, plant-insect interactions

Posted Date: April 13th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-399030/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Host plant range is arguably one of the most important factors shaping microbial communities associated with insect herbivores. However, it is unclear whether host plant specialization limits microbial community diversity or to what extent herbivores sharing a common host plant evolve distinct microbiomes. To investigate whether variation in host plant specialization influences the composition of herbivore symbiont populations we compared bacterial diversity across three milkweed aphid species (*Aphis nerii*, *Aphis asclepiadis*, *Myzocallis asclepiadis*) feeding on a common host plant (*Asclepias syriaca*) using 16S rRNA metabarcoding. Overall, bacterial species richness did not vary with degree of host plant specialization. However, aphid species harbored distinct bacterial communities that varied in composition and relative abundance of key symbionts. Differences in aphid microbiomes were primarily due to strain variation in the obligate symbiont *Buchnera* and facultative symbiont *Arsenophonus*, as most of the low-abundant taxa were found in all three species. Interestingly, *A. asclepiadis* harbored a greater diversity of unique strains of *Buchnera* and significantly higher *Arsenophonus* relative abundances compared to the other two aphid species. Although many low abundance microbes were shared across all milkweed aphids, key differences exist in symbiotic partnerships that could influence additional ecological variation, including variation in ant tending observed across milkweed aphid species via microbial induced changes to honeydew or defensive chemical profiles. This study suggests generalist and specialist herbivore microbiomes are similar when feeding on a common host plant and highlights an intriguing potential role for strain level variation of key aphid symbionts in host-plant interactions.

Introduction

Establishing partnerships with microbes is hypothesized to enhance the evolutionary and adaptive potential of the host organism [1–4]. In fact, both plants and insects harbor diverse microbial communities that provide unique advantages, including protection from stress and novel mechanisms for nutrient acquisition [4–7]. However, only recently have microbes come to be viewed as the architects of plant-insect interactions. From an insect perspective, microbes can be critical in facilitating or restricting the use of host plants by aiding in digestion of plant material and detoxification of anti-herbivore chemical defenses [3, 5, 8]. Soil and root-associated microbes also play essential roles in plant growth and defense against both above and belowground insect attackers [6, 7, 9, 10]. Although microbes unquestionably impact plant-insect interactions, the evolutionary forces shaping plant and insect-associated microbial communities remain poorly understood.

One factor thought to play a pivotal role in shaping insect herbivore microbiomes is host plant range or diet breadth, which can vary from a single species to hundreds of plant species [11–16]. Several plant associated factors, including nutritional quality and defensive chemistry, are known to influence insect microbial community dynamics by selecting for taxa that facilitate enhanced colonization and survival [17]. For example, associations with heritable bacterial symbionts have long been recognized as enabling phloem feeding insects to exploit an otherwise poor nutritional source [17, 18]. Microbes are also

hypothesized to contribute to variation in the capacity of insect herbivores to consume chemically defended plants [8]. Finally, feeding on different host plants can influence the composition of microbial communities associated with insect herbivores [11, 14, 19, 20]. However, aside from a handful of well-characterized nutritional symbionts it is unclear to what extent insect microbial communities either directly or indirectly contribute to host plant adaptation and diet breadth. Furthermore, the mechanisms contributing to variation in herbivore microbiomes and ecological implications for plant-insect coevolution are not well understood [14].

Insect microbiomes consist of varying combinations of heritable symbionts and flexible pools of environmentally acquired microbes, but how host plant range shapes the taxonomic and functional diversity of these communities is relatively unexplored [5, 21–23]. Microbial communities of specialist herbivores may be less diverse relative generalist species that possibly require a larger microbial repertoire to successfully colonize highly variable hosts or simply acquire greater environmental microbial diversity from feeding on a broader host plant range [5, 21]. Alternatively, overlapping host plant range could result in selection for common microbes needed for successful colonization across different herbivore species due to similarities in nutritional ecology or defensive chemistry. Currently, it is unknown whether broader patterns associated with host plant use exist, such as a gradient in microbiome diversity from specialist to generalist herbivores.

The current study aims to address the following question: How does the degree of host plant specialization shape insect herbivore symbiont community diversity and composition? Heritable bacterial symbionts are hypothesized to expand the diet breadth of sap-feeding insects, particularly aphids [23–25]. However, aphid studies have primarily focused on the role microbial symbionts play in host plant use by highly polyphagous species, while much less is known about monophagous or oligophagous species. Aphids colonizing milkweed species in the subfamily *Asclepiadaceae* show varying degrees of host specialization [26] and thus represent an excellent system in which to investigate the role diet breadth plays in shaping aphid microbial community composition. *Aphis nerii* is considered a broad specialist feeding on over 50 plant species, including oleander, milkweed, and periwinkle. *Aphis asclepiadis* is a narrow specialist, feeding on less than 10 *Asclepias* spp., and *Myzocalis asclepiadis* is monophagous, feeding only on the common milkweed *Asclepias syriaca* [26, 27]. It is thought that milkweed aphid species have few microbial symbionts besides the universal obligate symbiont *Buchnera*, possibly due to the toxicity of milkweed defensive chemicals (e.g. cardenolides) which when ingested could make the internal aphid environment inhospitable for sustained microbial growth (e.g. via direct anti-microbial effects). However, few populations have been screened for symbionts and thus the bacterial community of milkweed aphids remains largely unexplored [28]. We therefore characterized the diversity and composition of bacterial communities using targeted amplicon sequencing of field collected aphids found naturally colonizing common milkweed (*A. syriaca*).

Methods

Aphid Field Collection & Sample Preparation

Aphids were collected in July-Aug 2017 from a single host plant type, the common milkweed (*A. syriaca*), across 14 locations generally surrounding the Purdue campus (Online Resource 1 Table S1). Locations were chosen based on the presence of large patches of common milkweed in open grassy fields away from roadsides. Distance between locations ranged from a minimum of 1km to a maximum of 141km (Online Resource 1 Table S1). To avoid sampling heavily from single aphid clonal families (e.g. offspring from a single alate aphid colonizing a plant), aphids were collected from 3-5 separate plants (<20m apart) within a single location (e.g. from distinct milkweed patches when possible) and pooled together into a single sample. In some cases, 2 aphid species were found co-colonizing a single plant and thus single species versus mixed species samples were designated as such during data collection (Online Resource 1 Table S1). Individual aphids were removed from milkweed leaves by hand using a paint brush, identified and separated by species, and then placed in Eppendorf storage tubes containing 95% ethanol. All individuals collected from each separate plant at each location were stored at -20C until further processing. The total number of aphids collected per species at a single location ranged from 10 individuals to >50 individuals.

For bacterial community profiling, groups of 5 individual aphids per species were selected from the total pooled sample per location for DNA extraction and sequencing. Total DNA was extracted from groups of 5 whole aphids using the Qiagen DNeasy kit following standard protocols. Aphids were not surface sterilized beyond storage in 95% ethanol and therefore, while bacterial DNA recovered is expected to be primarily from internal communities, not all external or cuticular bacterial DNA was eliminated and is expected to contribute to overall diversity estimates [29]. Total DNA concentration was measured using a Nanodrop spectrophotometer for all samples. In total 65 pools of aphids were processed and sequenced for microbiome analysis. Targeted amplicon sequencing was used to characterize the bacterial communities associated with the three milkweed aphid species (*A. nerii*, *A. asclepiadis*, *M. asclepiadis*) across the 19 locations sampled. Library preparation and sequencing was performed at the University of Minnesota Genomics Core Facility on an Illumina MiSeq instrument (V3 cluster chemistry) following optimized methods [30] and using primers that target a 250bp segment of the V4 region of the 16s rDNA gene. Primers used were standard V4 region primers 515F-GTGCCAGCMGCCGCGGTAA and 806R-GGACTACHVGGGTWTCTAAT. All sequence data is available on NCBI SRA database under project number PRJNA635683.

Characterization & Analysis of Aphid Bacterial Communities

Sample demultiplexing was done by the University of Minnesota Genomics Center with Illumina software. Trimmomatic [v. 0.36; 32] and Cutadapt [v 1.13; 30] were used to remove adapters and primer sequences and low quality reads. All subsequent processing was performed in R (v 3.6.3) and Bioconductor (v 3.10). Trimmed reads were processed through the dada2 (v 1.14.1) pipeline by filtering and trimming based on read quality, inferring error rates, merging paired end reads, removing chimeras, and assigning taxonomy with the Silva reference database v. 132 [33]. Removal of very low abundance reads was done using a cut-off of fewer than 10 reads in 5% of the samples. Next, eukaryotic and mitochondrial sequences were

removed. Lastly, samples with fewer than 2500 reads were removed. A summary of raw sequencing results and processing steps can be found in Online Resource 1.

Following sequence processing, all downstream analyses were run in R (v 3.6.3). All code for statistical analyses and generation of figures can be found in the Purdue University Github (<https://github.itap.purdue.edu/LaramyEndersGroup/Milkweed-Aphid-Microbiome>). Specifically, we compared standard alpha and beta diversity metrics using the phyloseq [v 1.30.0; 32] and vegan [v 2.5-6; 33] packages in R to determine the extent to which microbiomes varied in taxonomic composition and structure across aphid species and sampling locations. To compare species richness and evenness we used the Kruskal-Wallis test to test the effect of aphid species on diversity metrics, followed by the Wilcoxon Rank Sum test to identify significant differences among aphid species. Differences in the structure of bacterial communities across aphid species was assessed through PERMANOVA analysis of beta diversity (Unifrac, weighted Unifrac, Bray-Curtis) and visualized using Non-Metric Multidimensional Scaling (NMDS). Homogeneity of dispersion across aphid species' samples was also tested using PERMDISP. To identify differentially abundant bacterial sequences across aphid species we applied generalized linear mixed models to normalized read counts using DeSeq2 [v 1.26.0; 34] and made pair-wise comparisons between each aphid species. Normalized read counts account for variation in library size and were used to estimate relative abundance of each ASV. Finally, we compared strain level genetic differences in *Buchnera* and *Arsenophonus* ASVs using a phylogenetic approach by aligning sequences with the DECIPHER package [v 2.14.0; [37] and building Maximum-likelihood trees with the `optim.pml(model="GTR", rearrangement = "stochastic")` function in the phangorn package in R [v 2.5.5; 35].

Results

Overall, most bacterial taxa identified through sequencing were found to occur in all three milkweed aphid species (Fig 1a). In total 49 amplified sequence variants (ASVs) were identified (Online Resource 1 Table S3), 42 of which occurred in each species when all sampling locations were considered. Bacterial communities in general were dominated by the primary aphid symbiont *Buchnera*, which was in high relative abundance compared to the remaining taxa (Fig 1b). Among the non-*Buchnera* ASVs identified several well-known aphid facultative symbionts were found, including *Arsenophonus*, *Serratia*, and *Hamiltonella*, which occurred in all milkweed aphid species to varying degrees (Fig 1b). In a few samples *Regiella*, *Rickettsiella*, and *Wolbachia* were identified in very low abundance (<2 reads in 10% of samples) and did not make the cutoff to be included in analysis (Online Resource 2 Fig S1). Much of the remaining low-abundance taxa identified are likely environmentally acquired from the plant host or surrounding soil (e.g. *Pantoea*, *Pseudomonas*, *Streptococcus*). The prevalence or proportion of samples that tested positive for each ASV also tended to vary across aphid species (Online Resource 1 Table S3). For example, *Buchnera* was found in all samples, but *Hamiltonella* ranged from 13% (2/15) prevalence in *A. asclepiadis* samples to 100% (6/6) in *M. asclepiadis* and 30% (13/43) in *A. nerii* samples. *Serratia* was also highly prevalent across all three species, ranging from 83% in *A. nerii* to 100% in *M. asclepiadis* samples. Interestingly, all three identified strains of *Lactobacillus* were found in 100% of *M. asclepiadis*

samples but were only present in 50% or less of samples of the other two species. Co-occurring strains of *Buchnera* and *Arsenophonus* were also found in all three species, although it should be noted that we tested groups of aphids and it remains unknown whether individual aphids harbor different strains (i.e. multiple infections).

Although most ASVs were shared at a global level, several key differences indicate that each milkweed aphid species harbors distinct bacterial communities that vary in composition and abundance (Figs 1 & 2, Online Resource 1 Table S4). NMDS of Bray-Curtis dissimilarity between samples indicates that bacterial community structure is unique to each aphid species ($p < 0.001$, Online Resource 1 Table S5), but did not vary by location (Fig 2 & 3). Hierarchical clustering of samples by differences in community structure and relative abundance of each taxa further shows that each aphid species forms a unique group (Fig 3). In addition to differences in overall community structure there was significant variation in the bacterial species richness and evenness of communities found across the three aphid species (Fig 4, Online Resource 1 Table S5). Interestingly, *A. nerii* had the lowest bacterial species richness and *A. asclepiadis* had the highest among the three species. Overall, results did not show a gradient in bacterial diversity (i.e. species richness) associated with diet breadth or degree of host-plant specialization across these three milkweed aphid species.

To further identify factors contributing to differences between aphid microbiomes we compared the relative abundance of individual taxa and variation in bacterial strain diversity (ASVs) (Online Resource 1 Table S5; Figures 3 & 5). Differences in community structure were primarily driven by 1) variation in the composition of facultative symbionts and other low-abundance (non-*Buchnera*) bacteria (Figs 1b & 3) and 2) the presence of *Buchnera* and *Arsenophonus* strains unique to each aphid species (Fig 5). Overall, *A. nerii* bacterial communities were dominated by *Buchnera* with few other symbionts in low abundance, which differs from the other two aphid species. *A. asclepiadis* has a greater diversity of unique *Buchnera* strains and overall higher *Arsenophonus* abundances. Each aphid species typically harbored a single dominant *Buchnera* strain in highest relative abundance, with some co-occurring strains at lower abundances (Figs 4 & 5; Online Resource 1 Table S3). Where communities differed the most was in *Arsenophonus* strain diversity and abundance (Figs 3 & 5). For example, there were 6 distinct *Arsenophonus* strains found within *A. asclepiadis* communities that ranged between 50 – 227 fold higher relative abundance compared to *A. nerii* and *M. asclepiadis*. *A. asclepiadis* also harbored 9 distinct *Buchnera* strains across the locations sampled. In addition, *Buchnera* strains varied from 91.6% sequence similarity to several strains differing by only 1-2 base pairs (>99.2% similarity) (Online Resource 1; Table S7a). *Arsenophonus* strains were even more similar to each other, varying from 94.7 – 99.6% similarity (Online Resource 1; Table S7b).

Discussion

The prevailing view of host-associated microorganisms and their role in insect-plant coevolution is rapidly changing. Acquisition of beneficial microbes serves not only to expand the ecological niche of the host, but can add novel weaponry in the adaptive battle between plants and the herbivores that prey upon

them [3, 5, 7, 8]. Furthermore, insects not only acquire microbes from host plants and the surrounding environment, but these microbial communities are intimately linked [9, 39, 40]. Disentangling the mechanisms driving variation in herbivore microbial communities and the ecological consequences for host plant specialization is therefore of great interest. In the current study, shifts were observed in the core microbiome of aphid species exhibiting varying degrees of host specialization within the milkweed family. Aphids feeding on milkweed share most bacterial taxa (Fig 1), but each species harbors unique symbiont populations that differ in strain diversity and relative abundance (Figs 1b, 3 & 5). However, broader patterns in bacterial symbiont diversity that scale with diet breadth were not observed (Figs 3 & 4).

Host plant specificity can influence herbivore associated microbial communities on multiple levels, including causing changes in functional, taxonomic or strain level diversity or by altering the abundance of individual taxa. The microbiomes of generalists and specialist herbivores could also vary simply due to differences in the contribution of heritable versus environmentally acquired microbes, the latter being more variable and transient. However, diet may in fact generally be a poor predictor of insect bacterial community composition [41], with exceptions only in certain groups of herbivores [42]. For example, Lepidopteran larva have variable gut microbiomes that are likely shaped by the host plant they feed on, but general patterns associated with diet breadth are not observed due to the high turnover rate of these microbial communities [14]. In contrast, recent work in Costa Rican rolled-leaf beetles (*Cephaloleia* spp.) shows diet breadth is linked to microbiome diversity and community structure [21]. Previous research also indicates the composition of aphid microbial communities are structured by host plant [11, 15, 16, 20, 22, 24, 43, 44] and heritable bacterial symbionts are involved in expanding diet breadth [23, 25].

Our results show milkweed aphids differing in host plant specialization and diet breadth have similar microbial communities on a taxonomic level but differ in strain diversity and relative abundance of key symbionts (e.g. *Arsenophonus*). One possible explanation is exposure to similar nutritional and chemical profiles could homogenize microbiomes of herbivore species feeding on the same host plants. The species in this study are specialists of the *Asclepias* (milkweed) family and therefore may have similar core symbionts due to exposure to closely related host plants. Additionally, phylogenetic relatedness can generally result in closely related species harboring more similar microbiota than distantly related species [45]. The current study is limited in that aphids were collected from a single host plant (i.e. common milkweed) and generalist species with a host range outside the milkweed family (e.g. *Myzus persicae*) were not characterized due to low occurrence in the field. Additional imbalance in sampling across aphid species (e.g. fewer *M. asclepiadis* samples, Table S1) could mean that some microbial variation was missed. Further research investigating the generalist-specialist gradient using herbivores that feed across multiple plant families is needed to clarify the extent to which diet breadth shapes microbial communities (e.g. [21]).

Although milkweed aphid microbiomes were overall similar, key differences in symbiotic partnerships could contribute to additional ecological variation (e.g. ant tending, parasitism rates, predation). Interestingly, variation in symbiont relative abundance and strain diversity contributed most to differences

observed across milkweed aphid microbiomes. In particular, *Arsenophonus* was found in higher abundance in *A. asclepiadis* compared to the other two species. *Arsenophonus* is a notorious shape-shifting insect symbiont, known best for reproductive manipulation of its host [46, 47]. Most aphid facultative symbionts are found in much lower abundances compared to the obligate nutritional symbiont *Buchnera*, suggesting the unusually high abundance observed in *A. asclepiadis* could be linked to symbiont complementarity, as has occurred in other aphid species [48]. It is also possible *Arsenophonus* provides a general fitness boost, similar to what has been observed in the soybean aphid [49]. Finally, differences in symbiont populations could shape milkweed aphid-ant mutualisms, possibly via microbial induced changes to honeydew or emission of chemical compounds that mediate partner attraction. Previous work shows insect social partnerships not only uniquely influence each host's symbiotic microbiome [50], but that volatile organic compounds produced by aphid-associated microbes play a role in attracting ant mutualists [51]. Among milkweed aphids, *A. asclepiadis* is consistently tended by ants and benefits from enhanced protection from predators, while *A. nerii* is occasionally ant tended and *M. asclepiadis* appears to be a loner lacking ant friends [52]. Based on the current study an intriguing question arises; Does *Arsenophonus* mediate milkweed aphid-ant interactions and thus contribute to observed differences in ant attendance? Although *Arsenophonus* does not appear to influence the intensity of ant attendance in cowpea aphids [53], it is possible this symbiont has evolved a different ecological role in the case of *A. asclepiadis*. However, given the 16s rRNA metabarcoding approach used in this study only provides relative abundances, *Arsenophonus* titer levels will need to be confirmed using additional methods such as quantitative PCR in order to take the first step towards addressing a potential role in aphid-ant interactions.

In summary, we did not find evidence for a gradient in bacterial diversity across generalist and specialist milkweed aphids. Instead, our results suggest overlapping host plant range and shared hosts can result in selection for common microbes and thus highly similar microbiomes across species. However, milkweed aphids do harbor unique bacterial populations that most notably vary in strain diversity and relative abundance of *Arsenophonus*, although a handful of other well known aphid symbionts were also found in low abundance. These findings suggest that while host plant specialization may not be a major driver of divergence in overall taxonomic composition of aphid symbiont communities, factors such as strain level variation and differences in abundance offer alternative routes to generating adaptive potential. Further research is needed to determine the functional or ecological role played by milkweed aphid facultative symbionts and different co-occurring strains.

Declarations

Acknowledgments: We would like to thank Purdue University for providing funding through faculty start-up funds to support this research, Purdue Entomology for hosting a “Paper in a Day” writing challenge that contributed to progress on this manuscript and the Purdue Entomology Women’s Writing Group for continued support, motivation and accountability.

Funding: This work was supported by start-up funds awarded to Dr. Laramy Enders from Purdue University.

Conflicts of interest/Competing interests: The authors have no conflicts of interest to declare that are relevant to the content of this article.

Ethics approval: Not applicable, research does not involve human subjects, their data or biological material

Consent to participate: Not applicable, research does not involve human subjects, their data or biological material

Consent for publication: Not applicable, research does not involve human subjects, their data or biological material

Availability of data and material: DNA sequencing data is available on the NCBI SRA database under project number PRJNA635683.

Code availability: All code for statistical analyses and figures are available at <https://github.itap.purdue.edu/LaramyEndersGroup/Milkweed-Aphid-Microbiome>.

Authors' contributions: L.E. wrote and revised all drafts of the manuscript, conceived of and designed the study, and contributed to data collection. T.H., K.B. and J.C. contributed to data collection and sample processing, provided conceptual guidance and critical revisions/edits of manuscript drafts prior to final submission. E.F. performed all statistical analyses, contributed to data interpretation, developed figures, and provided critical revisions of manuscript drafts prior to final submission. All authors read and approved the final manuscript.

References

1. Barbosa P, Krischik VA, Jones CG (1991) Microbial mediation of plant-herbivore interactions. John Wiley & Sons
2. Berenbaum MR (1988) Allelochemicals in insect-microbe-plant interactions; agents provocateurs in the coevolutionary arms race. *Nov Asp Insect-Plant Interact* 97–123
3. Mason CJ, Jones AG, Felton GW (2019) Co-option of microbial associates by insects and their impact on plant–folivore interactions. *Plant Cell Environ* 42:1078–1086
4. Sugio A, Dubreuil G, Giron D, Simon J-C (2015) Plant–insect interactions under bacterial influence: ecological implications and underlying mechanisms. *J Exp Bot* 66:467–478
5. Hansen AK, Moran NA (2014) The impact of microbial symbionts on host plant utilization by herbivorous insects. *Mol Ecol* 23:1473–1496
6. Mendes R, Garbeva P, Raaijmakers JM (2013) The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. *FEMS Microbiol Rev* 37:634–

7. Pineda A, Zheng S-J, van Loon JJ et al (2010) Helping plants to deal with insects: the role of beneficial soil-borne microbes. *Trends Plant Sci* 15:507–514
8. Hammer TJ, Bowers MD (2015) Gut microbes may facilitate insect herbivory of chemically defended plants. *Oecologia* 179:1–14
9. Liu H, Macdonald CA, Cook J et al (2019) An ecological loop: Host microbiomes across multitrophic interactions. *Trends Ecol Evol* 34:1118–1130
10. Grunseich JM, Thompson MN, Aguirre NM, Helms AM (2020) The Role of Plant-Associated Microbes in Mediating Host-Plant Selection by Insect Herbivores. *Plants* 9:6
11. Ferrari J, Darby AC, Daniell TJ et al (2004) Linking the bacterial community in pea aphids with host-plant use and natural enemy resistance. *Ecol Entomol* 29:60–65
12. McLean AH, Parker BJ, Hrčák J et al (2016) Insect symbionts in food webs. *Philos Trans R Soc B Biol Sci* 371:20150325
13. Giron D, Dedeine F, Dubreuil G et al (2017) Influence of microbial symbionts on plant–insect interactions. In: *Advances in botanical research*. Elsevier, pp 225–257
14. Jones AG, Mason CJ, Felton GW, Hoover K (2019) Host plant and population source drive diversity of microbial gut communities in two polyphagous insects. *Sci Rep* 9:1–11
15. Xu T-T, Jiang L-Y, Chen J, Qiao G-X (2020) Host Plants Influence the Symbiont Diversity of Eriosomatinae (Hemiptera: Aphididae). *Insects* 11:217. <https://doi.org/10.3390/insects11040217>
16. Qin M, Chen J, Xu S et al Microbiota associated with Mollitrichosiphum aphids (Hemiptera: Aphididae: Greenideinae): diversity, host species specificity and phyllosymbiosis. *Environ Microbiol* n/a: <https://doi.org/https://doi.org/10.1111/1462-2920.15391>
17. Douglas AE (2013) Microbial brokers of insect-plant interactions revisited. *J Chem Ecol* 39:952–961
18. Engel P, Moran NA (2013) The gut microbiota of insects–diversity in structure and function. *FEMS Microbiol Rev* 37:699–735
19. Chung SH, Scully ED, Peiffer M et al (2017) Host plant species determines symbiotic bacterial community mediating suppression of plant defenses. *Sci Rep* 7:1–13
20. Holt JR, Styer A, White JA et al (2020) Differences in Microbiota between two Multilocus Lineages of the sugarcane Aphid (*Melanaphis sacchari*) in the continental United States. *Ann Entomol Soc Am*
21. Blankenchip CL, Michels DE, Braker HE, Goffredi SK (2018) Diet breadth and exploitation of exotic plants shift the core microbiome of tropical herbivorous beetles. *PeerJ Prepr* 6:e26692v1
22. Gauthier J-P, Outreman Y, Mieuzet L, Simon J-C (2015) Bacterial communities associated with host-adapted populations of pea aphids revealed by deep sequencing of 16S ribosomal DNA. *PloS One* 10:e0120664
23. Wagner SM, Martinez AJ, Ruan Y-M et al (2015) Facultative endosymbionts mediate dietary breadth in a polyphagous herbivore. *Funct Ecol* 29:1402–1410

24. Guidolin AS, Cônsoli FL (2017) Symbiont diversity of *Aphis (Toxoptera) citricidus* (Hemiptera: Aphididae) as influenced by host plants. *Microb Ecol* 73:201–210
25. Leonardo TE, Muiru GT (2003) Facultative symbionts are associated with host plant specialization in pea aphid populations. *Proc R Soc Lond B Biol Sci* 270:S209–S212
26. Blackman RL, Eastop VF (2008) *Aphids on the world's herbaceous plants and shrubs*, 2 volume set. John Wiley & Sons
27. Züst T, Agrawal AA (2016) Population growth and sequestration of plant toxins along a gradient of specialization in four aphid species on the common milkweed *Asclepias syriaca*. *Funct Ecol* 30:547–556
28. Zytynska SE, Weisser WW (2016) The natural occurrence of secondary bacterial symbionts in aphids. *Ecol Entomol* 41:13–26
29. Binetruy F, Dupraz M, Buysse M, Duron O (2019) Surface sterilization methods impact measures of internal microbial diversity in ticks. *Parasit Vectors* 12:268
30. Gohl DM, Vangay P, Garbe J et al (2016) Systematic improvement of amplicon marker gene methods for increased accuracy in microbiome studies. *Nat Biotechnol* 34:942–949
31. Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120
32. Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* 17:10–12
33. Callahan BJ, McMurdie PJ, Rosen MJ et al (2016) DADA2: high-resolution sample inference from Illumina amplicon data. *Nat Methods* 13:581–583
34. McMurdie PJ, Holmes S (2013) phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PloS One* 8:e61217
35. Dixon P (2003) VEGAN, a package of R functions for community ecology. *J Veg Sci* 14:927–930
36. Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* 15:550
37. Wright ES (2016) Using DECIPHER v2. 0 to analyze big biological sequence data in R. *R J* 8
38. Schliep K, Potts AA, Morrison DA, Grimm GW (2016) Intertwining phylogenetic trees and networks. *PeerJ Preprints*
39. Hannula SE, Zhu F, Heinen R, Bezemer TM (2019) Foliar-feeding insects acquire microbiomes from the soil rather than the host plant. *Nat Commun* 10:1–9
40. Gomes SI, Kielak AM, Hannula SE et al (2020) Microbiomes of a specialist caterpillar are consistent across different habitats but also resemble the local soil microbial communities. *Anim Microbiome* 2:1–12
41. Jones RT, Sanchez LG, Fierer N (2013) A cross-taxon analysis of insect-associated bacterial diversity. *PLoS One* 8:e61218

42. Colman DR, Toolson EC, Takacs-Vesbach CD (2012) Do diet and taxonomy influence insect gut bacterial communities? *Mol Ecol* 21:5124–5137
43. Henry LM, Maiden MC, Ferrari J, Godfray HCJ (2015) Insect life history and the evolution of bacterial mutualism. *Ecol Lett* 18:516–525
44. Brady CM, White JA (2013) Cowpea aphid (*Aphis craccivora*) associated with different host plants has different facultative endosymbionts. *Ecol Entomol* 38:433–437.
<https://doi.org/https://doi.org/10.1111/een.12020>
45. McLean AH, Godfray HCJ, Ellers J, Henry LM (2019) Host relatedness influences the composition of aphid microbiomes. *Environ Microbiol Rep* 11:808–816
46. Jousselin E, Cø eur, d'Acier A, Vanlerberghe-Masutti F, Duron O (2013) Evolution and diversity of *Arsenophonus* endosymbionts in aphids. *Mol Ecol* 22:260–270
47. Nováková E, Hypša V, Moran NA (2009) *Arsenophonus*, an emerging clade of intracellular symbionts with a broad host distribution. *BMC Microbiol* 9:143
48. Chong RA, Moran NA (2018) Evolutionary loss and replacement of *Buchnera*, the obligate endosymbiont of aphids. *ISME J* 12:898–908
49. Wulff JA, White JA (2015) The endosymbiont *Arsenophonus* provides a general benefit to soybean aphid (Hemiptera: Aphididae) regardless of host plant resistance (Rag). *Environ Entomol* 44:574–581
50. Ivens AB, Gadau A, Kiers ET, Kronauer DJ (2018) Can social partnerships influence the microbiome? Insights from ant farmers and their trophobiont mutualists. *Mol Ecol* 27:1898–1914
51. Fischer CY, Lognay GC, Detrain C et al (2015) Bacteria may enhance species association in an ant–aphid mutualistic relationship. *Chemoecology* 25:223–232
52. Smith RA, Mooney KA, Agrawal AA (2009) Coexistence of Three Specialist Aphids on Common Milkweed, *Asclepias syriaca*. *Ecology* 89:2187–2196
53. Katayama N, Tsuchida T, Hojo MK, Ohgushi T (2013) Aphid Genotype Determines Intensity of Ant Attendance: Do Endosymbionts and Honeydew Composition Matter? *Ann Entomol Soc Am* 106:761–770

Figures

Figure 1

A) Venn Diagram showing the extent to which amplicon sequence variants (ASVs) are shared and which are unique to each milkweed aphid species from samples collected across all locations. B) log₁₀(x+1) normalized counts of bacterial taxa found across the three milkweed aphid species. If multiple strains were present they were grouped together under one genus. Very low abundant ASV (below 1% relative abundance) were grouped together into the “Other” category, except for known aphid symbionts (i.e. Hamiltonella) and significantly differentially abundant taxa (see Online Resource 1 & 2; Table S3 & S4; Fig S2 for details). Differing letters indicate significant differential abundance across species by DESeq2, with an adjusted p = 0.05. NS = not significant

Figure 3

Hierarchical clustering of all samples based on DESeq2 normalized read counts of each bacterial taxa

Figure 4

Species richness and evenness within aphid bacterial communities were compared across aphid species. Differing letters indicate significant differences at $p < 0.05$ by Wilcoxon Rank Sum Test (Online Resource 1; Table S5a)

Figure 5

Genetic variation and differences in relative abundance of *Buchnera* and *Arsenophonus* strains (ASVs) identified in the three milkweed aphid species. All *Buchnera* and *Arsenophonus* ASVs except ASV 49 were differentially abundant (adjusted $p < 0.05$) in at least one species comparison by DESeq2 (Online Resource 1 Table S6)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [MilkweedAphidMicrobiomeSupplementalTablesFinal032021.xlsx](#)
- [OnlineResource2.Supplemental.Figures.pdf](#)