The Impact of Tumor-Associated Macrophages on Chemoresistance via Angiogenesis in Colorectal Cancer

Masatsune Shibutani (✉ fbxbj429@ybb.ne.jp)

Department of Gastroenterological Surgery, Osaka City University Graduate School of Medicine

Shigetomi Nakao

Department of Gastroenterological Surgery, Osaka City University Graduate School of Medicine

Kiyoshi Maeda

Department of Gastroenterological Surgery, Osaka City General Hospital

Hisashi Nagahara

Department of Gastroenterological Surgery, Osaka City University Graduate School of Medicine

Shinichiro Kashiwagi

Department of Breast and Endocrine Surgery, Osaka City University Graduate School of Medicine

Kosei Hirakawa

Department of Gastroenterological Surgery, Osaka City University Graduate School of Medicine

Masaichi Ohira

Department of Gastroenterological Surgery, Osaka City University Graduate School of Medicine

Research Article

Keywords: tumor-associated macrophages, angiogenesis, chemoresistance, neovascularity, colorectal cancer

Posted Date: April 22nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-403212/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Anticancer Research on September 1st, 2021. See the published version at <https://doi.org/10.21873/anticancer.15253>.

Abstract

Background: The tumor microenvironment plays an important role in tumor progression. Tumor-associated macrophages (TAMs) have been reported to promote proliferation, invasion, metastasis, angiogenesis, and immunosuppression. Furthermore, angiogenesis has been reported to induce chemoresistance due to the inefficient distribution of drugs to cancer cells. However, the impact of TAMs on chemoresistance via angiogenesis in colorectal cancer (CRC) remains unclear. The aim of this study was to evaluate the impact of TAMs on the chemotherapeutic outcome in CRC.

Methods: We enrolled 54 patients who underwent chemotherapy for unresectable metastatic CRC after resection of the primary tumor. We evaluated the density of TAMs and the degree of angiogenesis by immunohistochemistry and then explored the correlation between the density of TAMs and the chemotherapeutic outcome. Furthermore, we also assessed any correlation between the density of TAMs and that of neovascularity.

Results: The high-TAMs group had a significantly worse progression-free survival ($p=0.0006$) and a poorer response rate ($p=0.0274$) than the low-TAMs group. In addition, a positive correlation was observed between the density of TAMs and the degree of neovascularity ($r=0.665$, $p=0.0004$).

Conclusions: TAMs were revealed to promote chemoresistance via angiogenesis in CRC.

Background

The tumor microenvironment has been recognized to play a crucial role in cancer progression^{1,2}. Most tumor-infiltrating macrophages (TAMs), which are immune cells present in the cancer microenvironment, have been reported to be differentiated into the M2 type, which promotes cancer progression via proliferation, invasion, metastasis, angiogenesis and suppression of anticancer immunity^{3–7}. Furthermore, angiogenesis induced by TAMs has been reported to be associated with not only cancer progression but also resistance to chemotherapy^{8,9}. These results have been proven in basic research using experimental animal models^{7,10}, but few studies have verified whether or not similar results can be obtained in the clinical setting.

The present study examined the correlation between the density of TAMs and the chemotherapeutic outcomes and evaluated the correlation between the density of TAMs and angiogenesis in clinical samples.

Methods

Patients

- Group I

To evaluate the correlation between the density of TAMs and the chemotherapeutic outcome, we retrospectively reviewed a database of 54 patients with stage IV colorectal cancer (CRC) who underwent palliative combination chemotherapy for unresectable metastatic tumors after resection of the primary tumor at the Department of Surgical Oncology of Osaka City University between 2007 and 2014.

- Group II

In addition to the above, to compare the differences in the cancer microenvironment between the primary tumor and the metastatic tumor, we retrospectively reviewed a database of 24 patients with stage IV CRC who underwent concurrent resection of the primary tumor and the metastatic liver tumor. Patients who underwent preoperative therapy, such as chemotherapy and radiotherapy, were excluded from this study.

This retrospective study was approved by the Ethics Committee of Osaka City University (approval number: 3853) and conducted in accordance with the Declaration of Helsinki. All patients provided their written informed consent.

Immunohistochemistry

CD31 has been used as a specific marker to identify vascular endothelial cells^{11,12}, and CD163 has been used as a specific marker to identify M2 macrophages^{7,13,14}. Surgically resected specimens were retrieved to perform immunohistochemistry. Sections 4 µm in thickness were deparaffined and rehydrated. The sections were then subjected to endogenous peroxidase blocking in 1% H₂O₂ solution in methanol for 15 min. Antigen retrieval was performed by autoclaving the sections at 105°C for 10 min in Dako Target Retrieval Solution (Dako, Glostrup, Denmark). Serum blocking was performed with 10% normal rabbit serum for 10 min. After H₂O₂ and serum blocking, the slides were incubated with primary mouse monoclonal anti-CD31 antibody (1:40 dilution; Dako, Glostrup, Denmark)/anti-CD163 antibody (1:200 dilution; Leica Biosystems, Newcastle Upon Tyne, UK) at room temperature for 1 h. The secondary antibody was biotin-labeled rabbit anti-mouse IgG (1:500; Nichirei, Tokyo, Japan). Detection was performed with a DAB kit (Histofine simple stain kit; Nichirei). The sections were counterstained with hematoxylin.

The evaluation of the density of microvessels

The morphometric analysis of microvessels was carried out on the immunohistochemical stained sections using “Image-J” (National Institutes of Health, Bethesda, MD, USA), an automated imaging software program. In tumor sections, 5 randomly selected fields at the invasive margin were selected with a light microscope at a magnification of 200x. The average area and number of microvessels obtained in five different fields was used to determine the density of microvessels for the data analysis according to the methods described in previous reports^{11,15–17} (Fig. 1).

The evaluation of the density of TAMs

The number of immunoreactive macrophages at the invasive margin was counted with a light microscope in a randomly selected field at a magnification of 400x (Fig. 2). The average number obtained in five different areas was used for the data analysis.

Statistical analyses

The relapse-free survival was defined as the interval between the date of operation and the date of the diagnosis of the first recurrence, death from any cause or last follow-up. Survival curves were made using the Kaplan-Meier method. Differences in the survival curves were assessed using the log-rank test. Response evaluations by computed tomography were performed according to the Response Evaluation Criteria in Solid Tumors¹⁸. Disease control was defined as a complete or partial response or stable disease. The association between the density of TAMs and that of microvessels was evaluated by Spearman's rank correlation coefficient. Similarly, the association between the density of TAMs in the primary tumor and that in the metastatic tumor was also evaluated by Spearman's rank correlation coefficient. All of the statistical analyses were conducted using the JMP software program, ver. 13.0.0 (SAS Institute Inc., Cary, NC, USA). $p < 0.05$ was considered to indicate a statistically significant difference.

Results

Patients' characteristics

- Group I

The characteristics of the patients with stage IV CRC who underwent palliative combination chemotherapy for unresectable metastatic tumor are listed in Table 1. A total of 26 men and 28 women were included. The median age of the patients was 63 years old (range: 40–83 years old). The most common metastatic organ was the liver. Twenty-four patients (44.4%) had multiple metastatic organs. Forty-six patients (85.2%) underwent oxaliplatin-based combination chemotherapy, and 8 (14.8%) underwent irinotecan-based combination chemotherapy as the first-line chemotherapy.

Table 1
Characteristics of Group I patients

Gender (n)	
Men	26
Women	28
Age (years)	
Median (range)	63 (40–83)
Location of the primary tumor (n)	
Colon	35
Rectum	19
Tumor depth (n)	
T1-3	20
T4	34
Histological Type (n)	
Well-, Moderately differentiated	50
Poorly differentiated, Mucinous	4
Liver metastasis (n)	
Negative	15
Positive	39
Peritoneal dissemination (n)	
Negative	43
Positive	11
Lung metastasis (n)	
Negative	33
Positive	21
Number of metastatic organs (n)	
1	30
≥ 2	24

FOLFOX: 5-fluorouracil + leucovorin + oxaliplatin; CapeOX: capecitabine + oxaliplatin; FOLFIRI: 5-fluorouracil + leucovorin + irinotecan; SOX: S-1 + oxaliplatin; TAMs: tumor-associated macrophages

Gender (n)	
Regimen of first-line chemotherapy (n)	
mFOLFOX6	31
CapeOX	14
FOLFIRI	8
SOX	1
Molecular-targeted therapy (n)	
Bevacizumab	21
Cetuximab	6
Panitumumab	2
None	25
The density of TAMs (/HPF)	
Median (range)	10.4 (3.2–28.0)
FOLFOX: 5-fluorouracil + leucovorin + oxaliplatin; CapeOX: capecitabine + oxaliplatin; FOLFIRI: 5-fluorouracil + leucovorin + irinotecan; SOX: S-1 + oxaliplatin; TAMs: tumor-associated macrophages	

- Group II

The characteristics of the patients who underwent concurrent resection of the primary tumor and the metastatic liver tumor are listed in Table 2. A total of 13 men and 11 women were included. The median age of the patients was 67 years old (range: 33–78 years old). Twenty-three patients (95.8%) had well- or moderately differentiated adenocarcinoma. The median number of metastatic liver tumors was 1 (range: 1–4).

Table 2
Characteristics of Group II patients

Gender (n)	
Men	13
Women	11
Age (years)	
Median (range)	67 (33–78)
Location of the primary tumor (n)	
Colon	15
Rectum	9
Tumor depth of the primary tumor (n)	
T1-3	12
T4	12
Histological type of the primary tumor (n)	
Well-, Moderately differentiated	23
Poorly differentiated, Mucinous	1
Lymphatic involvement of the primary tumor (n)	
Negative	1
Positive	23
Venous involvement of the primary tumor (n)	
Negative	10
Positive	14
Lymph node metastasis (n)	
Negative	8
Positive	16
Number of metastatic liver tumors (n)	
Median (range)	1 (1–4)

Classification according to the density of TAMs

We used the density of TAMs, which was a continuous variable, as the test variable and the objective response rate as the state variable. A receiver operating characteristic (ROC) curve analysis of the density of TAMs suggested the appropriate cut-off value was 14.9 with an area under the curve of 0.6484 (95% confidence interval: 0.475–0.822) (Fig. 3). Thus, 14.9 was adopted as the cut-off value of the density of TAMs (sensitivity: 100%, specificity: 40.9%), and the patients were classified into the high-TAMs group and the low-TAMs group based on this value.

Correlations between the density of TAMs and the clinicopathological factors

There was no marked relationship between the density of TAMs and the clinicopathological factors (Table 3).

Table 3
Correlations between the density of TAMs and the clinicopathological factors

Factors	The density of TAMs		
	High (n = 14)	Low (n = 40)	p-value
Gender (n)			
Men	5	21	
Women	9	19	0.276
Age (years) (n)			
< 63	7	22	
≥ 63	7	18	0.747
Location of the primary tumor (n)			
Colon	7	28	
Rectum	7	12	0.183
Histological type (n)			
Well-, Moderately differentiated	14	36	
Poorly differentiated, Mucinous	0	4	0.113
Number of metastatic organs (n)			
1	7	23	
≥ 2	7	17	0.543
Peritoneal dissemination (n)			
Negative	11	32	
Positive	3	8	0.909
Molecular-targeted therapy (n)			
Without	5	20	
With	9	20	0.356
TAMs: tumor-associated macrophages			

Correlations between the density of TAMs in the primary tumor and the chemotherapeutic outcome

The distribution of the chemotherapeutic response with reference to the TAMs subgroups is shown in Table 4. A high TAMs level was associated with a significantly lower objective response rate than a low TAMs level (50.0% vs. 82.5%, $p = 0.0274$). The progression-free survival was significantly worse in the high-TAMs group than in the low-TAMs group ($p = 0.0006$) (Fig. 4).

Table 4
The distribution of the chemotherapeutic response with reference to the TAMs subgroups

Response	The density of TAMs		
	High (n = 14)	Low (n = 40)	p-value
Complete response	0	2	
Partial response	7	31	
Stable disease	3	4	
Progressive disease	4	3	
Objective response rate	50.0%	82.5%	0.0274
TAM: tumor-associated macrophage			

Correlations between the density of TAMs and that of microvessels

Regarding the primary tumor, the number of TAMs was significantly associated with the number of microvessels and the area occupied by the microvessels ($r = 0.665$, $p = 0.0004$; $r = 0.633$, $p = 0.0009$, respectively) (Fig. 5AB). Furthermore, regarding the metastatic liver tumor, the density of TAMs was also significantly associated with the number of microvessels and the area occupied by the microvessels ($r = 0.859$, $p < 0.0001$; $r = 0.592$, $p = 0.0023$, respectively) (Fig. 5CD).

Correlations between the density of TAMs in the primary tumor and that in the metastatic liver tumor

The number of TAMs in the primary tumor was significantly associated with that in the metastatic liver tumor ($r = 0.781$, $p < 0.0001$) (Fig. 6).

Discussion

As a result of this research, it became clear that high TAMs infiltration was associated with worse chemotherapeutic outcomes than low infiltration in patients with unresectable metastatic CRC.

TAMs promote angiogenesis by producing vascular endothelial growth factor (VEGF) ¹⁹. Newly formed tumor blood vessels increased by angiogenesis differ from normal microvessels in that they have structural anomalies of the vessel wall, such as large gaps between endothelial cells, defects in pericyte

coverage and discontinuous basement membranes^{20,21}. Such structural abnormalities of the vessel wall increase the hyperpermeability of the tumor vessels, resulting in poor drug delivery to cancer cells. Furthermore, inadequate drug distribution diminishes the efficacy of chemotherapy. Based on these mechanisms, the chemotherapeutic effect is expected to decrease as the degree of TAMs infiltration increases. Although this relationship has been reported in basic research, it has rarely been verified using clinical samples. The significance of the present study is that it clinically shows that TAMs, as well as cancer cells, may promote chemoresistance via angiogenesis.

In this study, we examined the correlation between the density of TAMs in the primary tumor and the therapeutic outcomes. However, the relationship between the density of TAMs in the metastatic tumor, which was the target of treatment, and the chemotherapeutic outcome has not been investigated. Therefore, we examined the relationship between the density of TAMs in the primary tumor and that in the metastatic tumor, and the strong correlation between them became clear. Based on this result, the density of TAMs in the primary tumor was revealed to be a surrogate marker of the density of TAMs in the metastatic tumor. Because it is difficult to obtain a tissue sample of metastatic tumors in clinical practice, the findings revealed in this study were extremely significant.

Several limitations associated with the present study warrant mention. First, the current study was a retrospective study with a small cohort in a single center. Second, in addition to the TAMs, there are other factors, such as cancer cells, which promote angiogenesis by producing VEGF^{22,23}. Third, in this study, the density of microvessels was measured as a surrogate marker of neovascularization, according to previous reports^{11,15–17}, when angiogenesis was evaluated. However, it was difficult to determine whether or not the microvessels we counted were normal vessels that had originally existed or neovascularity that had newly developed in cancer microenvironment. In some studies, angiogenesis was assessed by the expression of Nestin or Ki-67 in vascular endothelial cells to assess the neovascularity strictly^{24,25}.

New drugs that block the differentiation of macrophages into the M2 type are being developed at the basic research level. In the future, these agents may be able to suppress angiogenesis and enhance the efficacy of chemotherapy.

Conclusions

TAMs were revealed to promote chemoresistance via angiogenesis in CRC.

Abbreviations

TAMs: tumor-associated macrophages; CRC: colorectal cancer; ROC: receiver operating characteristic; VEGF: vascular endothelial growth factor

Declarations

Ethics approval and consent to participate

This research conformed to the provisions of the Declaration of Helsinki. All patients were informed of the investigational nature of this study and provided their written informed consent. This retrospective study was approved by the ethics committee of Osaka City University (approval No.3853).

Consent for publication

Not applicable

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

No funding was acquired for this study.

Author's contributions

MS, SN and KM designed the study, performed the statistical analysis and draft the manuscript. HN collected the clinical data and revised the manuscript critically. SK, KH and MO designed the study and critically reviewed the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This research received no specific grants from any funding agency in the public, commercial or not-for-profit sectors. We thank Brian Quinn who provided medical writing services on behalf of JMC, Ltd.

References

1. Bissell MJ, Radisky D. Putting tumours in context. *Nat Rev Cancer*. 2001; 1: 46–54.
2. Li H, Fan X, Houghton J. Tumor microenvironment: the role of the tumor stroma in cancer. *J Cell Biochem*. 2007; 101: 805–15.
3. Allavena P, Sica A, Garlanda C, Mantovani A. The Yin-Yang of tumor-associated macrophages in neoplastic progression and immune surveillance. *Immunol Rev*. 2008; 222: 155–61.
4. Polverini PJ, Leibovich SJ. Induction of neovascularization in vivo and endothelial proliferation in vitro by tumor-associated macrophages. *Lab Invest*. 1984; 51: 635–42.

5. Mussoni L, Riganti M, Acero R, et al. Macrophages associated with murine tumours express plasminogen activator activity. *International journal of cancer*. 1988; 41: 227–30.
6. Mantovani A, Sozzani S, Locati M, Allavena P, Sica A. Macrophage polarization: tumor-associated macrophages as a paradigm for polarized M2 mononuclear phagocytes. *Trends Immunol*. 2002; 23: 549–55.
7. Sica A, Schioppa T, Mantovani A, Allavena P. Tumour-associated macrophages are a distinct M2 polarised population promoting tumour progression: potential targets of anti-cancer therapy. *Eur J Cancer*. 2006; 42: 717–27.
8. Wei C, Yang C, Wang S, et al. M2 macrophages confer resistance to 5-fluorouracil in colorectal cancer through the activation of CCL22/PI3K/AKT signaling. *Onco Targets Ther*. 2019; 12: 3051–63.
9. Larionova I, Cherdyntseva N, Liu T, Patysheva M, Rakina M, Kzhyskowska J. Interaction of tumor-associated macrophages and cancer chemotherapy. *Oncoimmunology*. 2019; 8: 1596004.
10. Yu H, Kortylewski M, Pardoll D. Crosstalk between cancer and immune cells: role of STAT3 in the tumour microenvironment. *Nat Rev Immunol*. 2007; 7: 41–51.
11. Des Guetz G, Uzzan B, Nicolas P, et al. Microvessel density and VEGF expression are prognostic factors in colorectal cancer. Meta-analysis of the literature. *British journal of cancer*. 2006; 94: 1823–32.
12. Uzzan B, Nicolas P, Cucherat M, Perret GY. Microvessel density as a prognostic factor in women with breast cancer: a systematic review of the literature and meta-analysis. *Cancer research*. 2004; 64: 2941–55.
13. Shabo I, Olsson H, Elkarim R, Sun XF, Svanvik J. Macrophage Infiltration in Tumor Stroma is Related to Tumor Cell Expression of CD163 in Colorectal Cancer. *Cancer Microenviron*. 2014; 7: 61–9.
14. Komohara Y, Jinushi M, Takeya M. Clinical significance of macrophage heterogeneity in human malignant tumors. *Cancer science*. 2014; 105: 1–8.
15. Weidner N, Semple JP, Welch WR, Folkman J. Tumor angiogenesis and metastasis—correlation in invasive breast carcinoma. *The New England journal of medicine*. 1991; 324: 1–8.
16. Badawi MA, Abouelfadl DM, El-Sharkawy SL, El-Aal WE, Abbas NF. Tumor-Associated Macrophage (TAM) and Angiogenesis in Human Colon Carcinoma. *Open Access Maced J Med Sci*. 2015; 3: 209–14.
17. Marech I, Ammendola M, Sacco R, et al. Tumour-associated macrophages correlate with microvascular bed extension in colorectal cancer patients. *J Cell Mol Med*. 2016; 20: 1373–80.
18. Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer*. 2009; 45: 228–247.
19. Ruffell B, Coussens LM. Macrophages and therapeutic resistance in cancer. *Cancer Cell*. 2015; 27: 462–72.
20. Jain RK. Normalizing tumor microenvironment to treat cancer: bench to bedside to biomarkers. *J Clin Oncol*. 2013; 31: 2205–18.

21. Huang D, Lan H, Liu F, et al. Anti-angiogenesis or pro-angiogenesis for cancer treatment: focus on drug distribution. *Int J Clin Exp Med*. 2015; 8: 8369–76.

22. Dvorak HF, Sioussat TM, Brown LF, et al. Distribution of vascular permeability factor (vascular endothelial growth factor) in tumors: concentration in tumor blood vessels. *J Exp Med*. 1991; 174: 1275–8.

23. Brown LF, Berse B, Jackman RW, et al. Expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in adenocarcinomas of the gastrointestinal tract. *Cancer research*. 1993; 53: 4727–35.

24. Gravdal K, Halvorsen OJ, Haukaas SA, Akslen LA. Proliferation of immature tumor vessels is a novel marker of clinical progression in prostate cancer. *Cancer research*. 2009; 69: 4708–15.

25. Nalwoga H, Arnes JB, Stefansson IM, Wabinga H, Foulkes WD, Akslen LA. Vascular proliferation is increased in basal-like breast cancer. *Breast Cancer Res Treat*. 2011; 130: 1063–71.

Tables

Table 1. Characteristics of Group I patients

Gender (n)	
Men	26
Women	28
Age (years)	
Median (range)	63 (40-83)
Location of the primary tumor (n)	
Colon	35
Rectum	19
Tumor depth (n)	
T1-3	20
T4	34
Histological Type (n)	
Well-, Moderately differentiated	50
Poorly differentiated, Mucinous	4
Liver metastasis (n)	
Negative	15
Positive	39
Peritoneal dissemination (n)	
Negative	43
Positive	11
Lung metastasis (n)	
Negative	33
Positive	21
Number of metastatic organs (n)	
1	30
≥2	24
Regimen of first-line chemotherapy (n)	
mFOLFOX6	31
CapeOX	14

FOLFIRI	8
SOX	1
Molecular-targeted therapy (n)	
Bevacizumab	21
Cetuximab	6
Panitumumab	2
None	25
The density of TAMs (/HPF)	
Median (range)	10.4 (3.2-28.0)

FOLFOX: 5-fluorouracil+leucovorin+oxaliplatin; CapeOX: capecitabine+oxaliplatin; FOLFIRI: 5-fluorouracil+leucovorin+irinotecan; SOX: S-1+oxaliplatin; TAMs: tumor-associated macrophages

Table 2. Characteristics of Group II patients

Gender (n)	
Men	13
Women	11
Age (years)	
Median (range)	67 (33-78)
Location of the primary tumor (n)	
Colon	15
Rectum	9
Tumor depth of the primary tumor (n)	
T1-3	12
T4	12
Histological type of the primary tumor (n)	
Well-, Moderately differentiated	23
Poorly differentiated, Mucinous	1
Lymphatic involvement of the primary tumor (n)	
Negative	1
Positive	23
Venous involvement of the primary tumor (n)	
Negative	10
Positive	14
Lymph node metastasis (n)	
Negative	8
Positive	16
Number of metastatic liver tumors (n)	
Median (range)	1 (1-4)

Table 3. Correlations between the density of TAMs and the clinicopathological factors

Factors	The density of TAMs		
	High (n=14)	Low (n=40)	p-value
Gender (n)			
Men	5	21	0.276
Women	9	19	
Age (years) (n)			
<63	7	22	0.747
≥63	7	18	
Location of the primary tumor (n)			
Colon	7	28	0.183
Rectum	7	12	
Histological type (n)			
Well-, Moderately differentiated	14	36	0.113
Poorly differentiated, Mucinous	0	4	
Number of metastatic organs (n)			
1	7	23	0.543
≥2	7	17	
Peritoneal dissemination (n)			
Negative	11	32	0.909
Positive	3	8	
Molecular-targeted therapy (n)			
Without	5	20	0.356
With	9	20	

TAMs: tumor-associated macrophages

Table 4. The distribution of the chemotherapeutic response with reference to the TAMs subgroups

Response	The density of TAMs		
	High (n=14)	Low (n=40)	p-value
Complete response	0	2	
Partial response	7	31	
Stable disease	3	4	
Progressive disease	4	3	
Objective response rate	50.0%	82.5%	0.0274

TAM: tumor-associated macrophage

Figures

Figure 1

The immunohistochemical detection of CD31, a vascular endothelial cell-specific marker (x200) (A). The area filled with green color is the vessel lumen (B).

Figure 2

The immunohistochemical detection of CD163, an M2 macrophage-specific marker (x400).

Figure 3

A receiver operating characteristic curve analysis of the CAR. Area under curve=0.6484, 95% confidence interval=0.475-0.822, $p=0.116$.

Figure 4

The Kaplan-Meier survival curves for the progression-free survival according to the density of tumor-associated macrophages (TAMs). The high-TAMs group was associated with a poorer prognosis than the low-TAMs group with regard to the progression-free survival ($p=0.0006$).

Figure 5

The correlation between the density of tumor-associated macrophages (TAMs) and that of microvessels. (A) The correlation between the number of TAMs in the primary tumor and the number of microvessels in the primary tumor. (B) The correlation between the number of TAMs in the primary tumor and the area occupied by microvessels in the primary tumor. (C) The correlation between the number of TAMs in the metastatic liver tumor and the number of microvessels in the metastatic liver tumor. (D) The correlation between the number of TAMs in the metastatic liver tumor and the area occupied by microvessels in the metastatic liver tumor.

Figure 6

The correlation between the density of tumor-associated macrophages (TAMs) in the primary tumor and that in the metastatic liver tumor. A significant correlation was observed ($r=0.781$, $p<0.0001$).