

First microsatellite markers for the European Robin (*Erithacus rubecula*) and their application in analysis of parentage and genetic diversity

Aleksandra Gwiazdowska

Museum and Institute of Zoology

Oliwia Karpńska

Warsaw University of Life Sciences

Katarzyna Kamionka-Kanclerska

Warsaw University of Life Sciences

Patryk Rowiński

Warsaw University of Life Sciences

Hanna Panagiotopoulou

Museum and Institute of Zoology

Jan J. Pomorski

Museum and Institute of Zoology

Richard K. Broughton

UK Centre for Ecology & Hydrology

Luis F. P. da Silva

CBIO-InBIO Campus Agrário de Vairão Rua Padre Armando Quintas

Robert Rutkowski (✉ robertrut@miiz.waw.pl)

Museum and Institute of Zoology

Research Article

Keywords: European Robin (*Erithacus rubecula*), genetic diversity, genetic markers, population genetics

Posted Date: April 14th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-404739/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

The European Robin is a small passerine bird associated with woodlands of Eurasia and North Africa. Despite being relatively widespread and common, little is known of the species' breeding biology and genetic diversity. We used Next Generation Sequencing (NGS) to develop and characterize microsatellite markers for the European Robin, designing three multiplex panels to amplify 14 microsatellite loci. The level of polymorphism and its value for assessing parentage and genetic structure was estimated based on 119 individuals, including seven full families and 69 unrelated individuals from Poland's Białowieża Primaeval Forest and an additional location in Portugal. All markers appeared to be highly variable. Analysis at the family level confirmed a Mendelian manner of inheritance in the investigated loci. Genetic data also revealed evidence for extra-pair paternity in one family. The set of markers that we developed are proven to be valuable for analysis of the breeding biology and population genetics of the European Robin.

Introduction

The European Robin *Erithacus rubecula*, a small (c.19 g) passerine bird, is a common and widespread species of wooded habitats in Europe, parts of western Asia and North Africa¹. Although it is a familiar and moderately well-known species in Western Europe, there is little information about the species' breeding biology across much of its range, and studies of its population genetics are mostly lacking²⁻⁹. This absence of genetic studies is probably related to the European Robin's elusiveness during the breeding season, when nests are notoriously difficult to locate¹.

Most of the available fragmentary data on the breeding biology of European Robins comes from highly transformed environments, such as gardens and secondary woodlands in Britain⁴. However, these studies may not be representative of the European Robin's ecology in more natural forests elsewhere in its range. This study bias and difficulty of observing breeding European Robins means that there is no information characterising the species' typical patterns of behaviour and breeding biology in primaeval forest habitats, to which it was originally adapted¹⁰.

The European Robin is considered to be socially monogamous and highly territorial in the breeding season, when pairs may have up to three consecutive breeding attempts between spring and early summer. Nests are typically located in recesses or cavities on standing or fallen trees within several meters of the ground, in banks or walls, or under overhanging vegetation on the ground itself; nests generally contain 4-5 eggs that are incubated for around 15 days before hatching altricial nestlings that leave after a further 15 days, becoming independent around 10 days later^{1,4,5,11,12}. Despite this basic knowledge there is no information on genetic relatedness or extra-pair paternity within or between broods and their parents.

Molecular methods have made an invaluable contribution to the study of the breeding biology and mating systems of birds¹³⁻¹⁷. These techniques have revealed that extra-partner mating is a much more frequent phenomenon among birds than was previously thought^{14,16,18}, and extra-pair nestlings occur frequently even among socially monogamous species¹⁴. Such studies are important in advancing the understanding of female mate choice and reproductive fitness^{17,19,20}. The benefits to females of adopting the extra-pair mating strategy are still debated, and there is no certain evidence that the advantages of such breeding activity outweigh the costs. Conversely, males obviously benefit from this behaviour through the increased reproductive success while engaging in the care of nestlings¹⁷.

Only three studies have applied molecular techniques to describe population genetics of the European Robin. The genetic structure of a resident population from the Canary Islands, which migrate short distances between breeding and wintering grounds, was characterized using polymorphism of mitochondrial DNA^{21,22}. This study revealed that birds from Gran Canaria island should be considered as a distinct subspecies that is genetically separated from birds on continental Europe and the western Canary Islands. Another resident population, from southern Italy, was investigated using cross-amplified microsatellites²³, which suggested significant genetic differentiation between populations with long- and short-distance migratory behaviour.

The use of cross-amplified microsatellites in genetic studies is known to have several potential disadvantages, including particular markers being less polymorphic in target species than in the species of origin²⁴⁻²⁶. Also, diversification of the species may lead to mutations in sequences of flanking regions of some alleles, preventing the primer annealing during PCR^{27,28}. This, in turn, may result in a high frequency of 'null alleles'. Moreover, the success rate of amplification frequently decreases in proportion with the genetic distance between species²⁹.

However, to date, there appears to have been no species-specific microsatellite markers described for the European Robin, including for the Central European population that represents the nominative and highly-migratory subspecies. Furthermore, this knowledge gap compounds the lack of data, supported by genetic information, for the diversity, ecology and biology of the European Robin's populations in natural forest ecosystems across Europe. Such studies from primary forest conditions are informative and highly valuable as a benchmark for understanding a species' biology in modified ecosystems and more outlying populations in the species' range.

In this study we provide the first genetic data for European Robins from the last significant expanse of European temperate primaeval forest, the Białowieża Forest in north-eastern Poland, and from a second novel location on the western margin of the species' European range, in Portugal. Information from the Białowieża Forest provides a unique opportunity to extend our knowledge of the European Robin's biology in the natural, unmodified forest that still retains an almost complete suite of flora and fauna³⁰. Such data provides an important reference system for further studies of evolutionary and ecological processes in this forest, in addition to those occurring in managed forests under strong anthropogenic pressure.

The main aim of the study is to describe the species-specific microsatellite markers for the European Robin, as a resource for population genetics and breeding biology studies. We then apply these molecular markers to investigate the paternity of multiple broods and individuals of European Robins from the Białowieża Forest, and additional samples from individuals resident approximately 2730 km away in Portugal, to estimate genetic diversity of the species across this geographical span.

Materials And Methods

This study was carried out in compliance with the ARRIVE guidelines. All methods were performed in accordance with the relevant guidelines and regulations. The methods used for animal care and handling were approved by 2nd Ethical Committee for Animal Experimentation at Warsaw University of Life Sciences, Poland (decision number: WAW2/93/2017, dated on 15.12.2017) and Portuguese Institute for Conservation, Nature and Forests (ICFN, decision number: 347/2020/CAPT).

Sample Collection And DNA Extraction

Samples of genetic material from European Robins were collected in two widely separated areas of Europe, in order to isolate microsatellite markers and investigate relatedness. In north-east Poland, birds were sampled between 2018–2019 in the Strict Reserve of Białowieża National Park (52° 46' 0" N, 23° 52' 0" E), which is characterized by features typical of pristine forest, namely a multi-storey profile of tree stands, abundant dead wood with many uprooted trees, tall tree heights and a diverse tree community of deciduous and coniferous species¹⁰. Genetic material was collected on three sample plots (about 30 ha each) at representative locations in the forest.

European Robins are summer migrants to the Białowieża forest, arriving in early spring in order to breed, after wintering in southern or western Europe. During the breeding season (late March to July), 95 individuals from the Białowieża forest were sampled for genetic analysis. Adult birds ($N=19$: 11 males, 8 females) were caught using mist-nets (fine sheets of netting suspended between poles) and a recording of songs and calls as a lure. Further samples of a single chick (8–10 days old) from each of 28 nests were also collected. These combined samples enabled genetic analysis for 47 presumably unrelated birds. Additionally, we collected samples from seven 'full' families (social father and mother plus all the nestlings) to verify Mendelian inheritance of the isolated microsatellites. These seven 'full families' included 14 of the 19 adult birds reported above. The material for genetic analysis was collected during standard capture and marking (with numbered metal leg-rings) procedures for adult birds, and during nest inspections for nestlings, and involved plucking a single tail feather from each individual. Samples were stored in Eppendorf tubes filled with 96 % pure ethyl alcohol.

In Portugal, where the breeding population of European Robins is resident all year round³¹, 24 samples of adult birds were collected from between April and June 2020 at five locations: (A) Fradelos (41° 21' 58" N, 8° 35' 55." W); (B) Azurara (41° 21' 02" N, 8° 43' 50" W); (C) Arvore (41° 19' 45" N, 8° 43' 59" W); (D) Larça (40° 18' 46" N, 8° 24' 13" W); and (E) Tapada de Mafra (38° 57' 40" N, 9° 17' 39" W).

Fradelos, Azurara and Arvore consist of mixed plantations of *Pinus pinaster* and *Eucalyptus globulus* with low management and an understory composed mostly of small native shrubs. Larça is a secondary native forest (tree height usually lower than 10 m) and a dense and diverse understory. These four locations are integrated in a landscape mainly composed by small forest patches, intertwined with agriculture and urbanized areas. Finally, Tapada de Mafra is a fenced property of more than 800 ha, managed for eco-tourism and big-game hunting with a mosaic of open areas and diverse forests dominated by *Quercus* spp. The birds were captured with mist nets placed along dirt tracks or with spring traps baited with mealworms (*Tenebrio molitor*). A blood sample was collected from each bird by puncturing the brachial vein near the joint between the humerus and ulna with a sterile needle (27G), following standard methods³². Approximately 75 µl of blood was collected with a capillary tube and then transferred to a 2 ml tube filled with 98% ethanol, which was stored at 4°C.

For the feather samples, the main source of DNA is cells attached to the base of the calamus (quill) when feathers are in the growth phase^{16,24}, or blood clots within the rachis of fully grown feathers^{16,33}. Hence, approximately 0.5 cm of the calamus was extracted and cut into smaller pieces using a sterile scalpel on sterile Petri dish. Cut parts of the calamus were then incubated with a Proteinase K overnight at 56°C.

For samples from Portugal, blood clots were removed from the alcohol and put into the buffer LT (Eurx Polska) and incubated with a Proteinase K overnight at 56°C. We then used a Tissue DNA Purification Kit (Eurx Polska) to extract DNA from the calamus and blood following standard protocols. Each extract was analysed using Nanodrop (Thermo Fisher Scientific) to check the purity and concentration of DNA. The extracted DNA was stored at -20°C.

Microsatellites Typing With Pacific Biosciences RSII

An amount of 7.4 µg of genomic DNA consisting of four equally (20 µl) mixed DNA samples was used for a 2 kb library construction. DNA was fragmented, using Covaris® System according to protocol, into sizes between 1.5 and 3 kb, applying the following modifications: intensity was set to 0.2, and time of sonication to 500 s. Fragmented DNA was purified using AMPure® PB Beads applying 0.6x volume ratio.

Subsequently, 750 ng of shared and concentrated DNA was used for library preparation with SMRTbell Template Prep Kit 1.0, applying the online Pacific Biosciences® protocol (PacBio). This procedure consisted of: 1. DNA damage and ends repair, 2. blunt ligation of hairpin adaptors at both ends of the DNA fragments, and 3. removal of failed ligation products with *ExoIII* and *ExoVII* enzymes. To separate the enzymatic reactions, DNA was purified with 0.6x volume ratio of AMPure® PB Beads. Size distribution of DNA fragments and its integrity was monitored on 0.5% agarose gels during and after completing the procedure of library preparation. The final library was double purified with 0.6x volume ratio of AMPure® PB Beads. Binding Calculator v.2.3.1.1 (PacBio) was applied for protocol generation in order to prepare the library for sequencing applying the DNA/Polymerase Binding Kit P6 V2 and MagBeads reagents. Sequencing was performed on a PacBio RS II sequencer running three SMRT Cells® in the 360-minute data collection mode.

Data generated through the PacBio sequencing platform was analysed to obtain Reads of Insert (ROI) according to the RS_ReadsofInsert.1 protocol applying default settings. The ROI reads were further analysed with the msatcommander v.1.08³⁴ program in order to identify microsatellite sequences and design primers. A threshold of at least five repetitions for tri- and tetra-nucleotide repeats, excluding mononucleotide repeats, was set in this analysis. Primer design parameters including the msatcommander option "combine loci" were as follows: length of the primers 18–22 bp, annealing temperature (Tm) 58°C–62°C, GC

content 30–70%, and amplicon product size 80–400 bp with minimum of 20–30 bp DNA sequence flanking from both sides of the tandem repeat of the microsatellite.

Microsatellite Marker Selection And Testing

Selection of microsatellite markers was performed in several steps. Firstly, based on information obtained for the set of microsatellite sequences, identified in the genome of the European Robin from the above analyses, loci with the highest number of tandem repeats were selected (total number of 281 sequences). In the second step, loci were filtered based on criteria calculated in silico: (i) maximal PCR efficiency, (ii) minimal penalty score, (iii) no dimer formation, (iv) a maximal difference in melting temperature of 2°C between a primer pair, and (v) sequence length of the microsatellite ranging from 100 to 350 bp. This gave 40 selected microsatellite loci, which were then divided based on product size into three groups: (1) short < 200 bp, (2) medium 200–300 bp, and (3) long > 300 bp. The reason for this division was optimization of designed multiplex PCR.

Finally, for further testing, we selected 11 short loci, 14 medium loci and 15 long loci. These loci were initially tested on 17 unrelated individuals following the method of Schuelke (2000)³⁵ and Austin et. al. (2011)³⁶. Briefly, the forward primers were tailed by a 5'-CACGACGTTGTAAAACGAC sequence tag, which was identical to four fluorescent labelled dyes: 6-Fam, Hex, Tamra and Rox. The PCR reaction mix contained: 1 µl of DNA, 7.5 µl of 2x Dream Taq Hot Start Green PCR Master Mix (Thermo Fisher Scientific), 0.02 M tagged forward locus specific primer, 0.2 M locus specific reverse primer and 0.2 M fluorescent labelled universal forward primers. The reaction was performed under the following conditions: initial denaturation for 15 min at 95°C, 35 cycles of 30 s at 95°C, 50 s at 57°C, 45 s at 72°C and final elongation of 5 min at 72°C. Microsatellite loci were amplified separately and visualised on 1.5% agarose gel. The genotyping analyses was performed using ABI 3500XL Genetic Analyzer (Applied Biosystems) and the PCR product sizes were read using GENEMAPPER v.4.1 software (Applied Biosystems). Based on the results obtained, 17 microsatellite loci were selected for further analysis and design of multiplex panels. These loci were characterized by a lack of amplification artefacts, a high number of alleles and no more than two peaks per locus on chromatograms.

The multiplex PCR reaction mix contained: 1 µl of DNA extract (containing 47–283 ng of DNA in the case of young birds) or 3 µl of DNA extract (containing 1–78 ng of DNA in the case of adult birds), 7.5 µl of PCR Master Mix (QIAGEN), 0.6 µl of the primer mix (forward and reverse for each locus; concentration: 0.85–1 µM) and 4.65 µl or 2.65 µl water for PCR depending on the amount of DNA extract used. This time forward primers were fluorescently labelled on their 5' end with one of the following dyes: 6-FAM, TAMRA, HEX and ROX (Sigma-Aldrich). The amplification conditions were as follows: initial denaturation for 15 min at 95°C, 40 cycles of 30 s at 94°C, 90 s at 57°C, 90 s at 72°C and final elongation for 10 min at 72°C.

Analysis of the amplification of the microsatellite loci in different combinations of multiplex sets was verified using ABI 3500XL Genetic Analyzer. As a size standard we used GeneScan™ 600 LIZ (Applied Biosystems). The genotyping was performed using a mixture of formamide with LIZ-600 and PCR product, which was denatured for 5 min. at 95°C. Genotypes were then identified using GENEMAPPER v.4.1.

Statistical analysis

All statistical analyses were performed on genotypes of 69 presumably unrelated individuals (45 from Poland and 24 from Portugal). For each locus we assessed the number of alleles (A), unbiased expected heterozygosity (H_E) and observed heterozygosity (H_O) using GenAEx version 6.5³⁷ and Hardy-Weinberg equilibrium (HWE), and also Genepop on the Web version 4^{38,39}. Additionally, the fixation index (F_{IS}) and F_{ST} were calculated using FSTAT version 2.9.3⁴⁰.

Parameters of usefulness of the selected microsatellites in paternity verification and population genetics studies were also estimated, using Cervus 3.0⁴¹. Specifically, we calculated the PIC index (polymorphic information content), NE-1P (average non-exclusion probability for the first parent), NE-2P (average non-exclusion probability for the second parent), NE-PP (average non-exclusion probability for a candidate parent pair), NE-I (average non-exclusion probability for identity of two unrelated individuals) and NE-SI (average non-exclusion probability for identity of two siblings). Probability of identity (PI) and probability of identity for siblings (PI_{Sibs}) were estimated using GenAEx version 6.5³⁷.

Results

RSII sequencing of the SMRT library generated 4,330 Mb of raw data covering 192,298 reads for the three SMRT Cells® analysed jointly. The average length of inserts in these runs equalled 1,959, 1,873 and 2,997 bp, respectively (with quality ranging between 0.91 and 0.96). The final output length of reads reached 3,800 Mb. Of these sequence sets, 65,162 reads of insert (ROI) were acquired and accumulated with 87,123,089 bp. The average ROI length equalled 1,337 bp and the average number of passes exceeded 16.

The final sequence list consisted of 62,872 DNA sequences that were subjected to microsatellite loci identification. Altogether 1,932 microsatellite loci covering tri- or tetra-nucleotide repeats in at least five repetitions were identified. For the *a priori* chosen parameters of microsatellite selection for amplification testing, primer pairs for 961 loci were successfully designed. From this set, 18 tri-nucleotide and 22 tetra-nucleotide markers (named ER1-ER40), with the number of repeats ranging between 37 and 10, were further chosen for amplification and polymorphism description (Supplementary Table S1) and then deposited in GenBank under accession numbers xxx–xxx.

Agarose gel electrophoresis indicated that 36 out of 40 tested primer pairs (90%) were successfully amplified. The amplification was evaluated as successful if the length of the band was similar to the size of alleles indicated by mstacommander. Genotyping using ABI 3500XL Genetic Analyzer (Applied Biosystems) showed that 23 primer pairs amplified microsatellite fragments, meaning that the peaks on chromatograms were sharp and the score of the peaks was similar to the expected product size. Finally, 17 primer pairs were selected for designing multiplex PCR. In the case of three primer pairs, amplification failed in multiplex reaction, despite applying several combinations. Hence, we finally designed three multiplex panels (Table 1), consisting of 14 primer pairs, as follows: Mix 1 (ER24, ER40, ER28, ER21, ER38, ER15), Mix 2 (ER32, ER7, ER4, ER5) and Mix 3 (ER39, ER13, ER25, ER31).

Table 1

The characterization of three multiplex PCR sets for the amplification of 14 microsatellite loci in 69 unrelated samples of European Robin. Column headings are as follows: Size range – size of fragments obtained during PCR, including flanking region; A – number of alleles; H_0 – heterozygosity observed; H_E – heterozygosity expected; F_{IS} – fixation index (* – F_{IS} value significant after Bonferroni correction, 200 randomizations, adjusted P -value = 0.00357), HWE – P -value of Hardy-Weinberg equilibrium exact test; PIC – polymorphic information content. Annealing temperature for all PCRs was 57°C

Locus	Primer sequence (5'-3')	Dye	Size range	A	H_0	H_E	F_{IS}	HWE
ER24	F: TGGTTGTCACCTTTGCACCTGG R: CTGCTTCACTCCTTGCACAG	TAM	128–185	16	0.594	0.696	0.154*	0.0104
ER40	F: GATTGCAGGGAACGCTGAC R: CAGGTGCTGCTGGAATGTTC	FAM	190–217	10	0.696	0.799	0.136	0.4742
ER28	F: TGGTTAATGAAGATGACGGCG R: AAGATCCCAGCCTCATTCCG	HEX	214–283	14	0.826	0.864	0.052	0.1114
ER21	F: ATGGATGCTGTGGGAGTCTC R: GTGATCACGACAGTCAGCAC	ROX	225–273	13	0.913	0.889	-0.020	0.4540
ER38	F: CGCCTGTGTAAATCTCGCTG R: TTGTTCCAACCTAGCCTGGG	FAM	283–358	12	0.797	0.799	0.010	0.3520
ER15	F: TACCTCCTGCTTGCTGTGAG R: TCCAACCAAGTCTCCAAGG	HEX	299–415	21	0.913	0.908	0.002	0.7325
ER32	F: CTCACCTGTGCTCTTAAGGC R: TGCAGTTCTTCAGTTTGGTGG	TAM	106–139	9	0.638	0.730	0.133	0.4697
ER7	F: CAACTGGTCAGCAACTCC R: TTGCAAGCTCAGGATCCTTG	ROX	262–294	10	0.899	0.851	-0.048	0.7983
ER4	F: TCAGAGGTGTGGCAGAAAGG R: TTGTTCTGCTGCTTTGGAC	FAM	204–292	17	0.696	0.850	0.188	0.0675
ER5	F: ACTCCCTACACTTGCCAAGG R: GTCTAAAGATGCCAGAACCTG	FAM	308–428	14	0.826	0.842	0.026	0.7992
ER39	F: CGGTGAACAAGAGCAGAAGG R: CCTTCTTCACTGCAAGCTGG	TAM	144–216	22	0.913	0.923	0.018	0.5100
ER13	F: AGTCTCCTTGCTGCTCTGTG R: TGCCCTTCTCTGACATGGG	ROX	180–256	19	0.913	0.885	-0.025	0.2387
ER25	F: GAAAGTATCGTGCAATGCAC R: TTCAGGTTGACAAAGAGGC	HEX	272–332	19	0.896	0.914	0.028	0.3522
ER31	F: GAAATGGGAAGTGTGCTGGC R: CTGCAGGCTTACTTCAACCAG	FAM	298–344	26	0.754	0.935	0.201*	0.0000

We failed to amplify loci ER25 in three samples, but the remaining 13 loci were amplified in all 119 samples. The resulting characterization of microsatellite polymorphism in 69 unrelated individuals of the European Robin indicated that the number of alleles at a single locus ranged from nine (locus ER32) to 26 (locus ER31).

The highest observed and expected values of heterozygosity were found in loci ER21, ER15, ER39, ER13 ($H_0 = 0.913$;) and ER31 ($H_E = 0.935$). The lowest value of heterozygosity was observed in locus ER24 ($H_0 = 0.594$; $H_E = 0.696$) (Table 1). Two loci (ER24 and ER31) significantly deviated from HWE (Table 1). Accordingly, in these loci, F_{IS} was significantly greater than zero, suggesting heterozygote deficiency. Overall, the observed heterozygosity was high ($H_0 = 0.805$) and significantly lower than the expected heterozygosity ($H_E = 0.849$, $P < 0.001$).

Considering the two separate locations, analysis of the 45 individuals from the population from Poland also indicated significant deviation from Hardy-Weinberg Equilibrium due to heterozygote deficiency ($H_0 = 0.819$; $H_E = 0.846$, $P < 0.002$), although F_{IS} was low and non-significant after Bonferroni correction ($F_{IS} = 0.04$; adjusted P -value = 0.004, 280 permutations). Similarly, in the group of samples from Portugal the observed heterozygosity was lower than expected

($H_0 = 0.780$; $H_E = 0.820$, $P = 0.063$), with an associated low value of F_{IS} ($F_{IS} = 0.07$). The mean number of alleles was higher in Poland (mean $A = 13.64$) than in Portugal (mean $A = 10.7$), although allelic richness was similar in both regions (Portugal $R = 10.71$; Poland $R = 11.60$).

Table 2

Informativeness and exclusion probability of analyzed loci, estimated based on genotypes of 69 unrelated individuals of European Robin from Poland and Portugal. PIC – polymorphic information content, PI – probability of identity; PI_{Sibs} – probability of identity for sibs; paternity non-exclusion probability (first second and pair parent, NE-1P, NE-2P and NE-PP respectively) and the combined non-exclusion probability of identity and sibling identity (NE-I and NE-SI) for each marker; overall – values for combination of 14 loci.

Locus	PIC	PI	PI_{Sibs}	NE-1P	NE-2P	NE-PP	NE-I	NE-SI
ER24	0.675	0.11	0.43	0.678	0.489	0.272	0.114	0.430
ER40	0.773	0.066	0.37	0.560	0.383	0.196	0.066	0.367
ER28	0.850	0.032	0.33	0.426	0.269	0.107	0.032	0.326
ER21	0.879	0.022	0.31	0.367	0.223	0.078	0.022	0.311
ER38	0.772	0.068	0.37	0.563	0.387	0.202	0.068	0.367
ER15	0.901	0.016	0.3	0.313	0.185	0.055	0.016	0.300
ER32	0.695	0.11	0.41	0.663	0.482	0.287	0.107	0.412
ER7	0.834	0.040	0.33	0.463	0.299	0.131	0.040	0.334
ER4	0.834	0.039	0.33	0.456	0.294	0.124	0.039	0.335
ER5	0.823	0.045	0.34	0.484	0.316	0.146	0.045	0.340
ER39	0.918	0.011	0.29	0.269	0.156	0.040	0.011	0.291
ER13	0.874	0.023	0.31	0.373	0.229	0.080	0.023	0.314
ER25	0.908	0.014	0.30	0.295	0.173	0.049	0.014	0.296
ER31	0.931	0.0079	0.28	0.231	0.131	0.029	0.008	0.285
Overall	0.8334	1,2E-21	2,1E-07	0.00000486	8.737E-0009	1.513E-0014	1.175E-0021	0.00000021

Genetic differentiation, assessed as F_{ST} , was low between both regions ($F_{ST} = 0.015$) but this was statistically significant ($P < 0.05$).

Parameters interlinked with the usefulness of identified loci in the parentage and relatedness analysis indicated that the described microsatellite panels can be successfully used in such studies. The polymorphic information content was high in all loci ($PIC > 0.5$; Table 2), and, accordingly, the analysis of 69 unrelated individuals showed a very low non-exclusion probability for the combination of 14 loci. The probability of identity and probability of sibling identity was the lowest in locus ER31 ($PI = 0.01$; $PI_{Sibs} = 0.28$) and the highest were in loci ER24 ($PI = 0.11$; $PI_{Sibs} = 0.43$) and ER32 ($PI = 0.11$; $PI_{Sibs} = 0.41$). The respective values for the combination of 14 loci were also very low (Table 2).

Paternity analysis in the seven families (with known social parents) indicated extra-pair paternity in only one case (Table 3, family WE8). In two of the nestlings, we didn't find the putative father's alleles in ten out of 14 investigated loci. Moreover, analysis of full families confirmed that all microsatellite loci had Mendelian inheritance (Table 3).

Table 3

Microsatellite genotypes of members of three full families of the European Robin. Example of families MO6 and WA6 confirm Mendelian manner of inheritance and paternity within the brood. Genotypes shown in bold suggest genetic contribution of an 'extra' male. F – female, M – son

MIX	MIX V		MIX IV																	M																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																														
Primer	ER24	ER40	ER28	ER21	ER38	ER15	ER32	ER7	ER4	ER5	ER6	ER11	ER12	ER13	ER14	ER16	ER17	ER18	ER19	ER20	ER22	ER23	ER25	ER26	ER27	ER29	ER30	ER31	ER33	ER34	ER35	ER36	ER37	ER39	ER41	ER42	ER43	ER44	ER45	ER46	ER47	ER48	ER49	ER50	ER51	ER52	ER53	ER54	ER55	ER56	ER57	ER58	ER59	ER60	ER61	ER62	ER63	ER64	ER65	ER66	ER67	ER68	ER69	ER70	ER71	ER72	ER73	ER74	ER75	ER76	ER77	ER78	ER79	ER80	ER81	ER82	ER83	ER84	ER85	ER86	ER87	ER88	ER89	ER90	ER91	ER92	ER93	ER94	ER95	ER96	ER97	ER98	ER99	ER100	ER101	ER102	ER103	ER104	ER105	ER106	ER107	ER108	ER109	ER110	ER111	ER112	ER113	ER114	ER115	ER116	ER117	ER118	ER119	ER120	ER121	ER122	ER123	ER124	ER125	ER126	ER127	ER128	ER129	ER130	ER131	ER132	ER133	ER134	ER135	ER136	ER137	ER138	ER139	ER140	ER141	ER142	ER143	ER144	ER145	ER146	ER147	ER148	ER149	ER150	ER151	ER152	ER153	ER154	ER155	ER156	ER157	ER158	ER159	ER160	ER161	ER162	ER163	ER164	ER165	ER166	ER167	ER168	ER169	ER170	ER171	ER172	ER173	ER174	ER175	ER176	ER177	ER178	ER179	ER180	ER181	ER182	ER183	ER184	ER185	ER186	ER187	ER188	ER189	ER190	ER191	ER192	ER193	ER194	ER195	ER196	ER197	ER198	ER199	ER200	ER201	ER202	ER203	ER204	ER205	ER206	ER207	ER208	ER209	ER210	ER211	ER212	ER213	ER214	ER215	ER216	ER217	ER218	ER219	ER220	ER221	ER222	ER223	ER224	ER225	ER226	ER227	ER228	ER229	ER230	ER231	ER232	ER233	ER234	ER235	ER236	ER237	ER238	ER239	ER240	ER241	ER242	ER243	ER244	ER245	ER246	ER247	ER248	ER249	ER250	ER251	ER252	ER253	ER254	ER255	ER256	ER257	ER258	ER259	ER260	ER261	ER262	ER263	ER264	ER265	ER266	ER267	ER268	ER269	ER270	ER271	ER272	ER273	ER274	ER275	ER276	ER277	ER278	ER279	ER280	ER281	ER282	ER283	ER284	ER285	ER286	ER287	ER288	ER289	ER290	ER291	ER292	ER293	ER294	ER295	ER296	ER297	ER298	ER299	ER300	ER301	ER302	ER303	ER304	ER305	ER306	ER307	ER308	ER309	ER310	ER311	ER312	ER313	ER314	ER315	ER316	ER317	ER318	ER319	ER320	ER321	ER322	ER323	ER324	ER325	ER326	ER327	ER328	ER329	ER330	ER331	ER332	ER333	ER334	ER335	ER336	ER337	ER338	ER339	ER340	ER341	ER342	ER343	ER344	ER345	ER346	ER347	ER348	ER349	ER350	ER351	ER352	ER353	ER354	ER355	ER356	ER357	ER358	ER359	ER360	ER361	ER362	ER363	ER364	ER365	ER366	ER367	ER368	ER369	ER370	ER371	ER372	ER373	ER374	ER375	ER376	ER377	ER378	ER379	ER380	ER381	ER382	ER383	ER384	ER385	ER386	ER387	ER388	ER389	ER390	ER391	ER392	ER393	ER394	ER395	ER396	ER397	ER398	ER399	ER400	ER401	ER402	ER403	ER404	ER405	ER406	ER407	ER408	ER409	ER410	ER411	ER412	ER413	ER414	ER415	ER416	ER417	ER418	ER419	ER420	ER421	ER422	ER423	ER424	ER425	ER426	ER427	ER428	ER429	ER430	ER431	ER432	ER433	ER434	ER435	ER436	ER437	ER438	ER439	ER440	ER441	ER442	ER443	ER444	ER445	ER446	ER447	ER448	ER449	ER450	ER451	ER452	ER453	ER454	ER455	ER456	ER457	ER458	ER459	ER460	ER461	ER462	ER463	ER464	ER465	ER466	ER467	ER468	ER469	ER470	ER471	ER472	ER473	ER474	ER475	ER476	ER477	ER478	ER479	ER480	ER481	ER482	ER483	ER484	ER485	ER486	ER487	ER488	ER489	ER490	ER491	ER492	ER493	ER494	ER495	ER496	ER497	ER498	ER499	ER500	ER501	ER502	ER503	ER504	ER505	ER506	ER507	ER508	ER509	ER510	ER511	ER512	ER513	ER514	ER515	ER516	ER517	ER518	ER519	ER520	ER521	ER522	ER523	ER524	ER525	ER526	ER527	ER528	ER529	ER530	ER531	ER532	ER533	ER534	ER535	ER536	ER537	ER538	ER539	ER540	ER541	ER542	ER543	ER544	ER545	ER546	ER547	ER548	ER549	ER550	ER551	ER552	ER553	ER554	ER555	ER556	ER557	ER558	ER559	ER560	ER561	ER562	ER563	ER564	ER565	ER566	ER567	ER568	ER569	ER570	ER571	ER572	ER573	ER574	ER575	ER576	ER577	ER578	ER579	ER580	ER581	ER582	ER583	ER584	ER585	ER586	ER587	ER588	ER589	ER590	ER591	ER592	ER593	ER594	ER595	ER596	ER597	ER598	ER599	ER600	ER601	ER602	ER603	ER604	ER605	ER606	ER607	ER608	ER609	ER610	ER611	ER612	ER613	ER614	ER615	ER616	ER617	ER618	ER619	ER620	ER621	ER622	ER623	ER624	ER625	ER626	ER627	ER628	ER629	ER630	ER631	ER632	ER633	ER634	ER635	ER636	ER637	ER638	ER639	ER640	ER641	ER642	ER643	ER644	ER645	ER646	ER647	ER648	ER649	ER650	ER651	ER652	ER653	ER654	ER655	ER656	ER657	ER658	ER659	ER660	ER661	ER662	ER663	ER664	ER665	ER666	ER667	ER668	ER669	ER670	ER671	ER672	ER673	ER674	ER675	ER676	ER677	ER678	ER679	ER680	ER681	ER682	ER683	ER684	ER685	ER686	ER687	ER688	ER689	ER690	ER691	ER692	ER693	ER694	ER695	ER696	ER697	ER698	ER699	ER700	ER701	ER702	ER703	ER704	ER705	ER706	ER707	ER708	ER709	ER710	ER711	ER712	ER713	ER714	ER715	ER716	ER717	ER718	ER719	ER720	ER721	ER722	ER723	ER724	ER725	ER726	ER727	ER728	ER729	ER730	ER731	ER732	ER733	ER734	ER735	ER736	ER737	ER738	ER739	ER740	ER741	ER742	ER743	ER744	ER745	ER746	ER747	ER748	ER749	ER750	ER751	ER752	ER753	ER754	ER755	ER756	ER757	ER758	ER759	ER760	ER761	ER762	ER763	ER764	ER765	ER766	ER767	ER768	ER769	ER770	ER771	ER772	ER773	ER774	ER775	ER776	ER777	ER778	ER779	ER780	ER781	ER782	ER783	ER784	ER785	ER786	ER787	ER788	ER789	ER790	ER791	ER792	ER793	ER794	ER795	ER796	ER797	ER798	ER799	ER800	ER801	ER802	ER803	ER804	ER805	ER806	ER807	ER808	ER809	ER810	ER811	ER812	ER813	ER814	ER815	ER816	ER817	ER818	ER819	ER820	ER821	ER822	ER823	ER824	ER825	ER826	ER827	ER828	ER829	ER830	ER831	ER832	ER833	ER834	ER835	ER836	ER837	ER838	ER839	ER840	ER841	ER842	ER843	ER844	ER845	ER846	ER847	ER848	ER849	ER850	ER851	ER852	ER853	ER854	ER855	ER856	ER857	ER858	ER859	ER860	ER861	ER862	ER863	ER864	ER865	ER866	ER867	ER868	ER869	ER870	ER871	ER872	ER873	ER874	ER875	ER876	ER877	ER878	ER879	ER880	ER881	ER882	ER883	ER884	ER885	ER886	ER887	ER888	ER889	ER890	ER891	ER892	ER893	ER894	ER895	ER896	ER897	ER898	ER899	ER900	ER901	ER902	ER903	ER904	ER905	ER906	ER907	ER908	ER909	ER910	ER911	ER912	ER913	ER914	ER915	ER916	ER917	ER918	ER919	ER920	ER921	ER922	ER923	ER924	ER925	ER926	ER927	ER928	ER929	ER930	ER931	ER932	ER933	ER934	ER935	ER936	ER937	ER938	ER939	ER940	ER941	ER942	ER943	ER944	ER945	ER946	ER947	ER948	ER949	ER950	ER951	ER952	ER953	ER954	ER955	ER956	ER957	ER958	ER959	ER960	ER961	ER962	ER963	ER964	ER965	ER966	ER967	ER968	ER969	ER970	ER971	ER972	ER973	ER974	ER975	ER976	ER977	ER978	ER979	ER980	ER981	ER982	ER983	ER984	ER985	ER986	ER987	ER988	ER989	ER990	ER991	ER992	ER993	ER994	ER995	ER996	ER997	ER998	ER999	ER1000
F	140	155	208	214	220	235	249	265	283	295	311	327	118	127	278	286	224	228	316	332	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
M	137	137	205	205	217	229	253	257	295	298	303	331	118	121	270	286	216	228	316	328	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	137	140	205	208	220	229	257	265	295	295	311	331	118	121	270	278	216	224	316	328	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
2	137	140	205	208	220	229	249	257	295	298	303	327	121	127	270	286	228	228	316	316	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
3	137	140	205	214	217	220	249	257	295	298	303	311	118	118	270	278	216	224	328	332	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
4	137	155	205	214	217	235	249	257	283	298	303	311	121	127	270	286	228	228	328	332	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
5	137	155	205	208	217	235	249	253	283	295	303	311	118	118	278	286	216	224	316	328	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
6	137	155	205	208	217	220	253	265	295	298	303	311	118	121	270	286	216	228	328	332	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
7	137	155	205	208	217	220	253	265	295	295	327	331	121	127	286	286	216	228	316	316	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
8	137	155	205	208	217	235	249	253	283	295	327	331	118	127	278	286	216	224	328	332	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
F	140	140	199	211	220	220	241	257	292	295	303	339	118	121	270	290	224	240	328	388	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
M	140	140	193	205	217	220	261	269	292	292	335	339	112	121	278	282	208	228	328	328	1																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
1	140	140	193	211	217	220	241	269	292	292	335	339	118	121	282	290	20																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																	

amplified well in the two populations that were separated by a large distance across Europe (c. 2730 km). Comparison of birds from these two populations showed that resident birds from Portugal have slightly lower genetic differentiation (in terms of heterozygosity, as well as allelic diversity) than the migratory population from Poland.

Higher genetic differentiation of migratory populations compared to resident ones has also been confirmed in North American Golden-crowned Kinglets (*Regulus satrapa*)⁴⁷. [I feel a few more examples should be included here]

We showed that genetic differentiation was low between the two populations, yet it was statistically significant. This supports previous studies that showed significant differentiation even within a limited geographical area, such as approximately 100 km² in one example²³. However, several other studies have suggested that migratory birds have weakly marked genetic differentiation, even over a large area⁴⁸⁻⁵⁰. Due to limited geographical sampling so far, the issue of genetic differentiation among populations of European Robins should be investigated further. In particular, for such a widespread and generally abundant species, it would be informative to investigate the relationships and genetic diversity across the very large western Palearctic range, where some populations are highly migratory and others are sedentary or highly insular (e.g. island populations)¹.

In the analysis of the genotypes of the seven full families of European Robins from Białowieża, we found strong evidence of extra-pair paternity in one nest. Although the sample size was low in this initial study (seven families, 48 nestlings), the result suggests a potentially low frequency of EPP in the study population (15% of sampled families, 4% of nestlings). However, this observation should be verified in a more extensive study, preferably by sampling a large number of representative families and populations across the species' range.

The rate of EPP that we found for European Robins was within the range of other European forest songbirds, such as the Wood Warbler *Phylloscopus sibilatrix* and Willow Warblers *P. trochilus*, which vary between zero and c. 50% of nests⁵¹⁻⁵⁴. These rates are also similar to many other bird species⁵⁵. Determining rates of EPP can be important for understanding strategies of breeding biology, settlement patterns and mate selection⁵⁶⁻⁵⁸.

In summary, our results are the first to provide the essential basis for similar ecological and genetic studies of European Robins, which is an iconic and culturally significant species in parts of Europe⁴, but has nevertheless been poorly studied across much of its range until the present time. We encourage further studies of the genetic diversity of European Robins across its broad geographical range, which may be valuable in understanding any differences between migratory and sedentary populations. Analysis of genetic diversity may also assist in the understanding of any differences between populations inhabiting the varying climatic spectrum in the European Robin's range (Mediterranean to Boreal) and the wide variety of modified habitats compared to the primeval forest of Białowieża. Comparisons in the genetic variability between other species across this range of environments and behaviours would also be valuable.

Declarations

Acknowledgements

The work was supported from the funds of the National Science Centre, Poland (research project No. 2016/23/N/NZ8/03374).

Author contributions

A.G. conducted all laboratory work, performed the statistical analysis, and drafted the article.

O.K. conceived the idea for this study, carried out the field work, and drafted the introduction and part of sample collection from Poland.

K.K.–K. carried out the field work.

P.R. conceived the idea for this study, and carried out the field work.

H.P. conducted molecular analysis with Pacific Biosciences RSII, and drafted the Microsatellites typing with Pacific Biosciences RSII and part of discussion about it.

J.P. performed bioinformatics analysis.

R.K.B. contributed to manuscript formulation and linguistics.

L.F.P.S. carried out the field work, and drafted part of sample collection from Portugal.

R.R. detailed the experimental plan, performed the statistical analysis, drafted the article, and coordinated the work.

References

1. Cramp, S. & Perrins, C. M. The birds of the Western Palearctic (ed. Cramp, S. & Perrins, C. M.) Vol. 7 (Oxford University Press, 1993).
2. Lack, D. Clutch and brood size in the Robin. *Birds*. **39**, 98-109, 130-135 (1946).
3. Lack, D. Further notes on clutch and brood size in the Robin. *Birds*. **41**, 98-104, 130-137 (1948).
4. Lack, D. The life of Robin. (Witherby, 1965).
5. Harper, D., G. C. Pairing strategies and mate choice in female Robins (*Erithacus rubecula*). *Behav.* **33**, 862-875 (1985).

6. Lebedeva, N. V. & Lomadze, N. H. The robin *Erithacus Rubecula* in the North-Western Caucasus. (ed. Matishov, G. G. & Lebedeva, N. V.) 252–277 (SSC RAS Publishing, Rostov-on-Don, 2007).
7. Knysh, N. P. Materials on the biology of Robin in forest-steppe deciduous forests of Sumy region. *Berkut* **17**, 41-60 (2008).
8. Zimin V. B. The robin in the north of the area. Volume 1. Distribution. Number. Reproduction. (ed. Zimin V.B.) 401 - 422 (Petrozavodsk: Karel'skiy nauchnyy centr RAN, 2009).
9. Baranovskiy, A. V. & Ivanov, E. S. Features of reproductive biology of robins (*Erithacus rubecula*) in anthropogenic habitats (for example, the city of Ryazan). *Principy èkologii* **6**, 17-25 (2017).
10. Wesołowski, T. Primeval conditions – What can we learn from them? *Ibis* **149**, 64-77 (2007).
11. Tobias, J. & Seddon, N. Territoriality as a paternity guard in the European robin *Erithacus rubecula*. *Behav.* **60**, 165-173 (2000).
12. Tobias, J. & Seddon, N. Female begging in European robins: do neighbors eavesdrop for extrapair copulations? *Ecol.* **13**, 637-642 (2002).
13. Lubjuhn, T., Strohbach, S., Brün, J., Gerken, T. & Epplen, J. T. EXTRA-PAIR PATERNITY IN GREAT TITS (PARUS MAJOR)-A LONG TERM STUDY. *Behaviour* **136**, 1157-1172 (1999).
14. Griffith, S. C., Owens, I. P. F. & Thuman K. A. Extra pair paternity in birds: a review of interspecific variation and adaptive function, *Ecol.* **11**, 2195-2212 (2002).
15. Cockburn, A. Prevalence of different modes of parental care in birds, *Proc Biol Sci.* **273**, 1375-1383 (2006).
16. Zagalska-Neubauer, M. & Dubiec, A. Techniki i markery molekularne w badaniach zmienności genetycznej ptaków. *Not. Ornith.* **48**, 193-206 (2007).
17. Brouwer, L. & Griffith, S. C. Extra-pair paternity in birds. *Ecol.* **28**, 4864-4882 (2019).
18. Petter, S. C., Miles, D. B. & White, M. M. Genetic evidence of mixed reproductive strategy in a monogamous bird. *The Condor* **92**, 702-708 (1990).
19. Jennions, M. D. & Petrie, M. Why do females mate multiply? A review of the genetic benefits. *Rev.* **75**, 21-64 (2000).
20. Akçay, E. & Roughgarden, J. Extra-pair paternity in birds: review of the genetic benefits. *Ecol Res.* **9**, 855-868 (2007).
21. Dietzen, C., Witt, H.-H. & Wink, M. The phylogeographic differentiation of the European robin *Erithacus rubecula* on the Canary Islands revealed by mitochondrial DNA sequence data and morphometrics: evidence for a new robin on Gran Canaria? *Avian Sci.* **3**, 115-131 (2003).
22. Rodrigues, P. et al. Phylogeography and genetic diversity of the Robin (*Erithacus rubecula*) in the Azores Islands: evidence of a recent colonisation. *J Ornithol.* **154**, 889-900 (2013).
23. Fulgione, D., Rippa, D., Manganiello, E., Caliendo, M. F. & Rastogi, R. K.. Seasonal Genetic Structure Analysis of a Resident Population of European Robin. *The Open Zoology Journal* **1**, 11-17 (2008).
24. Morin, P. A., Messier, J. & Woodruff, D. S. DNA extraction, amplification, and direct sequencing from hornbill feathers. *Sci. Soc. Thailand* **20**, 31-41 (1994).
25. Wright, T. F. et al. Microsatellite variation among divergent populations of stalk-eyed flies, genus *Cyrtodiopsis*. *Res.* **84**, 27-40 (2004).
26. Yue, G.-H., Kovacs, B. & Orban, L. A new problem with cross-species amplification of microsatellites: generation of non-homologous products. *Dongwuxue Yanjiu* **2**, 131-140 (2010).
27. Chapuis, M.-P. & Estoup, A. Microsatellite Null Alleles and Estimation of Population Differentiation. *Biol. Evol.* **24**, 621-631 (2007).
28. Dąbrowski, M. J., Bornelöv, S., Kruczyk, M., Baltzer, N. & Komorowski, J. 'True' null allele detection in microsatellite loci: a comparison of methods, assessment of difficulties and survey of possible improvements. *Ecol. Resour.* **15**, 477-488 (2015).
29. Primmer, C. R., Møller, A. P. & Ellegren, H. A wide-range survey of cross-species microsatellite amplification in birds. *Ecol.* **5**, 365-378 (1996).
30. Jaroszewicz B. et al. Białowieża Forest - a relic of the high naturalness of European forests. *Forests* **10**, 849 (2019).
31. Campos, A. R. et al. How do Robins *Erithacus rubecula* resident in Iberia respond to seasonal flooding by conspecific migrants? *Bird Study* **58**, 435-442 (2011).
32. Owen, J.C. Collecting, processing, and storing avian blood: a review. *Field Ornithol.* **82**, 339-354 (2011).
33. Horváth, M. B., Martínez-Cruz, B., Negro, J. J., Kalmár, L. & Godoy, J. A. An overlooked DNA source for non-invasive genetic analysis in birds. *Journal of Avian Biology* **36**, 84-88 (2005).
34. Faircloth, B. C. MSATCOMMANDER: detection of microsatellite repeat arrays and automated, locus-specific primer design. *Ecol. Resour.* **8**, 92-94 (2008).
35. Schuelke, M. An economic method for the fluorescent labeling of PCR fragments. *Biotechnol.* **18**, 233-234 (2000).
36. Austin, J. D. et al. Permanent genetic resources added to Molecular Ecology Resources Database 1 February 2011–31 March 2011. *Ecol. Resour.* **11**, 757-758 (2011).
37. Peakall, R. & Smouse, P. E. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics* **28**, 2537-2539 (2012).
38. Raymond, M. & Rousset, F. GENEPOP (version 1.2): population genetics software for exact tests and ecumenicism. *Heredity* **86**, 248-249 (1995).
39. Rousset, F. GENEPOP'007: a complete reimplementation of the GENEPOP software for Windows and Linux. *Ecol. Resour.* **8**, 103-106 (2008).
40. Goudet, J. FSTAT, a program to estimate and test gene diversities and fixation indices, version 2.9.3. <http://www.unil.ch/izea/software/fstat.html> (2001).
41. Kalinowski, S. T., Taper, M. L. & Marshall, T. C. Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. *Ecol.* **16**, 1099-1106 (2007).
42. Grohme, M. A., Soler, R. F., Wink, M. & Frohme, M. Microsatellite marker discovery using single molecule real-time circular consensus sequencing on the Pacific Biosciences RS. *Biotechniques* **55**, 253-256 (2013).

43. Liljegren, M. M., de Muinck, E. J. & Trosvik, P. Microsatellite length scoring by single molecule real time sequencing-effects of sequence structure and PCR regime. *PLoS ONE* **11**, e0159232 (2016).
44. Dutta, N. et al. Microsatellite marker set for genetic diversity assessment of primitive *Chitala chitala* (Hamilton, 1822) derived through SMRT sequencing technology. *Biol. Rep.* **46**, 41-49 (2018).
45. Selkoe, K. A. & Toonen, R. J. Microsatellites for ecologists: a practical guide to using and evaluating microsatellite markers. *Letts.* **9**, 615-629 (2006).
46. Corner, S., Yuzbasiyan-Gurkan, V., Agnew, D. & Venta, P.J. Development of a 12-plex of new microsatellite markers using a novel universal primer method to evaluate the genetic diversity of jaguars (*Panthera onca*) from North American zoological institutions. *Genet. Resour.* **11**, 487-497 (2019).
47. Graham, B. A., Carpenter, A. M., Friesen, V. L. & Burg, T. M. A comparison of neutral genetic differentiation and genetic diversity among migratory and resident populations of Golden-crowned-Kinglets (*Regulus satrapa*). *J Ornithol.* **161**, 509-519 (2020).
48. Bensch, S., Grahm, M., Müller, N., Gay, L. & Akesson, S. A. Genetic, morphological, and feather isotope variation of migratory willow warblers show gradual divergence in a ring. *Ecol.* **18**, 3087-3096 (2009).
49. Kralj, J., Procházka, P., Fainová, D., Patzenhauerová, H. & Tutiš, V. Intraspecific Variation in the Wing Shape and Genetic Differentiation of Reed Warblers *Acrocephalus scirpaceus* in Croatia. *Acta Ornithol.* **45**, 51-58 (2010).
50. Mettler, R. et al. Contrasting Patterns of Genetic Differentiation among Blackcaps (*Sylvia atricapilla*) with Divergent Migratory Orientations in Europe. *PLoS ONE* **8**, e81365 (2013).
51. Gyllensten, U., Jakonsson, S. & Temrin, H. No evidence for illegitimate young in monogamous and polygynous warblers. *Nature* **343**, 168-170 (1990).
52. Gil, D., Slater, P. J. B. & Graves, J. A. Extrapair paternity and song characteristics in the willow warbler *Phylloscopus trochilus*. *Avian Biol.* **38**, 291-297 (2007).
53. Moskalenko, V. N., Belokon, M. M., Belokon, Y. S. & Goretskaia, M. I. Extrapair young in nests of the Wood Warbler (*Phylloscopus sibilatrix*) in the Middle Russia (poster). 26th International Ornithological Congress; Tokyo, Japan (2014).
54. Grendelmeier, A., Arlettaz, R., Olano-Marin, J. & Pasinelli, G. Experimentally provided conspecific cues boost bird territory density but not breeding performance. *Behav Ecol* **28**, 174-185 (2017).
55. Petrie, M. & Kempnaers, B. Extrapair paternity in birds: explaining variation between species and populations. *Trends Ecol Evol.* **13**, 52-58 (1998).
56. Wagner, R. H. Hidden leks: sexual selection and the clustering of avian territories. *Ornithol Monogr.* **49**, 123-145 (1998).
57. Fletcher, R. J. & Miller, C. W. On the evolution of hidden leks and the implications for reproductive and habitat selection behaviours. *Behav.* **71**, 1247-1251 (2006).
58. Broughton, R. K., Bubnicki, J. W. & Maziarz, M. Multi-scale settlement patterns of a migratory songbird in a European primeval forest. *Behavioral Ecology and Sociobiology* **74**, 1-12 (2020).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryGwiazdowskaEtAlRobin11.04.21.xlsx](#)