

Adult HIV/AIDS Patients' Level of Satisfaction on Pharmaceutical Service: Institutional Prospective Cross Sectional Study

Adey Girmay

Addis Ababa University

Zelalem Tilahun

Addis Ababa University College of Health Sciences

Muluemebet Zinabu Akele

Addis Ababa University College of Health Sciences

solomon Assefa Huluka (✉ solomon.assefa@aau.edu.et)

Addis Ababa University

Research note

Keywords: ART, HIV/AIDS, Pharmaceutical Service, Satisfaction

Posted Date: October 17th, 2019

DOI: <https://doi.org/10.21203/rs.2.13343/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Objective : Patients are essential sources of information to assess the accessibility and effectiveness of pharmaceutical care. Patients' perception and satisfaction on the care they received has direct influence on treatment adherence. This study was thus aimed to assess HIV/AIDS patients' level of satisfaction on the pharmaceutical services provided in Zewditu Memorial Hospital, Addis Ababa, Ethiopia. An institution based cross sectional study was conducted from April 16, 2018 to May 16, 2018 in Zewditu Memorial Hospital. Result: Among the total 285 participants enrolled in the study, 172(60.4%) of them were females and the mean age of the participants was 42.5 ± 9.8 . The participants of this study received ART service for a mean duration of 7.3 ± 3.5 years. In this study, 26.7%, 13.7%, 38.6%, 13% and 8.1% of the study participants rated the overall satisfaction to the ART pharmaceutical service as excellent, very good, good, fair, and poor, respectively. Participants who were provided information about drug-drug and drug-food interaction were 2.72 times more likely to be satisfied than those who weren't provided (AOR=2.72, 95%CI : 1.13-6.54).

Introduction

HIV infection has remained a serious disease of public health importance since 1981. Since the start of the epidemic, it is claimed that more than 35.0 million people have died from AIDS related illnesses. Globally, it is estimated that approximately 36.7 million [34.0 million–39.8 million] people are living with HIV by the end of 2016 with 1.8 million new HIV infections, and 1 million AIDS related deaths [1].

Sub-Saharan Africa (SSA) countries continue to bear huge burden of HIV infection. The region is home to 25.7 million people living with HIV (PLWH) or 70% of world's PLWH [2]. Moreover, 75% of the total HIV/AIDS deaths in 2015 were from SSA [3]. Ethiopia is one of the sub-Saharan African countries with high burden of HIV [4]. In 2016, there were 720,000 PLWH and 27,104 newly diagnosed cases in Ethiopia. More than half (59%) of them were enrolled in highly active antiretroviral therapy (HAART) program [5]. Evidences suggest that improved access to antiretroviral therapy (ART) is helping to drive a decline in HIV related mortality [6]. In Ethiopia, ART service is available in 1047 health facilities [7].

Patient satisfaction on pharmaceutical care can be conceptualized as patient's evaluation of the pharmacist's performance in a variety of patient care activities [8]. It is one of the imperative components for the great success of any healthcare services, especially in the ART units [9]. Patient satisfaction has direct effect on adherence to HAART [10]. Certain patient characteristics are known to correlate with the global patient satisfaction rating. Older patients were significantly more satisfied than younger patients [11]. However, information provided pharmacist such as drug interaction were not known to be associated with patient satisfaction.

Patient's satisfaction is essentially an important issue in Africa particularly for countries like Ethiopia, where there is a dearth of adequate resources and skilled human capital to provide efficient health care services to the HIV/AIDS patients [9, 12]. We hypothesized that level of satisfaction in Ethiopia is low.

Hence, assessing the HIV/AIDS patients' level of satisfaction towards the pharmaceutical service in ART clinic is crucial for improving treatment outcome. This study is hence designed to identify the level of satisfaction of HIV/AIDS patients towards the pharmaceutical service provided at the ART clinic and its associated factors in Zewditu Memorial Hospital (ZMH), Addis Ababa, Ethiopia.

Methods

An institution-based prospective cross sectional study, using a quantitative data collection and key informant interview method, was conducted in ZMH, Addis Ababa, Ethiopia. The study setting was selected based on the virtue of being leading medical center in Ethiopia, for the treatment of HIV/AIDS. Currently, it provides ART pharmaceutical care for more than 7,000 PLWH. This study was carried out from April 16, 2018 to May 16, 2018.

In selecting the study participants, patients who were ART care beneficiaries for more than 1 year and above the age of 18 were included. The number of patients to be involved in the study was determined using the single proportion formula [13] considering the prevalence data (78.5%) taken from previous similar study done in Ethiopia [14]. Margin of error was set to be 5% (95% confidence interval). Accordingly, the sample size was:

$$n = (1.96)^2 \times 0.785(1-0.785) / (0.05)^2$$

$$n = 256$$

Assuming 10% non-response rate, the total sample size was 285.

For obtaining a representative of the population who has been visiting the ART pharmacy, a systemic random sampling technique was used. The average daily flow of patients to the hospital is approximately 100 patients. We had a 22 working days to collect data in the given study period. We expected to interview 13 patients per day. Accordingly, every eighth patient was approached to participate in the study. The first participant of the day was selected using a simple random sampling from one to eight. In place of patient refused to participate in the study, the next patient was enrolled in the study. None of the participants discontinued the study.

Data was collected using a tool adopted from Larson et al., [15]. Questionnaire was designed and modified appropriately after it was pretested on 25 clients prior to the actual data collection. Moreover, the content validity of the data collection tool was confirmed by team of expert including senior infectious disease physician in Zewditu memorial hospital and a public health personnel and clinical pharmacist from Addis Ababa University. Based on the pre-test we found that some of the information needed such as drug interaction was not included in the tool was modified. The aim of the study and the potential risk it might pose was explained to patients prior to data collection and their written consent was confirmed.

In collecting data on the patients' satisfaction towards the ART pharmaceutical service, study participants were asked to respond to a number of assessment questions classified on a Likert scale of 1–5, with “1” stood for rating of the item as “poor”, while “2”, “3”, “4”, and “5” stood for “fair”, “good”, “very good”, and “excellent”, respectively. These satisfaction assessment questions are regrouped in 8 themes.

Following checking its completeness and consistency, the data collected were entered to and analyzed using SPSS Statistics for Windows, version 20.0. Frequencies, percentages, *t*-test, and ANOVA with post hoc test were used to analyze different variables. *P*-value less than 0.05 were used as cut off points for determining statistical significance of associations among different variables.

Results

Socio-demographics Characteristics of Participants

Of the total 285 participants enrolled (100% response rate) in the study, majority (172; 60.4%) of them were females. More than half (53.3%) of the respondents were married. The mean age (\pm SD) of the participants was 42.5 ± 9.8 and the highest proportion was found between the age group of 31 and 40 years. More than one-third (34.4%) of them had attended secondary level education. Occupationally, 28.8% of them were self-employed. The participants had received the ART service for a mean duration of 7.3 ± 3.5 years (Table 1).

Table 1: Socio-demographic characteristics of adult HIV/AIDS patients in ZMH, Addis Ababa, 2018

Variable	Frequency (n=285)	Percent
Age groups, year		
18-30	33	11.6
31-40	111	38.9
41-50	77	27.0
51-60	57	20.0
>60	7	2.5
Sex		
Male	113	39.6
Female	172	60.4
Marital status		
Single	67	23.5
Married	152	53.3
Divorced	38	13.3
Widowed	28	9.8
Education status		
Cannot read or write	33	11.6
Primary	60	21.1
Secondary	98	34.4
Collage and above	94	33.0
Employment status		
Student	3	1.1
Civil servant	79	27.7
Private sector	63	22.1
Self-employed	82	28.8
House wife	50	17.5
Unemployed	8	2.8
Monthly income in ETB		
≤500	19	6.7
501-1000	20	7.0
1001-2000	59	20.7
2001-3000	67	23.5
3001-4000	46	16.1

4001-5000	39	13.7
>5000	35	12.3
Duration of ART use in year		
≤5	108	37.9
6-10	118	41.4
>10	59	20.7

Patients' Satisfaction towards the Service

More than half of the patients rated the time given by the pharmacist to listen what they are in need of as excellent (123; 43.2%) and very good (27; 9.5%). Moreover, majority (70.2%) of them claimed that there is excellent use of easy and understandable language by pharmacist for communication. Most participants (55.1%), however, rated the provision of written information by pharmacists as poor (Table 2).

As it is indicated in table 2 below, 156 (54.7%) of the participants rated the service provided about medication precaution and side effect by the pharmacist as poor. Similarly, more than half of the participants indicated that they are not satisfied with pharmacist effort to give information about drug-drug and drug-food interactions.

The arrangement of medication and the cleanliness of the environment was rated as excellent by 192 (67.4%) and 189(66.3%) participants, respectively. Of the total participants, 138(48.4%) of them rated the availability of medication as excellent and the overall appearance of the pharmacy was rated as excellent by 68.4% of the participants. As the overall satisfaction of the patients is indicated in table 2, more than a quarter (26.7%) of them rated the overall satisfaction as excellent; whereas, 13.7% , 38.6% ,13% and 8.1% of them rated their satisfaction as very good, good, fair and poor, respectively.

Table 2: patient satisfaction to words pharmaceutical service in ZMH, 2018 (n=285)

Questions	Mean (Likert scale)	Frequency		
		(%)	Excellent Good	Fair Very good Poor
Pharmacists' communication				
Pharmacist taking time to listen to you	3.47	123(43.2) 60(21.1) (22.5)	27(9.5) 11(3.9)	64
Pharmacists answers to your queries	3.85	168(58.9) (10.5) (16.1)	20(7.0) 21(7.4) 46(16.2)	30
Use of easy and understandable language	4.29	200(70.2) 15(5.3)	31(10.9) 15(5.3)	24(8.4)
labeling your medication with a readable and understandable instruction	2.07	50(17.5) 19(6.7) 180(63.2)	15(5.3) 21(7.4)	
Asked your concern about your medication	2.63	91(31.9) 28(9.8) 137(48.1)	8(2.8) 21(7.4)	
Provision of drug information in written form	2.35	68(23.9) 17(6.0) 157(55.1)	18(6.3) 25(8.8)	
Medication use information				
Pharmacist told you name of the medication	2.19	55(19.3) 33(11.6) 172(60.4)	14(4.9) 11(3.9)	
Brief explanation about medication administration	3.53	136(47.7) 36(12.6) 63(22.1)	28(9.8) 22(7.7)	
Provision of adequate information about your medication precaution and side effect	2.36	65(22.8) 29(10.2) 156(54.7)	18(6.3) 17(6.0)	
Information provided to you about your	2.19	54(18.9)	13(4.6)	

medication's drug-drug and drug-food interactions		30(10.5) 164(57.5)	24(8.4)
Pharmacist provides you information on how to solve medication related side effect	2.23	57(20.0) 25(8.8) 164(57.5)	17(6.0) 22(7.7)
Pharmacist provides you your medication with appropriate packaging	4.42	211(74.0) 14(4.9) 25(8.8)	34(11.9) 1(4.0)
The pharmacist provides you information about the proper storage of your medication	3.30	120(42.1) 31(10.9) 86(30.2)	33(11.6) 15(5.3)

Pharmacists' commitment and respect

The courtesy and respect shown to you by the Pharmacist

3.96	125(43.9) 74(26.6) 13(4.6)	62(21.8) 11(3.9)
------	----------------------------------	---------------------

Availability of the Pharmacist during the time of visit

3.96	136(47.7) 45(15.8) 27(9.5)	67(23.5) 10(3.5)
------	----------------------------------	---------------------

Pharmacist keeps your privacy during counseling

3.95	168(58.9) 30(10.5) 35(12.3)	29(10.2) 23(8.1)
------	-----------------------------------	---------------------

Time to receive pharmaceutical service

4.03	183(64.2) 20(7.0) 41(14.4)	25(8.8) 16(5.6)
------	----------------------------------	--------------------

Shorter waiting time to receive service

3.96	131(46) 0	10(35.4) 53(18.6)
------	--------------	----------------------

Pharmacists' efforts to solve medications problems

The pharmacist advice you how well your medical condition is controlled

2.27	63(22.1) 24(8.4)	17(6.0) 12(4.2) 169(59.3)
------	---------------------	------------------------------

The pharmacist advice you in detail how each medication supposed to help you

1.72	29(10.2) 15(5.3) 203(71.2)	10(3.5) 28(9.8)
------	----------------------------------	--------------------

The pharmacist Advise you about your disease and provide general advise on healthy life

2.16	57(20.0) 21(7.4) 181(63.5)	17(6.0) 9(3.2)
------	----------------------------------	-------------------

The pharmacist told you what to do when

2.24	55(19.3)	32(11.2)
------	----------	----------

you miss your dose		19(6.7)	0
The pharmacist offers you information source	1.56	179(62.8)	10(3.5)
		21(7.4)	21(7.4)
		13(4.6)	
		220(77.2)	
Infrastructure and appearance of the pharmacy			
How would you rate the arrangement of medication shelves and table	4.22	192(67.4)	34(11.9)
		17(6.0)	14(4.9)
		28(9.8)	
How would you rate the cleanliness of the environment and shelves	4.26	189(66.3)	33(11.6)
Is the space within the pharmacy adequate for all activities going in there	4.13	32(11.2)	9(3.2)
How would you rate the overall appearance of the pharmacy	4.02	182(63.9)	28(9.8)
Comfortable waiting hall	3.68	30(10.5)	21(7.4)
		172(68.4)	24(8.4)
		40(14.0)	23(8.1)
		24(8.4)	26(9.1)
		161(56.5)	18(6.3)
		22(7.7)	23(8.1)
			61(21.4)
Availability of medication			
How would you rate the availability of medication	3.68	138(48.4)	41(14.4)
		28(9.8)	34(11.9)
			44(15.4)
Overall satisfaction			
Participants overall satisfaction	3.38	76(26.7)	39(13.7)
		110(38.6)	37(13.0)
			23(8.1)

Factors Affecting Patient Satisfaction towards the Pharmaceutical service

Binary logistic regression analysis showed that there is significant ($p < 0.05$) association between provision of information about drug-drug/drug-food interaction and overall satisfaction of participants. Participants' who were provided information related to drug-drug and drug-food interaction were 2.72 times more likely to be satisfied than those who didn't receive information (AOR=2.72, 95% CI: 1.13-6.54). Participants who reported shorter time to receive service were 4.06 times more likely to be satisfied than those who did not (AOR=4.06, 95% CI: 1.69-9.74). However, no association was found between patients' socio-demographics variable and overall satisfaction (Table 3).

Table-3 Factors affecting overall satisfaction of patients attending ART service in ZMH, Addis Ababa, June 2018

Variables	Yes	No	Crudes odds ratio	Adjusted odds ratio
Age				
18-30	27	6	1.14[0.39-3.35]	4.06[0.95-17.28]
31-40	88	23	0.97[0.45-2.09]	1.86[0.67-5.10]
41-50	59	18	0.83[0.37-1.87]	1.43[0.51-3.98]
>51	51	13	1.00	1.00
Gender				
Male	93	20	1.40[0.77-2.56]	1.51[0.71-3.17]
Female	132	40	1.00	1.00
Marital status				
Single	50	17	0.63[0.21-1.94]	0.90[0.21-3.70]
Married	116	36	0.70[0.24-1.97]	1.11[0.31-3.98]
Divorced	36	2	3.91[0.70-21.88]	4.35[0.64-29.30]
Widowed	23	5	1.00	1.00
Educational status				
Can't read or write	27	6	0.92[0.32-2.59]	2.35[0.58-9.55]
Primary	48	12	0.82[0.35-1.88]	1.79[0.65-4.91]
Secondary	72	26	0.56[0.28-1.14]	1.09[0.44-2.68]
College	78	16	1.00	1.00
Information about missed dose				
Yes	90	135	1.83[0.97-3.44]	1.22[0.57-2.58]
No	16	44	1.00	1.00
Information about medication name				
Yes	87	138	1.89[0.99-3.59]	1.54[0.69-3.41]
No	15	45	1.00	1.00
Information about drug-drug and drug-food interaction				

Yes	87	10	3.15[1.51-6.54]*	2.72[1.13-6.54]*
No	138	50	1.00	1.00
Shorter waiting time				
Yes	198	27	5.60[2.92-10.74]*	4.06[1.69-9.74]*
No	34	26	1.00	1.00
Availability of medication				
Yes	176	49	3.36[1.84-6.10]*	2.03[0.96-4.27]
No	31	29	1.00	1.00

Discussion

In recent decades, determining patients' level of satisfaction is found to be the most useful tool for getting their views on how to provide them a care [16]. This is due to its ability to reflect the quality of health care delivery and provider's ability in successful delivery of care that meets patients' expectations and needs [17, 18]. A satisfied patient is more likely to develop a longer lasting rapport with their health care provider. An individual who fails to take the ART due to dissatisfaction will end up in developing resistant strains to the drugs in a matter of few minutes [19].

Present study showed that the overall mean satisfaction of the participants on ART pharmaceutical services provided at ZMH was 3.38 ± 1.23 . This finding is in line with a similar study done in Gondar on expectation and satisfaction of ART clients towards the pharmaceutical service [14]. This indicates the similarity of pharmacy practice in the settings.

In our study, 43.1% of participants rated the time taken by the pharmacist to listen to what they need as excellent; this was higher when compared with another study done in ART special unit in Addis Ababa [20]. This might be due to the effort exerted by the hospital to modernize the pharmacy service in recent years. The easy and understandable language utilization by the pharmacists was rated excellent (4.29). This was slightly higher compared to the report of the study in Gondar (3.90) by Abebe et al., [14].

The effort of pharmacists to solve patients' medication related problem was rated poor by majority of the participants. This finding is supported by a study done in Nigeria, where perception of patients regarding the pharmacists' efforts to solve patients' medication related problems were negative [21]. This could be because of lack of knowledge or willingness by the pharmacists side.

Several factors including patient's age, educational level, health status and the severity of illness influence satisfaction on care ART services [22]. However, in this study no significant association was found between patients' socio-demographics variable and overall satisfaction. This could be because the ART service is free of charge so could not be altered by of socio-demographic factors. In addition the

chronicity of the disease might make patient to have similar level of understanding about the service. Present study rather revealed that provision of information about ARV drug-drug and drug-food interaction, and shorter time to receive service influenced participants overall satisfaction towards the pharmaceutical service provided in the hospital. This indicate that improving the pharmaceutical service and develop pharmacist knowledge and skill can improve HIV/AIDS patients' satisfaction.

Conclusions

The adult HIV/AIDS patients' satisfaction with the overall ART pharmaceutical service was rated positively by participants with a mean value of 3.38. Patients claimed that they were better satisfied with infrastructure and the appearance of the pharmacy. In addition, the participants were better satisfied with the pharmacists' availability, commitment, confidentiality and for using easy and understandable language in the service. However, majority of the participants were not satisfied with the provision of medication related information and on the pharmacists' efforts to solve medications related problems.

Limitations

As the respondents were interviewed in the compound of the hospital, study bias could have occurred. Moreover, the study was conducted in a governmental hospital that provides care to the certain group of the population in the city and the sample size used is not that large enough. As such, the result may not be a reflective of the situation in the general population. The hospital manly provide service for patients who cannot afford to attend private hospitals. The finding of this study is interpreted in light of several limitations. The cross-sectional design gives only a snap shot of events.

Abbreviations

AOR: Adjusted Odds Ratio

CI: Confidence Interval

HAART: Highly Active Antiretroviral Therapy

PLWH: People Living with HIV

ZMH: Zewditu Memorial Hospital

Declarations

Ethical approval and consent to participate

Ethical approval was obtained from Ethics review committee of school of pharmacy and a formal letter was written from the Department of Pharmaceutics and Social Pharmacy to Zewditu Memorial Hospital to get permission for conducting the study. The data collectors clearly explained the aims of the study for

the study participants. Information was collected after obtaining written consent from each participant. To ensure confidentiality, participants' data were linked to a code number and registered. The right was given to the study participants to refuse or discontinue participation at any time they want and the chance to ask any thing about the study.

Availability of data and material

The data sets generated during the current study are available from corresponding author on a reasonable request.

Funding

The study was not funded

Acknowledgments

The authors acknowledge Zewditu memorial hospital for their support in providing a temporary office for the research team and facilitating the study.

Consent for Publication

Not applicable

Competing interest

The authors declare that they have no competing interests

Authors' Contributions

AG and ZT conceived the study and developed the proposal. AG, ZT and SA analyzed and interpreted the data. SA, ZT and MZ drafted the manuscript. MZ reviewed the article. All authors read and approved the final manuscript.

Author information

Department of Pharmaceutics and Social Pharmacy, School of Pharmacy, Addis Ababa University, PO Box 1176, Addis Ababa, Ethiopia

Adey Girmay & Zelalem Tilahun

Department of Pharmacology and Clinical Pharmacy, School of Pharmacy, Addis Ababa University, PO Box 1176, Addis Ababa, Ethiopia

Solomon Assefa Huluka

Muluemebet Zinabu Akele

Corresponding author

Correspondence to Solomon Assefa Huluka

References

1. Joint United Nations Programme on HIV/AIDS (UNAIDS). UNAIDS Data 2017. 2. Nash D, Yotebieng M, Sohn AH. Treating all people living with HIV in sub-Saharan Africa: a new era calling for new approaches. *J Virus Erad.* 2018; 4(Suppl 2): 1–4. 3. Wang H, Wolock TM, Carter A, Nguyen G, Kyu HH, Gakidou E, et al. Estimates of global, regional, and national incidence, prevalence, and mortality of HIV, 1980–2015: the Global Burden of Disease Study 2015. *Lancet HIV.* 2016; 3(8):e361–87. 4. UNAIDS. UNAIDS data 2017. Available from: http://www.unaids.org/sites/default/files/media_asset/20170720_Data_book_2017_en.pdf. Accessed on January 13th, 2018 5. Ethiopian Public Health Institute. HIV related estimates and projections for Ethiopia–2017. Addis Ababa: Ethiopian public health institute; 2017. 6. Phillips AN, Sabin C, Pillay D, Lundgren JD. HIV in the UK 1980–2006: reconstruction using a model of HIV infection and the effect of antiretroviral therapy. *HIV Med.* 2007; 8:536–46. 7. Federal Democratic Republic of Ethiopia HAPCO National HIV/AIDS Strategic Plan for 2015-2020; 2014. 8. Schommer JC, Kucukarslan SN. Measuring patient satisfaction with pharmaceutical services. *Am J Health Syst Pharm.* 1997; 54:2721–32. 9. Karunamoorthi K, Rajalakshmi M, Makesh SB, Yohannes A. HIV/AIDS patient's satisfactory and their expectations with pharmacy service at specialist antiretroviral therapy (ART) units. *Eur Rev Med Pharmacol Sci.* 2009; 13(5): 331-39. 10. Vahab SA, Madi D, Ramapuram J, Bhaskaran U, Achappa B. Level of Satisfaction Among People Living with HIV (PLHIV) Attending the HIV Clinic of Tertiary Care Center in Southern India. *JCDR.* 2016; 10(4): OC08. 11. Rahmqvist M, Bara A. Patient characteristics and quality dimensions related to patient satisfaction. *Int j qual heal care.* 2010; 22: 86-92. 12. Carlyn VI, Farris KB, Kassam R, Cox CE, Cave A. Pharmaceutical care research and education project: patient outcomes. *J Am Pharm Assoc.* 2001; 41(3): 411-20. 13. Lwanga SK, Lemeshow S. Sample size determination for health studies: A practical manual, World Health Organization, Geneva. 1991; pp. 1-5. 14. Abebe TM, Erku DA, Gebresillassie BM, Haile KT, Mekuria AB. Expectation and satisfaction of HIV/AIDS patients toward the pharmaceutical care provided at gondar University referral hospital, northwestern ethiopia: a cross-sectional study. *Patient Prefer and Adherence.* 2016; 10: 2073. 15. Larson LN, Rovers JP, Mackeigan LD. Patient satisfaction with pharmaceutical care: update of a validated instrument. *J Am Pharm Assoc.* 2002; 42:44-50. 16. Bahrapour A, Zolala F. Patient satisfaction and related factors in Kerman hospitals. *East Mediterr Health J.* 2005; 11(5/6): 905- 12. 17. Kagashe G, Rwebangila F. Patient satisfaction with health care services provided at HIV clinics at Amana and Muhimbili hospitals in Dar es Salaam. *Afr Health Sci.* 2011; 11(3): 60-66. 18. Mezemir R, Getachew D, Gebreslassie M. Patients' Satisfaction and its

determinants in Outpatient Department of Deberbirhan Referral Hospital, North Shoa, Ethiopia. *IJEMS*. 2014; 3(191): 2. 19. Wu AW, Gifford A, Asch S, Cohn SE, Bozzette SA, Yurk R. Quality-of-Care Indicators for HIV/AIDS: *Dis Manag Health Out*. 2000; 7:315. 20. Karunamoorthi K, Rajalakshmi M, Makesh SB, Yohannes A. HIV/AIDS patient's satisfactory and their expectations with pharmacy service at specialist antiretroviral therapy (ART) units. *Eur Rev Med Pharmacol Sci*. 2009; 13(5): 331-339. 21. Agu KA, Oqua D, Agada P, Ohiaeri SI, Adesina A, Abdulkareem MH, King RC, Wutoh AK. Assessment of satisfaction with pharmaceutical services in patients receiving antiretroviral therapy in outpatient HIV treatment setting. *Int J Clin Pharm*. 2014; 36(3): 636-647. 22. Getenet H, Haileamlak A, Tegegn A. Clients'satisfaction With Anti Retroviral Therapy Services At Jimma University Specialized Hospital. *J Health Sci*. 2008;2(18):21.