

Analysis of the efficacy of HIV protease inhibitors against SARS-CoV-2's main protease

mohamed mahdi (✉ mohamed@med.unideb.hu)

University of Debrecen <https://orcid.org/0000-0002-7959-5996>

János András Mótyán

Debreceni Egyetem

Zsófia Ilona Szojka

Debreceni Egyetem

Mária Golda

Debreceni Egyetem

Márió Miczi

Debreceni Egyetem

József Tózsér

Debreceni Egyetem

Research

Keywords: SARS-CoV-2, Protease, HIV Protease Inhibitors, In vitro assay, Inhibition profiling

Posted Date: July 10th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-40776/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on November 26th, 2020. See the published version at <https://doi.org/10.1186/s12985-020-01457-0>.

Abstract

Background

The pandemic caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has resulted in millions of infections worldwide. While the search for an effective antiviral is still ongoing, experimental therapies based on repurposing of available antivirals is being attempted, of which HIV protease inhibitors (PIs) have gained considerable interest. Inhibition profiling of the PIs directly against the viral protease has never been attempted *in vitro*, and while few studies reported an efficacy of lopinavir and ritonavir in SARS-CoV-2 context, the mechanism of action of the drugs remains to be validated.

Methods

We carried out an in-depth analysis of the efficacy of HIV PIs against the main protease of SARS-CoV-2 (M^{Pro}) in cell culture and *in vitro* enzymatic assays, using a methodology that enabled us to focus solely on any potential inhibitory effects of the inhibitors against the viral protease.

Results

Lopinavir, ritonavir, darunavir, saquinavir, and atazanavir were able to inhibit the viral protease in cell culture, albeit in concentrations much higher than their achievable plasma levels, given their current drug formulations. While inhibition by lopinavir was attributed to its cytotoxicity, ritonavir was the most effective of the panel, with IC_{50} of 13.7 μ M. Atazanavir on the other hand was the only PI to inhibit the viral protease both in cell culture and in our *in vitro* enzymatic assay.

Conclusion

Targeting of SARS-CoV-2 M^{Pro} by some of the HIV PIs might be of limited clinical potential, given the high concentration of the drugs required to achieve significant inhibition. Therefore, given their weak inhibition of the viral protease, any potential beneficial effect of the PIs in COVID-19 context might perhaps be attributed to acting on other molecular target(s), rather than SARS-CoV-2 M^{Pro} .

1. Background

In December 2019, a novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was identified as the etiological agent of viral pneumonia cases that occurred in Wuhan, Hubei Province, China. As of July 1st 2020, the pandemic has resulted in more than 10 million infections, and 500,000 deaths worldwide according to the World Health Organization.

There is currently no standardized treatment protocol, and there is no antiviral treatment of proven efficacy recommended for COVID-19. Clinical management of patients is mainly supportive, including supplementary oxygen and mechanical ventilation if needed. However, given the overwhelming burden of the pandemic on national healthcare systems and the global economy, experimental therapies have been attempted, which are predominantly based on the repurposing of FDA approved antivirals, antimalarials, arthritis drugs, and blood plasma derivatives [1].

The HIV protease inhibitors (PIs) lopinavir and ritonavir have gained particular interest, having shown documented *in vitro* activity against SARS-CoV and the Middle East respiratory syndrome coronavirus (MERS), however, these studies did not identify a molecular target for the drugs, since their efficacy was solely determined based on the inhibition of cytopathic effects or viral replication, respectively [2 3]. Given the 94.4% identity in amino acid sequence between SARS-CoV and SARS-CoV-2 [4], studying the efficacy of HIV protease inhibitors against SARS-CoV-2 would be of major relevance.

The genome of SARS-CoV-2 encodes for two viral cysteine proteases; nsp3 (papain-like protease) and nsp5 (main protease) [5]. The main protease (M^{pro}) of SARS-CoV-2; also named chymotrypsin-like protease (3CLpr), plays a crucial role in the viral life cycle, cleaving the initial polyproteins translated from the viral RNA at at least 11 of its 14 cleavage sites. M^{pro} of SARS-CoV-2 shares 96% sequence identity to that of SARS-CoV. The enzyme consists of three domains; two domains (I and II) which consist of antiparallel β -barrels, and an α -helical domain (I), which is responsible for dimerization and enzymatic activity [6 7]. Recent structure determination confirmed the similarities between the two enzymes [8].

One potential target for the HIV PIs is the M^{pro} . *In silico* screening identified nelfinavir as its potential inhibitor [9], while lopinavir and ritonavir were found to be potential inhibitors of the viral enzyme by molecular dynamics simulation [10].

It is important to note that the HIV protease is a C2-symmetric homodimeric aspartyl protease, composed of two identical subunits that are 99 amino acids each. The active site is located at the interface between the two monomers, and contains the catalytic Asp-Thr-Gly residues [11]. M^{pro} on the other hand, is a cysteine protease that can potentially be targeted by peptide mimetics. Given the structural difference between the two proteases, the efficacy of HIV protease inhibitors against SARS-CoV and SARS-CoV-2 is questionable.

Previous studies reported that a combination of lopinavir/ritonavir and ribavirin was effective against SARS-associated coronavirus, with concentrations of 4 $\mu\text{g}/\text{ml}$ and 50 $\mu\text{g}/\text{ml}$, respectively [12]. However, a recent clinical trial of 99 patients with laboratory-confirmed SARS-CoV-2 infection who were treated with lopinavir–ritonavir concluded that no significant benefit was observed in the treated group compared to those who received standard care [13]. Recently, a short communication reported that lopinavir inhibited SARS-CoV-2 replication in Vero E6 cells with IC_{50} of 26.63 μM , ritonavir, however, showed no inhibition of viral replication [14].

Early *in vitro* reports from China showed that darunavir inhibited SARS-CoV-2 replication, although at a very high concentration (300 μ M) [15]. Clinical trials are currently ongoing [16].

Our aim was to test the efficacy of a panel of HIV PIs against SARS-CoV-2 M^{Pro}, using a cell culture-based model. In this study, we determined the IC₅₀ of the PIs with the aid of a dark-to-bright GFP substrate system that had been developed and applied previously for the investigation of caspases [17]. Moreover, *in vitro* enzymatic inhibitory assays were also carried out using purified M^{Pro} and an oligopeptide substrate representing the AVLQ*SGFR cleavage site of SARS-CoV-2 polyprotein 1 ab (PP1ab).

2. Materials And Methods

2.1 Plasmids and inhibitors

Coding sequence of SARS-CoV-2 M^{Pro} (GenBank: MT291835.2) was cloned into pcDNA3.1(+) mammalian expression plasmid using BamHI/EcoRI restriction sites to create the SARS-CoV-2 M^{Pro} coding plasmid; thereafter referred to as CoV-2 M^{Pro}. The coding sequence of a dark-to-bright GFP reporter substrate; thereafter referred to as PR-Sub, was also cloned into pcDNA3.1(+) plasmid. The PR-Sub was designed to contain a sequence representing the N-terminal autoproteolytic cleavage site of SARS-CoV-2 M^{Pro} (TSAVLQ*SGFRKM); corresponding to the nsp4/nsp5 cleavage site, between the GFP and the Influenza A/M2 protein hydrophobic tail (CNDSSDPLVVAASIIIGILHLILWILDRL). For *in vitro* expression of the protease, the coding sequence of His₆-tagged M^{Pro} was cloned into pET11a bacterial expression plasmid using NdeI and BamHI enzymes. The above mentioned expression constructs were obtained using the gene synthesis service of GenScript.

The protease inhibitors darunavir, saquinavir, lopinavir, tipranavir, indinavir sulfate, and atazanavir sulfate were obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH. Ritonavir was obtained from Abbott laboratories, nelfinavir from Agouron, and atazanavir from Bristol-Myers Squibb.

A synthetic oligopeptide used in our *in vitro* enzymatic assay representing the N-terminal autoproteolytic cleavage site of SARS-CoV-2 M^{Pro} (AVLQ*SGFR) was obtained from a peptide synthesis service (BioBasic).

2.2 Analysis of transfection efficiency and proteolysis

293T human embryonic kidney cells (HEK-293T) (Invitrogen) were maintained in T-75 flask in 15 mL Dulbecco's modified Eagle's medium (DMEM) (Sigma-Aldrich) supplemented with 10% fetal bovine serum (FBS), 1% glutamine and 1% penicillin-streptomycin. Cells were transfected at 70% confluency with 5 μ g of either PR-Sub, or CoV-2 M^{Pro} plus PR-Sub plasmids using PEI method [18]. After 24 hours incubation, GFP fluorescence was analyzed by flow cytometry using FACS Calibur (BD Biosciences).

2.3 Inhibition profiling in cell culture

On the day of transfection, HEK-293T cells were split and transferred into a 48-well plates (30,000 cells/well) containing serial dilutions of the inhibitor ranging from 200 μM to 5 nM in a total volume of 200 μL DMEM/well, supplemented with 10% FBS, 1% glutamine and 1% penicillin-streptomycin. After 3 hours incubation at 37 °C, cells were transfected with 300–300 ng of CoV-2 M^{Pro} and PR-Sub plasmids using lipofectamine LTX reagent (Thermo Fisher Scientific), then the cells were incubated for 24 hours. GFP fluorescence was then measured by flow cytometry using FACS Calibur. The results were analyzed by FlowJo Software Version 10 (Becton, Dickinson and Company; 2019). Calculations of IC₅₀ were performed using GraphPad Prism 5.0 (GraphPad Software, Inc).

2.4 Cell viability assay

The day before the assay, HEK-293T cells were split into a 96-well plates (20,000 cells/well) containing serial dilutions of the inhibitor ranging from 200 μM to 100 nM in a total volume of 200 μL DMEM/well, supplemented with 10% FBS, 1% glutamine and 1% penicillin-streptomycin. The next day, the medium was replaced with 100 μL of OPTI-MEM culture media supplemented with 10% FBS, and 10 μL of the 12 mM 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) stock solution was added to the cells. After 4 hours incubation at 37 °C, 85 μL of supernatant was removed, and 50 μL of DMSO was added to the cells followed by incubation for 10 minutes at 37 °C. Absorbance was measured at 540 nm using Synergy H1 Hybrid Multi-Mode Reader (Agilent).

2.5 Expression and purification of His₆-M^{Pro}

His₆-M^{Pro} was expressed in BL21(DE3) cells, the expression was performed in Luria-Bertani (LB) medium supplemented with ampicillin (100 $\mu\text{g}/\text{ml}$ final concentration). Protein expression was induced by isopropyl β -D-1-thiogalactopyranoside (IPTG) (1 mM final concentration). Cells were pelleted by centrifugation at 4 °C for 20 min at 5,000 x g (Sorvall Lynx 4000, Thermo Fisher Scientific), the cell pellet was lysed in buffer A (20 mM Tris, 150 mM NaCl, 10 mM imidazole, pH 7.5) followed by sonication (Branson Sonifier 450). After a repeated centrifugation at 4 °C for 20 min at 10,000 x g, the pellet was discarded and His₆-M^{Pro} was purified from the supernatant by Ni-chelate affinity chromatography with the aid of His-Trap Column (GE Healthcare) using Äkta Prime instrument (Amersham Pharmacia Biotech). Afterwards, the purification buffer was changed to buffer B (20 mM Tris, 1 mM DTT, pH 8.0) using Amicon Ultra centrifugal filters (10K, Merck Millipore), and the protein was further purified by ion-exchange chromatography using HiTrap Q FF column (GE Healthcare). The high-purity fractions were dialyzed against buffer C (20 mM Tris, 150 mM NaCl, 1 mM EDTA, 1 mM DTT, pH 7.8).

2.6 *In vitro* protease assay

The AVLQ*SGFR oligopeptide was dissolved in distilled water and was used as substrate in activity measurements to test inhibitory potentials of the PIs.

The cleavage reactions contained 10 μL reaction buffer (20 mM Tris, 100 mM NaCl, pH 7.8), 4.8 μL oligopeptide substrate (1.37 mM final concentration), and 0.2 μL DMSO (in control samples) or 0.2 μL of the inhibitor (diluted in DMSO). For inhibitor screening, inhibitors were applied in 100 μM final

concentration. Reactions were initiated by the addition of 5 μL of His₆-M^{Pro} in a final total protein concentration of 0.12 μM , and the mixtures were incubated at 37 °C for 30 minutes. The reactions were terminated by the addition of 180 μL 1% trifluoroacetic acid (TFA). The cleavage products were detected using high performance liquid chromatography (HPLC), utilizing a 0-100% water-acetonitrile gradient in the presence of TFA using Merck Hitachi instrument. Relative activity was determined at less than 20% substrate hydrolysis. Activity measured in the presence of DMSO was considered to be 100%. While no potent inhibitor of M^{Pro} was available to perform active-site titration, 100% activity was assumed for the enzyme.

2.7 Modeling

Homology modeling of dark-to-bright GFP substrate was performed using Phyre2 web portal [19]. 97% of residues were modelled at > 90% confidence. Structural figures were prepared PyMol Molecular Graphics System (Version 1.3 Schrödinger, LLC).

3. Results

3.1 Inhibition profiling in cell culture

To measure M^{Pro} activity in cell culture experiments, we applied a modified version of a dark-to-bright GFP substrate [17]. The recombinant substrate consists of a GFP, followed by a proteolytic cleavage site and a C-terminal hydrophobic tail. Proteolysis at the inserted cleavage site releases the tail that serves as a hydrophobic quencher of fluorescence and facilitates tetramerization of GFP, which prevents chromophore maturation; the fluorescence is restored upon proteolysis (Fig. 1).

Firstly, we optimized the transfection of HEK-293T cells for the use of the SARS-CoV-2 M^{Pro} and the dark-to-bright GFP substrate. Transfection of cells with only the PR-Sub plasmid resulted in a maximum of 1% background fluorescence after 24 hours incubation. When cells were transfected with both the PR-Sub and CoV-2 M^{Pro} plasmids, GFP fluorescence ranged from 28–34%, indicating processing of the substrate and the activity of the protease (Fig. 2).

We then analyzed the inhibition efficacy of a panel of HIV PIs against SARS-CoV-2 M^{Pro}. While none of the inhibitors was able inhibit the viral protease in nanomolar concentration; which is expected for effective transition state analogs, in micromolar range, ritonavir was the most effective ($\text{IC}_{50} = 13.7 \pm 1.1 \mu\text{M}$). Saquinavir, darunavir, and atazanavir were also able to inhibit SARS-CoV-2 M^{Pro} at higher concentrations (Table 1).

Table 1
Results of inhibition profiling of HIV protease inhibitors against SARS-CoV-2 M^{Pro} in cell culture. Data are calculated from triplicate experiments. IC₅₀ for lopinavir was indeterminable accurately due to high cytotoxic effects in HEK-293T cells.

Inhibitor	IC ₅₀ (μM)	Standard error
Lopinavir + Ritonavir	10.9	± 1.1
Ritonavir	13.7	± 1.1
Saquinavir	31.4	± 1.2
Darunavir	36.1	± 1.2
Atazanavir	60.7	± 2.5
Lopinavir	Indeterminable	
Indinavir	No inhibition (up to 200 μM)	
Nelfinavir	No inhibition (up to 200 μM)	
Tipranavir	No inhibition (up to 200 μM)	

Although a combination of lopinavir and ritonavir resulted in better inhibition of the viral enzyme as compared to ritonavir alone (10.9 μM vs. 13.7 μM, respectively), when we carried out cell viability assays after treatment of the cells with the inhibitors, interestingly, inhibition by lopinavir was found to be a result of the high cytotoxicity observed at concentrations above 50 μM (90%) (Fig. 3).

In the case of ritonavir and saquinavir, cytotoxicity of > 50% was only observed at concentrations above 50 μM. High concentrations of darunavir and atazanavir on the other hand, were well tolerated by HEK-293T cells.

Indinavir, nelfinavir and tipranavir however, failed to inhibit SARS-CoV-2 M^{Pro}, even at 200 micromolar concentration of the inhibitors.

3.2 *In vitro* enzymatic assay

Following expression and purification of a His₆-tagged M^{Pro}, we determined its catalytic activity after incubation with AVLQSGFR oligopeptide substrate. Cleavage position within the substrate was determined using high-performance liquid chromatography coupled to electrospray ionization time-of-flight mass spectrometry (HPLC-ESI-TOF MS) (Supplementary Fig. 1). Inhibition profiling of the PIs was carried out after incubation of the inhibitors along with M^{Pro} and the substrate, and the relative efficacies of PIs were compared. Interestingly, of the panel, only atazanavir showed a significant inhibition of the M^{Pro} *in vitro* (p value = 0.04) (Fig. 4).

4. Discussion

To our knowledge, direct determination of the inhibition efficacy of HIV PIs against SARS-CoV-2 M^{Pro} in cell culture has not yet been attempted, although many *in silico* studies analyzing interaction between SARS-CoV-2 M^{Pro} and potential inhibitors were published [20–26] (Supplementary Table 1). Antiviral assays using lopinavir and ritonavir in Calu-3 cells were previously carried out for MERS-CoV, and the IC₅₀ for lopinavir, ritonavir and their combination was 11.6, 24.9, and 8.5 μM respectively [3]. In our analysis, we found that a combination of lopinavir plus ritonavir achieved the lowest IC₅₀, this however was due to the high cytotoxicity of lopinavir, and not as a result of direct inhibition of SARS-CoV-2 M^{Pro}. Ritonavir on the other hand was much more tolerable than lopinavir, and achieved the lowest IC₅₀. This should be taken into consideration, given the current drug formulation of lopinavir which is in combination with ritonavir, where ritonavir is used as a pharmacokinetic enhancer due to its inhibition of the cytochrome P450 3A4 isoenzyme, thereby increasing the bioavailability of lopinavir [27]. As a result, administration of ritonavir in combination with other PIs was found to decrease its minimum blood plasma concentration level, as compared to the generic formulation of the drug [28]. Also, while our result regarding ritonavir was in direct contrast to what Choy et.al. reported in their short communication [14], we believe that difference in methodologies is to blame for this discrepancy, as we examined the efficacy of ritonavir against the viral M^{Pro} protease per se.

It is important to note that the IC₅₀ of the inhibitors were in the micromolar range, which is not considered optimal for the inhibition of the viral enzyme. Previous studies have reported that the minimum concentrations (C_{min}) for lopinavir, darunavir, saquinavir, and atazanavir in patient's serum under antiretroviral treatment was found to be 9.3, 3.3, 3.8, and < 1 μM, respectively [29–32]. It would indeed be challenging to achieve such high plasma levels of the inhibitors in order to block the viral replication, moreover, the cytotoxic effects of some of the inhibitors, in addition to the side effects commonly observed with PIs questions the use of anti-HIV PIs in the context of SARS-CoV-2.

Additionally, using an *in vitro* enzymatic assay, we were able to directly analyze any potential inhibition of M^{Pro} by the HIV PIs. Of the panel, only atazanavir was able to significantly inhibit M^{Pro} *in vitro*, albeit at high concentrations. This finding however, should be interpreted with care, as even in the context of HIV protease, atazanavir had been known to have a variable and somewhat decreased efficacy on different HIV isolates [33 34].

5. Conclusion

In conclusion, to our knowledge, thorough analysis of the efficacies of PIs against SARS-CoV-2 remains scarce, and the targets of the drugs are yet to be verified. While few studies examined the efficacy of some PIs against the replication of SARS-CoV-2, we set out to study whether or not the inhibitors exert a direct effect on the viral protease. In our experiments, even though some of the PIs developed for the treatment of HIV were able to inhibit SARS-CoV-2 M^{Pro}, they were only able to do so at high

concentrations. The combination of lopinavir plus ritonavir resulted in the lowest IC₅₀ in cell culture, albeit at the cost of cellular viability. Although, darunavir and atazanavir required a much higher concentration to achieve the inhibition, cytotoxicity was not observed even at a concentration of 200 µM. It should be noted that there might be other molecular targets for the HIV PIs, as nelfinavir was recently shown to inhibit spike protein-mediated fusion of SARS-CoV-2 [35].

A drawback of this study is that we were not able to assess the interaction between the PIs and the papain-like protease of SARS-CoV-2, as our methodology only enabled us to study the viral main protease. Whether or not these exert any inhibitory effect on the papain-like protease is a subject for future studies. Also, while our cell culture assays were not performed in SARS-CoV-2 target cells, our methodology enabled us to directly examine any potential inhibition of the viral M^{pro} by the PIs, therefore, it is unlikely that different results will be obtained in target cells. Taking everything into consideration, the use of HIV PIs in the context of COVID-19 might be of limited clinical potential, beneficial effects of which might perhaps be attributed to acting on other molecular target(s), rather than M^{pro} itself. Data from clinical trials will indeed shed more light on their clinical efficacy.

Abbreviations

SARS-CoV-2: severe acute respiratory syndrome coronavirus 2

HIV: Human immunodeficiency virus

PI: Protease inhibitor

Mpro: Main protease

IC50: Half maximal inhibitory concentration

COVID-19: Coronavirus disease-2019

WHO: World health organization

FDA: Food and drug administration

MERS: Middle East respiratory syndrome

PR: Protease

GFP: Green fluorescent protein

FACS: Fluorescence-activated cell sorting

HEK-293T: Human embryonic kidney 293 cells

DMSO Dimethyl sulfoxide

MTT: 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide

IPTG: isopropyl β -D-1-thiogalactopyranoside

TFA: Trifluoroacetic acid

HPLC-ESI-TOF MS: high-performance liquid chromatography coupled to electrospray ionization time-of-flight mass spectrometry

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All data generated during this study are included in this article and its supplementary information files

Competing interests

The authors declare no competing interests

Funding

This work was supported in part by the Higher Education Institutional Excellence Programme (NKFIH-1150-6/2019) of the Ministry of Innovation and Technology in Hungary, within the framework of the Biotechnology thematic programme of the University of Debrecen.

The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results

Authors` contributions

M. Mahdi, J.A.M., and J.T. designed the experiments and interpreted the results; M. Mahdi, and Zs. Sz. carried out cell culture experiments, J.A.M., M.G., and M. Miczi carried out *in vitro* experiments. All authors contributed to the writing of the manuscript, and approved the final draft.

Acknowledgments

The authors would like to thank Szilvia Janics-Pető for her continuous help and technical assistance, and are grateful to the staff of the Laboratory of Retroviral Biochemistry. Authors are grateful to Tibor Nagy (Department of Applied Chemistry, Faculty of Science and Technology, University of Debrecen) for performing HPLC-ESI-TOF MS analysis.

References

1. Li G, De Clercq E. Therapeutic options for the 2019 novel coronavirus (2019-nCoV). *Nature reviews Drug discovery*. 2020;19(3):149–50. doi:10.1038/d41573-020-00016-0[published Online First: Epub Date].. (: do.
2. Su J, Rui Y, Lou M, et al. HIV-2/SIV Vpx targets a novel functional domain of STING to selectively inhibit cGAS-STING-mediated NF-kappaB signalling. *Nature microbiology*. 2019;4(12):2552–64. doi:10.1038/s41564-019-0585-4[published Online First: Epub Date]].
3. Chougui G, Munir-Matloob S, Matkovic R, et al. HIV-2/SIV viral protein X counteracts HUSH repressor complex. *Nature microbiology*. 2018;3(8):891–97. doi:10.1038/s41564-018-0179-6[published Online First: Epub Date]].
4. Manel N, Hogstad B, Wang Y, et al. A cryptic sensor for HIV-1 activates antiviral innate immunity in dendritic cells. *Nature*. 2010;467(7312):214–7. doi:10.1038/nature09337[published Online First: Epub Date]].
5. Chan JF, Kok KH, Zhu Z, et al. Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerging microbes infections*. 2020;9(1):221–36 doi. 2[published Online First: Epub Date].. (: 10.1080/22221751.2020.171990.
6. Anand K, Ziebuhr J, Wadhwani P, et al. Coronavirus main proteinase (3CLpro) structure: basis for design of anti-SARS drugs. *Science*. 2003;300(5626):1763–7. doi:10.1126/science.1085658[published Online First: Epub Date]].
7. Gentile D, Patamia V, Scala A, et al. Putative Inhibitors of SARS-CoV-2 Main Protease from A Library of Marine Natural Products: A Virtual Screening and Molecular Modeling Study. *Marine drugs* 2020;18(4) doi:10.3390/md18040225[published Online First: Epub Date]].
8. Dai W, Zhang B, Su H, et al. Structure-based design of antiviral drug candidates targeting the SARS-CoV-2 main protease. *Science* 2020 doi: 10.1126/science.abb4489[published Online First: Epub Date]].
9. Mittal L, Kumari A, Srivastava M, et al. Identification of potential molecules against COVID-19 main protease through structure-guided virtual screening approach. *Journal of biomolecular structure & dynamics* 2020:1–26 doi: 10.1080/07391102.2020.1768151[published Online First: Epub Date]..
10. 10.1021/acs.biochem.0c00160

- Nutho B, Mahalapbutr P, Hengphasatporn K, et al. Why Are Lopinavir and Ritonavir Effective against the Newly Emerged Coronavirus 2019? Atomistic Insights into the Inhibitory Mechanisms. *Biochemistry* 2020;**59**(18):1769-79 doi: 10.1021/acs.biochem.0c00160[published Online First: Epub Date]].
11. Wlodawer A, Miller M, Jaskolski M, et al. Conserved folding in retroviral proteases: crystal structure of a synthetic HIV-1 protease. *Science*. 1989;245(4918):616–21.
 12. Chu CM, Cheng VC, Hung IF, et al. Role of lopinavir/ritonavir in the treatment of SARS: initial virological and clinical findings. *Thorax*. 2004;59(3):252–6. doi:10.1136/thorax.2003.012658[published Online First: Epub Date]].
 13. Cao B, Wang Y, Wen D, et al. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19. *The New England journal of medicine* 2020 doi: 10.1056/NEJMoa2001282[published Online First: Epub Date]].
 14. Choy KT, Wong AY, Kaewpreedee P, et al. Remdesivir, lopinavir, emetine, and homoharringtonine inhibit SARS-CoV-2 replication in vitro. *Antiviral research*. 2020;178:104786. doi:10.1016/j.antiviral.2020.104786[published Online First: Epub Date]].
 15. Dong L, Hu S, Gao J. Discovering drugs to treat coronavirus disease 2019 (COVID-19). *Drug discoveries & therapeutics* 2020;**14**(1):58–60 doi: 10.5582/ddt.2020.01012[published Online First: Epub Date]].
 16. 10.1016/j.cell.2020.02.052
Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 2020;**181**(2):271 – 80 e8 doi: 10.1016/j.cell.2020.02.052[published Online First: Epub Date]].
 17. Nicholls SB, Chu J, Abbruzzese G, et al. Mechanism of a genetically encoded dark-to-bright reporter for caspase activity. *J Biol Chem*. 2011;286(28):24977–86. doi:10.1074/jbc.M111.221648[published Online First: Epub Date]].
 18. Baker A, Saltik M, Lehrmann H, et al. Polyethylenimine (PEI) is a simple, inexpensive and effective reagent for condensing and linking plasmid DNA to adenovirus for gene delivery. *Gene therapy*. 1997;4(8):773–82. doi:10.1038/sj.gt.3300471[published Online First: Epub Date]].
 19. Kelley LA, Mezulis S, Yates CM, et al. The Phyre2 web portal for protein modeling, prediction and analysis. *Nature protocols*. 2015;10(6):845–58. i: 10.1038/nprot.2015.053[published Online First: Epub Date]]. (: do.
 20. Shah B, Modi P, Sagar SR. In silico studies on therapeutic agents for COVID-19: Drug repurposing approach. *Life sciences*. 2020;252:117652. doi:10.1016/j.lfs.2020.117652[published Online First: Epub Date]].
 21. Sang P, Tian S-H, Meng Z-H, et al. Anti-HIV drug repurposing against SARS-CoV-2. *RSC Advances*. 2020;10(27):15775–83. i: 10.1039/D0RA01899F[published Online First: Epub Date]]. (: do.
 22. Pant S, Singh M, Ravichandiran V, et al. Peptide-like and small-molecule inhibitors against Covid-19. *Journal of biomolecular structure & dynamics* 2020:1–10 doi:

- 10.1080/07391102.2020.1757510[published Online First: Epub Date]..
23. Ortega JT, Serrano ML, Pujol FH, et al. Unrevealing sequence and structural features of novel coronavirus using in silico approaches: The main protease as molecular target. *EXCLI J.* 2020;19:400–09. doi:10.17179/excli2020-1189[published Online First: Epub Date].
 24. Calligari PB, Ricci S, Bocedi G. A.. Molecular Investigation of SARS–CoV-2 Proteins and Their Interactions with Antiviral Drugs. *Viruses.* 2020;12:445.
 25. Beck BR, Shin B, Choi Y, et al. Predicting commercially available antiviral drugs that may act on the novel coronavirus (SARS-CoV-2) through a drug-target interaction deep learning model. *Computational structural biotechnology journal.* 2020;18:784–90 doi. 5[published Online First: Epub Date].. :: 10.1016/j.csbj.2020.03.02.
 26. Fischer A, Sellner M, Neranjan S, et al. Potential Inhibitors for Novel Coronavirus Protease Identified by Virtual Screening of 606 Million Compounds. *Int J Mol Sci* 2020;21(10) doi:10.3390/ijms21103626[published Online First: Epub Date].
 27. Hull MW, Montaner JS. Ritonavir-boosted protease inhibitors in HIV therapy. *Annals of medicine.* 2011;43(5):375–88. i: 10.3109/07853890.2011.572905[published Online First: Epub Date].. (: do.
 28. van der Lugt J, Lange J, Avihingsanon A, et al. Plasma concentrations of generic lopinavir/ritonavir in HIV type-1-infected individuals. *Antivir Ther.* 2009;14(7):1001–4. i: 10.3851/IMP1410[published Online First: Epub Date].. (: do.
 29. Lopez-Cortes LF, Ruiz-Valderas R, Sanchez-Rivas E, et al. Lopinavir plasma concentrations and virological outcome with lopinavir-ritonavir monotherapy in HIV-1-infected patients. *Antimicrob Agents Chemother.* 2013;57(8):3746–51. doi:10.1128/AAC.00315-13[published Online First: Epub Date].
 30. Gutierrez-Valencia A, Torres-Cornejo A, BenMarzouk-Hidalgo OJ, et al. Darunavir minimum plasma concentration and ritonavir-boosted darunavir monotherapy outcome in HIV-infected patients. *Antivir Ther.* 2014;19(5):443–7. doi:10.3851/IMP2722[published Online First: Epub Date].
 31. Lotsch J, Harder S, Sturmer M, et al. Association of saquinavir plasma concentrations with side effects but not with antiretroviral outcome in patients infected with protease inhibitor-susceptible human immunodeficiency virus type 1. *Antimicrob Agents Chemother.* 2007;51(9):3264–72. doi:10.1128/AAC.00036-07[published Online First: Epub Date].
 32. Smith DE, Jeganathan S, Ray J. Atazanavir plasma concentrations vary significantly between patients and correlate with increased serum bilirubin concentrations. *HIV Clin Trial* 2006;7(1):34–8 doi: 10.1310/0KX0-H9VH-99EE-5D0L[published Online First: Epub Date]..
 33. Wood R. Atazanavir: its role in HIV treatment. *Expert review of anti-infective therapy* 2008;6(6):785 – 96 doi: 10.1586/14787210.6.6.785[published Online First: Epub Date].
 34. Mahdi M, Szojka Z, Motyan JA, et al. Inhibition Profiling of Retroviral Protease Inhibitors Using an HIV-2 Modular System. *Viruses* 2015;7(12):6152-62 doi: 10.3390/v7122931[published Online First: Epub Date].

35. Musarrat F, Chouljenko V, Dahal A, et al. The anti-HIV Drug Nelfinavir Mesylate (Viracept) is a Potent Inhibitor of Cell Fusion Caused by the SARS-CoV-2 Spike (S) Glycoprotein Warranting further Evaluation as an Antiviral against COVID-19 infections. Journal of medical virology. 2020. doi:10.1002/jmv.25985[published Online First: Epub Date].

Figures




Figure 1

The dark-to-bright reporter system utilized to investigate Mpro activity in cell culture. A) Sequences of Mpro and the reporter substrate (PR-Sub). B) Schematic representation of the dark-to-bright reporter system. Homology model structure of PR-Sub is also shown.



Figure 2

Optimization of HEK-293T cell transfection with SARS-CoV-2-Mpro and the dark-to-bright GFP substrate. A) Cells transfected with PR-Sub and CoV-2 Mpro under native microscopic light. B) Visualization of cells transfected with PR-Sub under fluorescent microscope. C) Visualization of cells transfected with PR-Sub and CoV-2 Mpro under fluorescent microscope. Co-transfection with both plasmids resulted in 28-34 % GFP fluorescence.




Figure 3

Determination of IC₅₀ in cell culture. Relative activity (%) is plotted on the left Y axis versus logarithmic transformation of the inhibitor's concentration (nM). Percentage of cytotoxicity is plotted on the right Y axis. Error bars represent SD (n=3).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementary.pdf](#)