

Nutrition-oriented reformulation of extruded cereals and associated environmental footprint: a case study

Namy Espinoza-Orias

Nestle Research Center

Antonis Vlassopoulos

Agricultural University of Athens <https://orcid.org/0000-0001-7136-1039>

Gabriel Masset (✉ gabriel.masset@nestle.com)

<https://orcid.org/0000-0001-6977-0247>

Research

Keywords: environmental impact, breakfast cereals, reformulation, life-cycle assessment

Posted Date: July 10th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-40806/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Foods on September 8th, 2020. See the published version at <https://doi.org/10.3390/foods9091260>.

Abstract

Background The global food system faces a dual challenge for the decades ahead: providing nutritious food to a growing population while reducing its environmental footprint. Reformulation of food has been identified as a strategy to improve the nutritional quality of products; but the environmental consequences of such reformulations have rarely been studied. The objective was to assess the evolution of five environmental impact indicators following reformulations of extruded breakfast cereals. **Methods** Recipe, nutritional composition and sourcing data for three extruded breakfast cereals were retrieved from the manufacturer, at three distinctive time points of a 15-year reformulation cycle: 2003, 2010, and 2018. Nutritional information was summarized using the Nutri-Score indicator. The environmental impact of all recipes was assessed by Life Cycle Assessment (LCA) using five indicators: climate change (CC), Freshwater consumption scarcity (FWCS), Abiotic resource depletion (ARD), Land use impacts on biodiversity (LUIB), and Impacts on ecosphere/ecosystems quality (IEEQ). Life cycle inventory (LCI) was both primary data from the manufacturer and secondary data from usual third-party LCI datasets. **Results** Reformulation led to improved nutritional quality for all three products. In terms of environmental impact, improvements were observed for the CC, ARD and IEEQ indicators, with average reductions of 12%, 14% and 2% between 2003 and 2018, respectively. Conversely, the FWCS and LUIB indicators were increased by 57% and 70%, respectively. For all indicators but ARD, ingredients contributed most to the environmental impact, while the absolute impacts of other stages of the supply chain (i.e. manufacturing, packaging, transport and end-of-life) reduced. A sensitivity analysis simulating no deforestation agricultural practices suggested that the CC indicator could be further reduced by 21 to 49% for the 2018 recipes. **Conclusions** This study highlighted the need to better account for the environmental consequences of changing food product recipes. While improvements can be achieved at all stages of products' life cycles, agricultural commodities need to be the focus for further improvement, in particular in shelf-stable grain-based products such as extruded cereals. This could be achieved through selection of less demanding ingredients and improvements in agricultural practices.

Background

The global food system faces a dual challenge for the decades ahead: providing nutritious food to a growing population while reducing its environmental footprint (1–3). Despite not being a new concept, the environmental sustainability of food is attracting renewed attention with the increased realization that sustainable food production will require extensive changes across the food system (e.g. agriculture, production, packaging and consumption) (1) and that a healthy and sustainable diet as proposed in the latest guidelines is likely too expensive for 1.58 billion people globally (4).

Public health interventions in nutrition have traditionally been designed along the principles of two complementary approaches: the issuing of food-based guidelines and the theory of incremental changes (5). The first encourages individuals to adopt healthier diets overall, i.e. changing their eating habits towards more plant-based products, and products with lower levels of saturated fats, sugar and sodium (1, 6, 7). The second focuses on the improvement of existing products and production methods, mainly

through product reformulation, and it is aimed as a short to medium term action that respects consumer expectations, avoids the inflation of a diet's cost and as such delivers improvements even among the lowest socioeconomic classes (8, 9).

While there is extensive evidence on the association between healthier and more environmentally friendly dietary choices overall (1, 6), recent evidence suggests that dietary shifts alone, i.e. the first approach, may not suffice to fit the global food system within the 'planetary boundaries' (10). Improvements of the production methods being needed to reduce further the environmental footprint of the food system. Studies looking at specific food products, their process and/or nutritional properties, have rarely combined environmental and nutritional dimensions together. Indeed, studies tend to focus on improvement of the food production per se, highlighting alternative methods to reduce the environmental footprint of products (11–14); or focusing on the evolution of the nutritional profile of specific food products, or group of food products, and their potential impact of nutritional intakes and health outcomes (8, 15–17). The question of whether such reformulations were compatible with a reduction of the environmental footprint of the food system has been rarely addressed, but the results are so far promising (18–21). However, the study of food sustainability presents challenges and would require the wider scientific community to familiarize with notions such as unit efficiency in food production and life-cycle assessments, as well as the understanding that environmental footprint analysis goes beyond the commonly used greenhouse gas emissions to include multiple indicators (22–24).

This paper aims to expand existing literature on food sustainability by presenting three cases of food reformulation in extruded breakfast cereals – carried out with the purpose of improving their nutritional value – and its impact on the products' environmental impacts. The latter were assessed through a complete Life Cycle Assessment including multiple sustainability indicators and presenting 15-years' worth (2003-2010-2018) of primary data from the three most produced recipes by Cereal Partners France. The aim of this analysis was to assess whether product reformulation could be an opportunity to improve a product's environmental footprint.

Methods

Cases selection

The three biggest extruded breakfast cereal recipes by volume of production by Cereal Partners France were selected. Cereal Partners France (CPF), a division of Cereal Partners Worldwide (CPW), manufactures breakfast cereals for domestic consumption at two facilities in France. Primary data was collected from CPF for 3 individual years over a 15 years period, namely: 2003, 2010 and 2018. Even though product reformulation cycles may be shorter, these three years represent key changes in nutritional profile improvements of the products, in particular the inclusion of whole grains in the recipes. Data from the 2003 recipes serve as the baseline while data from the next two major recipe changes in 2010 and 2018 serve as examples of the intermediate and current recipes. For all products only the main recipe was considered excluding any alternative products with different flavours or formats. The products

analysed are Chocapic® 430 g, Lion® 480 g, and Nesquik® 450 g, produced in the Rumilly and Itancourt factories of Cereal Partners France (hereafter mentioned as Cereals 1–3, respectively).

Life cycle Assessment Methodology

The functional unit chosen for the study is the “consumption of one serving size of breakfast cereal (30 g)”. Figure 1 describes the “cradle to grave” boundaries defined for the system under assessment. The system includes the agricultural production of ingredients, production of packaging materials, manufacture of breakfast cereals, distribution, retail, end of life of packaging materials, and all transportation in between life cycle stages. Two main life cycle stages are excluded from the assessment, namely consumption of cereals with milk or alternative products (plant-based beverages such as “soy milk”, “almond milk”, “oat milk”, or “pea milk”) and end of life management of post-consumer breakfast cereal waste. The reason to exclude the consumption stage lies in the fact that the nutrients provided by the combination of a serving size of breakfast cereal and a beverage (milk or a plant-based beverage) can differ greatly, rendering the potential scenarios for assessment not functionally comparable (understood as delivering the same nutrient-provision function). Similarly, the amount of post-consumer food waste depends on consumer behavior. Transportation of the product from retail store to home and of waste from home to end of life treatment facility are excluded because these are not significant contributors to the overall life cycle of the product.

Life cycle inventory included data on recipe formulation, packaging, sourcing of materials, manufacture, distribution, and retail as provided by the manufacturer. Secondary data coming from regional statistics was used to model end of life management of packaging materials in France. Life cycle inventory datasets are sourced from the following databases: ecoinvent v.3.4, allocation based on recycled content (2017); and World Food LCA Database (WFLDB) v.3.3 (2017). The LCA was carried out using the life cycle assessment software SimaPro v.9.0 (PRé).

Nutritional Data and Raw Material

The exact recipes and ingredients lists for the production of 100 g of finished product was provided by the manufacturer (Cereal Partners France) and closely matched the labelled ingredient list with the exception of humidity lost during production due to drying of the product (Supplementary Tables S1-S3). Region of origin was provided for all raw materials and it was assumed that agricultural production of the ingredients followed conventional agricultural practices. Cereals and sugar are sourced largely from France; palm oil and palm kernel oil are sourced from South East Asia; soybean oil and soy lecithin are sourced from South America; and the remaining ingredients are sourced from various European locations (Supplementary Table S4). For the purpose of this analysis, it was assumed that suppliers remained unchanged over the period of the study. The nutritional composition of the products was based on the nutritional declaration as labelled. The Nutri-Score algorithm was used to assess the nutritional quality of the products at each timepoint (25).

Environmental Footprint Assessment

Five environmental impact indicators were evaluated in the study, chosen because they provide a sufficient coverage of relevant environmental issues associated with packaged food products:

- Climate change [kg CO₂-eq] (100 a) (26). Biogenic CO₂ is assigned a characterization factor of 0. This is a midpoint indicator.
- Freshwater consumption scarcity [m³-eq] (27). This is a midpoint indicator.
- Abiotic resource depletion [kg Sb-eq] (CML 2001 method, v. 2.05) (28). This is a midpoint indicator.
- Land use impacts on biodiversity [PDF • m² • year] (IMPACT World+ / Land use method, v.0.05) (29). This is an endpoint indicator.
- Impacts on ecosphere/ecosystems quality [PDF • m² • year] (Impact 2002 + method v. Q2.27) (30, 31). This is an endpoint indicator. The method was modified to include aquatic eutrophication, aquatic acidification, terrestrial acidification and nutrients, and ecotoxicity (aquatic and terrestrial). Land occupation is addressed already by the Land use impacts on biodiversity indicator.

Inclusion of Packaging, Manufacture, Distribution and Retail, Packaging and of Life and Transportation in the Life cycle inventory

Primary data on packaging was used for the LCA calculated as g of material used per cereal box (Supplementary Table S5). In this case, primary packaging corresponds to the “bag in box” format, containing between 430 and 480 g of breakfast cereal. This packaging format is required to protect the brittle nature of the product. These boxes are then packed into secondary packaging made of corrugated board (between 12 and 22 boxes per case). The cases (between 16 and 24) are loaded in layers onto pallets and wrapped with stretch film (tertiary packaging). Plastic materials used include HDPE, high density polyethylene and LLDPE, linear low-density polyethylene sourced from the UK; folding board boxes (FBB) and corrugated board cases sourced from the UK and France. Virgin materials are used in the packaging system (plastic and fibre based). Other packaging, used for some raw ingredients and packaging materials is excluded from the assessment. Cereals and flours are delivered in bulk at factory gate.

Data corresponding to the overall annual operation at two factories in France for the three years (2003, 2010, 2018) were used to estimate the impact of manufacture on the environmental footprint of the products. The manufacturing process data were calculated per kg of produced products and as such annual production tonnage and investment on projects aiming to improve energy and water efficiency were taken into account (Supplementary Table S6). Waste is also generated during manufacturing, and detailed records exist for years 2010 and 2018, indicating the various waste streams and the respective end of life management options taken. However, after calculating the ratio of mass of waste to mass of product manufactured, it was decided not to include these datasets in the assessment due to them being non-material (ranging in the order of 10^{-5} to 10^{-4} kg per kg of final product).

For distribution and retail, the LCA model operated under the assumption that breakfast cereals are ambient, shelf-stable products, with a typical shelf life of 9 months. Once manufactured, products leave the factories and are taken to regional distribution centres where they are stored for 6 days. Then, they are taken to retail stores where the typical turn around period is 5 days. All transportation steps between life cycle stages were accounted for in the model.

The end of life assessment included only packaging material. Primary packing (the packaging used for the cereal boxes themselves) includes four waste channels: municipal solid waste landfilling and incineration, plastic packaging recycling and cardboard recycling. Data from national and European Union statistics were used to estimate how municipal solid waste and specific packaging materials were treated in France over the years of the assessment (2003, 2010 and 2018; full details in Supplementary Table S7) (32–34). In brief, it was assumed that in the years 2003 and 2010 plastic bags were not recycled and were instead treated as municipal solid waste and composting was not included as a disposal option in any timepoint. For secondary packaging (packaging used for transport and storage) it was assumed that secondary cardboard was recycled at a 100% rate while secondary plastic was treated under the same conditions as primary plastic.

Given the major contribution of land transformation to climate change and greenhouse gas emission, a sensitivity assessment was carried out to evaluate the effect of improved agricultural practices on this indicator. Conventional agricultural practices take into account statistical records of actual deforestation in the cultivation of three key commodities whose processed ingredients are part of the recipes of the breakfast cereals assessed: soybean oil, cocoa powder and palm oil. On the other hand, improved agricultural practices consider that no land use change would have taken place in the cultivation of said commodities. This was possible by using life cycle inventories from the World Food LCA Database (v.3.3), which incorporate country or commodity specific factors for land use change.

Statistical Analysis

Given that the sample selected for the analysis is a fixed sample of products (case study) and not meant to represent all products under the same category, only descriptive statistics were used to identify changes in key indicators.

Results

Raw materials and nutritional composition

In the previous 15 years the recipes of the products studied changed as a result of product reformulation (details in Supplementary Tables S1-S3). In terms of ingredient choices and recipe, the major changes include the introduction of wholegrains since 2010 as a substitute of the previously used refined grains. Between 2003–2018 the use of refined grains was reduced by 45% in average and wholegrains were introduced in amounts that covered the reduction in refined grains by 1.45 times. Similarly, there was an overall 60% reduction in the use of plant-based oils combined with a complete removal of palm oil by

2018 and a substitution by sunflower oil as the main plant-based oil used. Sugars from all sources were reduced by 10% in the 15 years studied and the use of cocoa powder or cocoa mass remained unchanged. The main cereals used include are wheat and corn with rice removed from the ingredients since 2010 (ingredient data not shown). As shown in Fig. 2 and supplementary Table S8, the main changes in the nutritional composition were a more than two-fold increase in the mean fiber content, followed by a nearly 70% reduction in mean saturated fat and a 37% reduction in mean sugar and sodium content. Protein was also increased by 31% in average while the energy content remained largely unchanged (< 6% reduction for all).

The changes in the nutritional composition were reflected in an improvement of the Nutri-Score rating of all three products (Fig. 3).

Environmental footprint assessment

In terms of changes in the environmental footprint of the cereal products studied, the analysis indicates a mixed picture (Fig. 4). On average the three products showed improvements for the climate change (mainly greenhouse gas emissions) and abiotic resource depletion indicators, with an average impact reduction in the range of 12% and 14%, respectively. On the other hand, smaller improvements were seen for impact on the ecosystem with a 2% reduction of the indicator and only one of the three products showing consistent improvements through the study period. Conversely, all three products showed higher needs for freshwater usage and a greater impact on the biodiversity due to land use with average increases of 57% and 70% respectively.

For all scenarios assessed, changes in freshwater usages were attributed to irrigation of crops in different countries (83–93% of the results of this indicator), and other uses account for the remaining 7–16% of impact (data not shown). Agriculture and especially the use of land for agricultural purposes explains between 85–97% of all impact in biodiversity. In particular, land use impacts on biodiversity are driven by occupation of land for the cultivation of annual crops such as cereals (52–94%), and perennial crops such as cocoa beans (3–32%).

A reduced use of resources such as solid fossil fuels (ranging between 25% and 30%), natural gas (ranging between 40% and 48%), or crude oil (ranging between 26% and 35%) explain changes in the abiotic resources' depletion. Similarly, there are four main drivers behind 90% of the impact on ecosystem quality in all scenarios. The largest contributor is ammonia (ranging between 61% and 67%) due to its characterization in aquatic and terrestrial ecotoxicity and aquatic and terrestrial acidification. The second largest contributor are nitrous oxides, ranging between 14% and 16% and characterized in terrestrial and aquatic acidification. The third largest contributor is phosphate, ranging between 9% and 10% and characterized only in aquatic eutrophication. Finally, sulfur dioxide contributes between 2% and 3% and is characterized in aquatic and terrestrial acidification.

On the contrary, the impact on climate change could be equally attributed to greenhouse gases (CO₂, CO, and CH₄) arising from land transformation and greenhouse gases arising from industrial processes (N₂O and fossil CO₂, CO and CH₄) for two out of three cases (46–55% land transformation to 43%-50% industrial processes), but not for the third case in which greenhouse gases from industrial processes explain 62–65% of the climate change impact (data not shown).

For all indicators, except Abiotic Resource Depletion, the choice of ingredients was the main contributor towards the environmental footprint of the average product explaining 64%-98% of all variability per indicator (Fig. 5 and Supplementary Tables S9-S11). For Abiotic Resource Depletion, both the choice of ingredient and manufacture had equal contribution of 25%-40%, followed by packaging that explained ~ 20% of the impact. Packaging was also responsible for 6.6– 12% of the impact on land use on biodiversity and both packaging and manufacture contributed ~ 10% of the total climate change impact. Interestingly, although the total impact on climate is reduced for all products from 2003 to 2018, the importance of ingredients choice as the main contributor towards their carbon footprint is increasing from 64,7% in 2003 to 71% in 2018. This could be linked to the reduced contribution of manufacturing towards the products' carbon footprint from 17–11.7% from 2003 to 2018.

Given that ingredients were shown to be the major contributor towards the environmental footprint for most indicators, a secondary analysis was carried out to measure the changes in all environmental indicators attributable only to ingredient selection (Table 1). Studying the changes in ingredients alone, the evolution of the environmental indicators was in the same direction as the complete LCA analysis – albeit to a smaller degree – with the exception of abiotic resource depletion and impact on the ecosphere quality. For those two indicators that are shown to have improved in the complete LCA analysis it would appear that changes in steps further down the products lifecycle have larger effects that mandate the net impact (likely manufacture and packaging). The same can be said for the climate change indicator whose 15-year improvement is 4-fold larger in the complete LCA as compared to when analyzing ingredients alone.

Table 1

Changes in the five environmental impact indicators in the 2003–2018 period for the product average-analysis of ingredients alone versus the complete LCA.

	Ingredients only					Complete LCA				
	ARD	CC	FWCS	IEEQ	LUIB	ARD	CC	FWCS	IEEQ	LUIB
2003	0%	0%	0%	0%	0%	0%	0%	0%	0%	0%
2010	-7%	-7%	-8%	-4%	18%	-13%	-10.3%	-8.7%	-6.3%	14.3%
2018	12%	-3%	62%	1%	78%	-14%	-11.7%	57%	-2.3%	69.7%
CC: Climate change; ARD: Abiotic resource depletion; LUIB: Land use impact on biodiversity; IEEQ: Impact on ecosphere/ecosystems quality; FWCS: Freshwater consumption scarcity.										

As ingredients are the main contributor to the carbon footprint and given the above stated results, a sensitivity analysis was carried out to measure the impact of changing agricultural practices on the carbon footprint of the products. As shown in Table 2 switching from ingredients that have been produced with conventional practices to those following practices that do not promote deforestation can reduce the estimated CO₂ emissions of a single serving by 21–55%, which highlight the significance of sourcing commodities from supply chains where increased traceability can demonstrate adherence to no deforestation commitments.

Table 2
Effect of introduction of no deforestation practices over climate change impact results [kg CO₂-eq/ 30 g serving size]

Product	Year	Conventional practices	No deforestation practices	Potential Change
Cereal 1	2003	9.4E-02	5.0E-02	-47%
	2010	9.6E-02	5.4E-02	-44%
	2018	8.9E-02	5.2E-02	-41%
Cereal 2	2003	1.2E-01	5.9E-02	-49%
	2010	1.1E-01	4.8E-02	-55%
	2018	1.0E-01	5.1E-02	-49%
Cereal 3	2003	9.0E-02	6.5E-02	-27%
	2010	6.7E-02	4.6E-02	-32%
	2018	7.6E-02	6.0E-02	-21%

Discussion

This study aimed to use real-life examples to estimate the potential to improve extruded breakfast cereals environmental impact via nutrition-focused reformulation.

The recipe changes that occurred during the 15-year reformulation cycle, i.e. switching from palm oil to sunflower oil and from purely refined cereals to a mix of wholegrain and refined cereals alongside a general reformulation to reduce sugar and fat content, led to substantial improvements in the products' nutritional value as measured by Nutri-Score. On the level of individual nutrients, the recipe changes were linked to reductions in energy, total sugars, saturated fats and sodium content with parallel improvements in their fiber and protein content.

The reformulation cycle, together with the changes that occurred in the other stages of the supply chain, was associated with improvements of three environmental impact indicators out of the five assessed in this study (climate change, abiotic resource depletion and impact on ecosphere quality); with increases observed for the land use impact on biodiversity and freshwater consumption scarcity.

Ingredients (i.e., the agricultural stage) contributed to more than 60% of the environmental impact for all indicators but abiotic resource depletion; at all phases of the reformulation cycle. In this specific study, the removal of rice, potentially one of the crops with the highest carbon footprints (35), and the switch from palm oil to sunflower oil could explain the changes in carbon footprint seen both at the ingredients level and for the full LCA (36, 37). However, despite the high irrigation needs of rice (38), its removal was not linked to improvements in freshwater scarcity. Similarly, the removal of palm oil was not associated with improvements in markers on biodiversity protection. A potential explanation for those changes could be linked to the hypothesis made in this analysis that all ingredients were produced following conventional practices. As shown in the sensitivity analysis, changes in the agricultural practices towards a no deforestation policy could have a high impact on the greenhouse gas emissions of these products (up to 55% reductions); and those same agricultural practices would promote greater biodiversity protection. The present results thereby strengthen the rationale for increasing the effort on improving farming practices to reduce the environmental impact of the food system (39, 40). The food industry has a key role to play in promoting those beneficial agricultural practices via its choice of suppliers and the promotion of agricultural transformation. While the choices of suppliers based on their environmental footprint may have attracted little attention so far, data indicate that even within the same country, the region of production for a raw material is linked with great differences in its carbon and water footprint, under the same agricultural practices (41). In reality though, the implementation of such environmental management strategy requires large resources in terms of establishing monitoring and auditing protocols, increasing the frequency of site visits, creating and strengthening partnerships with external verification organizations and local authorities, and investing in new technological solutions combining satellite remote sensing and big data analytics (42).

The contribution of ingredients to the environmental impact of products increased throughout the reformulation cycle. This was due to the reduction – in absolute terms – of the environmental impact of the other stages of the life-cycle, in particular manufacturing, packaging, and end-of-life. These improvements, which partly mitigated some negative effects of ingredient changes, highlight the increased rates of recycling that occurred in France in the 15 years analyzed, and the continuous progress that has been made by food manufacturers in terms of efficiency of manufacturing and packaging technology.

This analysis is not without caveats. Firstly, the product category studied is unique in terms of its environmental impact as the stage of transport, retail and food waste by the consumers are fairly small contributors in the environmental footprint given that the products are dried and shelf stable. The same analysis could yield substantially different results if carried out in products that require a cold-supply chain (43). Even among the cereal-based products, breakfast cereals have different LCAs than products like bread in which food waste and storage would play a larger role in the overall carbon footprint despite ingredients remaining the main contributor (44). Moreover, the choice of databases to retrieve the environmental data and the assumptions made for the suppliers and manufacturing methods of the historical recipes indicate that extrapolation of these results to other food categories should be made with caution (45–47). The lack of regularly updated life cycle inventory databases, which would include

good quality historical and statistical data and the reliance on the publicly available datasets, hinders the capacity to reflect on actual changes in the agricultural practices (48). On the other hand, this study is potentially the only one to present data on a full LCA analysis using a combination of publicly available and real-life data in all stages of the model and utilizing access to the manufacturers own databases to increase accuracy. In terms of carbon footprint, the current analysis estimated that greenhouse gas emissions of the analyzed products fall in the range of 67–100 g CO₂-eq for every 30 g portion after reformulation. Previous studies that followed a similar methodology and carried out a full LCA analysis (bar end of life analysis) indicate that greenhouse gas emissions for cereal products fall in the range of 80–117 g CO₂-eq with exception of two studies that indicated much lower emissions 21–30 g CO₂-eq, potentially due to the studies' limited scope (studying specific aspects of the cereal product production or use of different LCI databases) (11, 49–52).

Overall, the results add to an ongoing discussion among academia, food industry and the consumers on how food can become more sustainable and more nutritious in order to address future challenges. The results highlight that the supply chain as a whole and the raw materials in particular are key areas for future work. This observation also holds true for whole diets (53): the Mediterranean diet, with its short supply chains, high utilization of low environmental impact agriculture, and respect for seasonality being a prime example of the need for a systems approach on food sustainability (54). Consumers are increasingly concerned over the provenance and environmental performance of the food they consume, but they often seem to focus on one aspect of a product's environmental impact like its packaging (55) or the so-called food mile, a metric of the greenhouse gas emissions related to the transportation of a food from production to the kitchen cupboard. Often environmental sustainability is intertwined with the notion that a food is necessarily better for health or that it includes less unfamiliar ingredients and it is produced in a more 'natural' manner (56–59). The increased awareness of consumers around sustainability issues and the potential it has to impact consumer choices will require new research that expands beyond the simple greenhouse gas emissions metrics and the identification of new ways to communicate the complexity of an environmental impact LCA analysis to a non-expert audience (60).

Today multiple environmental impact indicators exist, each focusing on a certain aspect of the environmental impact equation. As previously shown in the nutrition world, in order to successfully drive consumer behavior and engage multiple stakeholders, the creation of composite markers that weigh the different aspects and serve as a simple to use metric to compare one food to another will be needed (61–63). The science of nutrient profiling was developed in nutrition for this purpose (64) and a similar discipline might soon be needed in the environmental sciences. Although primary reports exist on the conception and usefulness of such indicators (65), the creation of a unique metric that combines all aspects of a multi-indicator assessment will require a series of subjective choices which will eventually be challenged by the scientific community. The European Union Product Environmental Footprint methodology has attempted the creation of a framework to study simultaneously 15 environmental indicators, created a list of normalization factors and issued a series of pilot studies aiming in reducing

the complexity of multi-indicator assessment towards an easier assessment and communication of the impact of products on the environment (66, 67).

Conclusions

This study highlights that for the case of dry breakfast cereals, nutritional and environmental improvement do not necessarily go hand-in-hand. While ingredients contributed the most to the environmental impact of the analyzed products, the food industry has managed to reduce some of the environmental impact of such products mainly through improvements in post-agricultural life cycle stages. The results highlight the need for reformulation strategies that look beyond the factory gates and engage the agricultural sector in a way that promotes more sustainable production of raw material, improvements in its transportation and even reduction of food losses in the field.

List Of Abbreviations

ARD	Abiotic Resource Depletion
CC	Climate Change
CPF	Cereal Partners France
CPW	Cereal Partners Worldwide
FBB	Folding Board Boxes
FWCS	Freshwater Consumption Scarcity
HDPE	High Density Polyethylene
IEEQ	Impacts of Ecosphere/Ecosystems Quality
LCA	Life Cycle Assessment
LCI	Life Cycle Indicators
LLDPE	Linear Low-Density Polyethylene
LUIB	Land Use Impacts on Biodiversity

Declarations

Ethics approval and consent to participate

Not Applicable

Consent for Publication

Not Applicable

Availability of Data and Materials

The datasets generated during and/or analyzed during the current study are not publicly available as they include proprietary information of Société des Produits Nestlé S.A., Cereal Partners Worldwide and Cereal Partners France but may be available from the corresponding author on reasonable request.

Competing interests

NE, is employed by Nestlé Research, Switzerland, a member of the Société des Produits Nestlé S.A., a wholly owned affiliate of Nestlé S.A, which provides professional assistance, research and consulting services for food, dietary, dietetic and pharmaceutical products of interest to Nestlé S.A. GM is employed by Cereal Partners Worldwide. AV is a consultant to Cereal Partners Worldwide employed as a scientific writer.

Funding

The study was financed by Société des Produits Nestlé S.A., Nestlé France and Cereal Partners Worldwide.

Authors' contribution

NE acquired and analyzed the data, AV prepared the manuscript, GM & NE designed the study. All authors contributed to data interpretation and reviewed the manuscript. All authors have approved the submitted version of the manuscript and have agreed both to be personally accountable for the author's own contributions and to ensure that questions related to the accuracy or integrity of any part of the work, even ones in which the author was not personally involved, are appropriately investigated, resolved, and the resolution documented in the literature.

Acknowledgments

The authors would like to thank Guillaîne Kinimo for the preliminary data preparation and analyses, Adrian Cawsey for assisting in the primary data collection, and Jean-Manuel Bluet for the advice and support provided to conduct this study.

References

1. Willett W, Rockström J, Loken B, Springmann M, Lang T, Vermeulen S, et al. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. Vol. 393, *The Lancet*. Lancet Publishing Group; 2019. p. 447–92.
2. Intergovernmental Panel on Climate Change (IPCC). Climate Change and Land. An IPCC Special Report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems [Internet]. 2019 [cited 2020 Jul 6]. Available from: <https://www.ipcc.ch/srccl/>
3. Townsend RF, Jaffee S, Hoberg YT, Htenas A. Future of Food: Shaping the Global Food System to Deliver Improved Nutrition and Health [Internet]. Washington, DC: World Bank, Washington, DC; 2016 [cited 2020 Jul 6]. Available from: <https://openknowledge.worldbank.org/handle/10986/24104>
4. Hirvonen K, Bai Y, Headey D, Masters WA. Affordability of the EAT–Lancet reference diet: a global analysis. *Lancet Glob Heal* [Internet]. Elsevier Ltd; 2020 Jan 1 [cited 2020 Jun 29];8(1):e59–66. Available from: <http://www.thelancet.com/article/S2214109X19304474/fulltext>
5. Implementation I of M (US) C on DG, Thomas PR. Improving America’s Diet and Health [Internet]. Improving America’s Diet and Health. National Academies Press; 1991 [cited 2020 Jun 29]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK235261/>
6. Perignon M, Vieux F, Soler L-G, Masset G, Darmon N. Improving diet sustainability through evolution of food choices: Review of epidemiological studies on the environmental impact of diets. *Nutr Rev*. 2017;75(1).
7. Poore J, Nemecek T. Reducing food’s environmental impacts through producers and consumers. *Science* (80-). American Association for the Advancement of Science; 2018 Jun 1;360(6392):987–92.
8. Federici C, Detzel P, Petracca F, Dainelli L, Fattore G. The impact of food reformulation on nutrient intakes and health, a systematic review of modelling studies. *BMC Nutr* [Internet]. Springer Science and Business Media LLC; 2019 Dec 7 [cited 2020 Feb 18];5(1):2. Available from: <https://bmcnutr.biomedcentral.com/articles/10.1186/s40795-018-0263-6>
9. Gressier M, Privet L, Mathias KC, Vlassopoulos A, Vieux F, Masset G. Modeled dietary impact of industry-wide food and beverage reformulations in the United States and France. *Am J Clin Nutr* [Internet]. American Society for Nutrition; 2017 Jul 1 [cited 2020 Jun 29];106(1):225–32. Available from: <https://academic.oup.com/ajcn/article/106/1/225-232/4569820>
10. Springmann M, Clark M, Mason-D’Croz D, Wiebe K, Bodirsky BL, Lassaletta L, et al. Options for keeping the food system within environmental limits. *Nature*. Nature Publishing Group; 2018 Oct 25;562(7728):519–25.
11. Jeswani HK, Burkinshaw R, Azapagic A. Environmental sustainability issues in the food-energy-water nexus: Breakfast cereals and snacks. *Sustain Prod Consum*. Elsevier; 2015 Apr 1;2:17–28.
12. Kulak M, Nemecek T, Frossard E, Gaillard G. Eco-efficiency improvement by using integrative design and life cycle assessment. The case study of alternative bread supply chains in France. *J Clean Prod*. Elsevier Ltd; 2016 Jan 20;112:2452–61.

13. Noya LI, Vasilaki V, Stojceska V, González-García S, Kleynhans C, Tassou S, et al. An environmental evaluation of food supply chain using life cycle assessment: A case study on gluten free biscuit products. *J Clean Prod.* Elsevier Ltd; 2018 Jan 1;170:451–61.
14. Sieti N, Rivera XCS, Stamford L, Azapagic A. Environmental impacts of baby food: Ready-made porridge products. *J Clean Prod.* Elsevier Ltd; 2019 Mar 1;212:1554–67.
15. Masset G, Mathias KC, Vlassopoulos A, Mielenberg F, Lehmann U, Gibney M, et al. Modeled dietary impact of pizza reformulations in US children and adolescents. *PLoS One.* 2016;11(10).
16. Combris P, Goglia R, Henini M, Soler LG, Spiteri M. Improvement of the nutritional quality of foods as a public health tool. *Public Health [Internet].* 2011 Oct [cited 2019 Feb 4];125(10):717–24. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21890152>
17. Spiteri M, Soler LG. Food reformulation and nutritional quality of food consumption: An analysis based on households panel data in France. *Eur J Clin Nutr.* Nature Publishing Group; 2018 Feb 1;72(2):228–35.
18. Chaudhary A, Marinangeli C, Tremorin D, Mathys A. Nutritional Combined Greenhouse Gas Life Cycle Analysis for Incorporating Canadian Yellow Pea into Cereal-Based Food Products. *Nutrients [Internet].* MDPI AG; 2018 Apr 16 [cited 2020 Jun 29];10(4):490. Available from: <http://www.mdpi.com/2072-6643/10/4/490>
19. Van Mierlo K, Rohmer S, Gerdessen JC. A model for composing meat replacers: Reducing the environmental impact of our food consumption pattern while retaining its nutritional value. *J Clean Prod.* Elsevier Ltd; 2017 Nov 1;165:930–50.
20. Saget S, Costa M, Barilli E, Wilton de Vasconcelos M, Santos CS, Styles D, et al. Substituting wheat with chickpea flour in pasta production delivers more nutrition at a lower environmental cost. *Sustain Prod Consum [Internet].* Elsevier; 2020 Oct 1 [cited 2020 Jul 6];24:26–38. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S2352550920302815>
21. Heller MC, Keoleian G a, Willett WC. Toward a life cycle-based, diet-level framework for food environmental impact and nutritional quality assessment: a critical review. *Environ Sci Technol [Internet].* 2013 Nov 19;47(22):12632–47. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24152032>
22. Garnett T. Food sustainability: Problems, perspectives and solutions. In: *Proceedings of the Nutrition Society [Internet].* Cambridge University Press; 2013 [cited 2020 Jun 29]. p. 29–39. Available from: </core/journals/proceedings-of-the-nutrition-society/article/food-sustainability-problems-perspectives-and-solutions/B75C1F93146221F8EDD98A90CF9A67A2>
23. Williams P. Broadening influence on the food supply and environmental sustainability. *Nutr Diet [Internet].* Blackwell Publishing Ltd; 2019 Jul 1 [cited 2020 Jun 29];76(3):247–9. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1111/1747-0080.12554>
24. Lang T, Barling D. Nutrition and sustainability: An emerging food policy discourse. In: *Proceedings of the Nutrition Society [Internet].* Cambridge University Press; 2013 [cited 2020 Jun 29]. p. 1–12.

- Available from: /core/journals/proceedings-of-the-nutrition-society/article/nutrition-and-sustainability-an-emerging-food-policy-discourse/FC6900B6815969B0ACC9E93FD571257A
25. Santé Publique France. Usage regulation for the “Nutri-Score” logo [Internet]. 2019 [cited 2020 Jun 29]. Available from: <https://www.santepubliquefrance.fr/media/files/02-determinants-de-sante/nutrition-et-activite-physique/nutri-score/reglement-usage-en>
 26. IPCC. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, et al., editors. Cambridge, UK and New York, NY, USA: Cambridge University Press; 2013.
 27. Water Use in Life Cycle Assessment Group (WULCA). AWaRE (Available Water Remaining) Method [Internet]. 2016 [cited 2020 Jun 29]. Available from: <http://www.wulca-waterlca.org/aware.html>
 28. Guinée JB, Gorrée M, Heijungs R, Huppes G, Kleijn R, de Koning A, et al. Part III: Scientific background. In: Guinée JB, editor. Handbook on life cycle assessment Operational guide to the ISO standards. Dordrecht: Kluwer Academic Publishers; 2002.
 29. IMPACT World+ [Internet]. [cited 2020 Jun 29]. Available from: <http://www.impactworldplus.org>
 30. Jolliet O, Margni M, Charles R, Humbert S, Payet J, Rebitzer G, et al. IMPACT 2002 +: A New Life Cycle Impact Assessment Methodology. *Int J LCA*. 2003;8(6):324–30.
 31. Humbert S, De Schryver A, Bengoa X, Margni M, Jolliet O. IMPACT 2002+: User Guide. Draft for version Q2.21 [Internet]. 2012. Available from: http://www.quantis-intl.com/pdf/IMPACT2002_UserGuide_for_vQ2.21.pdf
 32. eurostat. Municipal waste by waste management operations. Updated on 24 February 2020 [Internet]. [cited 2020 Jun 29]. Available from: https://ec.europa.eu/eurostat/web/products-datasets/-/env_wasmun
 33. ADEME. Emballages ménagers. Données 2013. Faits et Chiffres [Internet]. 2014 [cited 2020 Jun 29]; Available from: <https://www.ademe.fr/emballages-menagers-donnees-2013>
 34. CITEO. La france de l'économie circulaire. Rapport annuel 2018 [Internet]. Paris; 2018 [cited 2020 Jun 29]. Available from: https://bo.citeo.com/sites/default/files/2019-07/CITEO_RA2018_web.pdf
 35. Zhang D, Shen J, Zhang F, Li Y, Zhang W. Carbon footprint of grain production in China. *Sci Rep* [Internet]. Nature Publishing Group; 2017 Dec 1 [cited 2020 Jul 3];7(1):1–11. Available from: www.nature.com/scientificreports
 36. Schmidt JH. United Plantation LCA Report [Internet]. 2014 [cited 2020 Jul 3]. Available from: [http://unitedplantations.com/About/United Plantation LCA Report.pdf](http://unitedplantations.com/About/United%20Plantation%20LCA%20Report.pdf)
 37. Spinelli D, Jez S, Basosi R. Integrated Environmental Assessment of sunflower oil production. In: *Process Biochemistry*. Elsevier; 2012. p. 1595–602.
 38. Bhatt R, Kukal SS, Busari MA, Arora S, Yadav M. Sustainability issues on rice–wheat cropping system. Vol. 4, International Soil and Water Conservation Research. International Research and Training Center on Erosion and Sedimentation and China Water and Power Press; 2016. p. 64–74.

39. Gan Y, Liang C, Chai Q, Lemke RL, Campbell CA, Zentner RP. Improving farming practices reduces the carbon footprint of spring wheat production. *Nat Commun* [Internet]. Nature Publishing Group; 2014 Nov 18 [cited 2020 Jul 3];5(1):1–13. Available from: www.nature.com/naturecommunications
40. Palhares JCP, Pezzopane JRM. Water footprint accounting and scarcity indicators of conventional and organic dairy production systems. *J Clean Prod*. Elsevier Ltd; 2015 Apr 15;93:299–307.
41. Nouri H, Stokvis B, Galindo A, Blatchford M, Hoekstra AY. Water scarcity alleviation through water footprint reduction in agriculture: The effect of soil mulching and drip irrigation. *Sci Total Environ*. Elsevier B.V.; 2019 Feb 25;653:241–52.
42. Vijay V, Pimm SL, Jenkins CN, Smith SJ. The Impacts of Oil Palm on Recent Deforestation and Biodiversity Loss. Anand M, editor. *PLoS One* [Internet]. Public Library of Science; 2016 Jul 27 [cited 2020 Jul 3];11(7):e0159668. Available from: <https://dx.plos.org/10.1371/journal.pone.0159668>
43. Gallo A, Accorsi R, Baruffaldi G, Manzini R. Designing Sustainable Cold Chains for Long-Range Food Distribution: Energy-Effective Corridors on the Silk Road Belt. *Sustainability* [Internet]. MDPI AG; 2017 Nov 8 [cited 2020 Jul 3];9(11):2044. Available from: <http://www.mdpi.com/2071-1050/9/11/2044>
44. Svanes E, Oestergaard S, Hanssen O. Effects of Packaging and Food Waste Prevention by Consumers on the Environmental Impact of Production and Consumption of Bread in Norway. *Sustainability* [Internet]. MDPI AG; 2018 Dec 21 [cited 2020 Jul 3];11(1):43. Available from: <http://www.mdpi.com/2071-1050/11/1/43>
45. O'Brien D, Capper JL, Garnsworthy PC, Grainger C, Shalloo L. A case study of the carbon footprint of milk from high-performing confinement and grass-based dairy farms. *J Dairy Sci*. Elsevier; 2014 Mar 1;97(3):1835–51.
46. Grant CA, Hicks AL. Comparative life cycle assessment of milk and plant-based alternatives. *Environ Eng Sci* [Internet]. Mary Ann Liebert Inc.; 2018 Nov 1 [cited 2020 Jul 3];35(11):1235–47. Available from: <https://www.liebertpub.com/doi/abs/10.1089/ees.2018.0233>
47. Winans KS, Macadam-Somer I, Kendall A, Geyer R, Marvinney E. Life cycle assessment of California unsweetened almond milk. *Int J Life Cycle Assess* [Internet]. Springer; 2020 Mar 1 [cited 2020 Jul 3];25(3):577–87. Available from: <https://doi.org/10.1007/s11367-019-01716-5>
48. Islam S, Ponnambalam SG, Lam HL. Review on life cycle inventory: methods, examples and applications. *J Clean Prod*. Elsevier Ltd; 2016 Nov 10;136:266–78.
49. Kramer GFH, Martinez E V, Espinoza-Orias ND, Cooper KA, Tyszler M, Blonk H. Comparing the Performance of Bread and Breakfast Cereals, Dairy, and Meat in Nutritionally Balanced and Sustainable Diets. *Front Nutr* [Internet]. Frontiers Media S.A.; 2018 Jun 7 [cited 2020 Jun 29];5:51. Available from: <https://www.frontiersin.org/article/10.3389/fnut.2018.00051/full>
50. Wallén A, Brandt N, Wennersten R. Does the Swedish consumer's choice of food influence greenhouse gas emissions? *Environ Sci Policy*. Elsevier; 2004 Dec 1;7(6):525–35.
51. Mertens E, Kaptijn G, Kuijsten A, van Zanten H, Geleijnse JM, van 't Veer P. SHARP-Indicators Database towards a public database for environmental sustainability. *Data Br* [Internet]. Elsevier Inc.; 2019 Dec 1 [cited 2020 Jun 29];27. Available from: <https://pubmed.ncbi.nlm.nih.gov/31656843/>

52. Masset G, Soler L-G, Vieux F, Darmon N. Identifying Sustainable Foods: The Relationship between Environmental Impact, Nutritional Quality, and Prices of Foods Representative of the French Diet. *J Acad Nutr Diet* [Internet]. 2014 Jun 4 [cited 2014 May 15];114(6):862–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24703928>
53. Clark MA, Springmann M, Hill J, Tilman D. Multiple health and environmental impacts of foods. *Proc Natl Acad Sci U S A* [Internet]. National Academy of Sciences; 2019 Nov 12 [cited 2020 Jul 3];116(46):23357–62. Available from: <https://www.pnas.org/content/116/46/23357>
54. CIHEAM, FAO. Mediterranean food consumption patterns: diet, environment, society, economy and health. A White Paper Priority 5 of Feeding Knowledge Programme [Internet]. Rome; 2015 [cited 2020 Jul 3]. Available from: <http://www.fao.org/3/a-i4358e.pdf>
55. Food Packaging Trends - UK - March 2018 - Market Research Report [Internet]. [cited 2020 Jul 6]. Available from: <https://reports.mintel.com/display/858877/>
56. Petrescu DC, Vermeir I, Petrescu-Mag RM. Consumer Understanding of Food Quality, Healthiness, and Environmental Impact: A Cross-National Perspective. *Int J Environ Res Public Health* [Internet]. MDPI AG; 2019 Dec 25 [cited 2020 Jul 6];17(1):169. Available from: <https://www.mdpi.com/1660-4601/17/1/169>
57. Asioli D, Aschemann-Witzel J, Caputo V, Vecchio R, Annunziata A, Næs T, et al. Making sense of the “clean label” trends: A review of consumer food choice behavior and discussion of industry implications. Vol. 99, *Food Research International*. Elsevier Ltd; 2017. p. 58–71.
58. Edwards-Jones G. Does eating local food reduce the environmental impact of food production and enhance consumer health? *Proc Nutr Soc* [Internet]. 2010 Nov [cited 2013 May 22];69(4):582–91. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20696093>
59. Aschemann-Witzel J. Consumer perception and trends about health and sustainability: Trade-offs and synergies of two pivotal issues. Vol. 3, *Current Opinion in Food Science*. Elsevier Ltd; 2015. p. 6–10.
60. Heller MC, Keoleian GA, Willett WC. Toward a life cycle-based, diet-level framework for food environmental impact and nutritional quality assessment: A critical review. *Environ Sci Technol* [Internet]. American Chemical Society; 2013 Nov 19 [cited 2020 Jul 6];47(22):12632–47. Available from: <https://pubs.acs.org/sharingguidelines>
61. Julia C, Etilé F, Hercberg S. Front-of-pack Nutri-Score labelling in France: an evidence-based policy [Internet]. Vol. 3, *The Lancet Public Health*. Elsevier Ltd; 2018 [cited 2020 Jul 3]. p. e164. Available from: <https://pubmed.ncbi.nlm.nih.gov/29483002/>
62. Julia C, Hercberg S. Nutri-Score: evidence of the effective-ness of the French front-of-pack nutrition label. *Ernahrungs Umschau*. 2017;64(12):181–7.
63. Finkelstein EA, Ang FJL, Doble B, Wong WHM, van Dam RM. A Randomized Controlled Trial Evaluating the Relative Effectiveness of the Multiple Traffic Light and Nutri-Score Front of Package Nutrition Labels. *Nutrients* [Internet]. MDPI AG; 2019 Sep 17 [cited 2020 Jul 3];11(9):2236. Available from: <https://www.mdpi.com/2072-6643/11/9/2236>

64. McColl K, Lobstein T, Brinsden H. Nutrient profiling could be used to transform food systems and support health-promoting food policies. *Public Heal Panor.* 2017;3(4).
65. Leach AM, Emery KA, Gephart J, Davis KF, Erisman JW, Leip A, et al. Environmental impact food labels combining carbon, nitrogen, and water footprints. *Food Policy.* Elsevier Ltd; 2016 May 1;61:213–23.
66. European Commission. Single Market for Green Products - The Product Environmental Footprint Pilots - Environment - European Commission [Internet]. [cited 2020 Jul 6]. Available from: https://ec.europa.eu/environment/eussd/smgp/ef_pilots.htm
67. European Commission. Single Market for Green Products - Communication - Environmental impacts - Environment - European Commission [Internet]. [cited 2020 Jul 6]. Available from: <https://ec.europa.eu/environment/eussd/smgp/communication/impact.htm>

Figures




Figure 1

Scope of the study and boundaries of the breakfast cereals product system.




Figure 2

Changes in the nutritional composition of the products studied over the 2003-2018 period




Figure 3

Changes in the Nutri-Score rating of the products studied over the 2003-2018 period




Figure 4

Changes in environmental indicators of the products studied over the 2003-2018 period




Figure 5

Contribution of each LCA stage on the overall environmental indicator (data for the average product, n=3)
 CC: Climate change; ARD: Abiotic resource depletion; LUIB: Land use impact on biodiversity; IEEQ: Impact on ecosphere/ecosystems quality; FWCS: Freshwater consumption scarcity.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplement.docx](#)