Early High-Dose Continuous Veno-Venous Hemofiltration Alleviates the Alterations of Cd4+ T Lymphocyte Subsets in Septic Patients Combined With Acute Kidney Injury

Beibei Jin

First Affiliated Hospital of Dalian Medical University

Da Cao

Southeast University Zhongda Hospital

Ning Yang

First Affiliated Hospital of Dalian Medical University

Ling Wang

First Affiliated Hospital of Dalian Medical University

Ruifang Li

First Affiliated Hospital of Dalian Medical University

Xiumei Liu

First Affiliated Hospital of Dalian Medical University

Ping Gong (✉ gongp828@sina.cn)

First Affiliated Hospital of Dalian Medical University <https://orcid.org/0000-0001-9910-7948>

Research

Keywords: Sepsis, CD4+ T lymphocyte subsets, Continuous veno-venous hemofiltration, Patient, Sequential organ failure assessment score

Posted Date: April 16th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-409007/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Artificial Organs on February 7th, 2022. See the published version at <https://doi.org/10.1111/aor.14199>.

Abstract

Background

This study aims to observe the changes in CD4⁺ T lymphocyte subsets in septic patients combined with acute kidney injury and the effect of early high-dose CVVH on CD4⁺ T lymphocyte subsets.

Methods

Enrolled septic patients combined with acute kidney injury were randomly divided into CVVH ($n = 50$) and conventional treatment (non-CVVH, $n = 53$) groups. Healthy volunteers ($n = 21$) were enrolled. CVVH was initiated within 12 h of intensive care unit (ICU) admission with the doses of 35 ~ 60 mL/kg/h and maintained for at least 72 h. Th1, Th2, Th17 and T_{reg} were measured by flow cytometry on days 1, 3 and 7 of ICU admission. Sequential organ failure assessment (SOFA) scores were calculated.

Results

Th1 percentages and Th1/Th2 ratios were lower, and Th2, Th17 and T_{reg} percentages and Th17/T_{reg} ratios were higher in patients with sepsis, when compared to healthy volunteers (all $P < 0.05$). Early high-dose CVVH treatment significantly increased Th1 percentages and Th1/Th2 ratios, and significantly decreased Th2, Th17 and T_{reg} percentages and Th17/T_{reg} ratios, when compared to non-CVVH treatment (all $P < 0.05$). Th1 percentages and Th1/Th2 ratios were negatively correlated with SOFA scores, while Th2, Th17 and T_{reg} percentages and Th17/T_{reg} ratios were positively correlated with SOFA scores (all $P < 0.05$). Patients with sepsis in the CVVH group had significantly lower SOFA scores on day 7 of ICU admission and a shorter ICU stay when compared to those in the non-CVVH group ($P < 0.05$). However, the difference in 28-day mortality was not significant between the CVVH and non-CVVH groups (22.0% vs. 37.7%, $P = 0.104$).

Conclusion

Septic patients combined with acute kidney injury exhibit the different alterations of CD4⁺ T lymphocyte subsets. Early high-dose CVVH treatment could alleviate the alterations, which may be one of factors associated with the improvement of sepsis severity, but do not significantly decrease the 28-day mortality.

Background

Sepsis is defined as a life-threatening organ dysfunction caused by a dysregulated host response to infection [1]. It is one of the leading causes of death in the intensive care unit (ICU). In spite of many therapeutic principles recommended by the international guidelines for the management of sepsis and septic shock, sepsis remains as a major health problem worldwide, with an ICU mortality rate of 25.8% in 84 countries in 2012 [2, 3].

In sepsis, the immune response initiated by an invading pathogen fails to return to homeostasis, which culminates in a pathological syndrome characterized by sustained excessive inflammation and immune suppression [4]. Sepsis has been considered as a result of a complex network of events that involve immune-inflammatory and anti-inflammatory processes, which contribute to the clearance of infection and tissue recovery, or organ injury [4, 5]. Recent studies have indicated that immune dysregulation (immune excess or suppression) might play a key role in the development of sepsis, and this dysregulation is closely associated with CD4⁺ T lymphocytes during both the early and late stage of sepsis [5]. Upon different stimulation in the microenvironment, naïve CD4⁺ T lymphocytes can be induced to differentiate into specific subtypes, such as T helper (Th) 1, Th2, Th17 and regulatory T (T_{reg}) cells. The effector functions of these cells are mediated by cytokines secreted by differentiated cells [6]. Th1 cells mainly secrete IFN- γ , TNF- β and IL-2, supporting cell-mediated immunity [6]. Th2 cells secrete the key effector-cytokines, such as IL-4, IL-5 and IL-13, and Th2 cells, mounting immune response to extracellular parasites and playing major roles in the induction and persistence of certain allergic diseases [6]. Th17 cells secrete key effector cytokines, including IL-17A, IL-17F, IL-21 and IL-22, and have a crucial role on the inflammatory response against parasites, and extracellular and fungal pathogens [7]. T_{reg} cells balance T lymphocyte homeostasis, activation and function via numerous different mechanisms, including the secretion of IL-10, TGF- β and IL-35 [4].

Clearly, CD4⁺ T lymphocytes represent a unique branch of the adaptive immune system, which is crucial for achieving a regulated effective immune response to pathogens, and their proper functioning is vital for the survival of patients infected with invading pathogens [6]. Higher Th1 and Th17 differentiation have been found to be associated with higher survival rate in severe sepsis [8]. However, pro-inflammatory Th1 and Th17 cells were found to be suppressed during human experimental endotoxemia [9], and the inversed proportion of Th17/T_{reg} may be associated with the aggravation of patients with sepsis [10]. Therefore, the adjustment of CD4⁺ T subtype may be one of promising options for the treatment of sepsis.

The direct removal of inflammatory mediators from circulation appears to be intuitively attractive, since this could efficiently interrupt the inflammatory cascade, affect the differentiation of CD4⁺ T subtypes, and attenuate inflammation. Continuous blood purification is a pivotal treatment strategy for renal failure in the intensive care setting [3, 11-13]. However, recent studies revealed that in addition to the removal of waste products, other molecules, such as inflammatory mediators/cytokines, were also the targets of removal in continuous blood purification, especially high-dose (namely high-volume) hemofiltration, for patients with sepsis [11, 12, 14-21]. Continuous blood purification has been generally applied for the treatment of sepsis as a strategy for the continuous removal of inflammatory mediators [3, 13]. However, to date, the effect of continuous venous-venous hemofiltration (CVVH), as a commonly used continuous blood purification mode, on the alterations of CD4⁺ T lymphocyte subsets in patients with sepsis, remains unclear. The present prospective study observed the changes in CD4⁺ T lymphocyte subsets in septic patients combined with acute kidney injury, and the effect of early high-dose CVVH on CD4⁺ T lymphocyte subsets alterations, disease severity and mortality.

Materials And Methods

Study design and population

The present study was conducted in accordance with the Declaration of Helsinki (2013 edition) adopted by the World Medical Association [22]. The study protocol was approved by the Ethics Committee of the First Affiliated Hospital of Dalian Medical University (PJ-KY-2018-12), Dalian, China. A written informed consent was obtained from all participants or legal guardians.

A total of 135 consecutive septic patients combined with acute kidney injury, who were admitted to the emergency ICU of our hospital from September 2016 to December 2018 and had the indication of CVVH treatment, were selected as subjects. The enrolled patients with sepsis were randomly divided into two groups: CVVH group, patients who received CVVH treatment; conventional treatment (non-CVVH) group. A total of 21 healthy volunteers were enrolled.

Inclusion and exclusion criteria

Patients with sepsis or septic shock were included in the present study if they met the requirement of the third international consensus definitions for sepsis and septic shock (Sepsis-3), accompanied by sepsis-induced acute kidney injury (diagnosed according to the KDIGO clinical practice guidelines) with increase in serum creatinine of more than 300 $\mu\text{mol/L}$, oliguria ($<100 \text{ mL/6 h}$) unresponsive to fluid resuscitation, severe acidemia of less than pH of 7.2, hyperkalemia of more than 6.5 mmol/L , or clinically significant organ edema (e.g., pulmonary edema) [1, 11, 23-25].

Patients were excluded from the present study based on the following criteria: patients < 18 years old; patients with immunodeficiency virus infection, neutropenia (defined as an absolute neutrophil count of $< 1,000$ neutrophils/ μL), acute bleeding, malignant tumor, pregnancy, or chronic kidney disease; patients who used corticosteroids or nephrotoxic drugs; patients who received any kind of renal replacement therapy prior to admission to the ICU; patients who refused to take part in the present study.

Treatments of the patients

The conventional therapy was based on standard sepsis treatments and according to Surviving Sepsis Campaign guidelines, which included intensive monitoring, fluid resuscitation, oxygen administration or mechanical ventilation, antimicrobial therapy (blood culture was conducted before the administration of antibiotics), vasopressor administration, glucose control, diuretics for oliguria with fluid overload, etc [3].

For the CVVH treatment, a 11.5Fr double lumen hemofiltration catheter (Lily Technology, Guangdong, China) was percutaneously inserted into the femoral vein. Hemofiltration was performed using the PrimaFlex apparatus (ALPRI, France) equipped with an acrylonitrile and sodium methallyl sulfonate copolymer + polyethylene (AN69ST) hollow-fiber high-flux hemofilter (1.5 m^2 , Braun Diapact, Germany), and was delivered in CVVH mode. The extracorporeal circuit and hemofilter were pre-flushed with heparinized normal saline (12500 U of unfractionated heparin added in 1000 mL of normal saline) to

prevent clotting. After circulation, the heparinized normal saline was flushed out of the hemofilter and circuit tubes prior to starting CVVH. Continuous intravenous pumping of unfractionated heparin (4 to 8 U/kg/h) was used to adjust the systemic activated partial thromboplastin time (APTT) between 35 and 45 s [26]. However, heparin-free CVVH was implemented if a patient had a bleeding tendency (e.g. low platelet count, prolonged activated partial thromboplastin time). CVVH was initiated within 12 h of ICU admission and maintained for at least 72 h with the doses of 35 ~ 60 mL/kg/h, and a blood flow rate of 150 ~ 200 mL/min. CVVH were stopped on basis of renal recovery and physician decision [24] The hemofilter was used till 72 h, or immediately changed if it did not work effectively because of clotting. The bicarbonate replacement solutions were commercially prepared by Qingshan Likang, Pharmaceutical Co. (Chengdu, China), and was pre-hemofilter diluted.

Variables

The following variables were collected for each patient on admission: age, sex, infection sites, results of blood culture, laboratory data (plasma creatinine, lactate, procalcitonin, leukocyte count, hemoglobin, platelet count and C-reactive protein), mean arterial pressure, number of patients using vasopressors, time from sepsis onset to ICU admission and number of patients using mechanical ventilation. On days 1, 3 and 7 of ICU admission, CD4⁺ T lymphocyte subsets (Th1, Th2, Th17 and T_{reg}) were measured, and Th1/Th2 and Th17/T_{reg} ratios were calculated; meantime, sequential organ failure assessment (SOFA) scores were evaluated.

Flow cytometry analysis

Two mL of peripheral venous blood was taken into EDTA-K3 coated tubes. The blood sample was first stimulated for 5 h with a cell stimulation cocktail (Sigma, St. Louis, MO, USA), containing 50 ng/mL of protein transport inhibitors, 1 mg/mL of ionomycin and 1.7 mg/mL of monensin, according to manufacturer's instructions.

T lymphocyte subsets, including Th1 (CD3⁺CD8-IFN- γ ⁺), Th2 (CD3⁺CD8-IL-4⁺), Th17 (CD4⁺IL-17⁺) and T_{reg} (CD4⁺CD25⁺Fox3⁺), were real-timely measured by flow cytometry (FACSVantage SE, BD Biosciences, San Jose, CA, USA). For the Th1 and Th2 analysis, the whole blood cells were extracellularly stained with FITC anti-human CD8 and PE-cy5 anti-human CD3 (BD Biosciences, San Jose, CA, USA) at 4°C for 30 min to determine the T lymphocyte population. Afterwards, the red blood cells in the whole blood were lysed using the FACS lysing solution (BD Biosciences, San Jose, CA, USA), and centrifuged (2,000×g) for five minutes. Then, the supernatant was abandoned, and the cell pellet was washed with phosphate buffered saline (PBS) and permeabilized with the Fix & Perm Reagent (BD Biosciences, San Jose, CA, USA). The permeabilized cells were further intracellular stained with either PE anti-human INF- γ antibody (Abcam, Cambridge, MA, USA) for the detection of Th1 cells (Fig.1), or PE anti-human IL-4 antibody (Abcam, Cambridge, MA, USA) for the detection of Th2 cells (Fig.1).

For the Th17 and T_{reg} analysis, the stimulated peripheral venous blood was extracellularly stained with the following combinations of antibodies: PE-cy5-conjugated anti-CD4 (Abcam, Cambridge, MA, USA) and PE anti-human IL-17 (BD Biosciences, San Jose, CA, USA) for Th17 (Fig.1); PE-cy5-conjugated anti-CD4 (Abcam, Cambridge, MA, USA), FITC-conjugated anti-CD25, and PE-conjugated anti-Fox3 (Abcam, Cambridge, MA, USA) for T_{reg} (Fig.1).

Statistical analysis

Data were analyzed using SPSS 22.0 (IBM, Armonk, NY, USA). Data were expressed as mean $\pm$ standard deviation (SD) for normal distribution, or median and interquartile range for skewed distribution. Pearson's chi-squared or Fisher's exact test was used, as appropriate, to compare the categorical variables. A Mann-Whitney *U* test was used to compare the variables between the CVVH and non-CVVH groups. Variables among time points and groups were compared using repeated-measures analysis of variance (ANOVA), followed by Bonferroni test for multiple comparisons. The differences were considered statistically significant when $P < 0.05$.

Results

Baseline characteristics of the enrolled subjects

A total of 135 eligible patients with sepsis or septic shock met the inclusion criteria (Fig.2). There were no significant differences between the CVVH ($n = 50$) and non-CVVH groups ($n = 53$) with regard to age, sex, infection sites, positive blood cultures, plasma creatinine, lactate and procalcitonin, leukocyte count, hemoglobin, platelet count, C-reactive protein, mean arterial pressure, the number of patients using vasopressors, time from sepsis onset to ICU admission and the number of patients receiving mechanical ventilation (all $P > 0.05$; Table 1). CVVH started at 5.5 (2.0, 8.5) hours after ICU admission. The parameters of CVVH were provided in Table 2.

The changes of T lymphocyte subsets in patients with sepsis on ICU admission

Patients with sepsis (both non-CVVH and CVVH groups) showed significantly lower Th1 percentages and Th1/Th2 ratios, and significantly higher Th2, Th17 and T_{reg} percentages and Th17/T_{reg} ratios on admission, when compared to healthy volunteers (all $P < 0.05$; Fig.3). However, the differences in the above variables on admission were not significant between the non-CVVH and CVVH groups (all $P > 0.05$).

The effect of high-dose CVVH treatment on Th1, Th2 and the Th1/Th2 ratios in patients with sepsis

After the one-week conventional therapy in the non-CVVH group, the low Th1 percentages and Th1/Th2 ratios were gradually improved, while the high Th2 percentages were gradually decreased (all $P < 0.05$; Fig. 3A, 3B and 3C). On day 7 of ICU admission, high-dose CVVH treatment exhibited more significant improvement in Th1 percentages and Th1/Th2 ratios, and more significant decrease in Th2 percentages

compared with non-CVVH treatment (all $P < 0.05$). However, on day 3 of ICU admission, the differences in the above variables were not significant between the non-CVVH and CVVH groups (all $P > 0.05$).

The effect of high-dose CVVH treatment on Th17, T_{reg} and Th17/T_{reg} ratios in patients with sepsis

After the one-week conventional therapy in the non-CVVH group, the increased Th17 and T_{reg} lymphocyte percentages and Th17/T_{reg} ratios could be efficiently reversed in patients with sepsis. On day 7 of ICU admission, high-dose CVVH treatment exhibited a more efficient effect on the reversion of these T lymphocyte subsets when compared to non-CVVH treatment (all $P < 0.05$; Fig. 3D, 3E and 3F). However, on day 3 of ICU admission, the differences in the above variables were not significant between the non-CVVH and CVVH groups (all $P > 0.05$).

The correlation between CD4⁺ T lymphocyte subsets and SOFA scores in patients with sepsis

The Spearman's correlation analysis indicated that the Th1 percentages and Th1/Th2 ratios were negatively correlated with the SOFA scores, while the Th2, Th17 and T_{reg} percentages, and Th17/T_{reg} ratios were positively correlated with the SOFA scores (all $P < 0.05$; Table 3).

The effect of high-dose CVVH on the severity and outcome of patients with sepsis

Patients with sepsis in the CVVH group had significantly lower SOFA scores on day 7 of ICU admission, and a shorter ICU stay, when compared to those in the non-CVVH group (both $P < 0.05$; Table 4). Patients with sepsis in the CVVH group had a tendency to a lower 28-day mortality when compared to those in the non-CVVH group (22.0% vs. 37.7%; Table 4), but the difference was not significant ($P = 0.104$).

Discussion

In the present study, we observed that septic patients combined with acute kidney injury exhibited higher Th2 percentages, lower Th1 percentages and Th1/Th2 ratios in the first week of ICU admission, when compared to healthy volunteers. This observation is similar to previous works, showing a shift from Th1 to Th2 cytokine profiles following infection [27, 28]. Higher Th17, T_{reg} and Th17/T_{reg} ratios were also observed in patients with sepsis in the present study when compared to healthy volunteers, which is consistent with the reports of other studies [29, 30]. These alterations of CD4⁺ T lymphocyte subsets indicate that T_{reg} and Th17 cells are different differentiations from Th1 and Th2 cells [28, 31, 32].

Although the underlying mechanisms of CD4⁺ T lymphocyte subset differentiation remain under investigation, it was speculated that these different differentiations may play crucial roles in disease progression in patients with sepsis [6, 28].

In addition, we observed that Th1 percentages and Th1/Th2 ratios were negatively correlated with the SOFA scores, while Th2, Th17, T_{reg} percentages and Th17/T_{reg} ratios were positively correlated with the SOFA scores. These observations further indicated that the differentiation and different alterations of CD4⁺ T lymphocyte subsets may be associated with the severity of sepsis, which [induced an](#)

immunoparalysis state [6, 28, 33]. T lymphocyte subsets, especially CD4⁺ T helper cells, have been demonstrated to contribute to the advancement of inflammatory diseases and microbial challenges [4, 6, 29-31, 34, 35]. The lack of the shift from Th1 to Th2 has been found to increase the survival among patients with sepsis [4]. Inhibiting a pro-inflammatory Th1 to an anti-inflammatory Th2 shift and attenuating the T_{reg} population can improve survival in secondary pneumonia infections [28]. In the present study, it was found that after one week of ICU admission, the proportions of T lymphocyte subsets in patients were getting close to those in healthy volunteers, which indicates the reversion of the CD4⁺ T lymphocyte subsets after conventional therapy.

Surprisingly, the early high-dose CVVH treatment exhibited a more efficient effect on this T lymphocyte subset reversion when compared to non-CVVH treatment, which further illustrates the correlation between CVVH treatment and the significantly decreased SOFA scores of patients. The mechanisms underlying the more efficient reversion of early high-dose CVVH treatment to the alterations of CD4⁺ T lymphocyte subsets have not yet been completely clarified. We speculated that this effect of early high-dose CVVH may be associated with the removal of high-dose CVVH to inflammatory mediators/cytokines, endotoxin, antigens and other solutes, which affected the differentiation, apoptosis and functions of CD4⁺ T lymphocyte subsets [5, 12, 14, 15, 17, 27, 33, 36]. AN69ST, utilized in the present study, is an effective cytokine-adsorbing hemofilter, and can also remove endotoxin [14, 37]. Ionic interactions have been demonstrated to play a role in cytokines (e.g. TNF- α , IL-6, and IL-8) adsorption by the AN69ST membrane [37]. Previous studies have shown that high-dose CVVH effectively removed serum inflammatory cytokines also via the mechanism of convection [19, 21]. Furthermore, cutting the peaks of soluble mediators through continuous hemofiltration, namely, a concept for the peak concentration hypothesis, might partly explain for the more efficient reversion of CVVH treatment to the alterations of CD4⁺ T lymphocyte subsets [17]. Thus, CVVH has been considered as an efficient immunomodulatory therapy for patients with sepsis [36]; the effect of the non-specific removal of unbound mediators and cytokines from the blood compartment is to reset to a lower level of immune dysregulation, both in the prevalently pro-inflammatory and counter-inflammatory phases of sepsis [17].

It was not surprising that the 28-day mortality of patients with sepsis failed to be significantly decreased by high-dose CVVH treatment in spite of the significantly decreased SOFA scores, which is inconsistent with the meta-analysis published by Putzu *et al.*, stating that low-quality evidence indicates the correlation between CVVH and the significant reduction in mortality in patients with severe sepsis [3, 13]. There are several potential explanations for this phenomenon. First, the single-center design and relatively small sample size in this study may not be adequate to reach a statistically significant reduction of mortality after CVVH treatment. Hence, a multi-center randomized trial with a large sample size would be needed in future trials. Second, the differentiations and functions of T lymphocyte subsets are complicate and dynamic, which might partly contribute to the severity of patients with sepsis and be insufficient to affect the outcome. Finally, the potential adverse effects of hemofilter during CVVH, such as being an inflammatory stimulus itself and activating leukocytes during hemofilter passage, excessive loss of electrolytes, micronutrients, vitamins or drugs (e.g., antibiotics), maybe weaken the beneficial

effects [18, 38]. In fact, to date the effects of CVVH treatment on mortality of patients with sepsis remains controversial. A recent meta-analysis also showed that high-volume hemofiltration did not change the mortality of patients with sepsis [16].

The present study had six main limitations. First, we failed to detect the levels of cytokines in plasma and filtered solution such as IL-4, IL-6, IL-1ra, IL-8, IL-10, IL-12 and tumor necrosis factor- α that may affect the differentiation and functions of these CD4⁺ T lymphocyte subsets. Second, CVVH is not a standardized technique. Hence, the settings (e.g. CVVH doses) and system used in the present study could result in different results from other studies [13, 39]. Third, the differences in subgroups were not assessed on the basis of sex and septic shock. However, the sex distribution and the number of patients using vasopressors on admission in the CVVH and non-CVVH groups was even. Fourth, although 35 ~ 60 mL/kg/h (> 35 mL/kg/h generally accepted as the definition of high dose in CVVH therapy) of CVVH doses used in the present study is not enough high, several meta-analyses showed that higher doses had no clinical advantage for blood purification purposes in the treatment of patients with sepsis [11, 16, 20, 40]. Finally, the association between CVVH and lower SOFA scores might be biased by the fact that hemofiltration eliminates creatinine and bilirubin, which itself are parameters of the SOFA scores.

Conclusions

Overall, the present study indicates that septic patients combined with acute kidney injury exhibit the different alterations of CD4⁺ T lymphocyte subsets in the first week of ICU admission. Early high-dose CVVH treatment could significantly alleviate the alterations of CD4⁺ T lymphocyte subsets, which may be one of factors associated with the improvement of sepsis severity, but do not significantly decrease the 28-day mortality.

Declarations

Ethics approval and consent to participate

The present study was conducted in accordance with the Declaration of Helsinki (2013 edition) adopted by the World Medical Association. The study protocol was approved by the Ethics Committee of the First Affiliated Hospital of Dalian Medical University (PJ-KY-2018-12), Dalian, China. A written informed consent was obtained from all participants or legal guardians.

Consent for publication

All the authors agree for publication.

Availability of data and materials

The datasets generated and analyzed during this study are available from the corresponding author upon reasonable request.

Competing interests

The authors declare that they have no conflicts of interest to disclose.

Funding

This work was supported by the National Natural Science Foundation of China (81571869).

Authors' contributions

PG and BJ conceived and designed the experiments. BJ, DC, NY, LW and RL carried out the experiments. BJ and DC analyzed the data. BJ and PG wrote the manuscript. PG took overall responsibility for the manuscript. All authors approved the final version of the manuscript.

Acknowledgements

The authors deeply appreciate Yuejian Liu for his technical assistance and the staff of the Emergency Intensive Care Unit for their helpful contributions.

References

1. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA*. 2016 Feb;315(8):801-10.
2. Vincent JL, Marshall JC, Namendys-Silva SA, Francois B, Martin-Loeches I, Lipman J, et al. Assessment of the worldwide burden of critical illness: the intensive care over nations (ICON) audit. *Lancet Respir Med*. 2014 May;2(5):380-6.
3. Rhodes A, Evans LE, Alhazzani W, Levy MM, Antonelli M, Ferrer R, et al. Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock: 2016. *Crit Care Med*. 2017 Mar;45(3):486-552.
4. van der Poll T, van de Veerdonk FL, Scicluna BP, Netea MG. The immunopathology of sepsis and potential therapeutic targets. *Nat Rev Immunol*. 2017 Jul;17(7):407-20.
5. Hotchkiss RS, Monneret G, Payen D. Sepsis-induced immunosuppression: from cellular dysfunctions to immunotherapy. *Nat Rev Immunol*. 2013 Dec;13(12):862-74.
6. Luckheeram RV, Zhou R, Verma AD, Xia B. CD4(+)T cells: differentiation and functions. *Clin Dev Immunol*. 2012;2012:925135.
7. Miossec P, Korn T, Kuchroo VK. Interleukin-17 and type 17 helper T cells. *N Engl J Med*. 2009 Aug;361(9):888-98.
8. Wu HP, Chung K, Lin CY, Jiang BY, Chuang DY, Liu YC. Associations of T helper 1, 2, 17 and regulatory T lymphocytes with mortality in severe sepsis. *Inflamm Res*. 2013 Aug;62(8):751-63.

9. Brinkhoff A, Sieberichs A, Engler H, Dolff S, Benson S, Korth J, et al. Pro-Inflammatory Th1 and Th17 Cells Are Suppressed During Human Experimental Endotoxemia Whereas Anti-Inflammatory IL-10 Producing T-Cells Are Unaffected. *Front Immunol*. 2018 May;9:1133.
10. Li J, Li M, Su L, Wang H, Xiao K, Deng J, et al. Alterations of T helper lymphocyte subpopulations in sepsis, severe sepsis, and septic shock: a prospective observational study. *Inflammation*. 2015;38(3):995-1002.
11. Romagnoli S, Ricci Z, Ronco C. CRRT for sepsis-induced acute kidney injury. *Curr Opin Crit Care*. 2018 Dec;24(6):483-92.
12. Villa G, Zaragoza JJ, Sharma A, Neri M, De Gaudio AR, Ronco C. Cytokine removal with high cut-off membrane: review of literature. *Blood Purif*. 2014;38(3-4):167-73.
13. Putzu A, Fang MX, Boscolo Berto M, Belletti A, Cabrini L, Cassina T, et al. Blood purification with continuous veno-venous hemofiltration in patients with sepsis or ARDS: a systematic review and meta-analysis. *Minerva Anesthesiol*. 2017 Aug;83(8):867-77.
14. Broman ME, Hansson F, Vincent JL, Bodelsson M. Endotoxin and cytokine reducing properties of the oXiris membrane in patients with septic shock: A randomized crossover double-blind study. *PLoS One*. 2019 Aug;14(8):e0220444.
15. Guo J, Tao W, Tang D, Zhang J. Th17/regulatory T cell imbalance in sepsis patients with multiple organ dysfunction syndrome: attenuated by high-volume hemofiltration. *Int J Artif Organs*. 2017 Oct;40(11):607-14.
16. Yin F, Zhang F, Liu S, Ning B. The therapeutic effect of high-volume hemofiltration on sepsis: a systematic review and meta-analysis. *Ann Transl Med*. 2020 Apr;8(7):488.
17. Ronco C, Tetta C, Mariano F, Wratten ML, Bonello M, Bordoni V, et al. Interpreting the mechanisms of continuous renal replacement therapy in sepsis: the peak concentration hypothesis. *Artificial organs*. 2003 Sep;27(9):792-801.
18. Junhai Z, Beibei C, Jing Y, Li L. Effect of High-Volume Hemofiltration in Critically Ill Patients: A Systematic Review and Meta-Analysis. *Med Sci Monit*. 2019 May;25:3964-75.
19. Park JT, Lee H, Kee YK, Park S, Oh HJ, Han SH, et al. High-Dose Versus Conventional-Dose Continuous Venovenous Hemodiafiltration and Patient and Kidney Survival and Cytokine Removal in Sepsis-Associated Acute Kidney Injury: A Randomized Controlled Trial. *Am J Kidney Dis* Oct. 2016;68(4):599-608.
20. Miao H, Wang F, Xiong X, Wang C, Zhang Y. Clinical Benefits of High-Volume Hemofiltration in Critically Ill Pediatric Patients with Severe Sepsis: A Retrospective Cohort Study. *Blood Purif*. 2018;45(1-3):18-27.
21. Ning B, Ye S, Lyu Y, Yin F, Chen Z. Effect of high-volume hemofiltration on children with sepsis. *Transl Pediatr*. 2020 Apr;9(2):101-7.
22. Hellmann F, Verdi M, Schlemper BR, Jr., Caponi S. 50th anniversary of the Declaration of Helsinki: the double standard was introduced. *Arch Med Res*. 2014 Oct;45(7):600-1.

23. Thomas ME, Blaine C, Dawnay A, Devonald MA, Ftouh S, Laing C, et al. The definition of acute kidney injury and its use in practice. *Kidney Int.* 2015 Jan;87(1):62-73.
24. Atan R, Peck L, Prowle J, Licari E, Eastwood GM, Storr M, et al. A Double-Blind Randomized Controlled Trial of High Cutoff Versus Standard Hemofiltration in Critically Ill Patients With Acute Kidney Injury. *Crit Care Med.* 2018 Oct;46(10):e988-e94.
25. Khwaja A. KDIGO clinical practice guidelines for acute kidney injury. *Nephron Clin Pract.* 2012;120(4):c179-84.
26. van de Wetering J, Westendorp RG, van der Hoeven JG, Stolk B, Feuth JD, Chang PC. Heparin use in continuous renal replacement procedures: the struggle between filter coagulation and patient hemorrhage. *J Am Soc Nephrol.* 1996 Jan;7(1):145-50.
27. Salomao R, Brunialti MK, Rapozo MM, Baggio-Zappia GL, Galanos C, Freudenberg M. Bacterial sensing, cell signaling, and modulation of the immune response during sepsis. *Shock.* 2012 Aug;38(3):227-42.
28. Luan YY, Yao YM, Sheng ZY. Update on the immunological pathway of negative regulation in acute insults and sepsis. *J Interferon Cytokine Res.* 2012 Jul;32(7):288-98.
29. Gupta DL, Bhoi S, Mohan T, Galwnkar S, Rao DN. Coexistence of Th1/Th2 and Th17/Treg imbalances in patients with post traumatic sepsis. *Cytokine.* 2016 Dec;88:214-21.
30. Venet F, Chung CS, Kherouf H, Geeraert A, Malcus C, Poitevin F, et al. Increased circulating regulatory T cells (CD4(+)/CD25 (+)/CD127 (-)) contribute to lymphocyte anergy in septic shock patients. *Intensive Care Med.* 2009 Apr;35(4):678-86.
31. Basu R, Hatton RD, Weaver CT. The Th17 family: flexibility follows function. *Immunol Rev.* 2013 Mar;252(1):89-103.
32. Zhang Y, Zhang Y, Gu W, He L, Sun B. Th1/Th2 cell's function in immune system. *Adv Exp Med Biol.* 2014;841:45-65.
33. Martin MD, Badovinac VP, Griffith TS. CD4 T Cell Responses and the Sepsis-Induced Immunoparalysis State. *Front Immunol.* 2020 Jul;11:1364.
34. Yu W, Yao D, Yu S, Wang X, Li X, Wang M, et al. Protective humoral and CD4(+) T cellular immune responses of *Staphylococcus aureus* vaccine MntC in a murine peritonitis model. *Sci Rep.* 2018 Feb;8(1):3580.
35. Venet F, Pachot A, Debard AL, Bohe J, Bienvenu J, Lepape A, et al. Increased percentage of CD4+CD25+ regulatory T cells during septic shock is due to the decrease of CD4+CD25- lymphocytes. *Crit Care Med.* 2004 Nov;32(11):2329-31.
36. Honore PM, Joannes-Boyau O, Gressens B. Blood and plasma treatments: the rationale of high-volume hemofiltration. *Contrib Nephrol.* 2007;156:387-95.
37. Moriyama K, Kato Y, Hasegawa D, Kurimoto Y, Kawaji T, Nakamura T, et al. Involvement of ionic interactions in cytokine adsorption of polyethyleneimine-coated polyacrylonitrile and polymethyl methacrylate membranes in vitro. *J Artif Organs.* 2020 Sep;23(3):240-246.

38. Lehner GF, Harler U, Feistritz C, Haller VM, Hasslacher J, Bellmann R, et al. Hemofiltration induces generation of leukocyte-derived CD31+/CD41- microvesicles in sepsis. *Annals of intensive care*. 2017 Sep;7(1):89.

39. Park JT, Lee H, Kee YK, Park S, Oh HJ, Han SH, et al. High-Dose Versus Conventional-Dose Continuous Venovenous Hemodiafiltration and Patient and Kidney Survival and Cytokine Removal in Sepsis-Associated Acute Kidney Injury: A Randomized Controlled Trial. *Am J Kidney Dis*. 2016 Oct;68(4):599-608.

40. Lehner GF, Wiedermann CJ, Joannidis M. High-volume hemofiltration in critically ill patients: a systematic review and meta-analysis. *Minerva Anesthesiol*. 2014 May;80(5):595-609.

Tables

Table 1 Baseline Characteristics of the Enrolled Subjects

	Healthy	Non-CVVH	CVVH	<i>P</i>
	(<i>n</i> = 21)	(<i>n</i> = 53)	(<i>n</i> = 50)	
Age (years)	59 (53 – 65)	58 (54 – 66)	57 (54 – 63)	NS
Male [<i>n</i> (%)]	12 (57.1%)	31 (58.5%)	28 (56.0%)	NS
Infection sites [<i>n</i> (%)]				
Abdomen	—	16 (30.2%)	14 (28.0%)	NS
Lung	—	19 (35.8%)	19 (38.0%)	NS
Kidney	—	7 (13.2%)	7 (14.0%)	NS
Skin/soft tissue	—	6 (11.3%)	5 (10.0%)	NS
Neurological	—	5 (9.5%)	6 (12.0%)	NS
Positive blood culture	—	15 (28.3%)	16 (32.0%)	NS
Escherichia coli	—	7 (13.2%)	8 (16.0%)	NS
Klebsiella pneumoniae	—	4 (7.5%)	4 (8.0%)	NS
Staphylococcus aureus	—	2 (3.8%)	1 (2.0%)	NS
Staphylococcus epidermidis	—	1 (1.9%)	1 (2.0%)	NS
Enterococcus faecium	—	1 (1.9%)	1 (2.0%)	NS
Laboratory findings				
Creatinine (μmol/L)	57.8 (48.2 – 94.5)	265.1 (189.2 – 394.0)	348.2 (256.1 – 419.6)	NS
Lactate (mmol/L)	0.8 (0.5 – 1.6)	3.7 (2.3 – 5.6)	3.6 (2.8 – 5.2)	NS
Procalcitonin (ng/mL)	0.02 (0.01 – 0.03)	16.17 (2.38 – 51.72)	16.96 (2.87 – 62.78)	NS
Leukocyte count (×10 ⁹)	7.72 (4.55 – 8.90)	14.72 (10.55 – 18.20)	15.72 (10.35 – 18.27)	NS
Hemoglobin (g/L)	133.5 (115.1 – 142.3)	122.1 (105.3 – 143.6)	123.4 (103.5 – 135.2)	NS
Platelet count (×10 ⁹)	210.6 (140.8 – 242.8)	130.6 (91.3 – 212.7)	110.9 (82.7 – 220.4)	NS
C-reactive protein (mg/L)	4.8 (2.4, 6.5)	94.8 (62.2, 155.9)	98.3 (63.4, 159.6)	NS
Mean arterial pressure (mmHg)	88.5 (78.5 – 93.7)	67.5 (62.0 – 75.8)	69.3 (63.0 – 72.7)	NS
Patients using vasopressors on	—	10 (18.9%)	8 (18.0%)	NS

admission [<i>n</i> (%)]				
Time from sepsis onset to ICU admission (days)	—	3.3 (2.0 – 7.5)	3.4 (2.1 – 6.5)	NS
Mechanical ventilation [<i>n</i> (%)]	—	20 (37.7%)	15 (30.0%)	NS

Notes: The values are expressed median and interquartile. AKI, acute kidney injury; CVVH, continuous veno-venous hemofiltration; ICU, intensive care unit; NS, no significance; SOFA, sequential organ failure assessment.

Table 2 The Parameters of CVVH

Blood flow (mL/min)	180.5 (170.2, 190.1)
Unfractionated heparin (U/kg/h)	5.2 (4.5, 6.8)
CVVH doses (mL/kg/h)	47.1 (40.5, 53.2)
Hemofilter survival (h)	35.6 (20.5, 42.0)

Notes: Values are expressed as median and interquartile. CVVH, continuous veno-venous hemofiltration.

Table 3 The Correlation among CD4⁺ T Lymphocyte Subsets and SOFA Scores

	SOFA scores	
	<i>R</i>	<i>P</i>
Th1	- 0.403	<0.001
Th2	0.387	<0.001
Th1/Th2	- 0.501	<0.001
Th17	0.376	<0.001
Treg	0.459	<0.001
Th17/Treg	0.564	<0.001

Table 4 Comparison of Clinical Severity and Outcome between the CVVH and Non-CVVH Groups

	Non-CVVH (<i>n</i> = 53)	CVVH (<i>n</i> = 50)	<i>P</i>
SOFA scores			
Day 1 of ICU admission	12 (10 – 14)	12 (9 – 14)	NS
Day 3 of ICU admission	11 (9 – 13)	9 (6 – 11)	NS
Day 7 of ICU admission	8 (6 – 11)	5 (3 – 9)	< 0.05
28-day mortality [<i>n</i> (%)]	20 (37.7%)	11 (22.0%)	0.104
ICU stay (days)	19 (7 – 22)	12 (9 – 16)	< 0.05

Notes: Values are expressed as median and interquartile. CVVH, continuous veno-venous hemofiltration; ICU, intensive unit; NS, no significance; SOFA, sequential organ failure assessment.

Figures

Figure 1

The flow cytometry of CD4⁺ T lymphocyte subsets from healthy volunteers and patients with sepsis. Th1 cells were identified with positive CD3 and IFN-γ, and negative CD8. Th2 cells were identified with positive

CD3 and IL-4, and negative CD8. Th17 cells were identified with positive CD3, CD4 and IL-17. Regulatory T cells were identified with positive CD4 and high CD25, and Foxp3.

Figure 2

The flow diagram of the participants. CVVH, continuous veno-venous hemofiltration.

Figure 3

Comparison of Th1 (A) and Th2 (B) percentages, Th1/Th2 ratio (C), Th17 (D) and Treg (E) percentages, and Th17/Treg ratio (F) in the healthy volunteers (n=21), non-CVVH (n=53) and CVVH (n=50) groups. The percentages of Th1 or Th2 meant the percentages of Th1 or Th2 in T lymphocytes; the percentages of Th17 or Treg meant the percentages of Th17 or Treg in CD4+ T lymphocytes. Day 1, the first day of intensive care unit (ICU) admission; Day 3, the third day of ICU admission; Day 7, the seventh day of ICU admission; CVVH, continuous veno-venous hemofiltration. Data were expressed as median and interquartile range.