

Analysis of Spatially Extended Excitable Izhikevich Neuron Model near Instability

Arnab Mondal

Indian Institute of Technology (Indian School of Mines): Indian Institute of Technology

Argha Mondal

University of Essex

S. Sharma

Indian Institute of Technology (Indian School of Mines): Indian Institute of Technology

Ranjit Kumar Upadhyay (✉ ranjit.chaos@gmail.com)

Indian Institute of Technology (Indian School of Mines): Indian Institute of Technology

<https://orcid.org/0000-0002-1210-7804>

Research Article

Keywords: Izhikevich model, firing activity, bifurcation scenarios, amplitude equations, diffusive instabilities

Posted Date: May 17th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-410280/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Nonlinear Dynamics on August 11th, 2021. See the published version at <https://doi.org/10.1007/s11071-021-06787-4>.

Analysis of spatially extended excitable Izhikevich neuron model near instability

Arnab Mondal · Argha Mondal · Sanjeev Kumar Sharma · Ranjit Kumar Upadhyay*

Received: date / Accepted: date

Abstract The article focuses on the issue of a spatiotemporal excitable biophysical model that describes the propagation of electrical potential called spikes to model the diffusion induced dynamics based on an analytical development of amplitude equations. Considering the Izhikevich neuron model consisting of coupled systems of ODEs, we demonstrate various results of spatiotemporal architecture (PDEs) using a suitable parameter regime. We analytically perform the saddle node bifurcation and Hopf bifurcation analysis with bifurcating periodic solutions that show the transition phases in the system dynamics. We study different types of firing patterns both analytically and numerically by the formation of Riccati differential equation. To examine the characteristics of diffusive instabilities, we use Turing amplitude equations by multiscaling method and then expansion in powers of a small control parameter. The instabilities and Turing bifurcation are established using the theoretical analysis and numerical simulations. The spatial system has potential effects on the deterministic system as a result of the

diffusive matrices with various couplings and the coupled oscillators with this nearest neighbor coupling show synchronization measured by the synchronization factor analysis. Our results qualitatively reproduce different phenomena of the extended excitable system based with an efficient analytical scheme.

Keywords Izhikevich model, firing activity, bifurcation scenarios, amplitude equations, diffusive instabilities

1 Introduction

To understand the various functional mechanisms in different brain areas (especially in cortical region), we definitely need the theoretical modeling and numerical simulations with experimental studies [1, 17, 21, 23, 30]. There are a certain number of models for excitable neural systems that are capable to reproduce the observed results by mathematical analysis and that are also computationally simple [19, 20, 33]. The models can generate rich dynamics often found in real neurons for electrophysiological activities. When a neuron receives a certain stimulus, it produces action potential (electrical impulses) that is propagated along the axon of the neuron [18]. Single neurons produce a spike when they are in the vicinity of a bifurcation point from resting state to firing activity. The following results show how the dynamics can be investigated for different characteristics. We are excited to study an excitable spatiotemporal system that supports the wave propagation of nerve impulses [7, 36, 38]. From the mathematical point of view, the model consists of a coupled nonlinear reaction diffusion systems of partial differential equations (PDEs) modeling the propagation of electrical potential. It is derived from a system of coupled ordinary differential equations (ODEs) modeling the cell membrane dynamics. The implementation of the proper analyti-

R.K. Upadhyay (✉)

Department of Mathematics & Computing, Indian Institute of Technology (Indian School of Mines) Dhanbad, 826004, Jharkhand, India.
E-mail: ranjit.chaos@gmail.com

Arnab Mondal · Sanjeev Kumar Sharma

Department of Mathematics & Computing, Indian Institute of Technology (Indian School of Mines) Dhanbad, 826004, Jharkhand, India.

Argha Mondal

Normandie Univ, UNIHAVRE, LMAH, FR-CNRS-3335, ISCN, 76600 Le Havre, France
Department of Mathematical Sciences, University of Essex, Wivenhoe Park, UK

School of Engineering, Amrita Vishwa Vidyapeetham, Amritapuri, Kollam 690525, India

* Corresponding author

cal and efficient numerical techniques is important to study such spatially extended systems.

Motivated by some recent articles, that were investigated on spatiotemporal patterns away from the equilibriums, traveling waves [6], spiral waves which frequently occur in biological, chemical and other physical systems [11, 13, 28, 29, 41, 42, 46]. We introduce the solution of the general amplitude equations and prove its well-posedness (for Hopf bifurcation [10, 12, 26, 27] and Turing instabilities [40, 45]). In our study, we consider zero flux boundary condition to show that the excitable system rest in an isolated condition [32]. Then, we will examine the corresponding system. We are interested to determine the variations in the firing potential as well as the stability analysis of the spatial system.

In this article, we are concerned with the analysis of spatiotemporal dynamics in the Izhikevich model [19, 20, 35] that plays a major role in the study of the cortical neurons. The famous mathematician and neuroscientist E. M. Izhikevich proposed an excitable two dimensional (2D) dynamical model that reproduces spiking-bursting activities [3, 20, 33, 39]. Here, we use this simple model that is biologically plausible and computationally efficient. The stability analysis can be observed from bifurcation scenarios where the predominant parameter is I (injected current stimulus). Then, we focus on the local nonlinear excitations in diffusively coupled Izhikevich model for different parameter regimes. Different parameter regimes corresponding to qualitatively diverse biophysical dynamics are investigated. Bifurcation scenarios have been widely used to analyze the dynamical responses that are modulated by uncoupled single model and coupled systems with various injected inputs such as synaptic coupling and current stimulus. We consider three firing regimes: fast spiking, tonic spiking, phasic spiking and verify the firing activities analytically using the formation of Riccati differential equation [14, 31]. Bifurcations are performed to show the transition phases of oscillations. Moreover, the dynamics behind it can generally be described through a phase diagram. We present a methodology to characterize the biophysics of this system, which takes into account the temporal structures of complex firings. This provides us to accurately distinguish the fundamental characteristics of the intrinsic dynamics and the modeling approaches enriches the spiking network level activities such as synchronization and its measure with synchronization factor [8, 25, 44]. For Hopf and Turing instabilities, responsible for close to the homogeneous fluctuations and the emergence of stationary and spatial patterns respectively, the corresponding amplitude equations are referred to as Turing amplitude equation (TAE) [43, 47] and complex Ginzburg-Landau equation (CGLE, it is responsible for the onset of Hopf instability in the occurrence of homogeneous fluctuations) [2, 22, 37].

We used TAE equation to get the insight into the nonequilibrium phenomenon in spatiotemporal systems. The analy-

sis scheme [9, 22, 37, 47] for the derivation of amplitude equations is based on the multiple space and time scales. The method provides a Taylor series expansion of the original nonlinear equations with many power operators and further it depends on the expansion of the linear and nonlinear terms of a small perturbation parameter close to the onset of diffusive instability. The small parameter measures the deviation from the diffusive instability region. It is used in our case as the reaction diffusion system has a diagonal diffusion matrices. The derivation of the procedure based on Kuramoto's method has been investigated to the systems. The coefficients of the amplitude equations obtained in the present article, may have their own values and it may consider to certain dynamical behavior in the diffusive system of experimental conditions. However, a systematic simulation is required to capture the dynamics of excitable models at a sufficient level of accuracy. The numerical solution of the coupled systems formed by the coupled nonlinear PDEs requires the suitable discretization scheme. The effects of diffusion in a biophysical extended system and then collective dynamical formation have been studied. The results show that the reaction diffusion system with different coupling characteristics that depend on the control parameters, participate in a collective behavior with diffusive instabilities.

The paper is organized as follows: In Section 2, we briefly describe the dynamics of the excitable model equations that shows the electrical potential called as spikes and its bifurcation scenarios. We also verify different types of oscillatory activities analytically. In Section 3, we verify our bifurcation results analytically. We demonstrate the Turing instability and spatiotemporal dynamics of the model system with some basic preliminaries. We derive the TAE, CGLE and find the coefficients. We report the main features with numerical simulations and analyzed the results to show the qualitative dynamics. Finally, we conclude the results with a discussion in Section 4.

2 Formulation of the excitable model and some preliminaries

This article focuses on the complex dynamics of the two variables excitable Izhikevich system over an appropriate range of parameters. The model construction does not account the biophysical behavior, however the output dynamics is very realistic and biophysically plausible. The time evolution of such a mathematical model is described by the following set of ODEs [19, 20]

$$\begin{aligned} \frac{dp}{dt} &= (0.04p^2 + 5p + 140 - q + I)/\varepsilon = f_1(p, q), \\ \frac{dq}{dt} &= a(bp - q) = f_2(p, q), \end{aligned} \quad (1)$$

where, p measures the membrane voltage dynamics and q , the recovery variable, that measures the activation of K^+ and

inactivation of Na^+ ionic currents, with an after-spike resetting constraint, i.e., when the voltage, p reaches the peak value, the following relation is applied: if $p \geq p_{\text{peak}} = 30$, then $p \leftarrow c$, $q \leftarrow q + d$. p is expressed in mV (millivolt) scale and time t is in ms (millisecond) scale [21]. The values of the parameters a, b, c and d determine spiking-bursting activities. The resting potential depends on the parameter b indicating the sensitivity of q to the subthreshold fluctuations of the voltage, p . The parameter a measures the timescale of the recovery variable, q . The parameters c and d control the afterspike reset values of p and q , respectively for the uncoupled dynamics. The function $(0.04p^2 + 5p + 140)$ was derived using the spike initiation dynamics of a cortical cell. The different suitable choices of parameters generate various types of oscillations such as regular spiking, various bursting, chattering and fast spiking, often found in neocortical and thalamic neurons [3, 24]. The stimulus currents are injected using the constant variable, I .

The equilibrium point, $E = (p_0, q_0)$ of system (1) can be found by solving $f_1(p_0, q_0) = 0$ and $f_2(p_0, q_0) = 0$, that gives $p_0 = \frac{-(5-b) \pm \sqrt{(5-b)^2 - 0.16(140+I)}}{0.08}$ and $q_0 = bp_0$. The Jacobian matrix of the system (1) at equilibrium point, $E = (p_0, q_0)$ is given by

$$J = \begin{pmatrix} J_{11} & J_{12} \\ J_{21} & J_{22} \end{pmatrix},$$

where, $J_{11} = \frac{0.08p_0+5}{\epsilon}$, $J_{12} = \frac{-1}{\epsilon}$, $J_{21} = ab$ and $J_{22} = -a$. The equilibrium point, $E = (p_0, q_0)$ of the system (1) is locally asymptotically stable if, $\text{Trace}(J) = J_{11} + J_{22} < 0$ and $\text{Det}(J) = J_{11}J_{22} - J_{12}J_{21} > 0$. We have considered the following parameter sets that produce different qualitative firing activities for the deterministic model [19, 20, 39] and depend on the dynamics with $\epsilon = 1$ (shown in Fig. 1 (a), (b) and (c) for parameter sets I and II) of the system (1). Set I : $a = 1$, $b = 1.5$, $c = -60$, $d = 0$, Set II : $a = -0.02$, $b = -1$, $c = -60$, $d = 8$.

The numerical simulations of the excitable systems are performed using the fourth-order Runge-Kutta method with a time step of 0.001, and the initial conditions are set to $p = -63$ and $q = bp$ [19, 20, 39]. The simulation results with a smaller time step do not show any significant differences. Bifurcation diagrams of the dynamical model are computed using the MatCont software package [4].

2.1 Oscillatory activities and Bifurcation analysis

We consider three different regimes i.e., fast spiking, tonic spiking and phasic spiking with the steady state. The bifurcation analysis of the system (1) is presented for parameter set I and II by considering current stimulus, I as a control parameter. The system (1) with parameter set I shows a sub-critical Hopf bifurcation and a saddle node bifurcation at $I = -65$ and -63.4375 respectively (see Fig. 1(d)).

For Saddle-node bifurcation, as the magnitude of the parameter, I changes, a stable equilibrium corresponding to the resting state condition is approached by an unstable fixed point; then they coalesce and annihilate each other. Since the resting state no longer exists (bistable regime exists see Fig. 1(d)) with the increase of I , the neuron starts to fire fast spiking (Fig. 1(a) solid blue line). There is a coexistence of resting and spiking states in the case of saddle-node bifurcation. At lower current stimulus ($I < 85$) with set II, it has two equilibrium points which collide at $I = 85$ (see Fig. 1(e) marked by SN1 point) and vanishes together for higher current stimulus $I > 85$. The system has a saddle node bifurcation at $I = 85$. Another interesting behavior is the system changes its stability from stable regime to unstable at $I = 78.9975$ (see Fig. 1(e) marked by HB1 point) with the transition from phasic spiking to tonic spiking regime (Figs. 1(b)-(c), solid blue lines). The system (1) shows supercritical Hopf bifurcation (HB1) at $I = 78.9975$. In Figs. 1(d)-(e), dashed and thick blue line represent the stable and unstable equilibrium regions.

Now, we are interested to find the solution of system (1) analytically to verify our numerical simulations i.e., different types of oscillatory activities. We begin with the recovery variable, q and find the solution of the last equation of system (1) by considering p as a constant between t_0 and t , where t_0 indicates initial time. The last equation of system (1) can be written as

$$\frac{dq}{dt} + aq = abp, \quad (2)$$

which is a first order linear ODE. In this context, we can find the solution of Eq. (2) using the integrating factor, I.F. = e^{at} and the solution is given by

$$q(t) = q(t_0)e^{-a(t-t_0)} + e^{-at} \int_{t_0}^t abpe^{a\tau} d\tau,$$

which can be also written as

$$q(t) = q(t_0)e^{-a(t-t_0)} + bp(1 - e^{-a(t-t_0)}).$$

For better result, we use small time step i.e., $\Delta t = t - t_0$ is very small and replace t by $t_0 + \Delta t$.

Now, we find the approximate solution of first equation of system (1) by considering q as a constant over some small time interval as described earlier. The first equation of system (1) can be written as

$$\frac{dp}{dt} = \frac{140-q+I}{\epsilon} + \frac{5}{\epsilon}p + \frac{0.04}{\epsilon}p^2, \quad (3)$$

which is a first order nonlinear ODE. It is impossible to find the exact solution for most of the nonlinear ODEs, fortunately above equation is a special type of ODE, which is known as Riccati differential equation [14, 31]. We can find the solution of Riccati differential equation using the transformation $r = \frac{1}{p-p_1}$, where p_1 is a particular solution to $\frac{dp}{dt}$. In this context, the particular solution p_1 can be found from $\frac{dp}{dt} = 0$, which leads to




Fig. 1 (a) Time series of the system (1) for parameter set I with $I = -64$ and set II (b)-(c) $I = 78$ and 80 are shown. Solid blue line indicates numerical simulation results, where dotted red lines are found analytically. (d)-(e) Bifurcation diagrams with respect to I for sets I and II respectively. Thick and dashed blue lines represent the stability and instability of the fixed points. HB and HB1 indicate sub-critical and super-critical Hopf bifurcation points. SN and SN1 indicate saddle node bifurcations.

$$p_1 = \frac{-5 \pm \sqrt{25 - 0.16(140 - q + I)}}{0.08}.$$

Here, we are interested to find only one particular solution, that's why we choose $p_1 = \frac{-5 + \sqrt{25 - 0.16(140 - q + I)}}{0.08}$. So, using the transformation $r = \frac{1}{p - p_1}$, the system (3) becomes

$$\frac{dr}{dt} + \frac{B}{\varepsilon} r = -\frac{0.04}{\varepsilon},$$

where $B = 5 + 0.08p_1$. To, find the solution of system (3), first we find the solution of above equation and then use the transformation $r = \frac{1}{p - p_1}$. So, the solution of system (3) is given by

$$p(t) = p_1 + \left(-\frac{0.04}{B} (1 - e^{-B(t-t_0)/\varepsilon}) + \frac{e^{-B(t-t_0)/\varepsilon}}{p(t_0) - p_1} \right) - 1,$$

provided p_1 is real i.e., B is real. Complex value of B indicates system (1) has no fixed point, then let us consider $B = iB_1$, $U = -B_1(t - t_0)/\varepsilon$ and $V = p(t_0) + \frac{5}{0.08}$. Then, the solution of system (3) is given by

$$p(t) = -\frac{5}{0.08} + \frac{V \cos U + (0.04V^2/B_1) \sin U - (B_1/0.16) \sin U}{\frac{1}{2}(1 + \cos U) + (0.08V/B_1) \sin U + 0.0032V^2/B_1^2(1 - \cos U)}.$$

Now, we plot the membrane voltage variable p with same time step as described earlier i.e., $\Delta t = 0.001$ and that are shown in Figs 1 (a) - (c) in dotted red lines. It is clear from Figs 1 (a) - (c) that analytical results are in good agreement with the numerical results.

3 Coupled Izhikevich model with 1D diffusion

The general 1D reaction diffusion system with the Izhikevich model is represented by the following two equations

$$\begin{aligned} \frac{\partial p}{\partial t} &= f_1(p, q) + D_{11} \frac{\partial^2 p}{\partial x^2}, \\ \frac{\partial q}{\partial t} &= f_2(p, q) + D_{22} \frac{\partial^2 q}{\partial x^2}, \end{aligned} \quad (4)$$

where $p = p(x, t)$ and $q = q(x, t)$ are the unknown functions to be evaluated and the subscripts t and x present the differentiation with respect to these variables. The initial conditions of the coupled PDEs are considered with the known functions $p(x, t = 0)$ and $q(x, t = 0)$ for $x \in \Omega$ and the boundary conditions show zero-flux boundary conditions $\frac{\partial p}{\partial n} = \frac{\partial q}{\partial n} = 0$, $x \in \partial\Omega$ and $t > 0$, where n measures the outward normal to $\partial\Omega$, the boundary of the interval and domain, Ω is the bounded interval or square domain for 1D and 2D diffusion respectively. In the 1D case, the length of the excitable cable is $x = 100$. D_{11} and D_{22} are the strengths of the synaptic couplings with positive values. One can explore the similar dynamics with the large finite length of the excitable cable by introducing more integrating space points in the simulation. We use a finite-difference scheme (with forward Euler method and central difference representation) with zero-flux boundary conditions for numerical simulation of a cable of finite length. The time step is $\delta t = 0.001$ and space step $\delta x = 0.5$, the number of nodes $N = x/\delta x = 200$

are considered. The second order partial derivative $\frac{\partial^2 p}{\partial x^2}$ and $\frac{\partial^2 q}{\partial x^2}$ are approximated as $\frac{1}{(\delta x)^2}(p_{i,j-1} - 2p_{i,j} + p_{i,j+1})$ and $\frac{1}{(\delta x)^2}(q_{i,j-1} - 2q_{i,j} + q_{i,j+1})$ respectively to find the numerical solution of the system (4). The initial conditions are considered with suitable periodic perturbations from the initial conditions as in the deterministic case. It is fixed for all the simulations. Biologically, the zero-flux boundary condition shows that the cell membranes are impermeable at the boundaries and it acts as an isolated cable [32].

To find the solution, we deviate the system (4) around the uniform steady state condition and linearize it around the nontrivial fixed point, we obtain the characteristic equation assuming the particular solution as $\begin{pmatrix} p \\ q \end{pmatrix} = \begin{pmatrix} p_0 \\ q_0 \end{pmatrix} + \varepsilon \begin{pmatrix} p_k \\ q_k \end{pmatrix} e^{\lambda_k t + ikx} + c.c. + o(\varepsilon^2)$ for model (4), where *c.c.* stands for the complex conjugate, λ_k is the wave length, k is the wave number along the x direction (x is the directional vector). The Jacobian matrix of Eq. (4) at the nontrivial equilibrium point $E = (p_0, q_0)$ is given by

$$J_D = J - k^2 D = \begin{pmatrix} J_{11} - D_{11}k^2 & J_{12} \\ J_{21} & J_{22} - D_{22}k^2 \end{pmatrix},$$

where $D = \begin{pmatrix} D_{11} & 0 \\ 0 & D_{22} \end{pmatrix}$. The stability conditions of system (4) at $E = (p_0, q_0)$ are given by $\text{Trace}(J_D) = J_{11} + J_{22} - D_{11}k^2 - D_{22}k^2 < 0$ and $\text{Det}(J_D) = (J_{11} - D_{11}k^2)(J_{22} - D_{22}k^2) - J_{12}J_{21} > 0$. The eigenvalues of the Jacobian matrix J_D are given by

$$\lambda = \frac{1}{2}(\text{Trace}(J_D) \pm \sqrt{(\text{Trace}(J_D))^2 - 4\text{Det}(J_D)}).$$

Now, we use the analytical approach to find the Hopf bifurcation point with its nature and the bifurcating periodic solution. In Andronov-Hopf bifurcation case [10, 12, 26], it produces a limit cycle from a fixed point or equilibrium solution of the dynamical model of ODEs, when the fixed point changes its stability through a pair of purely imaginary eigenvalues. The bifurcation can be supercritical or subcritical, resulting in stable or unstable limit cycle, respectively. For Supercritical Andronov-Hopf bifurcation, the stable equilibrium loses its stability and gives birth to a small-amplitude limit cycle attractor. When the magnitude of the parameter increases, the amplitude of the limit cycle increases and it becomes a full-size spiking limit cycle. First, consider the model (1) with the predominant parameter, I and the equilibrium state, $E = (p_0, q_0)$ depends on the values of I . Let the eigenvalues of the linearized system at the fixed point E becomes $\lambda(I)$, $\bar{\lambda}(I) = \alpha(I) \pm i\beta(I)$. Suppose the following conditions are satisfied for a particular critical value of I , say $I = I_H$.

- The Jacobian matrix has a simple pair of purely imaginary eigenvalues at the critical value of I , around the

fixed point, E , i.e., $\alpha(I = I_H) = 0$, $\beta(I = I_H) = w \neq 0$, where $\text{sgn}(w) = \text{sgn}[(\partial f_2(p, q)/\partial p)]|_{p_0, q_0}$ at $I = I_H$ (i.e., known as non-hyperbolicity condition). Then there exists a smooth curve of fixed point with the critical point of I and the eigenvalues vary smoothly.

- Moreover, if $\left. \frac{d\alpha(I)}{dI} \right|_{I=I_H} = m \neq 0$, (known as transversality condition, the eigenvalues cross the imaginary axis with non-zero speed).
- Finally, $n \neq 0$, where $n = (1/16)(f_{1ppp} + f_{1pqq} + f_{2ppq} + f_{2qqq}) + (1/16w)(f_{1pq}(f_{1pp} + f_{1qq}) - f_{2pq}(f_{2pp} + f_{2qq}) - f_{1pp}f_{2pp} + f_{1qq}f_{2qq})$ with $f_{1pq} = (\partial^2 f_1(p, q)/\partial p \partial q)|_{p_0, q_0}$ at $I = I_H$ (known as genericity condition). Then a unique curve of the periodic solutions bifurcates from the equilibrium solution E , into the parameter region $I > I_H$, if the condition $mn < 0$ holds or $I < I_H$, if $mn > 0$ holds. The fixed point E becomes stable if $I > I_H$ (respectively $I < I_H$) and unstable for $I < I_H$ (resp. $I > I_H$) for $m < 0$ (resp. $m > 0$) while the periodic solutions are stable (resp. unstable) if the equilibrium point is unstable (resp. stable) on the side of $I = I_H$, where the periodic solutions exist. It is called a supercritical Hopf bifurcation if the bifurcating periodic solutions are stable, and subcritical if they are unstable. To test the Hopf-bifurcation analysis, assume that the characteristic equation has a pair of purely imaginary roots $\lambda_{1,2} = \pm i\beta$ where ($\beta > 0$).

The system (4) in the absence of diffusion shows Hopf bifurcation if $\text{Im}(\lambda) \neq 0$ and $\text{Re}(\lambda) = 0$ with $k = 0$, i.e., $J_{11} + J_{22} = 0$, which gives $\frac{1}{\varepsilon}(0.08p_0 + 5) - a = 0$, where λ is the eigenvalue of the Jacobian matrix J . Thus, the critical value for the Hopf instability is given by

$$I_H = \frac{(5 - b)^2 - (a\varepsilon - b)^2}{0.16} - 140. \quad (5)$$

For parameter sets I and II, we obtain the critical value $I_H = -65$ and 78.9975 respectively, which are in good agreement with the numerical simulations. In the following, we study the stability of periodic solutions for Hopf bifurcation. The characteristic equation of the linearized matrix J is

$$\lambda^2 + (a - \frac{1}{\varepsilon}(0.08p_0 + 5))\lambda + \frac{a}{\varepsilon}(-0.08p_0 - 5) + ab = 0. \quad (6)$$

For the occurrence of Hopf bifurcation, $\frac{1}{\varepsilon}(0.08p_0 + 5) - a = 0$, then Eq. (6) becomes

$$\lambda^2 - a^2 + ab = 0. \quad (7)$$

Further simplification of Eq. (7) gives two purely imaginary roots as $\lambda = \pm i\sqrt{ab - a^2} = \pm i\beta$. Differentiating Eq. (6) with respect to I , we obtain

$$\lambda' = \frac{\frac{0.08}{\varepsilon}(\lambda + a)\frac{dp_0}{dI}}{2\lambda + a - \frac{1}{\varepsilon}(0.08p_0 + 5)}. \quad (8)$$

Now, substitute $\pm i\beta$ for λ in Eq. (8), we have $\text{Re}(\lambda') = 0.04 \frac{dp_0}{dt}$, where $\frac{dp_0}{dt} = \mp \frac{1}{\sqrt{(5-b)^2 - 0.16(140+I)}}$. Suppose η_1 and η_2 be the eigenvectors corresponding to eigenvalues $i\beta$ and $-i\beta$ respectively. Now, we define a non-singular matrix $A = \text{col}(\text{Re}(\eta_1), \text{Im}(\eta_1))$, so that

$$A^{-1} \begin{pmatrix} a & -1 \\ ab & -a \end{pmatrix} A = \begin{pmatrix} 0 & \beta \\ -\beta & 0 \end{pmatrix},$$

where $\beta = \sqrt{ab - a^2}$ and $\eta_1 = (1, a - i\sqrt{ab - a^2})$. Then $A = \begin{pmatrix} 1 & 0 \\ a - \sqrt{ab - a^2} & \end{pmatrix}$ and $A^{-1} = \begin{pmatrix} 1 & 0 \\ \frac{a}{\sqrt{ab - a^2}} & \frac{-1}{\sqrt{ab - a^2}} \end{pmatrix}$.

Let $Y = A^{-1}V$, where $Y = \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}$ and $V = \begin{pmatrix} p \\ q \end{pmatrix}$. We have $y_1 = p$ and $y_2 = \frac{a}{\sqrt{ab - a^2}}p - \frac{1}{\sqrt{ab - a^2}}q$. Then system (1) becomes

$$\begin{pmatrix} \dot{y}_1 \\ \dot{y}_2 \end{pmatrix} = \begin{pmatrix} 0 & \sqrt{ab - a^2} \\ -\sqrt{ab - a^2} & 0 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} + \begin{pmatrix} 0.04y_1^2 \\ \frac{0.04a}{\sqrt{ab - a^2}}y_1^2 \end{pmatrix}. \quad (9)$$

Writing F^i , $i = 1, 2$ for the nonlinear part of Eq. (9), we have $F^1 = 0.04y_1^2$ and $F^2 = \frac{0.04a}{\sqrt{ab - a^2}}y_1^2$. Again, we find $F_{11}^1 = 0.08$, $F_{11}^2 = \frac{0.08a}{\sqrt{ab - a^2}}$ and all other F_{jk}^i are zero. Thus, from the genericity condition, we have $n = \frac{1}{16\sqrt{ab - a^2}}(-F_{11}^1 F_{11}^2) = \frac{-0.04a}{ab - a^2}$. Therefore, $n < 0$ for $a > 0$, which indicates that the system (1) exhibits sub-critical Hopf bifurcation. The system (1) shows super-critical Hopf bifurcation if $n > 0$ i.e., $a < 0$. In our study, the system (1) shows a sub-critical and a super-critical Hopf bifurcation for parameter sets I and II respectively, which also satisfy our numerical result. Now, we study the Turing instability. In the presence of diffusion, Turing instability breaks the spatial symmetry i.e., the uniform steady state is stable for system (1) however, it is unstable for system (4). Thus the conditions for Turing instability are given by (i) $\text{Trace}(J) = J_{11} + J_{22} < 0$, (ii) $\text{Det}(J) = J_{11}J_{22} - J_{12}J_{21} > 0$, and (iii) $\text{Det}(J_D) = (J_{11} - D_{11}k^2)(J_{22} - D_{22}k^2) - J_{12}J_{21} < 0$.

We can find the threshold values of Turing instability and Turing bifurcation using $\text{Det}(J_D) = 0$ and $\text{Im}(\lambda) = 0$, $\text{Re}(\lambda) = 0$ at $k = k_{crit} \neq 0$ respectively. The threshold value of Turing bifurcation parameter is given by

$$I_{crit} = \frac{(5-b)^2 - (A-b)^2}{0.16} - 140, \quad (10)$$

where $A = \frac{\varepsilon(J_{22}D_{11} - 2\sqrt{-J_{12}J_{21}D_{11}D_{22}})}{D_{22}}$ and corresponding critical wave number is $k_{crit} = \left(\frac{\text{Det}(J)}{\text{Det}(D)}\right)^{1/4}$ i.e.,

$$k_{crit} = \left(\frac{ab - a(0.08p_0 + I)}{\varepsilon D_{11}D_{22}}\right)^{1/4}.$$

If we consider D_{22} as a Turing parameter, then the threshold value is given by

$$(D_{22})_{crit} = \frac{D_{11}}{J_{11}^2} (\text{Det}(J) - J_{12}J_{21} + 2\sqrt{-J_{12}J_{21}\text{Det}(J)}). \quad (11)$$

One can also evaluate the critical value of Turing instability with respect to D_{11} . Interestingly, it is clear from the above calculations that synaptic coupling D_{22} plays a major role for Turing instability. In the absence of D_{22} , only the membrane voltage spatially interacts in the excitable extended systems and as a result of it there is no Turing instability in the reaction diffusion system [5, 32]. Now, we derive the procedure to obtain the amplitude equations with the generalized TAE equations that explore the characteristics of the diffusive instability. We only consider constant diffusion scheme as it has interesting phenomenon in excitable spatial system. At the threshold of Turing point, the spatial symmetry of the reaction diffusion system is destroyed and arises a stationary form in time and oscillatory dynamics in space. It is noted that in the vicinity of a bifurcation point, the evolution of a spatial system generates critical slowing down, that can be described by an amplitude equation as following.

3.1 Turing amplitude equation (TAE)

For Turing instability, the dynamics of the system (4) can be described by the amplitude equations, which is known as TAE. The general form of TAE is given by [43, 47]

$$\frac{\partial W}{\partial t} = \eta_T W + g_T |W|^2 W + D_T \frac{\partial^2 W}{\partial x^2}, \quad (12)$$

where η_T , g_T and D_T are constants. The normalized form of this equation with $\eta_T = D_T = 1$ and $g_T = -1$ can be considered as the limiting condition of CGLE equation [47]. The TAE equation is a valid efficient tool in one spatial dimensional (1D) case. Here, we do not consider the case of wave instability [34]. In the following, we find the value of above parameters. At the onset of Turing instability, for critical wave number k_{crit} , the critical value ε_T is given by

$$\varepsilon_T = \frac{\text{Det}(D)(2b - \hat{J}_{11} + 2\sqrt{b(b - \hat{J}_{11})})}{aD_{11}^2},$$

where $\hat{J}_{11} = \varepsilon J_{11} = 0.08p_0 + 5$ and $\hat{J}_{12} = \varepsilon J_{12} = -1$. First, we introduce a control parameter μ_1 as normalized deviation from the critical value ε_T i.e., $\mu_1 = \frac{\varepsilon - \varepsilon_T}{\varepsilon_T}$. The right and left eigenvectors of the matrix $L_0 = (J_D)|_{\mu_1=0}$ are given by $PP = \begin{pmatrix} PP_p \\ PP_q \end{pmatrix} = \begin{pmatrix} 1 \\ \alpha \end{pmatrix} e^{ik_{crit}x}$ and $PP^* = (1 + \alpha\beta)^{-1} (1, \beta) e^{ik_{crit}x}$

respectively, where $\alpha = \frac{k_{crit}^2 D_{11} - \frac{\hat{J}_{11}}{\varepsilon_T}}{-\frac{1}{\varepsilon_T}}$ and $\beta = \frac{k_{crit}^2 D_{11} - \frac{\hat{J}_{11}}{\varepsilon_T}}{ab} =$

$\frac{-1}{k_{crit}^2 D_{22} + a} \frac{\varepsilon_T}{\varepsilon_T}$. The matrix $J^0 = J|_{\mu_1=0}$ is given by

$$J^0 = \begin{pmatrix} \frac{J_{11}}{\varepsilon_T} & -\frac{1}{\varepsilon_T} \\ ab & -a \end{pmatrix}.$$

Using the expressions $J_{ij}^1 = \frac{dJ_{ij}}{d\varepsilon} \frac{d\varepsilon}{d\mu_1} |_{\varepsilon=\varepsilon_T}$ and $\gamma_{ij} = -J_{ij}^0 + 4k_{crit}^2 D_{ij}$ the matrix J^1 and γ are given by

$$J^1 = \begin{pmatrix} -\frac{J_{11}}{\varepsilon_T} & \frac{1}{\varepsilon_T} \\ 0 & 0 \end{pmatrix},$$

and

$$\gamma = \begin{pmatrix} -\frac{J_{11}}{\varepsilon_T} + 4k_{crit}^2 D_{11} & \frac{1}{\varepsilon_T} \\ -ab & a + 4k_{crit}^2 D_{22} \end{pmatrix}$$

respectively. Finally, η_T , D_T and g_T are given by [47]

$$\eta_T = \frac{1}{1 + \alpha\beta} (J_{11}^1 + \alpha J_{12}^1 + \beta J_{21}^1 + \alpha\beta J_{22}^1) = \frac{\alpha - J_{11}}{\varepsilon_T(1 + \alpha\beta)}, \quad (13)$$

$$D_T = 4k_{crit}^2 \frac{(D_{21} + \alpha D_{22})(D_{12} + \beta D_{22})}{(1 + \alpha\beta)(J_{22}^0 - k_{crit}^2 D_{22})} = -\frac{4\alpha\beta k_{crit}^2 D_{22}^2}{(1 + \alpha\beta)(a + k_{crit}^2 D_{22})}, \quad \varepsilon_H = \frac{0.08p_0 + 5}{a}, \quad (14)$$

and

$$g_T = \frac{\bar{g}}{1 + \alpha\beta}, \quad (15)$$

where $\bar{g} = H_1^{pp}(-\frac{2}{DetJ^0}G_1J_{22}^0 + \frac{1}{Det\gamma}G_1\gamma_{22})$, $G_1 = \frac{H_1^{pp}}{2}$ and $H_1^{pp} = \frac{\partial^2 f_1(p,q)}{\partial p^2} = \frac{0.08}{\varepsilon_T}$. We can observe that effects of diffusion appear in the expressions of all the coefficients of the TAE, since the parameters α and β depend on diffusive couplings.

$g_T = 0$ line separates the region between supercritical and subcritical Turing instability. $g_T < (>)0$ indicates supercritical (subcritical) Turing instability. First, we consider parameter set I with $I = -68$ and $D_{11} = 1$. The deterministic system (1) is bistable around $I = -68$ (see Fig. 1(d)) i.e., stable and unstable equilibrium branch coexist. The stable equilibrium point $E = (-54.43, -81.645)$ becomes unstable for system (4) i.e., Turing instability occurs for suitable value of diffusion coefficient D_{22} . The threshold value for Turing instability is given by $(D_{22})_{crit} = 11.081$ (from Eq. (11)), i.e., stable equilibrium point becomes unstable for $D_{22} > 11.081$. For better understanding, we have shown the real part of the eigenvalue λ for different values of D_{22} in Fig. 2(a). In Fig. 2(a), $\text{Re}(\lambda)=0$ line separates the Turing and non-Turing domains.

As we already know that, system (4) shows Turing instability at parameter set I, with $I = -68$ and $D_{11} = 1$ for $D_{22} > (D_{22})_{crit} = 11.081$. Now, we are interested to find the nature of the Turing instability i.e., whether it's supercritical or subcritical. In Fig 2(b), we have shown the value of g_T is negative in Turing region. That indicates the instability is supercritical.

3.2 Complex Ginzburg-Landau equation (CGLE)

The dynamics of the spatiotemporal system (4) nearby to the Hopf instability can be described by the amplitude equations, which is known as CGLE [15, 16, 47]. The general form of CGLE is given by

$$\frac{\partial W}{\partial t} = (1 + iC_0)W + (1 + iC_1)\nabla^2 W - (1 + iC_2)|W|^2 W, \quad (16)$$

where i denotes the imaginary unit and ∇^2 is the one or two-dimensional Laplacian operator. When the real parameters C_0 , C_1 and C_2 vary, the complex amplitude, W shows rich dynamical behavior. First, we have to find the condition for Hopf instability and then the values of C_0 , C_1 and C_2 using Kuramoto's method [22, 47]. The manipulations are done using CGLE equation and only the coefficient C_1 depends on the diffusion. For the case of Hopf instability, $\text{Trace}(J) = J_{11} + J_{22} = 0$. The critical value ε_H of ε at the onset of Hopf instability is given by

Jacobian matrix J has purely imaginary eigenvalues $\pm i\omega_0$, in the case of Hopf instability, which gives $\omega_0^2 = J_{11}J_{22} - J_{12}J_{21}$ i.e.,

$$\omega_0^2 = -a^2 + \frac{ab}{\varepsilon_H}.$$

To find the values of C_0 , C_1 and C_2 , we introduce a small parameter μ as normalized deviation from the critical value ε_H i.e., $\mu = \frac{\varepsilon - \varepsilon_H}{\varepsilon_H}$. The matrix $J_0 = J|_{\mu=0}$ is given by

$$J_0 = \begin{pmatrix} a & \frac{-a^2 - \omega_0^2}{ab} \\ ab & -a \end{pmatrix}.$$

The left and right eigenvectors of the matrix J_0 are given by $P^* = \frac{1}{2} \left(-i\frac{a}{\omega_0}, \frac{1}{b} + i\frac{a}{b\omega_0} \right)$ and $P = \begin{pmatrix} P_p \\ P_q \end{pmatrix} = \begin{pmatrix} 1 + \frac{i\omega_0}{b} \\ a \end{pmatrix}$ respectively. The matrix $J_1 = \frac{dJ}{d\varepsilon} \frac{d\varepsilon}{d\mu} |_{\varepsilon=\varepsilon_H}$ is given by

$$J_1 = \begin{pmatrix} -a & \frac{1}{\varepsilon_H} \\ 0 & 0 \end{pmatrix}.$$

Next, we compute $P^* J_1 P$ to find the value of constant C_0 in Eq. (16), which gives $P^* J_1 P = -\frac{a+i\omega_0}{2} = \sigma_1 + i\omega_1$. Now,

$$C_0 = \frac{\omega_1}{\sigma_1} = \frac{\omega_0}{a}. \quad (17)$$

Next, we compute $P^* D P$ to find the value of constant C_1 in Eq. (16), which gives $P^* D P = -\frac{1}{2}(D_{11} + D_{22} + i\frac{a}{\omega_0}(D_{22} - D_{11})) = d_1 + id_2$. Now,

$$C_1 = \frac{d_2}{d_1} = \frac{a}{\omega_0} \frac{D_{22} - D_{11}}{D_{22} + D_{11}}. \quad (18)$$

To find $C_2 = \frac{g_2}{g_1}$, we calculate the vectors HXX and $NXXX$. HXX and $NXXX$, that are quadratic and cubic terms respectively [15, 47]. HXX is described by




Fig. 2 (a) Relation between the real part of the eigenvalue λ and k for parameter set I, with $I = -68$ and $D_{11} = 1$ for different diffusion coefficients, D_{22} . (b) The expression of g_T is shown graphically for set I, with $I = -68$ and $D_{11} = 1$ corresponding to D_{22} to determine the nature of the Turing instability. (c) Relation between $C_1 + C_2$ and D_{22} for set II, with $I = 78.9975$ and different D_{11} .

$$(HXX)_i = \frac{1}{2} \left[\frac{\partial^2 f_i(p,q)}{\partial p^2} \Big|_{p_0,q_0} P_p P_p + 2 \frac{\partial^2 f_i(p,q)}{\partial p \partial q} \Big|_{p_0,q_0} P_p P_q \right],$$

for $i = 1, 2$. $(HXX)_2 = 0$ as $f_2(p, q)$ is a linear function. $\frac{\partial^2 f_1(p,q)}{\partial p \partial q} = 0$ as in $f_1(p, q)$, no product term of p and q exists. Now, with $\mu = 0$ the first component $(H_0XX)_1$ is given by

$$(H_0XX)_1 = \frac{1}{2} \frac{\partial^2 f_1(p,q)}{\partial p^2} \Big|_{p_0,q_0} P_p P_p = \left(1 - \frac{\omega_0^2}{a^2}\right) \frac{H_1^{pp}}{2} + i \frac{\omega_0}{a} H_1^{pp},$$

where $H_1^{pp} = \frac{\partial^2 f_1(p,q)}{\partial p^2} = \frac{0.08}{\varepsilon_H}$. Similarly, the first component $(H_0X\bar{X})_1$ is given by

$$(H_0X\bar{X})_1 = \frac{1}{2} \frac{\partial^2 f_1(p,q)}{\partial p^2} \Big|_{p_0,q_0} P_p \bar{P}_p = \left(1 + \frac{\omega_0^2}{a^2}\right) \frac{H_1^{pp}}{2}.$$

As $f_1(p, q)$ is a quadratic function and $f_2(p, q)$ is a linear function, $(NXXX)_i = 0$, for $i = 1, 2$. Next, we compute the following expressions

$$(J_0 - 2i\omega_0 I)^{-1} = -\frac{1}{3\omega_0^2} \begin{pmatrix} -a - 2i\omega_0 & \frac{a^2 + \omega_0^2}{ab} \\ -ab & a - 2i\omega_0 \end{pmatrix},$$

$$Y_0 = -2J_0^{-1} H_0 X \bar{X} = \left(1 + \frac{\omega_0^2}{a^2}\right) \frac{H_1^{pp}}{\omega_0^2} \begin{pmatrix} a \\ ab \end{pmatrix}.$$

$$Y_+ = -(J_0 - 2i\omega_0 I)^{-1} H_0 X X = -\frac{1}{3\omega_0^2} \left(\left(1 - \frac{\omega_0^2}{a^2}\right) \frac{H_1^{pp}}{2} + i \frac{\omega_0}{a} H_1^{pp} \right) \begin{pmatrix} a + 2i\omega_0 \\ ab \end{pmatrix}.$$

Now, $(H_0XY_0)_1 = \frac{H_1^{pp} P_p Y_{0p}}{2}$, $(H_0XY_0)_2 = 0$, $(H_0\bar{X}Y_+)_1 = \frac{H_1^{pp} \bar{P}_p Y_{+p}}{2}$ and $(H_0\bar{X}Y_+)_2 = 0$. Next, we calculate g as $g = g_1 + ig_2 = -P^*(2H_0XY_0 + 2H_0\bar{X}Y_+ + 3N_0XX\bar{X})$, which gives

$$g_1 + ig_2 = -2P^* H_0XY_0 - 2P^* H_0\bar{X}Y_+. \quad (19)$$

Now,

$$-2P^* H_0XY_0 = \frac{1}{2\omega_0^3} (a^2 + \omega_0^2) (H_1^{pp})^2 \left(-\frac{\omega_0}{a} + i\right), \quad (20)$$

and

$$-2P^* H_0\bar{X}Y_+ = \frac{1}{12\omega_0^3 a^2} (3a\omega_0(a^2 + \omega_0^2) + i(a^2\omega_0^2 - a^4 + 2\omega_0^4)) (H_1^{pp})^2. \quad (21)$$

Using Eqs. (19), (20) and (21) with equating the complex and real parts, we obtain

$$g_1 = -\frac{1}{4\omega_0^2 a} (a^2 + \omega_0^2) (H_1^{pp})^2, \quad (22)$$

and

$$g_2 = -\frac{ag_1}{\omega_0} + \frac{1}{6\omega_0^3 a^2} (a^2 + \omega_0^2)^2 (H_1^{pp})^2.$$

Finally,

$$C_2 = \frac{g_2}{g_1} = -\frac{a}{\omega_0} - \frac{2}{3\omega_0 a} (a^2 + \omega_0^2). \quad (23)$$

The CGLE is valid if $g_1 > 0$. Hence, the Hopf bifurcation is super-critical if $g_1 > 0$ and sub-critical if $g_1 < 0$. The phase waves spreading away from the origin of perturbation and towards the origin of perturbation are known as Wave (W) and antiwave (AW) respectively [47]. W and AW domains are separated by the curve $C_1 + C_2 = 0$. W and AW domains are given by $C_1 + C_2 < 0$ and $C_1 + C_2 > 0$ respectively.

It is clear from the Eq. (22), that the sign of g_1 depends on a , super-critical Hopf bifurcation exists for $a < 0$, which is in good agreement with the above analytical and numerical simulations. Now, we consider parameter sets I and II with $I = -65$ and $I = 78.9975$ respectively. We derive the




Fig. 3 Spatiotemporal plots of the 2D Izhikevich cable for parameter set II, with $I=78$ and $D_{11} = 0.001$, (a),(d) $D_{22} = 0.00001$, (b),(e) $D_{22} = 0.01$ and (c),(f) $D_{22}=20$. (d)-(f) The blue, green and red lines indicate the corresponding time series of arbitrary oscillators in the reaction diffusion coupled systems (50th, 55th and 60th oscillators respectively).

following values $\varepsilon_H = 1$, $\omega_0 = 0.7071$, $C_0 = 0.7071$, $C_1 = -\frac{1.41421(D_{11}-D_{22})}{(D_{11}+D_{22})}$, $C_2 = -2.8284$, $g_1 = -0.0048$ and $\varepsilon_H = 1$, $\omega_0 = 0.14$, $C_0 = -7$, $C_1 = \frac{0.142857(D_{11}-D_{22})}{(D_{11}+D_{22})}$, $C_2 = 4.9048$, $g_1 = 0.0816 > 0$ for parameter sets I and II respectively. In Fig. 2(c), we have shown the value of $C_1 + C_2$ is positive for different values of D_{11} , corresponding to D_{22} for set II. Therefore, only antiwave exists.

3.3 Synchronization factor

Next, we have considered parameter set II with $I = 78$ and $D_{11} = 0.001$. At lower diffusion $D_{22} = 0.00001$, the system (4) produces a de-synchronized firing pattern (see Fig. 3(a)). For better understanding, we have presented time-series of different oscillators in Fig. 3(d). At higher diffusion $D_{22} = 0.01$, we observed more synchronized firing pattern (see Figs. 3(b)-(e)). Finally, at $D_{22} = 20$, all nodes are perfectly synchronized i.e., all nodes show same firing behavior. To understand the synchronization behavior, we derive a statistical measure of synchronization i.e, the synchronization factor R , which is given by the following equation

$$R = \frac{\langle E^2 \rangle - \langle E \rangle^2}{\frac{1}{N} \sum_{i=1}^N [\langle p_i^2 \rangle - \langle p_i \rangle^2]},$$

where,

$$E = \frac{1}{N} \sum_{i=1}^N p_i.$$

The symbol $\langle * \rangle$ indicates the mean value of the variable over time. R takes the value between 0 and 1 [8, 44], where the value one indicates that all nodes are perfectly synchronized. We have shown the effect of the diffusion coefficient D_{22} on synchronization factor R in Fig 4 . Further, if we consider a bursting regime, we obtained similar results with no significant changes. All the nodes are not synchronized for weak coupling. However, with the increase of diffusion coefficient D , the value of R tends to 1, which indicates all the nodes are perfectly synchronized for higher coupling.

4 Conclusions

The effects of diffusion in a biophysical extended system and then collective dynamical formation have been an ongoing research theme. The diffusion can enhance the spatial characteristics and even induce different diffusive instabilities. We consider diverse excitabilities of the deterministic system and verify the analytics of the model using Riccati differential equation. Next, the diffusion induced characteristics can be evaluated on the basis of amplitude equations that provide a mathematical analysis of reaction diffusion systems nearly to the onset of instabilities [34, 37, 47]. We report the effects of the diffusion on the system dynamics in the vicinity of Hopf and Turing instability. The 1D cable, which is in steady state for uncoupled system can produce ir-




Fig. 4 Distribution for the synchronization factor of the 2D Izhikevich cable with diffusion for the parameter set II with $I=78$ and $D_{11} = 0.001$.

regular oscillatory patterns. We also study the effects of diffusion on synchronization in oscillatory regime. For the TAE equation, it is classified the Turing instability and divided it into two domains. We verify the numerical simulations with proper analytical treatment for different firing patterns, Hopf bifurcation and Turing instability. The diffusive coupling affects the coefficient C_1 , whereas the term C_2 is independent of diffusion for CGLE equation. This approach will require the modification to observe the effects on spatially extended fast-slow excitable dynamical system. Note that, the value of the diffusion coefficient has strong effects on k_T and hence on the diffusive instabilities. The diffusive couplings contribute to all the coefficients of the amplitude equations. The approach demonstrates flexibility of our methodology for various firing regions supported by Hopf bifurcation analysis. The present article reports the emergence of two types of bifurcations in the temporal dynamics that control different oscillations exhibited by the system. Neural oscillations reflect rhythmic activities and can be generated either by mechanisms in single neurons or by network of neurons connected in a complex fashion. The results further show the dynamical behavior of the transition phases in single model and the diffusion induced system.

Acknowledgements This work is supported by University Grants Commission (UGC), Govt. of India under NET-JRF scheme to the coauthor SKS.

Compliance with ethical standards

Conflicts of interest The authors declare that there is no conflict of interests regarding the publication of this article.

Ethical standard The authors state that this research com-

plies with ethical standards. This research does not involve either human participants or animals.

References

1. Aïnseba, B., Bendahmane, M., Ruiz-Baier, R.: Analysis of an optimal control problem for the tridomain model in cardiac electrophysiology. *J. Math. Anal. Appl.* **388**(1), 231–247 (2012)
2. Aranson, I.S., Kramer, L.: The world of the complex ginzburg-landau equation. *Rev. Mod. Phys.* **74**(1), 99 (2002)
3. Connors, B.W., Gutnick, M.J.: Intrinsic firing patterns of diverse neocortical neurons. *Trends Neurosci.* **13**(3), 99–104 (1990)
4. Dhooge, A., Govaerts, W., Kuznetsov, Y.A., Meijer, H.G.E., Sautois, B.: New features of the software matcont for bifurcation analysis of dynamical systems. *Math. Comput. Model. Dyn. Syst.* **14**(2), 147–175 (2008)
5. Ermentrout, B., Lewis, M.: Pattern formation in systems with one spatially distributed species. *Bulletin of mathematical biology* **59**(3), 533–549 (1997)
6. Feng, Z., Chen, G.: Traveling wave solutions in parametric forms for a diffusion model with a nonlinear rate of growth. *Discrete Contin. Dyn. Syst. Ser. A* **24**(3), 763–780 (2009)
7. Gani, M.O., Ogawa, T.: Instability of periodic traveling wave solutions in a modified fitzhugh–nagumo model for excitable media. *Appl. Math. Comput.* **256**, 968–984 (2015)
8. Ge, M., Jia, Y., Xu, Y., Lu, L., Wang, H., Zhao, Y.: Wave propagation and synchronization induced by chemical autapse in chain hindmarsh–rose neural network. *Appl. Math. Comput.* **352**, 136–145 (2019)
9. Gong, Y., Christini, D.J.: Antispiral waves in reaction-diffusion systems. *Phys. Rev. Lett.* **90**(8), 088302 (2003)
10. Guo, H., Chen, Y.: Supercritical and subcritical hopf bifurcation and limit cycle oscillations of an airfoil with cubic nonlinearity in supersonic/hypersonic flow. *Nonlin. Dyn.* **67**(4), 2637–2649 (2012)
11. Hou, Z., Shi, B., Chai, Z.: A lattice boltzmann based local feedback control approach for spiral wave. *Comput. Math. with Appl.* **74**(10), 2330–2340 (2017)
12. Hsü, I.D., Kazarinoff, N.D.: Existence and stability of periodic solutions of a third-order non-linear autonomous system simulating immune response in animals. *Proc. R. Soc. Edinb. A* **77**(1-2), 163–175 (1977)
13. Hu, B., Ma, J., Tang, J.: Selection of multiarmed spiral waves in a regular network of neurons. *PLoS One* **8**(7), e69251 (2013)

14. Humphries, M.D., Gurney, K.: Solution methods for a new class of simple model neurons. *Neural computation* **19**(12), 3216–3225 (2007)
15. Ipsen, M., Hynne, F., Sørensen, P.: Systematic derivation of amplitude equations and normal forms for dynamical systems. *Chaos* **8**(4), 834–852 (1998)
16. Ipsen, M., Kramer, L., Sørensen, P.G.: Amplitude equations for description of chemical reaction–diffusion systems. *Phys. Rep.* **337**(1-2), 193–235 (2000)
17. Iqbal, N., Wu, R., Liu, B.: Pattern formation by superdiffusion in fitzhugh–nagumo model. *Appl. Math. Comput.* **313**, 245–258 (2017)
18. Izhikevich, E.M.: Neural excitability, spiking and bursting. *Int. J. Bifurcation and Chaos* **10**(06), 1171–1266 (2000)
19. Izhikevich, E.M.: Simple model of spiking neurons. *IEEE Trans. Neural Netw.* **14**(6), 1569–1572 (2003)
20. Izhikevich, E.M.: Which model to use for cortical spiking neurons? *IEEE Trans. Neural Netw.* **15**(5), 1063–1070 (2004)
21. Izhikevich, E.M.: *Dynamical systems in neuroscience*. MIT press (2007)
22. Kuramoto, Y.: *Springer series in synergetics*. In: *Chemical oscillations, waves and turbulence*, vol. 19, p. 130. Springer Berlin (1984)
23. Labouriau, I.S., Murza, A.C.: Periodic solutions in an array of coupled fitzhugh–nagumo cells. *J. Math. Anal. Appl.* **412**(1), 29–40 (2014)
24. Lau, H.T., Yu, M., Fontana, A., Stoeckert, C.J.: Prevention of islet allograft rejection with engineered myoblasts expressing fasl in mice. *Science* **273**(5271), 109–112 (1996)
25. Li, C.H., Yang, S.Y.: Eventual dissipativeness and synchronization of nonlinearly coupled dynamical network of hindmarsh–rose neurons. *Appl. Math. Model.* **39**(21), 6631–6644 (2015)
26. Liao, X., Li, S., Wong, K.w.: Hopf bifurcation on a two-neuron system with distributed delays: a frequency domain approach. *Nonlin. Dyn.* **31**(3), 299–326 (2003)
27. Lv, Y., Pei, Y., Yuan, R.: Hopf bifurcation and global stability of a diffusive gause-type predator–prey models. *Comput. Math. with Appl.* **72**(10), 2620–2635 (2016)
28. Ma, J., Gao, J.H., Wang, C.N., Su, J.Y.: Control spiral and multi-spiral wave in the complex ginzburg–landau equation. *Chaos Solit. Fract.* **38**(2), 521–530 (2008)
29. Ma, J., Hu, B., Wang, C., Jin, W.: Simulating the formation of spiral wave in the neuronal system. *Nonlin. Dyn.* **73**(1-2), 73–83 (2013)
30. Ma, J., Tang, J.: A review for dynamics in neuron and neuronal network. *Nonlin. Dyn.* **89**(3), 1569–1578 (2017)
31. Mokhtarzadeh, M., Pournaki, M., Razani, A.: A note on periodic solutions of riccati equations. *Nonlin. Dyn.* **62**(1), 119–125 (2010)
32. Mondal, A., Sharma, S.K., Upadhyay, R.K., Aziz-Alaoui, M., Kundu, P., Hens, C.: Diffusion dynamics of a conductance-based neuronal population. *Phys. Rev. E* **99**(4), 042307 (2019)
33. Mondal, A., Upadhyay, R.K.: Diverse neuronal responses of a fractional-order izhikevich model: journey from chattering to fast spiking. *Nonlin. Dyn.* **91**(2), 1275–1288 (2018)
34. Mondal, A., Upadhyay, R.K., Mondal, A., Sharma, S.K.: Dynamics of a modified excitable neuron model: Diffusive instabilities and traveling wave solutions. *Chaos* **28**(11), 113104 (2018)
35. Montani, F., Baravalle, R., Montangie, L., Rosso, O.A.: Causal information quantification of prominent dynamical features of biological neurons. *Philos. Trans. R. Soc. A* **373**(2056), 20150109 (2015)
36. Murza, A.C.: Oscillation patterns in tori of modified fhn neurons. *Appl. Math. Model.* **35**(3), 1096–1106 (2011)
37. Newell, A.C.: *Nonlinear Wave Motion* [proceedings of the Summer Seminar in Applied Mathematics Sponsored by American Mathematical Society and Society for Industrial and Applied Mathematics and Held at Potsdam, New York, 1972]. American Mathematical Society (1974)
38. Taliaferro, S.D.: Stability and bifurcation of traveling wave solutions of nerve axon type equations. *J. Math. Anal. Appl.* **137**(2), 396–416 (1989)
39. Teka, W.W., Upadhyay, R.K., Mondal, A.: Spiking and bursting patterns of fractional-order izhikevich model. *Commun. Nonlinear Sci. Numer. Simul.* **56**, 161–176 (2018)
40. Turing, A.M.: The chemical basis of morphogenesis. *Bull. Math. Biol.* **52**(1-2), 153–197 (1990)
41. Vanag, V.K., Epstein, I.R.: Inwardly rotating spiral waves in a reaction-diffusion system. *Science* **294**(5543), 835–837 (2001)
42. Vanag, V.K., Epstein, I.R.: Pattern formation in a tunable medium: The belousov-zhabotinsky reaction in an aerosol or microemulsion. *Phys. Rev. Lett.* **87**(22), 228301 (2001)
43. Walgraef, D.: The hopf bifurcation and related spatio-temporal patterns. In: *Spatio-Temporal Pattern Formation*, pp. 65–85. Springer (1997)
44. Wang, C., Lv, M., Alsaedi, A., Ma, J.: Synchronization stability and pattern selection in a memristive neuronal network. *Chaos* **27**(11), 113108 (2017)
45. Wu, R., Zhou, Y., Shao, Y., Chen, L.: Bifurcation and turing patterns of reaction–diffusion activator–inhibitor model. *Physica A: Statistical Mechanics and its Applications* **482**, 597–610 (2017)

-
46. Wu, X., Ma, J.: The formation mechanism of defects, spiral wave in the network of neurons. *PLoS One* **8**(1), e55403 (2013)
 47. Zemskov, E.P., Vanag, V.K., Epstein, I.R.: Amplitude equations for reaction-diffusion systems with cross diffusion. *Phys. Rev. E* **84**(3), 036216 (2011)

Figures




Figure 1

(a) Time series of the system (1) for parameter set I with $I = -64$ and set II (b)-(c) $I = 78$ and 80 are shown. Solid blue line indicates numerical simulation results, where dotted red lines are found analytically. (d)-(e) Bifurcation diagrams with respect to I for sets I and II respectively. Thick and dashed blue lines represent the stability and instability of the fixed points. HB and HB1 indicate sub-critical and super-critical Hopf bifurcation points. SN and SN1 indicate saddle node bifurcations.




Figure 2

(a) Relation between the real part of the eigenvalue $\text{Re}(\lambda)$ and k for parameter set I, with $\text{Re}(\lambda) = -68$ and $D_{11} = 1$ for different diffusion coefficients, D_{22} . (b) The expression of $\text{Re}(\lambda)$ is shown graphically for set I, with $\text{Re}(\lambda) = -68$ and $D_{11} = 1$ corresponding to D_{22} to determine the nature of the Turing instability. (c) Relation between $C_1 + C_2$ and D_{22} for set II, with $\text{Re}(\lambda) = 78.9975$ and different D_{11} .




Figure 3

Spatiotemporal plots of the 2D Izhikevich cable for parameter set II, with $\text{Re}(\lambda) = 78$ and $D_{11} = 0.001$, (a),(d) $D_{22} = 0.00001$, (b),(e) $D_{22} = 0.01$ and (c),(f) $D_{22} = 20$. (d)-(f) The blue, green and red lines indicate the corresponding time series of arbitrary oscillators in the reaction diffusion coupled systems (50th, 55th and 60th oscillators respectively).




Figure 4

Distribution for the synchronization factor of the 2D Izhikevich cable with diffusion for the parameter set II with $I=78$ and $D_{11} = 0.001$.