Characteristics and prognostic outcome factors in elderly peritoneal dialysis patients:a prospective observational study

Xiaoying Ma

Cangzhou central hospital

HaiPing Xu

Cangzhou central hospital

FuYun Sun (✉ 13315777305@163.com)

Research article

Keywords: Elderly, peritoneal dialysis, death, prognosis

Posted Date: July 23rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-41395/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background This study aimed to analyze the characteristics, outcomes and prognosis factors of survival in elderly peritoneal dialysis patients, so as to better understand the status of elderly peritoneal dialysis patients and improve their quality of life.

Methods This study was a prospective, observational study that included peritoneal dialysis (PD) patients. Categorizing by age, elderly group is aged ≥ 65 , younger group is aged < 65 . Clinical characteristics, survival and transferring to hemodialysis were compared between two groups. Meanwhile, risk factors of death in elderly PD patients were explored.

Results A total of 202 PD patients were enrolled, including 61 in elderly group and 141 in younger group. The comparison of baseline data between two groups: in elderly group serum albumin and normalized protein catabolic rate (nPCR) decreased, the incidence of previous cardiovascular and cerebrovascular diseases and Charlson Comorbidity Index (CCI) were higher, the major primary disease in elderly patients was diabetic nephropathy, significant differences were found between elderly and younger group ($P < 0.01$). The mortality in elderly group was substantially higher, 27 patients (44.3%) died in elderly group and 21 patients (14.3%) died in younger group. The 1-year, 2-year, 3-year, 4-year survival rate were 81.97%, 70.49%, 60.66%, 55.74%, respectively. Cardiovascular disease was the main cause of death in elderly PD patients. Higher BMI, CCI, and previous ischemic heart disease (IHD) were risk factors for long-term survival of elderly PD patients. Compared with the younger group, elderly patients were less likely to transfer to hemodialysis: 2 cases (3.3%) in elderly group and 23 cases (16.3%) in younger group. Peritonitis was primary reason for converting to hemodialysis (HD) in both two group.

Conclusions Poor nutrition, more complications and diabetic nephropathy were characteristics of elderly PD patients. High BMI, CCI and previous IHD were independent predictors of death in elderly PD patients. Cardiovascular disease was the main cause of death in elderly PD patients, while the chief reason of transferring to hemodialysis was peritonitis.

1. Background

Chronic kidney disease is a global public health problem and becoming increasingly common. A survey in China found that patients with chronic kidney disease stage 3 (CKD3) were about 1.2 million in 2012, and average age was 63.6 years [1]. With the development of CKD to the end stage, peritoneal dialysis has become one of the main renal replacement therapy and better choice of elderly patients with end stage renal disease, because of its small influence on hemodynamics, protection of residual renal function, low cost and simple operation [2].

Elderly PD patients have unique clinical characteristics, and incline to have multiple comorbidities, including diabetes, hypertension, cardiovascular and cerebrovascular diseases [3]. Many factors can influence the clinical outcomes of elderly PD patients such as comorbidities, complications, age and nutritional deficiency [4, 5]. However, previous studies were mainly retrospective studies, and the influencing factors of death were variously in different countries. Some studies have shown that death is the main reason for patients to withdraw from peritoneal dialysis, age and peritonitis are the risk factors for patients to withdraw from

peritoneal dialysis [6, 7]. However, there have been few reports about the causes of withdrawal from PD in China, especially transferring to hemodialysis.

In this study, We intended to analyze the characteristics, prognosis (death, transfer to hemodialysis) of elderly PD patients by comparing with younger group, assess the influencing factors of mortality in elderly patients.

2. Methods

Patients and groups

The study was approved by the ethics committee of the Cangzhou Central Hospital in Cangzhou City, Hebei Province, China. Written informed consent was attained from each patient. 202 patients who received continuous ambulatory peritoneal dialysis from January 2016 to January 2017 in the Department of Nephrology of Cangzhou Central Hospital were enrolled in this prospective observational study. They did the exchange themselves only after successful training by primary nurses. Conventional PD solutions (Dianeal 1.5%, 2.5%, or 4.25% dextrose; Baxter Healthcare, Guangzhou, China), Y-sets, and twin-bag systems were used. According to age, PD patients were divided into elderly group (age ≥ 65) and younger group. Elderly was defined as ≥ 65 years old at PD initiation. Patients should meet the following criteria: age > 18 years old, first peritoneal dialysis treatment and dialysis time more than 3 months, complete data. The exclusion criteria included incomplete data, loss of follow-up, those who did not cooperate with follow-up, transferred to other dialysis centers. Additionally, we did not include patients who were diagnosed as acute renal failure, multiple organ failure. End of follow-up was defined as the day of the outcome (death or transfer to hemodialysis) or end of follow-up (January 2020).

Demographic and clinical data collection

The demographic data including gender, age, body mass index (BMI), etiological diagnosis (diabetic nephropathy, hypertensive renal damage, primary glomerular disease), peritonitis or not, bacterial culture results (gram-positive or gram-negative bacilli) were collected. We also recorded outcomes of patients: continue PD, death, transfer to hemodialysis, kidney transplantation and time of death, reason of death, cause of transferring to hemodialysis. The Charlson index of comorbidity (CCI) was used to measure comorbidity [8]. The following variables were recorded: white blood cells (WBC), red blood cells (RBC), hemoglobin (HGB),

platelets (PLT), albumin, alanine aminotransferase (ALT), aspartate Aminotransferase (AST), total bilirubin, direct bilirubin, serum creatinine, urea nitrogen, serum calcium, potassium, sodium, ferritin, total Kt/V value, residual Kt/V, weekly creatinine clearance rate (WCcr), residual creatinine clearance rate, nPCR, dialysis time; whether the patients used angiotensin-converting enzyme inhibitors/angiotensin receptor blocker (ACEI / ARB) drugs, ever had cerebrovascular disease, IHD, heart failure or not.

Statistical analysis

Data analyses were performed using SPSS 23.0 and R 3.6.1 version. Results were expressed as mean $\pm$ standard deviation (SD), median and interquartile ranges, or numbers and percentages (%), as appropriate. Kolmogorov–Smirnov test was used to examine the normality of all numeric continuous variables. Independent sample t-tests were used if with a normal distribution, while nonparametric tests (Mann–Whitney

U test) were used to examine variables without a normal distribution,when comparing between groups. Categorical data were compared between elderly and younger groups by chi-square test. When there was only less than 5 observations in an group-outcome combination, Fisher's test was used.The Kaplan – meier survival curve was performed to investigate the difference of survival and transferring to hemodialysis between the elderly and the younger group. Multivariate Cox regression analysis was used to screen risk factor of mortality in elderly patients. Using the backward method, the entry probability was set to 0.05, and the removal probability was set to 0.10. R 3.6.1 version was constructed to draw nomogram map for building model to predict survival. All tests were two-sided, and P values < 0.05 were considered statistically significant.

3. Results

General characteristics of elderly patients

We enrolled 202 PD patients, including 61 in elderly group and 141 in younger group, with a median follow-up time of 44 months. Compared with the younger group, elderly patients were more likely to have lower albumin and nPCR, more comorbidity, especially with cardiovascular and cerebrovascular diseases .Diabetic nephropathy was common in elderly group, the difference was statistically significant ($P < 0.01$). Table 1 showed the baseline characteristics of patients grouped by age(see below).

Table 1
Basic characteristics of PD patients according to age

	All patients N = 202	Elderly group N = 61	Younger group N = 141	P
Demography				
Age,years	57.00(45.00,65.00)	72.00(66.50,76.00)	50.00(37.00,57.50)	< 0.001
Male,n (%)	121 (59.9)	34 (55.7)	87 (61.7)	.427
Laboratory parameter				
BMI,kg/m ²	23.97 ± 4.33	23.92 ± 3.54	24.00 ± 4.64	.903
WBC,x10 ⁹ /L	6.64(5.30,8.11)	7.13(5.33,9.32)	6.40(5.26,7.77)	.158
RBC,x10 ¹² /L	3.46 ± 0.59	3.36 ± 0.69	3.50 ± 0.54	.159
HGB,g/L	102.00(91.00,116.00)	100.00(85.00,115.50)	103.00(93.00,116.00)	.279
PLT, x10 ⁹ /L	209.50(163.00,251.25)	202.00(146.50,252.00)	212.00(165.00,251.00)	.281
ALB,g/L	33.70(37.20,40.10)	35.60(32.00,38.00)	38.40(34.95,41.00)	< 0.001
ALT,u/L	12.60(9.15,17.15)	13.00(9.80,16.90)	12.45(9.10,17.38)	.153
AST,u/L	15.80(12.00,20.30)	16.00(12.10,20.45)	15.45(12.00,20.15)	.171
TBIL,umol/L	4.10(4.95,6.60)	3.90(4.50,6.10)	4.38(5.10,6.60)	.151
DBIL,umol/L	1.70(1.20,2.30)	1.70(1.20,2.30)	1.70(1.20,2.30)	.759
Scr,umol/L	815.50(587.50,1039.75)	747.00(541.00,979.00)	607.00(828.00,1042.00)	.253
BUN,mmol/L	19.50(14.78,22.93)	18.20(13.90,22.20)	19.90(15.50,23.50)	.133
serum Ca,mmol/L	2.16(2.01,2.31)	2.12(1.96,2.29)	2.17(2.02,2.33)	.135
Serum K,mmol/L	4.20(3.70,4.80)	4.20(3.60,4.75)	4.20(3.80,4.80)	.414

PD = peritoneal dialysis;BMI = body mass index (weight/height²); WBC = white blood cell;

RBC = red blood cell;PLT = platelet;ALB = albumin;TBIL = total bilirubin;

DBIL = direct bilirubin;Scr = serum creatinine;BUN = blood urea nitrogen;

WCcr = weekly creatinine clearance rate;nPCR = normalized protein catabolic rate;

DM = diabetes mellitus;HRD = hypertensive renal damage;IHD = ischemic heart disease;

HF = heart failure;DN = Diabetic Nephropathy;CG = chronic glomerulonephritis;HD = hemodialysis

	All patients N = 202	Elderly group N = 61	Younger group N = 141	P
Serum Na,mmol/L	141.00(138.00,142.15)	141.00(137.00,143.00)	141.00(139.00,142.00)	.309
Ferritin,ug/L	269.70(137.70,573.50)	300.00(138.00,514.45)	257.00(136.25,584.80)	.725
Total kt/v	1.72(1.45,2.04)	1.69(1.50,2.04)	1.73(1.42,2.04)	.784
Residual kt/v	0.08(0.00,0.33)	0.10(0.01,0.32)	0.06(0.00,0.34)	.343
Total WCcr,ml/min	45.67(39.89,53.94)	46.09(40.77,55.15)	45.17(39.58,53.68)	.769
Residual Ccr,ml/min	2.00(0.00,9.79)	2.82(0.07,15.07)	1.19(0.00,10.29)	.235
nPCR,g/kg/day	0.82(0.71,0.92)	0.74(0.62,0.85)	0.85(0.76,0.93)	.003
Dialysis time	26.00(13.00,49.00)	21.00(9.50,46.00)	27.00(14.00,51.00)	.279
Charlson score	9.00(5.00,13.00)	11.00(6.50,15.00)	9.00(5.00,11.00)	.003
Medical and diseases history				
ISD,n(%)	58(28.7)	34(55.7)	24(17.0)	< .001
HF,n(%)	73(36.1)	27(44.3)	46(32.6)	.114
Cerebrovascular Disease,n(%)	40(19.8)	19(31.1)	21(14.9)	.008
peritoneal dialysis-related peritonitis n(%)	88(43.6)	21(34.4)	67(47.5)	.085
Gram positive, n(%)	32(15.8)	10(16.4)	22(15.6)	.888
Gram negative, n(%)	20(9.9)	8	12	.314
ACEI/ARB Use,n,(%)	74(36.6)	18(29.5)	56(39.7)	.167

PD = peritoneal dialysis;BMI = body mass index (weight/height²); WBC = white blood cell;

RBC = red blood cell;PLT = platelet;ALB = albumin;TBIL = total bilirubin;

DBIL = direct bilirubin;Scr = serum creatinine;BUN = blood urea nitrogen;

WCcr = weekly creatinine clearance rate;nPCR = normalized protein catabolic rate;

DM = diabetes mellitus;HRD = hypertensive renal damage;IHD = ischemic heart disease;

HF = heart failure;DN = Diabetic Nephropathy;CG = chronic glomerulonephritis;HD = hemodialysis

	All patients N = 202	Elderly group N = 61	Younger group N = 141	P
Primary disease				
DN,n(%)	46(22.8)	22(36.1)	24(20.6)	.003
HRD,n(%)	27(13.4)	11(18.0)	16(10.9)	.200
CG,n(%)	30(14.9)	6(9.8)	24(14.8)	.187
PD = peritoneal dialysis;BMI = body mass index (weight/height ²); WBC = white blood cell;				
RBC = red blood cell;PLT = platelet;ALB = albumin;TBIL = total bilirubin;				
DBIL = direct bilirubin;Scr = serum creatinine;BUN = blood urea nitrogen;				
WCcr = weekly creatinine clearance rate;nPCR = normalized protein catabolic rate;				
DM = diabetes mellitus;HRD = hypertensive renal damage;IHD = ischemic heart disease;				
HF = heart failure;DN = Diabetic Nephropathy;CG = chronic glomerulonephritis;HD = hemodialysis				

Outcomes Of Elderly Patients

To the end of the study, 121 (59.9%) patients continued to receive PD, 48 (23.8%) died, 25 (12.4%) were transferred to hemodialysis, 8 (3.9%) received a kidney transplantation. Among elderly group, 27 (44.3%) died, 2 (3.3%) were transferred to hemodialysis, and 1 (1.6%) received a kidney transplantation. Due to small number of kidney transplantation patients in elderly group, the following is not further discussed. Table 2 showed the outcomes of all peritoneal dialysis patients.

Table 2
Outcomes of patients

Outcomes	All patients N = 202	Elderly group N = 61	Younger group N = 141
Stay on PD,n(%)	121(59.9)	31(50.8)	90(63.8)
Death, n (%)	48 (23.8)	27 (44.3)	21 (14.9)
Transferred to HD, n (%)	25(12.4)	2(3.3)	23(16.3)
Kidney transplant, n (%)	8(3.9)	1(1.6)	7(5.0)

Mortality Of Elderly Patients

27 (44.3%) of elderly patients and 21 (14.3%) of younger patients died.(Fig. 1Kaplan–Meier survival curve between elderly and younger patients)The survival time of elderly group was significantly lower than that of younger group. The 1-year 2-year3-year and 4-year survival rate were 81.97%, 70.49%, 60.66% and 55.74%,respectively. The main cause of death in elderly PD patients was cardiovascular disease, while in younger group was infection.(Table 3)

Table 3
Causes of Death

Death	Elderly group	Younger group
	Cases,n(%)	Cases,n(%)
Cardiovascular disease	10(37.0)	4(19.0)
Cerebrovascular disease	4(14.8)	4(19.0)
Infection	3(11.1)	6(28.6)
Cachexia	3(11.1)	2(9.5)
Other	7(25.9)	5(23.8)

Transferring To Hd In Elderly Patients

2(3.3%)elderly PD patients, 23(16.3%) younger group transferred to HD. Elderly patients with PD were less likely to convert to HD. Peritonitis was the chief reason for both elderly and younger PD patients to transfer to HD. (Table 4)

Table 4
Causes of transferring to HD

Transfer to HD	Elderly group	Younger group
	Cases,n(%)	Cases,n(%)
Peritonitis	2(3.3)	18(12.8)
Inadequate dialysis	0	2(1.4)
Dialysis failure	0	2(1.4)
Others	0	1(0.7)

prognostic mortality factors in elderly PD patients

For elderly PD patients, univariate analysis Cox regression analysis indicated that BMI, CCI, previous IHD, peritonitis, dialysis time were the risk factors for the long-term survival of the elderly PD patients. BMI, CCI and IHD remained independent significant predictors of long-term survival of elderly PD patients by multivariate Cox regression analysis (Table 5 and Fig. 2). The impact of peritonitis,dialysis time disappeared in multivariate

analysis. The survival model of elderly PD was established by R 3.6.1 version (Fig. 3). CCI were related to peritonitis ($r = 0.471$, $P < 0.001$) and dialysis time ($r = 0.260$, $P = 0.043$), these may be a confounding factor.

Table 5
Independent relative risk of death in elderly peritoneal dialysis patients

univariate analysis					multivariate analysis				
95%CI					95%CI				
Factor	P	HR	Lower Limit	Upper Limit	P	HR	Lower Limit	Upper Limit	
BMI	.016	1.139	1.025	1.265	.005	1.0	1.0	1.1	
CCI	.018	1.073	1.012	1.137	.022	1.1	1.0	1.2	
ISD	.011	3.084	1.301	7.315	.042	2.5	1.0	6.1	
Peritonitis	.042	2.305	1.031	5.154	.284	N/A	N/A	N/A	
Dialysis Time	.089	1.014	.998	1.030	.310	N/A	N/A	N/A	

4. Discussion

With the aging of the population, spiraling rates of end stage renal failure have increased and the elderly needing for PD is gradually increasing [9]. In order to improve the prognosis and quality of life in elderly patients with PD, it is especially important to explore the survival status of elderly patients and actively prevent and control the risk factors.

We analyzed the general condition of the elderly patients, the reasons for withdrawal from PD therapy, and the risk factors for survival of patients, to provide guidance for improving prognosis and prolonging the survival time of patients on PD with ESRD.

Compared with the younger group, elderly group of PD patients tended to have hypoalbuminemia and low nPCR, both of which are important indicators to evaluate the nutrition of patients [10], this result was consistent with the Kholshali et al study [11]. Wang XX et al also reported that serum albumin remained lower in elderly patients [12]. In the course of PD treatment, a large number of nutrients are lost in dialysis solution, and the absorption capacity of gastrointestinal function is decreased in elderly patients, leading to malnutrition. Peritoneal dialysis does not need to establish vascular access, has little impact on hemodynamics, so elderly patients with diabetes tend to choose PD. Arteriosclerosis and the incidence of cardiovascular/cerebrovascular events is often occurred in diabetic nephropathy [13, 14]. Age is also a risk factor of vascular calcification [15]. Some studies have shown that elderly PD patients are inclined to develop peritonitis [16], but there was no difference in the incidence of peritonitis between elderly and younger group in this study, which was consistent with a Canada study. It was reported that elderly PD patients who were 70 years old or more had no relationship with peritonitis [17], considering that elderly patients had better family/social support and was treated with advanced antibiotics.

In this study, the mortality of elderly patients with PD was significantly lower than that of younger patients. Age is a risk factor for survival of PD patients [4, 18]. The 1-, 3-, and 5-year survival rates of elderly PD patients were as follows: 81.97%, 70.49%, 60.66%, 55.74%. A study in China found that the 1-, 3-, and 5-year mortality of elderly patients with PD respectively were 79%, 56% and 30% [19], which is generally consistent with our results. Cardiovascular disease is the leading cause of death, 53.4% of elderly PD patients existed with cardiovascular complications in a retrospective study [19]. The USRDS data showed that cardiovascular events were the main cause of death in dialysis patients [20], which was related with the metabolic abnormalities and the increased incidence of diabetic nephropathy in elderly PD patients.

In this study, we provided evidence that high BMI, high CCI and IHD were independent risk factors for predicting death in elderly PD patients. Patients with high BMI is inclined to have high risk of dialysis [21], and obesity can increase cardiovascular and cerebrovascular events, thus increasing the risk of death [22]. However, the relationship between BMI and survival in elderly patients with PD is controversial. Some studies found that obese PD patients lived longer than those with lower BMI [23, 24], considering that patients with lower BMI will increase protein consumption and inflammation, and high BMI can increase hemodynamic stability and the isolation of fat tissue from toxins [25]. Another study found that baseline BMI was not related to the prognosis of peritoneal dialysis [26], so more studies are needed to explore the relationship between BMI and death in elderly PD patients. As mentioned, CCI is an important index to evaluate the complications. Sandrine genestier et al showed that CCI was positively correlated with the mortality of elderly patients [27]. A retrospective analysis of 292 patients found that 2-year survival rate of elderly patients with CCI score > 9 was 38%, and that of patients with CCI < 7 reached 69% [28]. Elderly PD patients often suffer with IHD, and the risk of recurrence with cardiovascular and cerebrovascular events increases [29], so elderly PD patients are at high risk of death.

In this study, peritonitis was main cause of transferring to hemodialysis in both elderly and younger patients. Minli et al. [30] and Sakaci et al. [31] also found that the main cause of turning to hemodialysis was peritonitis. Elderly patients turning to hemodialysis was smaller than that of younger patients. Elderly patients had a higher risk of peritonitis-related and all-cause mortality. In a recent study from Brazil, the only factor associated with death during peritoneal infection was older age [32], so elderly patients had no chance to transfer to HD.

This study elucidated the characteristics of elderly PD patients, the prognosis of elderly PD patients and the risk factors for predicting death, which help nephrologists better understand the characteristics of elderly PD patients and improve the prognosis of patients. However, this study has some limitations. First of all, this study is a single center, relatively small sample study, which can not represent the situation of elderly PD patients in China, additional multiple studies are needed in larger populations to confirm the conclusion of our study. In addition, we did not analyze economic and social situation of each patient.

5. Conclusions

In conclusion, compared with younger group, elderly PD patients had poor nutritional status, more complications, majority of diabetic nephropathy and high mortality. High BMI, high CCI and previous IHD events were the independent risk factors to predict mortality of elderly PD patients. Cardiovascular disease was main cause of death in elderly group. Peritonitis was major reason for transferring to HD in PD patients.

Abbreviations

PD=peritoneal dialysis;HD=hemodialysis; nPCR=normalized protein catabolic rate;CCI= Charlson Comorbidity Index; BMI=body mass index; WBC= white blood cell; RBC=red blood cell;PLT = platelet;ALB =albumin;TBIL=totalbilirubin;DBIL=direct bilirubin;Scr=serumcreatinine;BUN=blood urea nitrogen;WCcr=weekly creatinine clearance rate; DM=diabetesmellitus;HRD=hypertensive renal damage;IHD=ischemic heart disease;HF=heart failure;DN =Diabetic Nephropathy;CG =chronic glomerulonephritis;SD=standard deviation

Declarations

Ethics approval and consent to participate

The study was approved by the ethics committee of the Cangzhou Central Hospital in Cangzhou City, Hebei Province, China. Written informed consent was attained from each patient.

Consent for publication

Not applicable

Availability of data and materials

The datasets generated and/or analyzed during the current study are not publicly available due to local regulations on the management of medical records but are available from the corresponding author on reasonable request.

Conflict of interest

None declared

Funding

Not applicable

Authors' Contributions

SF and MX: project development, data collection, and management, data analysis, manuscript writing and revising; XH: data collection, data analysis; XH: manuscript editing and revising; All authors have read and approved the manuscript.

Acknowledgements

We would like to thank the nursing staff of the PD unit for inspiration and helpful discussions.

References

1. Zhang L, Wang F, Wang L, Wang W, Liu B, Liu J, Chen M, He Q, Liao Y, Yu X *et al*: **Prevalence of chronic kidney disease in China: a cross-sectional survey**. *Lancet* 2012, **379**(9818):815-822.

2. Brown EA, Johansson L: **Epidemiology and management of end-stage renal disease in the elderly.** *Nat Rev Nephrol* 2011, **7**(10):591-598.
3. Dimkovic NB, Prakash S, Roscoe J, Brissenden J, Tam P, Bargman J, Vas SI, Oreopoulos DG: **Chronic peritoneal dialysis in octogenarians.** *Nephrol Dial Transplant* 2001, **16**(10):2034-2040.
4. Hung CC, Chang CT, Lee CC, Chen KH, Yu CC, Wu CH, Huang JY, Wu MS, Yang CW: **Prognostic predictors of technique and patient survival in elderly Southeast Asian patients undergoing continuous ambulatory peritoneal dialysis.** *Int J Clin Pract* 2009, **63**(2):254-260.
5. Weinhandl ED, Foley RN, Gilbertson DT, Arneson TJ, Snyder JJ, Collins AJ: **Propensity-matched mortality comparison of incident hemodialysis and peritoneal dialysis patients.** *J Am Soc Nephrol* 2010, **21**(3):499-506.
6. Mizuno M, Ito Y, Tanaka A, Suzuki Y, Hiramatsu H, Watanabe M, Tsuruta Y, Matsuoka T, Ito I, Tamai H *et al*: **Peritonitis is still an important factor for withdrawal from peritoneal dialysis therapy in the Tokai area of Japan.** *Clin Exp Nephrol* 2011, **15**(5):727-737.
7. Luo Q, Xia X, Lin Z, Lin J, Yang X, Huang F, Yu X: **Very early withdrawal from treatment in patients starting peritoneal dialysis.** *Ren Fail* 2018, **40**(1):8-14.
8. Beddhu S, Zeidel ML, Saul M, Seddon P, Samore MH, Stoddard GJ, Bruns FJ: **The effects of comorbid conditions on the outcomes of patients undergoing peritoneal dialysis.** *Am J Med* 2002, **112**(9):696-701.
9. Disney AP: **Demography and survival of patients receiving treatment for chronic renal failure in Australia and New Zealand: report on dialysis and renal transplantation treatment from the Australia and New Zealand Dialysis and Transplant Registry.** *Am J Kidney Dis* 1995, **25**(1):165-175.
10. Grzegorzewska A, Dobrowolska-Zachwieja A: **[Comparison of indices of kinetic modelling of urea and creatinine as well as evaluation of nutritional status of patients treated with continuous ambulatory peritoneal dialysis].** *Pol Arch Med Wewn* 1994, **91**(6):417-426.
11. Khoshhali M, Kazemi I, Hosseini SM, Seirafian S: **Relationship between trajectories of serum albumin levels and technique failure according to diabetic status in peritoneal dialysis patients: A joint modeling approach.** *Kidney Res Clin Pract* 2017, **36**(2):182-191.
12. Wang X, Han Q, Wang T, Tang W: **Serum albumin changes and mortality risk of peritoneal dialysis patients.** *Int Urol Nephrol* 2020, **52**(3):565-571.
13. Yao Q, Lindholm B, Heimbürger O: **Peritoneal dialysis prescription for diabetic patients.** *Perit Dial Int* 2005, **25 Suppl 3**:S76-79.
14. Hadjadj S, Cariou B, Fumeron F, Gand E, Charpentier G, Roussel R, Kasmi AA, Gautier JF, Mohammedi K, Gourdy P *et al*: **Death, end-stage renal disease and renal function decline in patients with diabetic nephropathy in French cohorts of type 1 and type 2 diabetes.** *Diabetologia* 2016, **59**(1):208-216.
15. Ma D, Yan H, Yang X, Yu Z, Ni Z, Fang W: **Abdominal aortic calcification score as a predictor of clinical outcome in peritoneal dialysis patients: a prospective cohort study.** *BMC Nephrol* 2020, **21**(1):151.
16. Wu H, Ye H, Huang R, Yi C, Wu J, Yu X, Yang X: **Incidence and risk factors of peritoneal dialysis-related peritonitis in elderly patients: A retrospective clinical study.** *Perit Dial Int* 2020, **40**(1):26-33.
17. Nessim SJ, Bargman JM, Austin PC, Story K, Jassal SV: **Impact of age on peritonitis risk in peritoneal dialysis patients: an era effect.** *Clinical journal of the American Society of Nephrology : CJASN* 2009,

4(1):135-141.

18. Iqbal MM, Islam MN, Mansur MA, Naeem GM, Sattar H, Hossain RM, Mohsin M, Rahman MH, Rashid HU: **Outcome of peritoneal dialysis and hemodialysis in elderly patients with diabetes: early experience from Bangladesh.** *Adv Perit Dial* 2005, **21**:85-89.
19. Joshi U, Guo Q, Yi C, Huang R, Li Z, Yu X, Yang X: **Clinical outcomes in elderly patients on chronic peritoneal dialysis: a retrospective study from a single center in china.** *Perit Dial Int* 2014, **34**(3):299-307.
20. **US Renal Data System 2016 Annual Data Report: Epidemiology of Kidney Disease in the United States.** *Am J Kidney Dis* 2017, **69**(3s1):A4.
21. Kalantar-Zadeh K, Kopple JD: **Body mass index and risk for end-stage renal disease.** *Ann Intern Med* 2006, **144**(9):701; author reply 701-702.
22. Goodkin DA, Mapes DL, Held PJ: **The dialysis outcomes and practice patterns study (DOPPS): how can we improve the care of hemodialysis patients?** *Semin Dial* 2001, **14**(3):157-159.
23. Chung SH, Lindholm B, Lee HB: **Influence of initial nutritional status on continuous ambulatory peritoneal dialysis patient survival.** *Perit Dial Int* 2000, **20**(1):19-26.
24. McCusker FX, Teehan BP, Thorpe KE, Keshaviah PR, Churchill DN: **How much peritoneal dialysis is required for the maintenance of a good nutritional state? Canada-USA (CANUSA) Peritoneal Dialysis Study Group.** *Kidney Int Suppl* 1996, **56**:S56-61.
25. Park J, Ahmadi SF, Streja E, Molnar MZ, Flegal KM, Gillen D, Kovesdy CP, Kalantar-Zadeh K: **Obesity paradox in end-stage kidney disease patients.** *Prog Cardiovasc Dis* 2014, **56**(4):415-425.
26. Badve SV, Paul SK, Klein K, Clayton PA, Hawley CM, Brown FG, Boudville N, Polkinghorne KR, McDonald SP, Johnson DW: **The association between body mass index and mortality in incident dialysis patients.** *PLoS One* 2014, **9**(12):e114897.
27. Genestier S, Meyer N, Chantrel F, Alenabi F, Brignon P, Maaz M, Muller S, Faller B: **Prognostic survival factors in elderly renal failure patients treated with peritoneal dialysis: a nine-year retrospective study.** *Perit Dial Int* 2010, **30**(2):218-226.
28. Vrtovsnik F, Porcher R, Michel C, Hufnagel G, Queffeuilou G, Mentz F, Mignon F: **Survival of elderly patients on peritoneal dialysis: retrospective study of 292 patients, from 1982 to 1999.** *Perit Dial Int* 2002, **22**(1):73-81.
29. Abu-Assi E, Lopez-Lopez A, Gonzalez-Salvado V, Redondo-Díaz A, Peña-Gil C, Bouzas-Cruz N, Raposeiras-Roubín S, Riziq-Yousef Abumuaileq R, García-Acuña JM, Gonzalez-Juanatey JR: **The Risk of Cardiovascular Events After an Acute Coronary Event Remains High, Especially During the First Year, Despite Revascularization.** *Rev Esp Cardiol (Engl Ed)* 2016, **69**(1):11-18.
30. Li M, Yan J, Zhang H, Wu Q, Wang J, Liu J, Xing C, Zhou Y: **Analysis of outcome and factors correlated with maintenance peritoneal dialysis.** *J Int Med Res* 2019, **47**(10):4683-4690.
31. Sakac T, Ahbap E, Koc Y, Basturk T, Ucar ZA, Şengül A, Şevinc M, Kara E, Akgöl C, Kayalar AO *et al*: **Clinical outcomes and mortality in elderly peritoneal dialysis patients.** *Clinics (Sao Paulo)* 2015, **70**(5):363-368.
32. de Moraes TP, Olandoski M, Caramori JC, Martin LC, Fernandes N, Divino-Filho JC, Pecoits-Filho R, Barretti P: **Novel predictors of peritonitis-related outcomes in the BRAZPD cohort.** *Perit Dial Int* 2014, **34**(2):179-

Figures

Figure 1

Kaplan–Meier survival curve according to age Time means follow-up time(months)

Figure 2

Independent relative risk of death in elderly peritoneal dialysis patients by multivariate analysis

Figure 3

Joint modeling for prognostic predictors of survival in elderly patients

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [STROBEchecklist.docx](#)