Complete Genome Sequence of a Novel Mitovirus From the Phytopathogenic Fungus *Fusarium Oxysporum*

Jing wang

Henan Academy of Agricultural Sciences <https://orcid.org/0000-0001-8247-6085>

Rui Qiu

Henan Academy of Agricultural Sciences

Chengjun Li

Henan Academy of Agricultural Sciences

Shujun Li (✉ 13603749396@126.com)

Henan Academy of Agricultural Sciences <https://orcid.org/0000-0003-0778-3186>

Xiaojie Li

Henan Academy of Agricultural Sciences

Jun Zhao

Henan Academy of Agricultural Sciences

jingke Bai

Henan Academy of Agricultural Sciences

Yuguo Chen

Henan Academy of Agricultural Sciences

Research Article

Keywords: Fusarium oxysporum, mycovirus, Mitovirus, genomic structure

Posted Date: April 28th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-416498/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Fusarium oxysporum is a cosmopolitan plant pathogen causing Fusarium wilt and Fusarium root rot in many economically important crops. There is still limited information about mycoviruses that infect *F. oxysporum*. Here, a novel mitovirus tentatively named Fusarium oxysporum mitovirus 1 (FoMV1) was identified from *F. oxysporum* strain B2-10. The genome of FoMV1 is 2,453 nt in length with a predicted AU content of 71.6%, and contains one large open reading frame (ORF) using the fungal mitochondrial genetic code. The ORF encodes RNA-dependent RNA polymerases (RdRp) of 723 amino acids with a molecular mass of 84.98 kDa. The RdRp domain of FoMV1 shares 29.01–68.43% sequence similarity to the members of the family *Mitoviridae*. Phylogenetic analysis further suggested that FoMV1 is a new member of a distinct species in the genus *Mitovirus*.

Introduction

Fusarium oxysporum is one of the most destructive pathogen in agricultural production that can infect more than 120 plant species including some economically important crops such as cotton, tomato, banana, and tobacco [1, 2]. This species complex not only causes vascular disease in a large number of economically important crops, but also causes Fusarium root rot in different *Solanaceae* species [3, 4]. The typical symptoms of Fusarium root rot in tobacco include the chlorosis of lower leaves, decaying of root and vascular discoloration. Generally, stem and taproot showed reddish to brown vascular discoloration, and the whole plant died in the end [5]. This pathogen was soil bored and could be detected both inside and exterior of the seed. Moreover, it can interact with nematodes of parasitic root knot and cyst nematodes to increase disease, and also survive in the soil for years in the absence of a host crop [6]. Although the fumigants were the most consistently effective management tactics against *F. oxysporum*, the application of methyl bromide and chloropicrin resulted in poor wrapper leaf quality and burn [6]. Hence, resistance varieties are also the most effective and economical means of reducing disease [7].

Mycoviruses are viruses that infect fungi, whose hosts include phytopathogenic and entomopathogenic fungi [8-12]. Most of the mycoviruses are cryptic infections and seldom produce symptoms [9]. Some mycoviruses belong to different viral families who possess variable effects on the morphological changes of their host, including reduced mycelia growth and increased pigmentation [13, 14]. Mycovirus-induced hypovirulence is very attractive that appeals to pathologists most. The successful case of applying hypovirulent strains of *Cryphonectria parasitica* to combat chestnut blight in Europe has provided impetus for exploiting mycoviruses as virocontrol agents [15]. The other mycoviruses with significant biocontrol effects include *Sclerotinia sclerotiorum* hypovirulence-associated DNA virus 1 (SsHADV1) and *Rosellinia necatrix* megabirnavirus 1 (RnMBV1) [16-19]. With the development of high-throughput sequencing technology, more and more novel mycoviruses could be identified. Hence, there will be more chances to screening for novel fungal viruses that could be used as biological control agents.

The knowledge of mycoviruses in phytopathogenic species *Fusarium oxysporum* is still very limited. Four dsRNA segments were first detected in the hypovirulent strain of *F. oxysporum* isolated from soybean seedling, but the mycoviruses corresponding to dsRNA have not been identified [20]. Two mycoviruses *Fusarium oxysporum* f. sp. *dianthi* mycovirus 1 (FodV1) and *Fusarium oxysporum* ourmia-like virus 1 (FoOuLV1) associated with hypovirulence were identified and characterized in *F. oxysporum* isolated from diseased carnation plant and bitter melon, respectively [21, 22]. After that, some novel mycoviruses have been reported, such as *Fusarium oxysporum* f. sp. *dianthi* hypovirus 2 (FodHV2) and *Fusarium oxysporum* f. sp. *dianthi* mitovirus 1 (FodMV1). FodHV2 belongs to the family *Hypoviridae* but did not alter the vegetative growth or virulence of the host [23, 24]. A novel polomyco-related virus termed Hadaka virus 1 (HadV1) identified from *F. oxysporum*. HadV1 has 11(+) RNA genomic segments and lacks a typical structural protein that reveals a potential novel lifestyle of multi-segmented RNA viruses [25]. All these results indicated that the mycoviruses harbored by *F. oxysporum* have a rich diversity, and some of them have potential as biological control agents. Here, a novel mitovirus was identified in strain B2-10 of *F. oxysporum*, and described the characterization of the genome, and tentatively named as “*Fusarium oxysporum* mitovirus 1” (FoMV1).

Provenance and sequencing of strains

The strain B2-10 of *F. oxysporum* was isolated from the diseased root of a tobacco plant with typical symptoms of *Fusarium* root rot, which was collected from Xuchang city, Henan province, China, in 2019. The strain was purified by single spore isolation technique. The specific primers of translation elongation factor 1-alpha (EF-1α) and *F. oxysporum* were used for *Fusarium* species identification [26]. Mycelial agar plugs of strain B2-10 were transferred to a PDA plate (9 cm diam) covered with cellophane membrane and cultured at 25 °C in the dark for four to six days. Mycelial mats were harvested using a sterilized medicine spoon and stored at -70 °C until use. Total RNA was extracted from 1.0 g of mycelium using an RNAiso kit (TaKaRa, China). Total RNA of strains were sent to Shanghai Bohao Biotechnology Co., Ltd. for high-throughput sequencing on an Illumina HiSeq 2500 platform. The rRNA was depleted using a Ribo-Zero™ rRNA Removal Kit (Illumina, CA, USA). Detailed parameters used in the bioinformatics pipeline according to the procedures described by Wang et al. (2020) [27]. Finally, the contigs that were identical or complementary to the viral genomic sequence were identified as putative viral sequences.

The cDNA of strain B2-10 was synthesized using the PrimerScript™1st Strand cDNA Synthesis Kit (TaKaRa, China) following the manufacturer's instruction. The specific primers (F1: 5'-AGGTCAACCAATGGGAAC-3'; R1: 5'-CCAACAAAAGTAGGATAG-3') were designed to verify the presence of newly discovered mitovirus FoMV1 with the PCR amplicon of 756 bp in length. The system and step of RT-PCR were conducted refer to the procedure described by Wang et al. (2020) [27]. A SMARTer^R RACE 5'/3' Kit (TakaRa, China) was used to complete the 5'- and 3'-terminal genomic sequence with the help of gene-specific primers (GSPs). GSP-763F1 (5'-GGAATACCAGTAAGAGGGATA-3') and GSP-763F2 (5'-GTAAAAGCCTAGAGGTTGGT-3') were used for 3'-RACE as inner and outer primers, respectively. Similarly, GSP-763R1 (5'-AAAGAGAAAAGGGTTTGAAC-3') and GSP-763R2 (5'-CTTGCAGTCCTATGGTCTAC-3') were

used for 5'-RACE as inner and outer primers, respectively. The procedure was performed following the kit manufacturer's instruction. The target PCR product was gel-purified by MiniBEST Agarose Gel DNA Extraction Kit (TaKaRa, China), and cloned into the pMD19-T vector (Sangon Biotech, China), then introduced into *Escherichia coli* Trelief 5α (TSINGKE Biotech, China). The target clones were sent to Sangon Biotech for sequencing.

Putative open reading frames (ORFs) were predicted using ORF Finder in the NCBI database (<https://www.ncbi.nlm.nih.gov/orffinder/>). The potential secondary structures for the 5'-terminal and 3'-terminal nucleotide sequences of FoMV1 (positive strand) were predicted using RNAfold WebServer (<http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi>). Multiple sequence alignment of the RdRp sequences encoded by FoMV1 and other mitoviruses were performed using DNAMAN software (Version 9) and CLUSTALX program (version 2.1). A phylogenetic tree was constructed using the neighbor-joining method with a bootstrap value of 1000 replicates through the MEGA-X program (Version 10.1.8).

Sequence properties

The complete genome sequence of FoMV1 obtained by assembling partial-length cDNAs that amplified from the total RNA, the 5'- and 3'-terminal sequences were determined Using RT-PCR with oligo (dT) primers and rapid amplification of cDNA ends. Then, the full-length sequence of FoMV1 has been submitted to the GenBank database with the accession number MW690927 (not yet released).

The full-length cDNA sequence of FoMV1 is 2,453 nt long and is relatively high A+U content of 71.6%. The AU-rich (71.6%) of FoMV1 is higher than the AU-rich of *Fusarium andiyazi* mitovirus 1 and *Fusarium oxysporum* f. sp. *dianthi* mitovirus 1 with 70.5% and 58.8%, respectively [28, 24]. AU-richness is a common characteristic of fungal and plant mitochondrial mitovirus [29]. The nucleotide sequence of FoMV1 was examined by ORF Finder using the fungal mitochondrial codon usages. The positive strand was found to contain one major large ORF (nt 241-2,412) and one small ORF (nt 83-172). The untranslated regions (UTRs) at the 5'- and 3'- ends are 240 nt and 41 nt long, respectively (Fig. 1A). The large ORF encodes a 723-aa protein with a calculated molecular weight of 84.98 KDa and a predicted pI of 9.83. A total of 13 UGA-encoded tryptophans were found in this ORF [30]. A small ORF near the 5'-terminus encodes a protein of 29-aa with a calculated molecular weight of 3.31 KDa and a predicted pI of 8.94. The 5'- and 3'- UTRs of FoMV1 were predicted for potential secondary structure. The result indicated that the 5'-terminal sequence (nt 4-47) could be folded into three stem-loop structure with a ΔG value of -15.60 kcal/mol (Fig. 1B). The 3'-terminal sequence (nt 2,415-2,453) could be folded into a stable stem-loop structure with a ΔG value of -8.70 kcal/mol (Fig. 1B). Moreover, the 5'- and 3'- UTRs of the positive strand of FoMV1 had an inverted complementarity, which a potentially stable panhandle structure was predicted with a ΔG value of -11.00 kcal/mol (Fig. 1B).

A search of BLASTX in NCBI indicated that the internal 2,150 nt region (241-2,391) of the FoMV1 sequence is 74.41% identical (E-value 0.0) to the full-length genome sequence (229-2,386) of *Fusarium andiyazi* mitovirus 1 (GenBank Acc. No. QIQ28423). A homology search with BLASTP showed that this amino acid sequence of RdRp of FoMV1 is most closely related to the RdRp of *Fusarium andiyazi*

mitovirus 1, *Fusarium circinatum* mitovirus 2, *Fusarium sacchari* mitovirus 1, and *Fusarium poae* mitovirus 1 with the identity of 45.19%-68.43% (E-value 0.0). A conserved domain database (CDD) search confirmed that position 925-2,136 nt of the RdRp of FoMV1 contained a conserved domain Mitovir_RNA_pol (pfam05919). Furthermore, a multiple alignment of amino acid sequences of RdRp between FoMV1 and other mitoviruses suggested that it contained six conserved motifs (Fig. 2). That is a characteristic of the RdRps of mitochondrial viruses [31, 32]. Therefore, according to the rules of species demarcation criteria about mitoviruses defined by the ICTV (<https://talk.ictvonline.org/>), suggesting that FoMV1 should be considered a novel mitovirus. It was a tentative member of a new species that belonged to the family Mitoviridae. However, there were no proteins or polypeptides homologous to the putative protein sequence of the small ORF. In addition, the presence of small ORF coding for polypeptides with unknown function near the 5'-terminal region has been previously reported in other mycovirus, including *Botrytis cinerea* mitovirus 1 [33], *Helminthosporium victoriae* 190S virus [34], *Macrophomina phaseolina* victorivirus 1 [35], and *Rosellinia necatrix* megabirnavirus 1 [18].

A phylogenetic tree based on the full length amino acid sequence of the viral RdRp of FoMV1 and 25 other selected mitoviruses were constructed using the Neighbor-Joining method (Fig. 3). FoMV1 clustered with *sacchari* mitovirus 1, *Fusarium andiyazi* mitovirus 1, and *Fusarium circinatum* mitovirus 2-1 to form a clade that clustered with *Fusarium circinatum* mitovirus 1, *Fusarium globosum* mitovirus 1, *Fusarium coeruleum* mitovirus 1, and *Fusarium poae* mitovirus 2 to form a large clade that were all mitochondrial viruses infecting *Fusarium* species. *Fusarium oxysporum* f. sp. *dianthi* mitovirus 1 (FodMV1) is the first mitovirus that identified in *F. oxysporum* [24]. However, FoMV1 didn't cluster with FodMV1, but with mitoviruses that infects *Fusarium* sp. This result of the phylogenetic analysis of RdRp sequences suggested that FoMV1 is a novel member of the family *Mitoviridae*.

Declarations

Funding:

This research work is supported by the Science and Technology Project of Henan Provincial Tobacco Company (2020410000270012). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Availability of data and material

The datasets used or analyzed during the current study are available from the corresponding author on reasonable request.

Conflicts of Interest/Competing interests

The authors declare no conflict of interest.

Author Contributions:

J.W., and S.J.L., conceived and designed the experiments; J.W. performed the experiments; R.Q., C.J.L., X.J.L., J.Z., J.K.B., and Y.G.C., collected the samples; J.W. wrote the paper; S.J.L., and R.Q., revised the paper.

Ethical approval

This article does not contain any studies with human participants or animals performed by any of the authors.

References

1. Michielse CB, Rep M (2009) Pathogen profile update: *Fusarium oxysporum*. Mol Plant Pathol 10:311-324. <https://doi.org/10.1111/j.1364-3703.2009.00538.x>
2. Pietro AD, Madrid MP, Caracuel Z, Delgado-Jarana J, Roncero MIG (2003) *Fusarium oxysporum*: exploring the molecular arsenal of a vascular wilt fungus. Mol Plant Pathol 4:315-325. <https://doi.org/10.1046/j.1364-3703.2003.00180.x>
3. Armstrong GM, Armstrong JK (1968) Formae speciales and races of *Fusarium oxysporum* causing a tracheomycosis in the syndrome diseases. Phytopathology 58:1242-1246. <https://doi.org/10.1094/phyto-68-19>
4. Chehri K, Salleh B, Zakaria L (2015) Morphological and phylogenetic analysis of *Fusarium solani* species complex in Malaysia. Microbial Ecol 69:457-471. <https://doi.org/10.1007/s00248-014-0494-2>
5. Yang M, Cao JD, Zheng YX, Zhou HF, He MD, Duan J, Tong WJ, Deng XP, Chen XL (2020) First Report of *Fusarium* Root Rot of Tobacco Caused by *Fusarium solani* in Lincang, China. Plant Dis 104:1541. <https://doi.org/10.1094/PDIS-08-19-1711-PDN>
6. LaMondia JA (2015) *Fusarium* wilt of tobacco. Crop Protection 73:73-77. <https://doi.org/10.1016/j.cropro.2015.03.003>
7. Snow D, Lucas GB (1966) Diseases of Tobacco. Journal of Applied Ecology 3:409. <https://doi.org/10.2307/2401265>
8. Ghabrial S, Suzuki N (2009) Viruses of plant pathogenic fungi. Annu Rev Phytopathol 47:353-384. <https://doi.org/10.1146/annurev-phyto-080508-081932>
9. Kotta-Loizou I, Coutts RHA (2017) Studies on the virome of the entomopathogenic fungus *Beauveria bassiana* reveal novel dsRNA elements and mild hypervirulence. PLoS Pathog 13:e1006183. <https://doi.org/10.1371/journal.ppat.1006183>
10. Son M, Yu J, Kim KH (2015) Five questions about mycoviruses. PLoS Pathog. <https://doi.org/10.1371/journal.ppat.1005172>
11. Herrero N, Dueñas E, Quesada-Moraga E, Zabalgoitia I (2012) Prevalence and diversity of viruses in the entomopathogenic fungus *Beauveria bassiana*. Appl Environ Microbiol 78:8523-8530. <https://doi.org/10.1128/AEM.01954-12>

12. Ghabrial SA, Casto'n JR, Jiang D, Nibert ML, Suzuki N (2015) 50-plus years of fungal viruses. *Virology* 479:356-368. <https://doi.org/10.1016/j.virol.2015.02.034>
13. Deng F, Xu R, Boland GJ (2003) Hypovirulence-associated double-stranded RNA from *Sclerotinia homoeocarpa* is conspecific with Ophiostoma novo-ulmi mitovirus 3a-Ld. *Phytopathology* 93:1407-1414. <https://doi.org/10.1094/PHYTO.2003.93.11.1407>
14. Xie J, Jiang D (2014) New insights into mycoviruses and exploration for the biological control of crop fungal diseases. *Annu Rev Phytopathol* 52:45-68. <https://doi.org/10.1146/annurev-phyto-102313-050222>
15. Hillman BI, Suzuki N (2004) Viruses of the chestnut blight fungus, *Cryphonectria parasitica*. *Adv Virus Res* 63:423-472. [https://doi.org/10.1016/S0065-3527\(04\)63007-7](https://doi.org/10.1016/S0065-3527(04)63007-7)
16. Xiao Y, Li B, Fu Y, Jiang D, Ghabrial SA, Li G, Peng Y, Xie J, Cheng J, Huang J, Yi X (2010) A geminivirus-related DNA mycovirus that confers hypovirulence to a plant pathogenic fungus. *Proc Natl Acad Sci USA* 107:8387-8392. [https://doi.org/10.1016/S0065-3527\(04\)63007-7](https://doi.org/10.1016/S0065-3527(04)63007-7)
17. Xiao Y, Li B, Fu Y, Xie J, Cheng J, Ghabrial SA, et al. (2013) Extracellular transmission of a dna mycovirus and its use as a natural fungicide. *Proc Natl Acad Sci USA* 110(4):1452-1457. <https://doi.org/10.1073/pnas.1213755110>
18. Chiba S, Salaipeth L, Lin YH, Sasaki A, Kanematsu S, Suzuki N (2009) A novel bipartite double-stranded RNA mycovirus from the white root rot fungus *Rosellinia necatrix*: molecular and biological characterization, taxonomic considerations, and potential for biological control. *J Virol* 83:12801-12812. <https://doi.org/10.1128/JVI.01830-09>
19. Salaipeth L, Eusebio-Cope A, Chiba S, Kanematsu S, Suzuki N (2014) Biological properties and expression strategy of *Rosellinia necatrix* megabirnavirus 1 in an experimental host *Cryphonectria parasitica*. *J Virol* 95:740-750. <https://doi.org/10.1099/vir.0.058164-0>
20. Kilic O, Griffin GJ (1998) Effect of dsRNA-containing and dsRNA-free hypovirulent isolates of *Fusarium oxysporum* on severity of fusarium seedling disease of soybean in naturally infested soil. *Plant & Soil* 201:125-135. <https://doi.org/10.1023/A:1004319614390>
21. Lemus-Minor CG, Carmen CM, García-Pedrajas MD, Pérez-Artés E (2019) Horizontal and vertical transmission of the hypovirulence-associated mycovirus *Fusarium oxysporum* f. sp. *dianthi* virus. *Eur J Plant Pathol* 153:645-650. <https://doi.org/10.1007/s10658-018-1554-0>
22. Zhao Y, Zhang Y, Wan X, She Y, Li M, Xi H, Xie J and Wen C (2020) A Novel Ourmia-Like Mycovirus Confers Hypovirulence-Associated Traits on *Fusarium oxysporum*. *Front Microbiol.* . <https://doi.org/10.3389/fmicb.2020.569869>
23. Torres-Trenas A, Cañizares MC, García-Pedrajas MD, Pérez-Artés E (2020) Molecular and biological characterization of the first hypovirus identified in *Fusarium oxysporum*. *Front Microbiol* 10:3131-3143. <https://doi.org/10.3389/fmicb.2019.03131>
24. Torres-Trenas A, Pérez-Artés E (2020) Characterization and Incidence of the First Member of the Genus Mitovirus Identified in the Phytopathogenic Species *Fusarium oxysporum*. *Viruses* 12:279-293. <https://doi.org/10.3390/v12030279>

25. Sato Y, Shamsi W, Jamal A, Bhatti MF, Suzuki N (2020) Hadaka virus 1: a capsidless eleven-segmented positive-sense single-stranded rna virus from a phytopathogenic fungus, *Fusarium oxysporum*. mBio 11:20-38. <https://doi.org/10.1128/mBio.00450-20>
26. Mishra PK, Fox RT, Culham A (2003) Development of a PCR-based assay for rapid and reliable identification of pathogenic Fusaria. FEMS Microbiol Lett 218:329-332. [https://doi.org/10.1016/S0378-1097\(02\)01174-6](https://doi.org/10.1016/S0378-1097(02)01174-6)
27. Wang J, Ni Y, Liu X, Zhao H, Xiao Y, Xiao X, Li S, Liu H (2020) Divergent RNA Viruses In *Macrophomina phaseolina* exhibit potential as virocontrol agents. Virus Evolution. <https://doi.org/10.1093/ve/veaa095>
28. Yao Z, Zou C, Peng N, Zhu Y, Bao Y, Zhou Q, Wu Q, Chen B and Zhang M (2020) Virome Identification and Characterization of *Fusarium sacchari* and *andiyazi*: Causative Agents of Pokkah Boeng Disease in Sugarcane. Front Microbiol 11:240. <https://doi.org/10.3389/fmicb.2020.00240>
29. Paquin B, Laforest MJ, Forget L, Roewer I, Wang Z, Longcore J, Lang BF (1997) The fungal mitochondrial genome project: Evolution of fungal mitochondrial genomes and their gene expression. Curr Genet 31:380-395. <https://doi.org/10.1248/yakushi.126.1179>
30. Nibert ML (2017) Mitovirus UGA (Trp) codon usage parallels that of host mitochondria. Virology 507:96-100. <https://doi.org/10.1016/j.virol.2017.04.010>
31. Hong Y, Dover SL, Cole TE, Brasier CM, Buck KW (1999) Multiple mitochondrial viruses in an isolate of the Dutch elm disease fungus *Ophiostoma novo-ulmi*. Virology 258:118-127. <https://doi.org/10.1006/viro.1999.9691>
32. Hillman BI, Cai G (2013) The family Narnaviridae: simplest of RNA viruses. Adv Virus Res 86:149-176. <https://doi.org/10.1016/B978-0-12-394315-6.00006-4>
33. Wu M, Zhang L, Li G, Jiang D, Ghabrial SA (2010) Genome characterization of a debilitation-associated mitovirus infecting the phytopathogenic fungus *botrytis cinerea*. Virology 406(1):117-126. <https://doi.org/10.1016/j.virol.2010.07.010>
34. Huang S, Ghabrial SA (1996) Organization and expression of the double-stranded RNA genome of Helminthosporium victoriae 190S virus, a totivirus infecting a plant pathogenic filamentous fungus. Proc Natl Acad Sci U S A 93:12541-12546. <https://doi.org/10.2307/40654>
35. Wang J, Xiao Y, Liu X, Ni Y, Zhao H, Zhao X, Liu H, Xiao X (2020) Complete genome sequence of a novel victorivirus isolated from the sesame charcoal rot fungus *Macrophomina phaseolina*. Arch Virol 165(2): 509-514. <https://doi.org/10.1007/s00705-019-04497-2>