

Pulmonary Embolism Risk Factors in the General Trauma Population: A Nested Case-Control Study Using Nationwide Trauma Registry Data in Japan

Hiroki Iriyama

Juntendo University

Akira Komori

Juntendo University

Takako Kainoh

Juntendo University

Yutaka Kondo

Juntendo University Urayasu Hospital

Toshio Naito

Juntendo University

Toshikazu Abe (✉ abetoshi111@gmail.com)

University of Tsukuba

Research Article

Keywords: pulmonary embolism, venous thromboembolism, acute respiratory failure, trauma, surgery, coagulopathy.

Posted Date: April 29th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-418103/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Data assessing specific risk factors for post-traumatic pulmonary embolism (PE) are scarce.

Methods: This was a nested case-control study using the Japan Trauma Data Bank between 2004 and 2017. We enrolled patients aged ≥ 16 years, Injury Severity Score ≥ 9 , and length of hospital stay ≥ 2 days, with PE and without PE, using propensity score matching. We conducted logistic regression analyses to examine risk factors for PE.

Results: We included 719 patients with PE and 3,595 patients without PE. Of these patients, 1,864 [43.2%] were male, and their median Interquartile Range (IQR) age was 73 [55–84] years. The major mechanism of injury was blunt (4,282 [99.3%]). Median [IQR] Injury Severity Score (ISS) was 10 [9–18]. In the multivariate analysis, the variables spinal injury [odds ratio (OR), 1.40 (1.03–1.89)]; long bone open fracture in upper extremity and lower extremity [OR, 1.51 (1.06–2.15) and OR, 3.69 (2.89–4.71), respectively]; central vein catheter [OR, 2.17 (1.44–3.27)]; and any surgery [OR, 4.48 (3.46–5.81)] were independently associated with PE.

Conclusions: Spinal injury, long bone open fracture in extremities, central vein catheter placement, and any surgery were risk factors for post-traumatic PE.

Background

Pulmonary embolism (PE) poses a great risk of mortality and morbidity for patients after trauma^{1,2}, among whom prophylaxis for PE is an important consideration. However, patients with lower extremity trauma were unable to receive mechanical prophylaxis. Neither could pharmacologic prophylaxis be applied to patients with high risk of bleeding—especially soon after trauma or surgery. It is thus important to carefully assess risk of PE for each individual patient.

Previous studies evaluated the risk of post-traumatic PE. Aging^{3–10} and trauma severity^{5–9,11–14} are established risk factors. Other, more controversial risk factors of post-traumatic PE are injury sites, blood transfusion, and timing of definitive surgery. Previous studies were not designed to assess specific risk factors for post-traumatic PE, rather investigating the overall outcome of venous thromboembolism (VTE), including deep vein thrombosis (DVT). Few risk assessment studies have focused only on post-traumatic PE. Patients with post-traumatic PE, known as de novo PE, do not often have DVT. The risk of PE and DVT may be different among patients with trauma^{15–17}.

We therefore aimed to investigate specific unknown risk factors for acute PE such as injury sites and timing of definitive surgery in trauma patients using a nationwide trauma registry in Japan.

Methods

Design and setting

We conducted a nested case–control study between January 2004 and December 2017 utilizing a nationwide trauma registry, the Japan Trauma Data Bank (JTDB) database, established in 2003. In 2017, a total of 264 hospitals, including 95% of all tertiary emergency medical centers in Japan, participated in the database¹⁸, which includes patient demographics, Abbreviated Injury Scale (AIS) scores, Injury Severity Score (ISS), emergency procedures, in-hospital complications, and clinical outcomes. Data collection was performed as a part of routine clinical patient management.

Participants

We included patients who had blunt or penetrative trauma, were aged ≥ 16 years old, had ISS ≥ 9 , and were admitted to the intensive care unit (ICU) or general ward. In addition, we included only patients who survived for more than 2 days after hospital admission in order to exclude the impact of early trauma deaths^{14,19}. We excluded patients who were pregnant, underwent chronic dialysis²⁰, and received anticoagulation drugs (Fig. 1).

Data definitions

The primary outcome was occurrence of PE during hospital stay. We categorized AIS codes representing fractures into open, closed, or unclassifiable, using AIS 90 Update 98 and AIS 2005 Update 2008. We categorized AIS codes such as 752604.3 (humerus fracture), 752804.3 (radius fracture), 753204.3 (ulna fracture), 853422.3 (tibia fracture), and 852604.3 (pelvic fracture) in AIS 90 Update 98 as open fractures in this study because almost all were open, although these codes include several other kinds of fractures. We defined blood transfusion as any blood product within the first 24 hours—the same method as that used in previous studies^{20–22}. The database recorded central vein catheter placements only at the emergency department.

Statistical analyses

We included patients from the database without PE as a control, using propensity score (PS) matching to ensure that the PE group and the control group were equally balanced on baseline characteristics and severity of trauma. We used the variables age, gender, vital signs at emergency department (Glasgow Coma Scale, systolic blood pressure, heart rate, and respiratory rate), mechanism of injury, transport type, ISS, and admission ward, to calculate PS. We carefully selected these variables based on clinical relevance and previous research^{18,23}. We employed nearest neighbor matching without replacement in a 1:5 manner. We used a caliper of 0.01 standard deviation of the logit of the PS. We evaluated standardized mean difference of the variables over 0.1 as a meaningful imbalance after PS matching. We calculated descriptive statistics comparing the PE group and the control group, using the Wilcoxon signed-rank test for continuous variables and the chi-square or Fisher exact test for categorical variables, having first evaluated the normality of all continuous variables using the Kolmogorov-Smirnov test. No continuous variables had normal distributions.

We used conditional logistic regression models to identify specific risk factors for post-traumatic PE. In the 1st model, the candidates of risk factors were body regions with AIS ≥ 3 . In the 2nd model we

included long bone and pelvic fracture instead of AIS ≥ 3 in the upper and lower extremities. In the 3rd model, we categorized long bone and pelvic fracture into open, closed, or unclassifiable fractures. In the final model, we added procedures such as blood transfusion, central vein catheter placement, and any surgery, to the 3rd model. We adjusted all the models for comorbidities (see Supplementary Table S1). We did not adjust all the models for in-hospital complications such as pneumonia and sepsis because the database had no data on whether PE or these in-hospital complications occurred antecedently. For all models we did not use covariates which had 10 or less patients in the PE group or control group. When we conducted conditional logistic regression models, we calculated the variance inflation factor of the covariates to evaluate multicollinearity, evaluating a variance inflation factor > 5 as meaningful multicollinearity. However, we did not confirm any meaningful multicollinearity. All p values were two-sided; we considered p values of < 0.05 statistically significant. We performed statistical analyses using R software (Version 3.6.2) ²⁴.

Subgroup analysis

We conducted a subgroup analysis to investigate the influence of time to bone fixation. We included only patients who had received primary bone fixation. We used a non-conditional logistic regression model with covariates time to bone fixation and the same covariates as those used in the final model. We defined time to bone fixation as the time between hospital arrival and surgery. We did not use covariates which had 10 or less patients in the PE group or control group for the model, in the same way as in the main analysis.

Sensitivity analysis

We conducted a sensitivity analysis to exclude the possibility that mechanical and pharmacological prophylaxis could be potential confounding elements. Because prophylaxis is well adhered to in ICU in Japan, we included only patients admitted to the ICU. We used a non-conditional logistic regression model and included the same covariates as those in the final model. Again, we did not use covariates which had 10 or less patients in the PE group or the control group for the model, using the same method as in the main analysis.

Imputation

Before PS matching, we replaced missing values concerning vital signs at the emergency department with vital signs at the injury site (we replaced no missing records for the Glasgow Coma Scale, 18 missing for systolic blood pressure, 82 missing for heart rate, and 318 missing for respiratory rate among all the 4,314 patients in this study). We defined missing values for blood transfusion as no blood transfusion (82 missing among the 4,314 patients in this study). We replaced missing values for time of hospital arrival with time when an emergency medical technician had contacted patients at the injury site (67 missing among 1,997 patients who received primary bone fixation).

Ethics approval

The Ethics Committee of the Juntendo University approved this study (IRB No. 19 – 010). We confirmed that all methods were performed in accordance with the relevant guidelines. We confirmed that the need for informed consent was waived by The Ethics Committee of the Juntendo University. The JTDB administrators also provided permission for use of the data from their database.

Results

A total of 155,525 patients among the patients who were registered in the JTDB database were eligible for this study after inclusion and exclusion criteria. A total of 772 (0.5%) patients developed post-traumatic PE. After PS matching, the PE group comprised 719 patients and the control group comprised 3,595 patients (Fig. 1). We detected no meaningful imbalance of the variables for the PS matching between the two groups (Table 1).

Table 1
Baseline characteristics of patients with and without pulmonary embolism

Characteristics		PE group (n = 719)	Control group (n = 3,595)	SMD	P value
Age, median (IQR) ^a		74 (55–84)	73 (56–84)	0.008	
Male, No (%) ^a		321 (44.6)	1543 (42.9)	0.04	
Mechanism of injury, No (%) ^a	Blunt / Penetrate	713/6 (99.2/0.8)	3569/26 (99.3/0.7)	0.01	
Transport type, No (%) ^a	Ambulance without physician	647 (90.0)	3205 (89.2)	0.04	
	Ambulance/Helicopter with physician	58 (8.1)	296 (8.2)		
	Other	14 (1.9)	94 (2.6)		
Vital signs at ED, median (IQR) ^a	GCS	15 (14–15)	15 (14–15)	0.005	
	SBP	140 (119–159)	140 (120–161)	0.04	
	HR	82 (71–95)	82 (72–95)	0.02	
	RR	20 (16–24)	20 (18–24)	0.009	
Admission ward, No (%) ^a	General ward / ICU	393/326 (54.7/45.3)	2011/1584 (55.9/44.1)	0.03	
ISS, median (IQR) ^a		10 (9–20)	10 (9–18)	0.02	
AIS ≥ 3, No (%)	Head	119 (16.6)	1050 (29.2)		< 0.001
	Thorax	137 (19.1)	768 (21.4)		0.18
	Abdomen	46 (6.4)	194 (5.4)		0.33
	Spine	75 (10.4)	387 (10.8)		0.84
	Upper extremity	59 (8.2)	186 (5.2)		0.002
	Lower extremity (including pelvis)	479 (66.6)	1690 (47.0)		< 0.001
Upper extremity (detail), No (%)	Long bone fracture	60 (8.3)	187 (5.2)		0.001

Characteristics		PE group (n = 719)	Control group (n = 3,595)	SMD	P value
	Open	57 (7.9)	165 (4.6)		< 0.001
	Closed	3 (0.4)	23 (0.6)		0.61
Lower extremity & pelvis (detail), No (%)	Long bone & pelvic fracture	490 (68.2)	1670 (46.5)		< 0.001
	Open	199 (27.7)	276 (7.7)		< 0.001
	Closed	56 (7.8)	239 (6.6)		0.31
	Unclassifiable ^b	283 (39.4)	1306 (36.3)		0.14
Blood transfusion, No (%)		132 (18.4)	511 (14.2)		0.006
Central vein catheter, No (%)		50 (7.0)	105 (2.9)		< 0.001
Surgery, No (%)	Any	623 (86.6)	2,051 (57.1)		< 0.001
	Orthopedic	550 (76.5)	1,650 (45.9)		< 0.001
	Bone fixation	521 (72.5)	1,476 (41.1)		< 0.001
	Abdominal	48 (6.7)	181 (5.0)		0.09
	Head	32 (4.5)	184 (5.1)		0.51
	Chest	11 (1.5)	34 (0.9)		0.16
	Others ^c	23 (3.2)	103 (2.9)		0.72

^aVariables included in the propensity score matching

^bAIS90 Update 98 included lower extremity and pelvic fracture codes which represent both open and closed fracture.

^cOther surgeries include facial, neck, and skin surgeries.

Definition of abbreviations: PE = pulmonary embolism; SMD = standardized mean difference; IQR = interquartile range; ED = emergency department; GCS = Glasgow Coma Scale; SBP = systolic blood pressure; HR = heart rate; RR = respiratory rate; ICU = intensive care unit; ISS = Injury Severity Score; AIS = abbreviated injury scale score

Missing data: None

With regard to baseline characteristics, the PE group had head injury with AIS ≥ 3 less frequently and lower extremities and pelvic injury with AIS ≥ 3 more frequently than those of the control group (16.6% vs. 29.2%, 66.6% vs. 47.0%, respectively) (Table 1). Specifically, the PE group had long bone and open fractures in the upper and lower extremities more frequently than the control group. We found no consistent pattern of comorbidities between the two groups (e-Table 1). However, we noted several obvious concomitant complications that occurred more frequently in the PE group than in the control group (see Supplementary Table S2). Such complications included pneumonia (47.4% vs. 2.9%), osteomyelitis (44.6% vs. 0.1%), sepsis or multiple organ failure (30.7% vs. 0.6%), and wound infection (51.5% vs. 1.1%).

Regarding treatments, the PE group frequently received blood transfusion (18.4% vs. 14.2%, $p = 0.006$), central vein catheter placement (7.0% vs. 2.9%, $p < 0.001$), and surgery (86.6% vs. 57.1%, $p < 0.001$). In both groups, bone fixation was the most frequent surgery (72.5% and 41.4%). Among patients who received bone fixation, a lower proportion of the PE group than that of the control group received bone fixation within 24 hours (22.4% vs. 27.5%) (see Supplementary Table S3).

The overall in-hospital mortality was higher in the PE group than in the control group (5.8% vs. 3.4%, $P = 0.003$) (Table 2). In terms of survivor dispositions, the PE group were more likely to be transferred to other facilities than the control group (68.9% vs. 58.4%, $P < 0.001$). The PE group had longer hospital stays than the control group (32 [interquartile range (IQR): 19–60] days vs. 23 [IQR: 13–36] days, $P < 0.001$).

Table 2
Outcomes of patients with and without pulmonary embolism

Outcomes	PE group (n = 719)	Control group (n = 3,595)	P value
In-hospital mortality, No (%)	42 (5.8)	122 (3.4)	0.003
Place after discharge, No (%)	Home	1365 (39.7)	< 0.001
	Other facilities	2006 (58.4)	
	Other	64 (1.9)	
Length of hospital stay, median (IQR)	32 (19–60)	22 (13–36)	< 0.001

Definition of abbreviations: PE = pulmonary embolism; IQR = interquartile range.

Missing data: In-hospital mortality = 32; Place after discharge = 203

Table 3 shows conditional logistic regression models to identify risk factors for PE. Head injury with AIS ≥ 3 was associated with a lower occurrence of PE in the 1st, 2nd, and 3rd model. However, in the final model, head injury was not associated with PE (odds ratio (OR) [95% confidence interval (CI)]: 0.78 [0.61–1.01]). Orthopedic injury such as spine injury with AIS ≥ 3 (OR [95% CI]: 1.40 [1.03–1.89]), open fracture in the upper extremities (OR [95% CI]: 1.51 [1.06–2.15]), and open fracture in the lower extremities (OR [95% CI]: 3.69 [2.89–4.71]) were independently associated with PE in the final model. Regarding treatments, central vein catheter placement (OR [95% CI]: 2.17 [1.44–3.27]) and any surgery (OR [95% CI]: 4.48 [3.46–5.81]) were associated with PE in the final model.

Table 3
Association between pulmonary embolism and various characteristics from the models under consideration

Variable	OR (95% CI)			
	1st model (n = 4,314)	2nd model (n = 4,314)	3rd model (n = 4,314)	Final model (n = 4,314)
Head AIS \geq 3	0.77 (0.60–0.98)	0.76 (0.60–0.97)	0.67 (0.53–0.86)	0.78 (0.61–1.01)
Thorax AIS \geq 3	1.20 (0.95–1.52)	1.13 (0.90–1.42)	1.09 (0.87–1.38)	1.22 (0.95–1.56)
Abdomen AIS \geq 3	1.48 (1.03–2.11)	1.47 (1.03–2.11)	1.42 (0.99–2.04)	1.00 (0.68–1.48)
Spine AIS \geq 3	1.50 (1.12–2.02)	1.51 (1.13–2.02)	1.26 (0.94–1.69)	1.40 (1.03–1.89)
Upper extremity AIS \geq 3	2.26 (1.62–3.16)			
Long bone fracture		2.27 (1.63–3.15)		
Open			2.24 (1.59–3.15)	1.51 (1.06–2.15)
Closed			a	a
Lower extremity AIS \geq 3	2.61 (2.08–3.26)			
Long bone & pelvic fracture		2.79 (2.25–3.46)		
Open			4.82 (3.82–6.10)	3.69 (2.89–4.71)
Closed			1.18 (0.85–1.63)	1.03 (0.73–1.44)
Blood transfusion				0.84 (0.64–1.09)
Central vein catheter				2.17 (1.44–3.27)
Any Surgery				4.48 (3.46–5.81)

All the models were also controlled for comorbidities.

The 3rd model and the final model were adjusted for unclassifiable fracture in lower extremity and pelvis.

^aCovariate which was not used for the model because of less than 10 patients in PE group

Definition of abbreviations: OR = odds ratio; CI = confidence interval; AIS = Abbreviated Injury Scale score

In the subgroup analysis, we included 1,997 patients treated with primary bone fixation. Our non-conditional logistic regression model showed that time to primary bone fixation of 24–120 h—compared to that within 24 h—was associated with PE (Fig. 2).

Sensitivity analysis showed that the following covariates were independently associated with PE, which is consistent with the final model: spine AIS ≥ 3 (OR [95% CI]: 1.62 [1.15–2.27]); open fracture in lower extremity (OR [95% CI]: 2.11 [1.50–2.96]); central vein catheter placement (OR [95% CI]: 1.91 [1.27–2.86]); and any surgery (OR [95% CI]: 3.53 [2.57–4.85]) (see Supplementary Table S4). Only Open fracture in the upper extremities was not associated with PE (OR [95% CI]: 1.30 [0.80–2.10]), a finding which differed from that of the final model.

Discussion

This nested case–control study of general trauma population revealed that long bone open fracture in extremities, spinal injury, central vein catheter placement, and any surgery were risk factors for PE. Bone fixation was the most frequent surgery, and delayed bone fixation also could be a risk factor for PE.

Long bone open fractures in upper and lower extremities emerged as high risk trauma for PE. Lower extremity fracture is an established risk factor for PE; however, in previous studies, upper extremity fracture has not been risk factor for PE^{3,8,10,21}. Whether open or closed fracture was a risk factor for PE was also unknown. In this study, long bone open fracture in the lower extremities was the most relevant trauma with PE, and long bone open fracture in the upper extremities was the second most relevant trauma. The possible underlying pathophysiology may be that open fracture has greater soft tissue damage involving the venous endothelium and has more bleeding from open wounds, resulting in more severe coagulopathy, than closed fracture^{25–27}. Venous stasis caused by immobility could also play a role in patients with lower extremity fracture²⁸. Open fracture in the extremities could be related to occurrence of PE.

We confirmed that spinal injury^{9,12,21,29–31}, central vein catheter placement^{6,32,33}, and surgery^{3,10,12,21} were risk factors for PE, which is consistent with previous studies. Spinal injury is related to venous stasis because of long-term immobility. Vascular-related procedures such as central vein catheter placement and surgery could be related to endothelial injury and hypercoagulability.

Head, thoracic, and abdominal trauma were not risk factors for PE in this study, although whether these injury sites are risk factors for PE has been controversial in previous studies^{3,4,7,9,11}. Compared with this study, previous studies were limited by small sample size and uncontrolled potential confounders such as comorbidities, severity, and surgery. Regarding head trauma, recent studies focusing on the timing of

post-traumatic PE showed that head trauma was associated with late onset of PE^{16,34–36}. Our cohort might have captured only early PE.

Early definitive surgery for bone fracture could reduce risk for PE. Previous studies showed that delayed definitive surgery for bone fracture (after 24 to 48 h) led to pulmonary complications such as pneumonia and acute respiratory distress syndrome because post-traumatic and post-surgical inflammation act as “two hit model”^{22,37–39}. However, few studies have showed an association between timing of definitive surgery and post-traumatic PE because it is a relatively infrequent complication⁴⁰. A subgroup analysis in this study showed that bone fixation during 24–120 h after injury was associated with a higher risk for PE. Our study suggests that early total care is better than damage control orthopedics regarding PE among patients with bone fracture.

In this study, patients with spinal injury, open fracture in the extremities, central vein catheter placement, and any surgery were at high risk of post-traumatic PE. Patients with these characteristics should be probably treated with pharmacological prophylaxis as soon as bleeding risk is adequately controlled.

The current study has several important limitations that warrant discussion. First, neither mechanical nor pharmacological prophylaxis was recorded in this database and could be potential confounding factors. Therefore, we performed a sensitivity analysis among ICU patients because prophylaxis is well adhered to in ICU in Japan (see Supplementary Table S4). The sensitivity analysis showed similar results, four of five risk factors for PE in the main analysis were independently associated with PE. Second, because they were risk factors for PE in previous studies, pneumonia and sepsis also could be potential confounding factors^{13,32,41}. However, we did not include these complications to logistic regression analyses because we did not have onset-time data of complications in our database. Third, we did not include patients who did not survive < 48 hours to exclude the impact of early trauma deaths. Some patients with PE might have been lost in our study. However, because VTE prophylaxis such as anticoagulants could not have been used in early phase of trauma care, these patients were presumably small number and their deaths were unpreventable. Fourth, there might be an information bias. The diagnosis of PE was based on the reports of the physician in charge, and it might have been an underdiagnosis. However, the incidence of PE among patients with eligible criteria was 0.5% (772/155,525) in this study, which is consistent with previous studies (0.1–2.6%)^{3,5–7,9–11,40}. Diagnosis was likely to have been correct. Moreover, physicians in Japan, compared with those in other countries, have relatively easier access to the use of computed tomography, because there are many computed tomography scanners in Japan⁴².

Conclusions

Among general trauma population, long bone open fracture in extremities, spinal injury, central vein catheter placement, and any surgery could be risk factors for post-traumatic PE. Among patients with bone fracture, delayed bone fixation also could be a risk factor for PE.

Declarations

Data availability

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Acknowledgments

The authors would like to thank Enago (www.enago.jp) for the English language review.

Author contributions

TA is the guarantor of the paper. HI, AK, TK, YK, and TA conceived the study. HI and TA performed analyses for the study. HI and TA wrote the first draft of the manuscript. All authors provided critical input into manuscript drafting and revisions.

Competing interests

The authors declare no competing interests.

Ethics approval and consent to participate

The Ethics Committee of the Juntendo University, which did not mandate obtaining consent from patients in observational studies using anonymous data, approved this study (IRB No. 19-010).

References

1. Knudson, M. M., Collins, J. A., Goodman, S. B. & McCrory, D. W. Thromboembolism following multiple trauma. *J. Trauma*. **32**, 2–11 (1992).
2. Gudipati, S. *et al.* A cohort study on the incidence and outcome of pulmonary embolism in trauma and orthopedic patients. *BMC Med*. **12**, 39 (2014).
3. Knudson, M. M., Ikossi, D. G., Khaw, L., Morabito, D. & Speetzen, L. S. Thromboembolism after trauma: an analysis of 1602 episodes from the American College of Surgeons National Trauma Data Bank. *Ann. Surg.* **240**, 490–498 (2004).
4. Schultz, D. J. *et al.* Incidence of asymptomatic pulmonary embolism in moderately to severely injured trauma patients. *J. Trauma*. **56**, 727–733 (2004).
5. Tuttle-Newhall, J. E., Rutledge, R., Hultman, C. S., Fakhry, S. M. & Statewide population-based, time-series analysis of the frequency and outcome of pulmonary embolus in 318,554 trauma patients. *J. Trauma*. **42**, 90–99 (1997).
6. O'Malley, K. F. & Ross, S. E. Pulmonary embolism in major trauma patients. *J. Trauma*. **30**, 748–750 (1990).
7. Spencer Netto, F. *et al.* Pulmonary emboli after blunt trauma: timing, clinical characteristics and natural history. *Injury*. **43**, 1502–1506 (2012).

8. Rogers, F. B. *et al.* Determining venous thromboembolic risk assessment for patients with trauma: The Trauma Embolic Scoring System. *J. Trauma. Acute Care Surg.* **73**, 511–515 (2012).
9. Jeremitsky, E., St Germain, N., Kao, A. H., Ong, A. W. & Smith, R. S. Risk of pulmonary embolism in trauma patients: Not all created equal. *Surgery.* **154**, 810–815 (2013).
10. Meizoso, J. P. *et al.* A simplified stratification system for venous thromboembolism risk in severely injured trauma patients. *J. Surg. Res.* **207**, 138–144 (2017).
11. Hoyt, D. B. *et al.* A risk analysis of pulmonary complications following major trauma. *J Trauma.* **35**, 524–531 (1993).
12. Azu, M. C., McCormack, J. E., Huang, E. C., Lee, T. K. & Shapiro, M. J. Venous thromboembolic events in hospitalized trauma patients. *Am. Surg.* **73**, 1228–1231 (2007).
13. Paffrath, T. *et al.* Venous thromboembolism after severe trauma: incidence, risk factors and outcome. *Injury.* **41**, 97–101 (2010).
14. Lichte, P. *et al.* Post-traumatic thrombo-embolic complications in polytrauma patients. *Int. Orthop.* **39**, 947–954 (2015).
15. Brakenridge, S. C. *et al.* Comparing clinical predictors of deep venous thrombosis versus pulmonary embolus after severe injury: a new paradigm for posttraumatic venous thromboembolism? *J. Trauma. Acute Care Surg.* **74**, 1231–1238 (2013).
16. Bennis, M., Reilly, P. & Kim, P. Early pulmonary embolism after injury: a different clinical entity? *Injury.* **45**, 241–244 (2014).
17. Kim, K. S. & Brophy, G. M. Symptomatic venous thromboembolism: incidence and risk factors in patients with spontaneous or traumatic intracranial hemorrhage. *Neurocrit. Care.* **11**, 28–33 (2009).
18. Abe, T. *et al.* Trauma complications and in-hospital mortality: failure-to-rescue. *Crit. Care.* **24**, 223 (2020).
19. Knudson, M. M., Gomez, D., Haas, B., Cohen, M. J. & Nathens, A. B. Three thousand seven hundred thirty-eight posttraumatic pulmonary emboli: a new look at an old disease. *Ann. Surg.* **254**, 625–632 (2011).
20. Selby, R. *et al.* Hypercoagulability after trauma: hemostatic changes and relationship to venous thromboembolism. *Thromb. Res.* **124**, 281–287 (2009).
21. Geerts, W. H., Code, K. I., Jay, R. M., Chen, E. & Szalai, J. P. A prospective study of venous thromboembolism after major trauma. *N. Engl. J. Med.* **331**, 1601–1606 (1994).
22. Vallier, H. A., Cureton, B. A., Ekstein, C., Oldenburg, F. P. & Wilber, J. H. Early definitive stabilization of unstable pelvis and acetabulum fractures reduces morbidity. *J. Trauma.* **69**, 677–684 (2010).
23. Haas, B., Gomez, D., Hemmila, M. R. & Nathens, A. B. Prevention of complications and successful rescue of patients with serious complications: characteristics of high-performing trauma centers. *J. Trauma.* **70**, 575–582 (2011).
24. Ihaka, R. & Gentleman, R. R. A Language for Data Analysis and Graphics. *J. Comput. Graph. Stat.* **5**, 299–314 (1996).

25. Blann, A. D. How a damaged blood vessel wall contributes to thrombosis and hypertension. *Pathophysiol. Haemost. Thromb.* **33**, 445–448 (2003).
26. Johansson, P. I., Stensballe, J., Rasmussen, L. S. & Ostrowski, S. R. A high admission syndecan-1 level, a marker of endothelial glycocalyx degradation, is associated with inflammation, protein C depletion, fibrinolysis, and increased mortality in trauma patients. *Ann. Surg.* **254**, 194–200 (2011).
27. Engelman, D. T., Gabram, S. G., Allen, L., Ens, G. E. & Jacobs, L. M. Hypercoagulability following multiple trauma. *World J. Surg.* **20**, 5–10 (1996).
28. Lowe, G. D. O. Virchow's triad revisited: abnormal flow. *Pathophysiol. Haemost. Thromb.* **33**, 455–457 (2003).
29. Fujii, Y. *et al.* Thrombosis in spinal cord injury. *Thromb. Res.* **68**, 357–368 (1992).
30. Aito, S., Pieri, A., D'Andrea, M., Marcelli, F. & Cominelli, E. Primary prevention of deep venous thrombosis and pulmonary embolism in acute spinal cord injured patients. *Spinal cord.* **40**, 300–303 (2002).
31. Merli, G. J., Crabbe, S., Paluzzi, R. G. & Fritz, D. Etiology, incidence, and prevention of deep vein thrombosis in acute spinal cord injury. *Arch. Phys. Med. Rehabil.* **74**, 1199–1205 (1993).
32. Bahl, V. *et al.* A validation study of a retrospective venous thromboembolism risk scoring method. *Ann. Surg.* **251**, 344–350 (2010).
33. Levy, M. M., Albuquerque, F. & Pfeifer, J. D. Low incidence of pulmonary embolism associated with upper-extremity deep venous thrombosis. *Ann. Vasc. Surg.* **26**, 964–972 (2012).
34. Brakenridge, S. C., Toomay, S. M., Sheng, J. L., Gentilello, L. M. & Shafi, S. Predictors of early versus late timing of pulmonary embolus after traumatic injury. *Am. J. Surg.* **201**, 209–215 (2011).
35. Coleman, J. J. *et al.* Factors associated with pulmonary embolism within 72 hours of admission after trauma: a multicenter study. *J Am Coll Surg.* **220**, 731–736 (2015).
36. Gambhir, S. *et al.* Early versus late pulmonary embolism in trauma patients: not all pulmonary embolisms are created similarly. *J. Surg. Res.* **239**, 174–179 (2019).
37. Brundage, S. I., McGhan, R., Jurkovich, G. J., Mack, C. D. & Maier, R. V. Timing of femur fracture fixation: effect on outcome in patients with thoracic and head injuries. *J. Trauma.* **52**, 299–307 (2002).
38. Nahm, N. J., Como, J. J., Wilber, J. H. & Vallier, H. A. Early appropriate care: definitive stabilization of femoral fractures within 24 hours of injury is safe in most patients with multiple injuries. *J. Trauma.* **71**, 175–185 (2011).
39. Harvin, J. A. *et al.* Early femur fracture fixation is associated with a reduction in pulmonary complications and hospital charges: a decade of experience with 1,376 diaphyseal femur fractures. *J. Trauma Acute Care Surg.* **73**, 1442–1449 (2012).
40. Byrne, J. P., Nathens, A. B., Gomez, D., Pincus, D. & Jenkinson, R. J. Timing of femoral shaft fracture fixation following major trauma: A retrospective cohort study of United States trauma centers. *PLoS Med.* **14**, e1002336 (2017).

41. Gangireddy, C. *et al.* Risk factors and clinical impact of postoperative symptomatic venous thromboembolism. *J. Vasc. Surg.* **45**, 335–341 discussion 341 – 332 (2007).
42. Organisation for Economic Co-operation and Development. *Computed tomography (CT) scanners (indicator)*. <https://data.oecd.org/healthqct/computed-tomography-ct-scanners.htm#indicator-chart> (2021).

Figures




Figure 1

Patient selection Definition of abbreviations: PE = pulmonary embolism; ISS = Injury Severity Score; GCS = Glasgow Coma Scale; SBP = systolic blood pressure; HR = heart rate; RR = respiratory rate; ICU = intensive care unit.




Figure 2

Association between pulmonary embolism and time to primary bone fixation This figure is the result of subgroup analysis using non-conditional logistic regression model. This model was adjusted for the same covariates as in the final model. Definition of abbreviations: OR = odds ratio; CI = confidence interval.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [PEsuppv25scientificreport.docx](#)