

Clinical Variables Responsible for Early and Late Diagnosis of Foreign Body Aspiration in Pediatrics age Group

Samarth Goyal

Manipal University Kasturba Medical College Manipal

Shubhika Jain

Manipal University Kasturba Medical College Manipal

Guruprasad Rai (✉ guru.rai@manipal.edu)

Kasturba Hospital Manipal <https://orcid.org/0000-0002-8785-0803>

Rajkamal Vishnu

Manipal Academy of Higher Education

Ganesh Sevagur Kamath

Manipal Academy of Higher Education

Arvind Kumar Bishnoi

Manipal Academy of Higher Education

Yogesh Gaude

Manipal Academy of Higher Education

Vijaya Kumara

Manipal Academy of Higher Education

Harshil Joshi

Manipal Academy of Higher Education

Revanth Reddy

Manipal Academy of Higher Education

Research article

Keywords: Bronchus, Foreign Body, Foreign Body Aspiration (FBA), Rigid Bronchoscopy

Posted Date: July 22nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-41881/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on September 29th, 2020. See the published version at <https://doi.org/10.1186/s13019-020-01314-9>.

Abstract

Background

Incidence of foreign body aspiration (FBA) has been noticed maximally in age group ranging from 12 months-3 years. Foreign body in the trachea is a medical emergency as presentation is of respiratory distress. Obstruction of only one main or distal bronchus, leads to severe cough, choking sensation and breathlessness. Without early intervention it can lead to collapse, consolidation and pneumonia of the affected lung.

Methods

We retrospectively analysed 37 pediatric case records who presented from January 2014-December 2018 with FBA. Our primary aim was to assess the parameters responsible for early and late diagnosis of FBA. We concluded with a diagnostic algorithm for management of FBA on the basis of this outcomes.

Results

Around 32.5% came with a history of aspiration, 43% were referred from primary centers with a suspicion for the same and the rest came our tertiary care hospital directly. Those who presented within a week came with complaints of wet cough, wheeze and tachypnea where as those who came in after a week had dry cough and fever as their main complaint. Majority of ingested foreign body was vegetative type (80%) as compared to the non -vegetative material (20%).

Conclusion

Unlike adults, FBA in children is most commonly diagnosed on history, suspicion and clinical findings. Chest x ray has been the primary investigation of choice but in majority of the cases it was normal with subtle changes. Early diagnosis is the key to avoid complication.

Background:

Foreign body aspiration is a uncommon medical emergency ⁽¹⁾. Both adults and children have tendency to inhale foreign bodies. The object enters into the trachea and usually goes further down and lodges into one of the bronchus. Children have a tendency to swallow whatever comes into their hands which include wide variety of objects like coin, parts of toys, seeds, nuts etc and when they cry or laugh or jump with the mouth full, the Fb can enter the airway in deep inspiration. In adult's the main cause for aspiration is trying to swallow food when they are under intoxication and common Fb are unchewed food, fish bone, etc. FBA is said to be a primary cause of accidental death in children under 12 months to 3 years of age

(2). literatures suggests sex discrepancy of 60% and majority of the patients being males⁽³⁾. Foreign body (FB) in airway can lead to choking and ultimately death due to asphyxia if the diagnosis is delayed⁽⁴⁾. National Safety Council of America in 2016 proposed the rate of fatal choking in children < 5 years of age in American general population was 0.43 per 100,000.⁽⁵⁾

We aimed to analyze retrospectively and determine clinical predictors like history of presentation, physical and radiological findings which help in early diagnosis. We also intend to study the parameters which lead to delayed diagnosis and draw an algorithm for management on the basis of our experience.

Methods:

We conducted this retrospective study at Kasturba Medical College Hospital, Manipal between January 2014 and December 2018. Institutional ethical committee clearance was obtained before the initiation of the study. A retrospective analysis of the medical records of 37 consecutive patients of pediatric age group who were treated in our tertiary care hospital for FBA aspiration were included in the study.

We analyzed the following details from the medical records: demographic profile (age and gender), time of onset of symptoms and history of treatment at local hospital, time of referral to our hospital and the treatment given. The duration from the onset of symptoms and the referral. The clinical features, physical examination and radiological findings at presentation were noted. Bronchoscopy findings were also tabulated. Final result and complications were documented.

Data was entered and analyzed using Statistical Package for Social Sciences (SPSS) version 15. The results are summarized as percentages and proportions.

Results:

Total of 37 case files were analysed. The socio-demographic data of the patients is depicted in Table 1.

Table 1
Socio-Demographic details.

	n = 37	%
Age		
1–3 years	32	86.4
> 3 years	5	13.6
Gender		
Male	25	67.5
Female	12	32.5

Out of 37 cases studied, 12 (32.4%) came with history of foreign body aspiration. Sixteen cases (43.2%) were referred to the hospital from primary centers with suspicion of foreign body whereas remaining presented to the hospital directly.

Table 2
Time duration between onset of symptom and presentation to hospital.

	n = 37	%
Within 24 hrs.	9	24.32
1 day to 1 week	8	21.6%
1 week to 1 month	10	27.02
> 1 month	10	27.02

Delay in presentation was more in patient who were initially treated symptomatically in a primary health center. Of 20 patients who came to the hospital after 1 week, 13 (65%) were treated locally whereas who presented before 1 week of onset of symptoms only 3 (17.7%) out of 17 underwent previous treatment.

Figure 1: Relation of duration of stay in the hospital and local treatment.

Most common presenting features in cases that presented early (< 1 week) were Tachypnea (58.8%), Wheeze (47.05%) and wet cough (41.1%). Cases that presented late (> 1 week) came mainly with complain of fever (65%) and dry cough (45%). History of foreign body aspiration was present in 47.05% of early cases and 20% of late cases.

Eighty percent of foreign bodies were of vegetative type, peanut being most common (43.24%) and 18.9% included non-vegetative material like metal screw, pin, needle and crayon. Details on type of foreign body has been depicted in Fig. 3.

Table 3
Location of foreign body and related examination and radiological findings

Location on the basis of bronchoscopy	Respiratory examination		Radiological findings (x ray)	
Left bronchus			X ray done	100%
	Decreased left sided chest movements	26.3%	X ray showed left sided FB	42.1%
	Decreased air entry on left side	47.4%	Left lung collapse	21.1%
	Decreased air entry on right side	5.3%	Mediastinal shift to left	5.3%
	Rhonchi	36.9%	Mediastinal shift to right	5.3%
	Crepitus	31.6%		
Right bronchus			X ray done	100%
	Decreased right sided chest movements	5.9%	X ray showed right sided FB	41.4%
			Right lung collapse	17.7%
	Decreased air entry on right side	64.7%	Mediastinal shift to right	5.9%
	Decreased air entry on left side	5.9%	Mediastinal shift to left	11.8%
	Rhonchi	47.1%	Pericardial pneumonitis	11.8%
	Crepitus	17.7%		

Table 4
Bronchoscopy data of the cases

		n = 37	%
Route of bronchoscopy	Orotracheal	35	94.6%
	Nasotracheal	2	5.4%
Method of bronchoscopy	Rigid	36	97.3%
	Fiberoptic	1	2.7%
No. of episodes	One	33	89.2%
	Two	4	10.8%

All the cases underwent rigid bronchoscopy once. Flexible Bronchoscopy done in 4 cases was repeated in 4 cases due to; incomplete removal, slippage of some part into segmental bronchuse and impaction of FB to the mucosa.

Out of 37 cases, 35 cases (94.6%) have undergone rigid bronchoscopic removal. Adjuvant treatment that was given along with bronchoscopic removal was antibiotics (72.97%), bronchodilators (29.73%) and steroids (18.90%).

Discussion:

Foreign-body aspiration accounts for high morbidity if the diagnosis is delayed or missed. Mortality also is reported in children, especially between ages 12 months to 3 years.⁽²⁾ Most children under the age of 3 years tend to mull over most particles in their mouths, they also have flawed nibbling habits and premature swallowing coordination which makes them more prone for a FBA. Children are more susceptible for a FBA complications due to immature defense mechanisms.⁽⁶⁾ Often the presentation and initial radiological findings are vague which impedes the early diagnosis. An alleged episode of choking and severe cough is a critical evidence in the diagnosis of FB aspiration.

In our study, choking (5.40%) and hemoptysis (2.70%) were seen in limited cases. Most common symptoms in patients who presented early (< 1 week) were Tachypnea (58.8%), Wheeze (47.05%) and wet cough (41.1%). In cases of delayed presentation (> 1 week) majority complained of fever (65%) and dry cough (45%). It was found out that delay in presentation mostly due to unintentional aspiration by child unnoticed by parents, vague clinical history, lack of respiratory symptoms, unwillingness of the treating physician for getting an chest X ray and delayed referral. It was noted that the duration of treatment at the local hospital y was directly proportional to the duration of admission later for effective management.

Unlike adults, babies do not have clear recall of FBA, thus diagnosis is most often dependent on the mothers history, clinical findings such as onset and duration of symptoms and suspicion. Chest x -ray is the most common investigation done in these cases and majority of the time chest x rays are normal. The typical findings in chest radiograph which is diagnostic is hyperinflation, collapse, consolidation of one lung and mediastinal shift. Many authors in their study have reported rates of normal chest X -ray around 20–42%^(7, 8). In our study normal chest x-ray was found in 39% patients. Normal chest X ray are accepted in early phase as most aspirated foreign bodies are organic in nature and cannot be seen on chest x ray. Later when the organic substance swells with inflammation it occludes the bronchus completely and typical chest xray findings appear.

Recently CT has been introduced as a noninvasive procedure in diagnosis of FBA, but it is known to show false positive and has radiations hazards associated.^(9, 10, 11)

Various diagnostic modalities have been reported to be effective, rigid bronchoscopic inspection being the gold standard⁽¹²⁾. It is an invasive procedure done under general anaesthesia and has some

morbidity.⁽⁷⁾ Some authors suggested that flexible bronchoscopy is a useful procedure for foreign body retrieval from infants and children with 91.3% success rate⁽¹³⁾ .with availability of proper bronchoscopic accessories. Rigid or flexible bronchoscopy especially in pediatric patients remains a controversy and depends on the institute protocol.. In our study all patients underwent rigid bronchoscopy initially. Only in 4 cases were incomplete retrieval was done and flexible bronchoscopy was used for inspection followed by rigid bronchoscopy for retrieval.

In our cases nature of foreign body aspirated were mostly organic 81.1% and inorganic foreign body accounted for 18.9%. Routinely found FB was peanut being present in 43.24% patients.

Many authors have reported the gap between aspiration and hospital admission was more than 24 hours and even after that there was a delay in diagnosis⁽¹⁴⁾. Some of the larger reviews have documented delays ranging from 20–40% of the patients reviewed⁽¹⁵⁾. In this present study 54.04% of the patients reviewed had delayed presentation. Delay in presentation were due to unintentional aspiration by child unnoticed by parents ,vague clinical history, lack of respiratory symptoms, unwillingness of the treating physician for getting an chest X ray and late referral. The severity of symptoms depends on the degree of obstruction of the tracheobronchial tree so it can be complete or incomplete and site of obstruction. If there is complete obstruction these patients are referred early. Delay usually occurs in patients with incomplete obstruction with normal chest X ray. Lack of history of inhalation most commonly led to delayed presentation and resulted in various respiratory complications like Pneumonia ,Obstructive emphysema ,Atelectasis ,Pneumothorax and Pneumomediastinum^(16–118). Literature reveals there is a 7% incidence of pre hospital deaths in cases of foreign body aspiration⁽¹⁶⁾.

Another factor that causes preoperative complications associated with FBA is lack of standard treatment guidelines and management procedures.⁽⁶⁾

Based on our study we have developed an algorithm for management on the basis of outcomes.

Conclusion

Experience from our institute suggests that majority of complication occurs only when there is a major airway obstruction. Early diagnosis and referral reduces complication and mortality. Initial chest ray is a useful tool and in a crunch situation never be indecisive for doing a rigid bronchoscopy. Prophecy of negative bronchoscopy will probably counterbalance when it gets to saving the child.

List Of Abbreviations

Foreign body aspiration – FBA

Foreign body- FB

Declarations

Acknowledgement:

The Medical Records Department at Kasturba Hospital for providing access to case records.

Authors contribution :

Conception or design of the work: Samarth Goyal , Shubhika Jain

Data collection: Vijaya Kumara , Harshil Joshi , Revanth Reddy

Data analysis and interpretation: Guruprasad Rai

Drafting the article: Yogesh Gaude , Rajkamal Vishnu

Critical revision of the article: Ganesh Sevagur Kamath

Final approval of the version to be published: Arvind Kumar Bishnoi

Funding : None.

Availability of data and materials: The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request

Ethics approval and consent to participate : This study was approved by the Kasturba Medical College and Kasturba Hospital Institutional Ethics Committee (IEC :350/2019)

Consent for publication: All authors reviewed and approved the final version of the manuscript and consented for publication.

Competing interests : The authors declare that they have no Competing interests.

Author details:

Department of Cardiothoracic and vascular surgery, Kasturba Medical College, Manipal Academy of Higher Education, Manipal ,Karnataka, India– 576 104

References

1. Janahi IA, Khan S, Chandra P, Al-Marri N, Saadoon A, Al-Naimi L, Al-Thani M, Greer W. A new clinical algorithm scoring for management of suspected foreign body aspiration in children. BMC pulmonary medicine. 2017 Dec;17(1):61.
2. National Center for Injury Prevention and Control. WISQARS Leading Causes of Death Reports, 1999–2001.

3. Foltran F, Ballali S, Rodriguez H, van As AB, Passali D, Gulati A, Gregori D. Inhaled foreign bodies in children: a global perspective on their epidemiological, clinical, and preventive aspects. *Pediatric pulmonology*. 2013 Apr;48(4):344–51.
4. Lifschultz BD, Donoghue ER. Deaths due to foreign body aspiration in children: the continuing hazard of toy balloons. *Journal of Forensic Science*. 1996 Mar 1;41(2):247 – 51.
5. Hanba C, Cox S, Bobian M, Svider PF, Gonik NJ, Shkoukani MA, Sheyn A. Consumer product ingestion and aspiration in children: A 15-year review. *Laryngoscope*. 2017 May;127(5):1202–7.
6. Yang XJ, Zhang J, Chu P, Guo YL, Tai J, Zhang YM, Tang LX, Ni X. Pneumomediastinum secondary to foreign body aspiration: clinical features and treatment exploremment in 39 pediatric patients. *Chinese medical journal*. 2016 Nov 20;129(22):2691.
7. Zerella JT, Dimler M, McGill LC, Pippus KJ. Foreign body aspiration in children: value of radiography and complications of bronchoscopy. *Journal of pediatric surgery*. 1998 Nov 1;33(11):1651-4.
8. Metrangolo S, Monetti C, Meneghini L, Zadra N, Giusti F. Eight years' experience with foreign-body aspiration in children: what is really important for a timely diagnosis?. *Journal of pediatric surgery*. 1999 Aug 1;34(8):1229-31.
9. Haliloglu M, Ciftci AO, Oto A, Gumus B, Tanyel FC, Senocak ME, Buyukpamukcu N, Besim A. CT virtual bronchoscopy in the evaluation of children with suspected foreign body aspiration. *European journal of radiology*. 2003 Nov 1;48(2):188–92.
10. Hong SJ, Goo HW, Roh JL. Utility of spiral and cine CT scans in pediatric patients suspected of aspirating radiolucent foreign bodies. *Otolaryngology—Head and Neck Surgery*. 2008 May;138(5):576–80.
11. Kosucu P, Ahmetoglu A, Koramaz I, Orhan F, Özdemir O, Dinç H, Ökten A, Gümele HR. Low-dose MDCT and virtual bronchoscopy in pediatric patients with foreign body aspiration. *Am J Roentgenol*. 2004 Dec;183(6):1771–7.
12. Adaletli I, Kurugoglu S, Ulus S, Ozer H, Elicevik M, Kantarci F, Mihmanli I, Akman C. Utilization of low-dose multidetector CT and virtual bronchoscopy in children with suspected foreign body aspiration. *Pediatric radiology*. 2007 Jan 1;37(1):33–40.
13. Ramírez-Figueroa JL, Gochicoa-Rangel LG, Ramírez-San Juan DH, Vargas MH. Foreign body removal by flexible fiberoptic bronchoscopy in infants and children. *Pediatric pulmonology*. 2005 Nov;40(5):392–7.
14. Losek JD. Diagnostic difficulties of foreign body aspiration in children. *The American journal of emergency medicine*. 1990 Jul 1;8(4):348 – 50.
15. Kim IG, Brummitt WM, Humphry A, Siomra SW, Wallace WB. Foreign body in the airway: a review of 202 cases. *Laryngoscope*. 1973 Mar;83(3):347–54.
16. Hidaka H, Obara T, Kuriyama S, Kurosawa S, Katori Y, Kobayashi T. Logistic regression analysis of risk factors for prolonged pulmonary recovery in children from aspirated foreign body. *International journal of pediatric otorhinolaryngology*. 2013 Oct 1;77(10):1677-82.

17. Shafi M, Suhail Z, Ashrafi SK, Sajjad SM. Frequency of tracheobronchial foreign bodies and their management in urban population of Sindh. J Pak Med Assoc. 2012 Sep 1;62(9):896-9.
18. Ciftci AO, Bingöl-Koloğlu M, Şenocak ME, Tanyel FC, Büyükpamukçu N. Bronchoscopy for evaluation of foreign body aspiration in children. Journal of pediatric surgery. 2003 Aug 1;38(8):1170-6.

Figures




Figure 1

Relation of duration of stay in the hospital and local treatment.




Figure 2

Clinical manifestation at time of presentation




Figure 3

Types of foreign body.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Chart.jpg](#)