SARS-CoV-2 Infection Elicits a Greater Humoral Immune Response in the Second Wave Than the First Wave

Sayeh Ezzikouri (✉ sayeh.ezzikouri@pasteur.ma)

Institut Pasteur du Maroc <https://orcid.org/0000-0002-3982-6163>

Hind Majidi

Gouvernement du Royaume du Maroc Ministere de la Sante

Soad Redwane

Ministère de la Santé: Gouvernement du Royaume du Maroc Ministere de la Sante

Nadia Meziane

Centre Régional de transfusion sanguin, Casablanca

Abdellatif Zahir

Centre de Transfusion Sanguine, Eljadida

Islam Abbadi

Institut Pasteur du Maroc

Chaimaa Zerrad

Institut Pasteur du Maroc

Asmaa Haddaji

Institut Pasteur du Maroc

Karima Abounouh

Institut Pasteur du Maroc

Soufiane Hilmi

Institut Pasteur du Maroc

Safaa Aqillouch

Institut Pasteur du Maroc

Mohcine Elmessaoudi-Idrissi

Institut Pasteur du Maroc

Oumaima Laazaazia

Institut Pasteur du Maroc

Abdelhakim Ainahi

Institut Pasteur du Maroc

Nabila Rmili

Institut Pasteur du Maroc

Abderrahmane Maaroufi

Research Article

Keywords: COVID-19, SARS-CoV-2, antibody, B cells, vaccine

Posted Date: April 21st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-419367/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Abstract

Purpose

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is the causative agent of the ongoing pandemic of coronavirus disease 2019 (COVID-19). One year has passed since the first COVID-19 case, and concerns have emerged regarding new variants, meaning that characterizing antibody-mediated immune response dynamics is of paramount importance. Here, we evaluated the humoral immune response against SARS-CoV-2 in subjects during the first and second waves of the pandemic in Morocco.

Methods

We assessed humoral immune response in samples from 94 seropositive individuals in the first wave (February to August 2020) and 596 seropositive individuals in the second wave (December 2020 to January 2021). Plasma samples were collected from volunteer blood donors and their levels of serum IgG to SARS-CoV-2 nucleoprotein (NP) were determined using architect SARS-CoV-2 IgG chemiluminescence microparticle immunoassay.

Results

Our results revealed an increase of humoral immunity during the second wave than first wave (3.897 ± 0.079 vs. 2.842 ± 0.153 , respectively, $p < 0.0001$). Notably, we found an age-related gradient in antibody level, with higher antibody index in subjects at 45 years old and above ($p = 0.01451$). However, no significant difference was found according to gender ($p = 0.8629$).

Conclusions

Our data highlighted an important issue regarding antibody-mediated immune response against SARS-CoV-2 infection during the second wave and this issue might have arisen due to the dynamics of different strains circulating during the progress of the pandemic.

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) —the causative agent of the ongoing pandemic of coronavirus disease 2019 (COVID-19)—emerged one year ago [1], and is now posing further challenges in many countries worldwide as new variants are identified. These variants include those identified in the United Kingdom (UK) (variant 20I/501Y.V1, VOC 202012/01, or B.1.1.7), South Africa (variant 20H/501Y.V2 or B.1.351), and Brazil (variant P.1), and have raised many questions around severity of disease, increased risk of death, vaccine efficacy, and the speed at which COVID-19 may proliferate [2–8].

The North African kingdom of Morocco has adopted a range of measures to combat the COVID-19 pandemic, with surveillance for new variants including diagnostic and screening measures with nucleic

acid amplification tests, antigen-based tests, serologic assays and genome sequencing among the approaches that have been applied. Previously, Morocco appeared to achieve a degree of control against the disease since the country's Ministry of Health first reported a case, on March 2nd, 2020. The 5,443,945 tests carried out in Morocco as of March 28th, 2021 yielded a total of 494,659 positive results, with 8,798 deaths and 482,352 patients recorded as recovering from COVID-19. In January 28th, 2021, the kingdom launched the national vaccination campaign and as of March 28th, 2021, 4,302,183 persons received their first dose and 3,332,292 second dose of a COVID-19 vaccine. As with many other countries, Morocco appears to have experienced waves of infection, with the first wave regarded as commencing with the first reported case in March 2020, and the second wave around September 2020, just weeks after the traditional Eid al-Adha festivities, which are marked by social and religious gatherings. The two waves are separated by a trough in daily recorded cases (Fig. 1). However, on January 18th, 2021, Morocco's health ministry confirmed its first imported case of the UK variant, which was detected in the northern port of Tangier in a Moroccan national returning from Ireland via Marseille.

With recently described variants in SARS-CoV-2, and characterizing antibody (Ab)-mediated immune response dynamics of SARS-CoV-2 is fundamental to understand COVID-19 epidemiology, the path to protection against SARS-CoV-2 infection or re-infection, and for effective vaccine development [9]. Data of this nature are not yet available for SARS-CoV-2 infection. To investigate the humoral immune response against SARS-CoV-2 infection in the first and second waves in Morocco, we subjected blood samples from donors who recovered from COVID-19 during the first wave, or seroconverted during the second wave to a comparative evaluation.

Materials And Methods

Study participants

This is an observational study in healthy volunteers. We enrolled two groups of donors in two different periods, between 14th of February and 7th of August 2020 (for the first wave), or 23th of December 2020 through January 8, 2021 (for the second wave). Blood donors satisfying the donation eligibility criteria set by the Moroccan Ministry of Health and testing positive for SARS-CoV-2 could be included in the study. Individuals underwent a health screening procedure before blood donation. All study participants provided written informed consent after they had been briefed on the purposes of the study. After enrolment, participants were grouped into three age categories: 18–25 years, 26–45 years and more than 45 years. The samples were collected from Regional Blood Transfusion Center in Casablanca and Blood Transfusion Center in El Jadida. The study protocol was in accordance with the Helsinki declaration and received an approval from the ethics Committee of the Faculty of Medicine of Rabat (N°17/20).

Serological testing

Serologic tests to ascertain the presence of the antibody against SARS-CoV-2 (IgG) were conducted at the Viral Hepatitis Laboratory of the Pasteur Institute of Morocco. We used Architect™ SARS-CoV-2 IgG

chemiluminescence microparticle immunoassay (CMIA) (Abbott Park, Illinois, USA) to detect serum IgG Abs directed against the nucleocapsid protein of SARS-CoV-2. Interpretation of results was based on manufacturer's criteria: negative when the cut-off index (S/C) < 1.4 and positive when the cut-off index (S/C) $\geq$ 1.4.

Statistical Analysis

Data are presented as mean values (with corresponding standard errors of mean). Two-tailed Student's t-test or one-way ANOVA were used to analyze the differences between groups. Correlations between antibody index and age were assessed using Pearson's correlation coefficient. Statistical analyses were performed using GraphPad PRISM version 6.0e (GraphPad Software, San Diego, CA, USA).

Results

Among 14,000 blood donors screened in the first wave, 94 individuals who tested positive for SARS-CoV-2- IgG, and were included in the study population as the first-wave cohort. Among 1,986 blood donors in the second wave, 596 individuals were positive for SARS-CoV-2- IgG, and were included in the study population as the second-wave cohort.

The 94 COVID-19-recovered subjects in the first wave had a mean age of 38 ± 10 years and sex ratio (male/female) of 1. The 596 COVID-19- recovered subjects in the second wave had a mean age of 36 ± 10 years and sex ratio (male/female) of 2.16. No significant difference was observed regarding age between these cohorts ($p = 0.2325$). However, significant difference was observed regarding sex between these cohorts ($p = 0.0002$). Antibody index analyses revealed that the second-wave cohort had an elevated humoral immunity relative to the first-wave cohort (3.897 ± 0.079 vs. 2.842 ± 0.153 , $p < 0.0001$, respectively) (Fig. 2A).

We tested for associations of age and sex with antibody index among subjects in the second wave cohort. Notably, we found an age-related gradient in antibody levels ($p = 0.01451$) (Fig. 2B & Fig. 2C). Interestingly, SARS-CoV-2 antibody index was higher in individuals aged 45 years or above (Fig. 2B). However, no significant difference was observed between men and woman regarding antibody index ($p = 0.8629$) (Fig. 2D).

Discussion

Herein, we report a comparative evaluation of humoral immune response in SARS-CoV-2-IgG positive blood donors covering two waves of infection over a period of more than 11 months. We found that antibody response was greater in the second than the first wave. However, we are unable to assess the functionality of antibodies during the first and second waves due to lack of virus neutralization assay. However, the N protein is predicted to serve as a surrogate marker for the S protein-neutralizing antibody complex [10]. While all viral proteins are vulnerable to mutations that could impact diagnostic test performance, the S protein is under more selective pressure [2].

This atypical humoral immunity is particularly relevant in Morocco, where the second wave commenced around September 2020 and was probably linked to multiple distinct variants having accumulated mutations that differed from those in variants circulating during the first wave. Indeed, the SARS-CoV-2 strains circulating during the early epidemic phase in Morocco are different to those circulating in the second wave. Analysis of Moroccan SARS-CoV-2 sequences from the first wave clustered SARS-CoV-2 strains in clades G, B11, and A2 with the presence of amino acid change at D614G, which resembles strains seen in Asia, Europe, North America, Australia, and Africa, this indicating multiple SARS-CoV-2 introductions into Morocco [11–13]. Similarities are also evident with European strains seen at the beginning of the pandemic, and SARS-CoV-2 strains isolated in America and Africa later in the pandemic [14]. In addition, in Morocco, SARS-CoV-2 strains identified during December 2020 revealed an increased number of mutations relative to those identified in the first wave [15, 16]. This pattern appears to be in line with previous reports showing that viral genomes from the first and second waves belong to different clades/lineages [17–19]. Moreover, several recent reports of reinfection, as indicated by viral sequence differences, clinical data and data on potential exposure, have raised critical questions about whether and how well a first infection protects against reinfection [9, 17–20]. In addition, a relationship between a humoral immune response to SARS-CoV-2 infection and protection against reinfection with the identical SARS-CoV-2 strain has been shown in rhesus macaques, but has yet to be established in humans [21]. However, with the emergence of new variants more studies are needed to determine whether the SARS-CoV-2 variants reflect antigenic differences between strains circulating in the first and second waves, or waning immunity against SARS-CoV-2 [17].

In this study, older age was associated with greater IgG responses to SARS-CoV-2 but gender did not affect this association. Previous studies showed that older age is associated with greater antibody production and that males consistently had greater antibodies levels [22, 23].

In conclusion, our data revealed an important issue regarding antibody-mediated immune response against SARS-CoV-2 infection during the second wave. This issue may have arisen due to the dynamics of differences between circulating strains as the pandemic progressed and/or between specific human genetic variants that contribute to inter-individual variability in serum immunoglobulin levels [24].

There are limitations associated with this study. The antibody response induced by COVID-19 tends to be linked to the severity of the disease. However, the disease severity could not be determined in the recovering subjects. The study was also limited by the lack of assessment of neutralizing antibody levels. Further investigations are warranted to determine whether common or divergent factors are driving this drift in humoral immunity during the COVID-19 pandemic.

Declarations

Declaration of Competing Interest

All authors declare that there are no conflicts of interests.

Funding

This study was supported by Ministry of Health, Morocco.

Conflicts of interest

None

Availability of data and material

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Code availability

Not applicable

Ethics approval and consent to participate

The study protocol was in accordance with the Helsinki declaration and received an approval from the ethics Committee of the Faculty of Medicine of Rabat (N°17/20). The need for informed consent was waived because this was an observational retrospective study, and all patient data were analyzed anonymously.

Author Contribution

Sayeh Ezzikouri, Soumaya Benjelloun, Abderrahmane Maaroufi, Nabila Rmili, Hind Majidi, and Soad Redwane participated in the conception and design of the study. Nadia Meziane and Abdellatif Zahir collected samples and data. Islam Abbadi, Chaimaa Zerrad, Asmaa Haddaji, Soufiane Hilmi, Karima Abounouh, Oumaima Laazaazia, and Abdelhakim Ainahi conducted the serological assay. Sayeh Ezzikouri, Soumaya Benjelloun, Abderrahmane Maaroufi, Hind Majidi, and Soad Redwane participated in acquisition and interpretation of the data. Sayeh Ezzikouri and Mohcine Elmessaoudi-Idrissi analyzed the data. Sayeh Ezzikouri wrote the manuscript with assistance from all coauthors and approved the version of paper.

Consent for publication

All authors approved the final version for publication.

Acknowledgments

We acknowledge the contributions of other clinical and technical staff of the Regional Blood Transfusion Centers in Casablanca and El Jadida and from Ministry of Health.

References

- [1] Wang C, Horby PW, Hayden FG, Gao GF (2020) A novel coronavirus outbreak of global health concern. *Lancet* 395 (10223):470-473
- [2] Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, Hengartner N, Giorgi EE, Bhattacharya T, Foley B, Hastie KM, Parker MD, Partridge DG, Evans CM, Freeman TM, de Silva TI, Sheffield COVID-19 Genomics Group, McDanal C, Perez LG, Tang H, Moon-Walker A, Whelan SP, LaBranche CC, Saphire EO, Montefiori DC (2020) Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. *Cell* 182 (4):812-827.e819
- [3] Rambaut A, Loman N, Pybus O, Barclay W, Barrett J, Carabelli A, Connor T, Peacock T, Robertson LD, Volz E, UK C-GC (2020) Preliminary genomic characterisation of an emergent SARS-CoV-2 lineage in the UK defined by a novel set of spike mutations. *Virological.org*. Posted December 16, 2020. Accessed January 19, 2021. <https://virological.org/t/preliminary-genomic-characterisation-of-an-emergent-sars-cov-2-lineage-in-the-uk-defined-by-a-novel-set-of-spike-mutations/563>.
- [4] Tegally H, Wilkinson E, Giovanetti M, Iranzadeh A, Fonseca V, Giandhari J, Doolabh D, Pillay S, San EJ, Msomi N, Mlisana K, Gottberg AV, Walaza S, Allam M, Ismail A, Mohale T, Glass AJ, Engelbrecht S, Zyl GV, Preiser W, Petruccione F, Sigal A, Hardie D, Marais G, Hsiao M, Korsman S, Davies MA, Tyers L, Mudau I, York D, Maslo C, Goedhals D, Abrahams S, Laguda-Akingba O, Alisoltani-Dehkordi A, Godzik A, Wibmer CK, Sewell BT, Lourenço J, Alcantara LCJ, Pond SLK, Weaver S, Martin D, Lessells RJ, Bhiman JN, Williamson C, de Oliveira T (2020) Emergence and rapid spread of a new severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in South Africa. Preprint at medRxiv.
- [5] Kirby T (2021) New variant of SARS-CoV-2 in UK causes surge of COVID-19. *Lancet Respir Med* S2213-2600 (21):00005-00009
- [6] Callaway E (2021) Could new COVID variants undermine vaccines? Labs scramble to find out. *Nature* 589 (7841):177-178
- [7] Xie X, Zou J, Fontes-Garfias CR, Xia H, Swanson KA, Cutler M, Cooper D, Menachery VD, Weaver S, Dormitzer PR, Shi PY (2021) Neutralization of N501Y mutant SARS-CoV-2 by BNT162b2 vaccine-elicited sera. *bioRxiv*. .
- [8] Resende PC, Bezerra JF, de Vasconcelos RHT, Arantes I, Appolinario L, Mendonça AC, Paixao AC, Rodrigues ACD, Silva T, Rocha AS, Pauvolid-Corrêa A, Motta FC, Teixeira DLF, de Oliveira Carneiro TF, Neto FPF, Herbster ID, Leite AB, Riediger IN, Debur MDC, Naveca FG, Almeida W, Livorati M, Bello G, Siqueira MM (2021) Spike E484K mutation in the first SARS-CoV-2 reinfection case confirmed in Brazil, 2020. *Virological.org*. Posted January 10, 2021 Accessed January 20, 2021. <https://virological.org/t/spike-e484k-mutation-in-the-first-sars-cov-2-reinfection-case-confirmed-in-brazil-2020/584>.
- [9] Overbaugh J (2020) Understanding protection from SARS-CoV-2 by studying reinfection. *Nat Med* 26 (11):1680-1681

- [10] Batra R, Olivieri LG, Rubin D, Vallari A, Pearce S, Olivo A, Prostko J, Nebbia G, Douthwaite S, Rodgers M, Cloherty G (2020) A comparative evaluation between the Abbott Panbio™ COVID-19 IgG/IgM rapid test device and Abbott Architect™ SARS CoV-2 IgG assay. *J Clin Virol* 132:104645
- [11] Lemriss S, Souiri A, Amar N, Lemzaoui N, Mestoui O, Labioui M, Ouhaariba N, Jibjibe A, Yartaoui M, Chahmi M, El Rhouila M, Sellak S, Kandoussi N, El Kabbaj S (2020) Complete Genome Sequence of a 2019 Novel Coronavirus (SARS-CoV-2) Strain Causing a COVID-19 Case in Morocco. *Microbiol Resour Announc* 9 (27):e00633-00620
- [12] Jouali F, Marchoudi N, El Ansari FZ, Kasmi Y, Chenaoui M, El Aliani A, Azami N, Loukman S, Ennaji MM, Benhida R, Fekkak J (2020) SARS-CoV-2 Genome Sequence from Morocco, Obtained Using Ion AmpliSeq Technology. *Microbiol Resour Announc* 9 (31):e00690-00620
- [13] Laamarti M, Chemaoui-Elfihri MW, Kartti S, Laamarti R, Allam L, Ouadghiri M, Smyej I, Rahoui J, Benrahma H, Diawara I, Alouane T, Essabbar A, Siah S, Karra M, El Hafidi N, El Jaoudi R, Sbabou L, Nejari C, Amzazi S, Mentag R, Belyamani L, Ibrahim A (2020) Genome Sequences of Six SARS-CoV-2 Strains Isolated in Morocco, Obtained Using Oxford Nanopore MinION Technology. *Microbiol Resour Announc* 9 (32)::e00767-00720
- [14] Song S, Ma L, Zou D, Tian D, Li C, Zhu J, Chen M, Wang A, Ma Y, Li M, Teng X, Cui Y, Duan G, Zhang M, Jin T, Shi C, Du Z, Zhang Y, Liu C, Li R, Zeng J, Hao L, Jiang S, Chen H, Han D, Xiao J, Zhang Z, Zhao W, Xue Y, Bao Y (2020) The global landscape of SARS-CoV-2 genomes, variants, and haplotypes in 2019nCoV. *bioRxiv*:2020.2008.2030.273235
- [15] Fournier PE, Colson P, Levasseur A, Devaux C, Gautret P, Bedotto M, Delerce J, Brechard L, Pinault L, Lagier JC, Fenollar F, Raoult D (2021) Emergence and outcome of the SARS-CoV-2 “Marseille-4” variant. Preprint IHU doi: <https://doi.org/10.35081/xcrm-6t77>.
- [16] Colson P, Levasseur A, Gautret P, Fenollar F, Thuan Hoang V, Delerce J, Bitam I, Saile R, Maaloum M, Padane A, Bedotto M, Brechard L, Bossi V, Khedher MB, Chaudet H, Million M, Tissot-Dupont H, Lagier JC, Mboup S, Fournier PE, Raoult D (2021) Introduction into the Marseille geographical area of a mild SARS-CoV-2 variant originating from sub-Saharan Africa: an investigational study. *Travel Med Infect Dis* 101980
- [17] To KK, Hung IF, Chan KH, Yuan S, To WK, Tsang DN, Cheng VC, Chen Z, Kok KH, Yuen KY (2020) Serum antibody profile of a patient with COVID-19 reinfection. *Clin Infect Dis* ciaa1368
- [18] Colson P, Finaud M, Levy N, Lagier JC, Raoult D (2020) Evidence of SARS-CoV-2 re-infection with a different genotype. *J Infect* S0163-4453 (20):30706-30704
- [19] Colson P, Levasseur A, Delerce J, Chaudet H, Bossi V, Ben Khedher M, Fournier PE, Lagier JC, Raoult D (2020) Dramatic increase in the SARS-CoV-2 mutation rate and low mortality rate during the second epidemic in summer in Marseille. Preprint IHU 2020;doi:<https://doi.org/10.35088/68c3-ew82> .

- [20] Tillett RL, Sevinsky JR, Hartley PD, Kerwin H, Crawford N, Gorzalski A, Laverdure C, Verma SC, Rossetto CC, Jackson D, Farrell MJ, Van Hooser S, Pandori M (2021) Genomic evidence for reinfection with SARS-CoV-2: a case study. *Lancet Infect Dis* 21 (1):52-58
- [21] Deng W, Bao L, Liu J, Xiao C, Liu J, Xue J, Lv Q, Qi F, Gao H, Yu P, Xu Y, Qu Y, Li F, Xiang Z, Yu H, Gong S, Liu M, Wang G, Wang S, Song Z, Liu Y, Zhao W, Han Y, Zhao L, Liu X, Wei Q, Qin C (2020) Primary exposure to SARS-CoV-2 protects against reinfection in rhesus macaques. *Science* 369 (6505):818-823
- [22] Klein SL, Pekosz A, Park HS, Ursin RL, Shapiro JR, Benner SE, Littlefield K, Kumar S, Naik HM, Betenbaugh MJ, Shrestha R, Wu AA, Hughes RM, Burgess I, Caturegli P, Laeyendecker O, Quinn TC, Sullivan D, Shoham S, Redd AD, Bloch EM, Casadevall A, Tobian AA (2020) Sex, age, and hospitalization drive antibody responses in a COVID-19 convalescent plasma donor population. *J Clin Invest* 130 (11):6141-6150
- [23] Zhang B, Zhou X, Zhu C, Song Y, Feng F, Qiu Y, Feng J, Jia Q, Song Q, Zhu B, Wang J (2020) Immune Phenotyping Based on the Neutrophil-to-Lymphocyte Ratio and IgG Level Predicts Disease Severity and Outcome for Patients With COVID-19. *Front Mol Biosci* 7:157
- [24] Scepanovic P, Alanio C, Hammer C, Hodel F, Bergstedt J, Patin E, Thorball CW, Chaturvedi N, Charbit B, Abel L, Quintana-Murci L, Duffy D, Albert ML, Fellay J, Milieu Intérieur Consortium (2018) Human genetic variants and age are the strongest predictors of humoral immune responses to common pathogens and vaccines. *Genome Med* 10 (1):59

Figures

Montly COVID-19 Cases in Morocco

Montly COVID-19 Deaths in Morocco

Figure 1

SARS-CoV-2 epidemiological dynamics in Morocco. A). The epidemic curve of new confirmed COVID-19 cases. B) Daily confirmed COVID-19 deaths.

Figure 2

Antibody index of seropositive subjects. A) Comparison between first and second waves. Data are presented as mean and standard error of mean (SEM). B) Box and whiskers plot of antibody index with regard to age. The line crossing the box represents the mean. The whiskers extend to the minimal and maximal observed in each cohort. C) Age is associated with antibody responses to SARS-CoV-2. D) The graphic shows aligned dot plot of antibody index according to gender. Data are presented as mean $\pm$ SEM. Data were compared using Student's two-tailed t-test or one-way ANOVA.