

Statistical Analysis of Ionospheric Total Electron Content (TEC): Long-Term Estimation of Extreme TEC in Japan

Michi Nishioka (✉ nishioka@nict.go.jp)

National Institute of Information and Communications Technology <https://orcid.org/0000-0001-7147-5933>

Susumu Saito

Electronic Navigation Research Institute, Japan

Chihiro Tao

National Institute of Information and Communications Technology, Japan

Daikou Shiota

National Institute of Information and Communications Technology

Takuya Tsugawa

National Institute of Information and Communications Technology

Mamoru Ishii

National Institute of Information and Communications Technology

Full paper

Keywords: total electron content (TEC), extreme TEC, long-term ionosonde observation, manually scaled foF2, slab thickness

Posted Date: July 20th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-41992/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Version of Record: A version of this preprint was published on February 15th, 2021. See the published version at <https://doi.org/10.1186/s40623-021-01374-8>.

Abstract

Ionospheric total electron content (TEC) is one of the key parameters for users of radio-based systems, such as the Global Navigation Satellite System, high-frequency communication systems, and space-based remote sensing systems, since total ionospheric delay is proportional to TEC through the propagation path. It is important to know extreme TEC values in readiness for hazardous ionospheric conditions. The purpose of this study is to estimate extreme TEC values with occurrences of once per year, ten years, and hundred years in Japan. In order to estimate the extreme values of TEC, a cumulative distribution function of daily TEC is derived using 22 years of TEC data from 1997 to 2018. The extreme values corresponding to once per year and ten years are 90 and 130 TECU, respectively, in Tokyo, Japan. On the other hand, the 22-year data set is not sufficient to estimate the once-per-hundred-year value. Thus, we use the 62-year data set of manually scaled ionosonde data for the critical frequency of the F-layer ($f_{\text{OF}2}$) at Kokubunji in Tokyo. First, we study the relationship between TEC and $f_{\text{OF}2}$ for 22 years and investigate the slab thickness. Then the result is applied to the statistical distribution of $f_{\text{OF}2}$ data for 62 years. The result shows that the once-per-100-year TEC is about 190 TECU at Tokyo. The value is also estimated to be 230 TECU in Kagoshima and 150 TECU in Hokkaido, in the southern and northern parts of Japan, respectively.

1. Introduction

The ionospheric condition is one of the most important space weather features for users of radio-based systems, such as navigation systems based on the Global Navigation Satellite System (GNSS), high frequency (HF) communication systems, and space-based remote sensing systems. Radio waves propagating in the ionosphere experience a delay due to the electrons in the ionosphere. The ionospheric delay is proportional to the ionospheric total electron content (TEC) along the propagation path. The easiest way to correct the ionospheric delay is to utilize broadcast ionospheric delay models based on simple empirical TEC models such as the Klobuchar and NeQuick models. The TEC value is determined by many factors, such as solar activity, the season, local time, and geomagnetic activity. TEC variations caused by solar activity, the season, and local time may be estimated using these simple models but those caused by geomagnetic storms and other phenomena cannot be fully removed from these models. Therefore, users of radio-based systems may be affected by positive and/or negative ionospheric storms. During negative ionospheric storms, TEC is greater than or equal to 0 TECU even if the negative storm is extremely severe. On the other hand, extreme TEC values during positive storms are not unknown and should be studied.

For the design and operation of systems that may be impacted by space weather phenomena, it is important to know the possible extent of the impact and how often such events are likely to occur. Thus, it is important to study extreme values related to various space weather phenomena. For users of trans-ionosphere radio-based systems, the extreme TEC value is a key value.

Extreme values of some space weather parameters have been studied. For example, that of the Dst index was investigated using extreme value modeling (Tsubouchi and Omura, 2007). Those of the solar flare X-ray flux, speed of coronal mass ejection, Dst index, and proton energy in proton events were studied by Riley (2012) using complementary cumulative distribution functions. More recently, that of short-wave fadeout by a solar flare was examined on the basis of long-term ionosonde observation data (Tao et al., in this issue).

However, extreme TEC values of once per long period of time have not yet been quantitatively estimated. Several countries have prepared documents with space weather benchmarks. The US White House published "Space Weather Phase 1 Benchmarks" in June 2018 (US White House, 2018). Although it lists three factors that cause ionospheric disturbances, such as geomagnetic storms, quantitative benchmarks were not provided because the ionospheric effects of geomagnetic storms on the ionosphere largely differ from event to event and even their mechanism is not completely understood.

Another reason why extreme TEC values have not been fully studied is that only 20 years has passed since the start of fully fledged TEC observations. TEC observations started with measurements of the Faraday rotation or Doppler effect many decades ago. Since these observations were conducted by a few transmitters and receivers, it is difficult to study TEC behavior statistically. With the spread of GNSS and its ground-based receivers, the number of TEC observations dramatically increased. Thanks to the GNSS-TEC observation systems, we have learned a lot about TEC behavior during the last 20 years (for example, Foster, 2007; Nishioka et al., 2009 Maruyama et al., 2013). The purpose of this study is to estimate extreme values of TEC with their occurrence rates. We investigate the occurrence rates of extreme values of TEC in Japan in the short, mid-, and long term, which are once per year, ten years, and hundred years, respectively.

To evaluate TEC corresponding to an occurrence rate of once per hundred years, 20 years of data is obviously insufficient. Furthermore, solar activity in the last 20 years has on average been moderate, although several intense geomagnetic storms occurred during solar cycle 24. Compared with GNSS-TEC observation, ionosonde observation has a much longer history. This technique was developed in the late 1920s and began to be implemented in the 1940s in order to monitor shortwave propagation. In Japan, ionosonde observation began in 1931. After going through various changes, routine ionosonde observation was started by the predecessor of National Institute of Information and Communications Technology (NICT) in 1951 using an automatic system. Ionospheric parameters derived from the long-term ionosonde observation are archived by World Data Center for the Ionosphere at NICT (<http://wdc.nict.go.jp/IONO/wdc/>). Long-term ionosonde data have been used for various studies such as a study of the long-term trends of the ionosphere (Xu et al., 2004) and for the development of empirical models (Bilitza, 2018; Yue et al., 2006; Maruyama, 2011). As the TEC and the maximum density of the F region derived from ionosonde observation (NmF2) are known to be correlated, NmF2 can be a proxy of TEC. In this study, about 70 years of data of ionospheric parameters derived from the long-term ionosonde observation are used. Although the data period is still shorter than one hundred years, we

investigate statistical characteristics of extreme TEC values in order to estimate the ionospheric once-per-hundred-year condition.

The TEC value over Japan depends on the latitude, normally with a larger value in southern Japan. Japan is mainly located in the lower mid-latitude region with a latitudinal range of about 20 degrees. The southern part of Japan is located at the poleward slope of the equatorial ionospheric anomaly (EIA) crest. On the other hand, the northern part is hardly affected by EIA variation and may rather be affected by phenomena originating from the polar region (Cherniak et al., 2015). Therefore, extreme TEC values should also differ among the center, southern, and northern parts of Japan.

Details of the data set used in this study and the analysis method are described in Sections 2 and 3, respectively. Analysis results are presented in Section 4. In Section 4, the result obtained using about 20 years of TEC data collected in Tokyo, which is almost in the center of Japan, is shown as the first step. Then long-term ionosonde data are analyzed. On the basis of the result, extreme TEC values with probabilities of once per year, ten years, and hundred years are estimated for Tokyo. In the last part of Section 4, the extreme TEC values in southern and northern Japan are also estimated. In Section 5, the results are discussed in comparison with those of case studies of geomagnetic storms in previous papers. Section 6 provides the conclusions of this study.

2. Data Set

In this study, we use TEC data derived from the nationwide GNSS network over Japan, which is called the GNSS Earth Observation Network System (GEONET) and operated by the Geospatial Information Authority of Japan, and ionosonde observation data collected over Tokyo.

GNSS-TEC data derived from GEONET have been archived by NICT since 1997. Using the network data, the slant TEC along the line of sight between the receiver and the satellite was derived from pseudo-range and carrier-phase measurements by dual-frequency GPS receivers (Saito et al., 1998). The instrumental bias of the TEC associated with the inter-frequency bias of the satellite and receiver was obtained by a technique proposed by Otsuka et al. (2002), in which the daily bias values are derived by assuming that hourly averaged TEC values are uniform within the field of view of a given GNSS receiver. The slant TEC is converted to the vertical TEC after removing the instrumental bias. The median value of the vertical TEC whose ionospheric pierce point is located within 100 km from a given location over one hour is derived as an hourly TEC. The largest hourly TEC in a given day is noted as the daily TEC in this paper. The daily TECs of 22 years from 1997 to 2018 are used in this study and studied in Section 4.1.

Ionospheric conditions have been monitored for about 70 years by NICT using ionosondes in Kokubunji, Tokyo (36.7°N, 139.5°E, 26.8°E in Mag.Lat) and other stations. Ionospheric parameters have been manually scaled from ionograms. In order to ensure uniform quality of data, the scalers have discussed and established scaling rules, although automatic scaling tools have been developed in recent years. Thanks to the substantial efforts of the scalers, ionospheric parameters from the 1950's to the present are now available. In this study, the manually scaled critical frequency of the F-layer (f_{OF2}), which

corresponds to the peak density of the F-layer, is used. In order to study foF2 with the daily TEC, we refer to the maximum foF2 in a given day as the daily foF2. In Section 4.2, a 22-year data set of daily foF2 values from 1997 to 2018 is used. In the same section, a 62-year data set of daily foF2 values from 1957 to 2018 is also used.

3. Method

In order to find extreme values of TEC corresponding to an occurrence frequency of once every certain number of years, the cumulative distribution function (CDF) of daily TEC occurrence is investigated. The CDF of the daily TEC occurrence is a distribution function of daily TEC values that are greater than or equal to a critical TEC, which is denoted as $TEC_{critical}$. One of the advantages of investigating the CDF instead of a simple occurrence probability is that it is easy to find TEC values with an occurrence frequency of once per long period (Riley, 2012). In other words, the CDF of the daily TEC occurrence provides an occurrence probability of a daily TEC that is greater than or equal to a certain value, while a normal distribution provides the occurrence probability of a daily TEC between two values.

Although a data set of TEC values over 22 years may be sufficient to investigate TEC values with occurrence frequency of once per year and ten years, it would not be sufficient to investigate the TEC value with an occurrence frequency of once per hundred years.

To compensate the insufficient TEC data set, we utilized a 62-year data set of foF2 values and study a property of the relationship between TEC and foF2. The relationship between TEC and foF2 is given by the following equation:

$$TEC = S \cdot NmF2,$$

where S is the slab thickness. In this study, characteristics of slab thickness are studied using the 22-year data set of TEC and foF2 values. By utilizing the characteristics of the slab thickness and the 62 years of foF2 data, we deduce CDFs of TEC values over 62 years, from which we estimate the TEC value corresponding to occurrence frequency of once per hundred years.

4. Results

4.1 Statistical analysis of TEC over 22 years

Figure 1 shows the CDF of the daily TEC occurrence at Tokyo. The occurrence rate is shown on the left axis. The occurrence rate on the left-hand axis of the ordinate is days per hundred years, that is, an occurrence rate of one day means an occurrence rate of once per hundred years. The occurrence rate is converted to the occurrence percentage and shown on the right-hand axis of the ordinate. An occurrence probability of 0.3%, which corresponds to a frequency of once per year, is shown as a solid horizontal line. It is found that the daily TEC can reach about 90 TECU with a frequency of once per year. The occurrence probabilities of once per ten years and once per hundred years correspond to 0.03% and

0.003% and are shown with dotted and dashed horizontal lines, respectively. It is found that a daily TEC of more than 100 TECU occurs with a frequency of once per ten years. The TEC values with frequencies of once per year and once per ten years are summarized in Table 1. On the other hand, the daily once-per-hundred-year TEC value cannot be appropriately estimated from Figure 1 because the distribution is based on only 22 years of data.

The colors in the histograms in Figure 1 represent the classifications based on solar and geomagnetic activity: red, pink, blue, and light blue represent days of high solar activity and high geomagnetic activity (HSHG), high solar activity and low geomagnetic activity (HSLG), low solar activity and high geomagnetic activity (LSHG), and low solar activity and low geomagnetic activity (LSLG), respectively. Solar and geomagnetic activities are respectively defined on the basis of the solar sunspot number (SSN) and disturbance storm-time (DST) index, which are provided as sunspot data from the World Data Center SILSO, Royal Observatory of Belgium, Brussels (<http://sidc.be/silso/datafiles>) and WDC for Geomagnetism, Kyoto (<http://wdc.kugi.kyoto-u.ac.jp/dstdir/index.html>), respectively. HS (LS) days are defined as days for which the average daily SSN for the previous 27 days is greater than or equal to (less than) 50. HG (LG) days are defined as days for which the average daily DST of the current day and the previous day is less than or equal to (greater than) -50 nT. It can be seen that a TEC of 60 TECU or larger is most likely to be observed when either the solar activity or the geomagnetic activity is high, while those exceeding 100 TECU are observed when the solar activity is high.

4.2 Statistical analysis of foF2 over 22 and 62 years

Here, CDFs of the daily foF2 occurrence are studied in order to estimate once-per-hundred-year values. First, a CDF of the daily foF2 occurrence over the same period as in Figure 1, from 1997 to 2018, were examined in comparison with that of the 22 years of TEC data in Figure 1. Figure 2 shows a CDF of the daily foF2 occurrence, that is, the distribution of the daily foF2 that is greater than or equal to some critical foF2, $foF2_{critical}$. As in Figure 1, the occurrence rate per hundred years is shown on the left-hand axis of the ordinate and the occurrence rate in percentage is shown on the right axis. The occurrence frequencies of once per year, ten years, and hundred years of 0.3%, 0.03%, and 0.003% are shown as solid, dotted, and dashed horizontal lines, respectively. The colors in Figure 2 represent solar and geomagnetic activities similarly to in Figure 1; red, pink, blue, and light blue represent days of HSHG, HSLG, LSHG, and LSLG, respectively. The largest foF2 was about 16.7 MHz. It is found that foF2 was higher than 15 MHz for only HSHG and HSLG days, which is similar to the result in Figure 1.

The same analysis is carried out for the 62-year foF2 data set from 1957 to 2018. The result is shown in Figure 3 in the same format as Figure 2. The maximum observed foF2 is about 18.7 MHz, which is slightly larger than that obtained from the 22-year data set in Figure 2. Moreover, the occurrence rate of daily foF2 values larger than 16.8 MHz in Figure 3, which corresponds to the rightmost bar in the histogram, is about twice of that in Figure 2.

4.3 Estimation of extreme TEC from slab thickness

As the characteristics of the CDFs of the daily foF2 occurrence are different for the 22- and 62-year data sets, the once-per-hundred-year TEC value cannot be estimated by extrapolating the CDF of the daily TEC occurrence obtained from the 22-year data set. In this section, we estimate the once-per-hundred-year TEC value by using the 62-year foF2 data set.

The value of foF2 is proportional to the square root of the maximum ionospheric density, NmF2. NmF2 is given by the following equation.

$$\text{NmF2}[\text{m}^{-3}] = 1.24 \times 10^{10} \times \text{foF2}^2[\text{MHz}]$$

Figure 4 shows the correlation between daily TEC and NmF2 derived from the daily foF2. All data collected over 22 years are shown in this scatter plot. It can be seen that TEC and NmF2 have a strong correlation. The red line is the least-squares linear approximation of all data. The slope, which is about 250 km, is equivalent to the thickness of the ionosphere that gives a TEC value with a density of NmF2. This parameter, which is called the ionospheric slab thickness, is used to deduce TEC from NmF2 because of the strong correlation between daily TEC and daily foF2.

Since the slab thickness is known to have seasonal dependence, a single value of the slab thickness is not appropriate for estimating TEC from foF2 or equivalently NmF2. Figure 5 shows the slab thickness against the day of the year for 22 years from 1997 to 2018. Data are sparser from June to August compared with other months, because foF2 values often cannot be obtained owing to masking by the sporadic E-layer, which often appears in these months. The red polyline is the monthly mean of the slab thickness. The monthly mean slab thickness is about 180 km in winter and 250 km in summer. Blue and red vertical lines indicate the ranges of $\pm 3\sigma$ and $\pm 4.2\sigma$, which are equivalent to a probabilities of once per ten and hundred years, respectively, when the estimated slab thickness is assumed to have a normal distribution.

Here we estimate the daily TEC from the daily NmF2 data, assuming the slab thickness has only seasonal dependence. Figure 6 shows the CDFs of the estimated daily TEC occurrence obtained using the monthly mean slab thickness and observed NmF2 from 1957 to 2018. The black histograms are distributions of the daily TEC estimated with the monthly mean slab thickness, which is shown with a red polyline in Figure 5. The number of days per 100 years and the occurrence rate are shown on the left- and right-hand axes of the ordinate, respectively. The black solid, dotted, and dashed horizontal lines correspond to 0.3% (once a year), 0.03% (once every ten years), and 0.003% (once every hundred years), respectively. The blue histograms in Figure 6 are the distribution of TEC estimated with the average + 3σ slab thickness (upper value of the blue vertical line in Figure 5), which corresponds to a slab thickness with a frequency of once per ten years. According to this histogram, the TEC with a frequency of once per ten years is 130 TECU or more. Furthermore, the red histograms in Figure 6 are derived from the average + 4.2σ slab thickness (upper limit of the red vertical line in Figure 3). This result indicates that TEC values of more than 190 TECU can be observed with a frequency of once per hundred years. These TEC values are summarized in Table 1.

4.4 Latitudinal dependence of extreme TEC

Figures 1–6 are results based on data obtained in Tokyo. Here we estimate extreme TEC values for southern and northern Japan because TEC behavior is expected to be different at different magnetic latitudes. Figure 7 shows the correlations of daily TEC between Tokyo and Kagoshima (31.2°N , 130.6°E , 21.7°N in Mag. Lat) and between Tokyo and Hokkaido (45.2°N , 141.8°E 36.4°N in Mag. Lat) for 22 years from 1997 to 2018. Basically, the TEC in Tokyo is smaller than that in Kagoshima and larger than that in Hokkaido. The red line represents the linear approximation of these data and reveals that the TECs in Kagoshima and Hokkaido are, on average, 1.2 and 0.8 times that in Tokyo, respectively. From these results, the TEC values with probabilities of once per year, ten years, and hundred years are estimated as 110, 160, and 230 TECU (70, 105, and 150 TECU), respectively, in Kagoshima (Hokkaido) as summarized in the second and third rows in Table 1.

5. Discussion

It is important to estimate the occurrence rates of extreme values of TEC in Japan in the short, mid-, and long term, which are once per year, ten years, and hundred years, respectively, in readiness for hazardous ionospheric conditions. “Space Weather Phase 1 Benchmarks”, which was published by the USA White House in June 2018, lists three factors that cause ionospheric disturbances: solar flares, proton events, and geomagnetic storms. However, quantitative benchmarks are difficult to derive because the effects of geomagnetic storms largely differ from event to event. Furthermore, the mechanism of ionospheric storms is not yet completely understood. Although the results in this paper are limited to the region around Japan, they are a starting point for evaluating benchmarks in other regions.

In this study, we estimated extreme TEC values by assuming that the slab thickness has only seasonal dependence. The seasonal dependence of the slab thickness shown in Figure 5 is consistent with the results of previous studies (Jin et al., 2007; Huang et al., 2016). Another factor determining the slab thickness is the dynamics and/or composition change caused by geomagnetic disturbances. According to Stankov and Warnant (2009), the slab thickness is systematically enhanced during geomagnetic disturbances for both positive and negative ionospheric storms. Extreme values of TEC during geomagnetic storms can be obtained by using a slab thickness of mean+ 4.2σ to estimate CDFs of the daily TEC occurrence in Figure 6.

Large values of TEC have mainly been recorded during geomagnetic storms. The largest reported TEC is about 330 TECU to our knowledge, which was recorded by a GPS receiver onboard the CHAMP satellite at an altitude of about 400 km during the October 2003 Halloween storm (Mannucci et al., 2005). Extreme positive storms are thought to be caused by a geomagnetic disturbance that induces prompt penetration of the electric field (Tsurutani et al., 2004). The TEC value of 330 TECU is much higher than the results in this paper. Here we discuss possible reasons for the difference between these values. One possibility is differences in observation opportunities. The characteristics of ionospheric storms are not always similar among geomagnetic storms, with their magnitude varying greatly from event to event. Mannucci et al.

(2008) analyzed four intense geomagnetic storms in 2003 including the event for which the extreme value of 330 TECU was observed by the CHAMP satellite. A dramatic increase in TEC was observed in only one event. The observed TEC on the other three storm days was around 100 TECU or less. If the event-to-event difference is too large, 70 years of data might not be enough to estimate TEC values for once-per-hundred-year or once-per-thousand-year events.

Another possibility accounting for the difference between the extreme value of 330 TECU in Mannucci et al. (2005) and our result is the longitude dependence of the ionospheric influence on geomagnetic storms. Immel and Mannucci (2013) analyzed global TEC maps during geomagnetic storms over seven years. Their analysis confirmed that on average the American sector exhibits larger TEC enhancements regardless of the onset UT. Greer et al. (2017) used the Global Ionosphere–Thermosphere Model to carry out an experiment on a geomagnetic storm by modifying the storm arrival UT. The result indicated that the strongest enhancements of TEC during storms are found in the American and Pacific longitude sectors. They suggested that the longitudinal dependences were due to Earth's asymmetrical geomagnetic topology in the American and Pacific sectors. The difference between our results and that of Mannucci et al. (2005) may originate from the difference between the Japanese and American/Pacific sectors. In order to clarify whether the longitudinal dependence results in the large difference between the results of this study and that of Mannucci et al. (2008), long-term observational data in addition to data over oceans are necessary.

This study focuses on positive ionospheric storms, which may significantly affect GNSS users. On the other hand, the effect of negative storms on space weather users may also be significant, particularly for HF communicators, who may experience blackouts during negative ionospheric storms. In addition, parameters other than TEC, such as maximum usable frequency (MUF) and scintillation indices, should be studied for extreme cases.

6. Summary

In this study, extreme values of TEC with frequencies of once per year, ten years, and hundred years were investigated. The results are summarized as follows:

- The CDF of daily TEC values was studied for a 22-year data set observed in Tokyo in order to estimate TECs with frequencies of once per year and ten years. The obtained once-per-year and once-per-ten-year TECs were 90 and 130 TECU, respectively.
- In order to estimate the once-per-hundred-year TEC value, 62 years of manually scaled ionosonde data were used to augment the insufficient observation period of TEC. The slab thickness was assumed to have only seasonal variation and was used to estimate TEC from 62 years of foF2 data. The obtained once-per-hundred-year TEC value was 190 TECU in Tokyo.
- Extreme TEC values were also studied for Kagoshima and Hokkaido in southern and northern Japan, respectively. Once-per-hundred-year values of 230 and 150 TECU were obtained, respectively.

Abbreviations

EIA	equatorial ionospheric anomaly
EUV	solar extreme ultraviolet (EUV)
foF2	critical frequency of the F-layer
GEONET	GNSS Earth Observation Network System
GNSS	Global Navigation Satellite System
HF	high frequency
HSHG	high solar and high geomagnetic activity
HSLG	high solar and low geomagnetic activity
LSHG	low solar and high geomagnetic activity
LSLG	low solar and low geomagnetic activity
MUF	maximum usable frequency
NICT	National Institute of Information and Communications Technology
NmF2	maximum density of the F2 layer
TEC	total electron content

Declarations

Availability of data and materials

The TEC data used in this study are archived on NICT's homepage (<https://aer-nc-web.nict.go.jp/GPS/GEONET/>). Manually scaled ionosonde parameters are also archived on NICT's homepage (http://wdc.nict.go.jp/IONO/HP2009/ISDJ/manual_txt.html).

Competing interests

The authors declare that they have no competing interests.

Funding

Extreme TEC values in once-per-year, once-per-ten-year, and once-per-hundred year are estimated to be 90 TECU, 130 TECU, and 190 TECU at Tokyo, Japan.

Extreme TEC values in once-per-hundred years in Kagoshima and Hokkaido, which is southern and northern Japan, respectively, are estimated to be 230 and 150 TECU, respectively.

Authors' contributions

MN conducted the research and has responsibility for the results presented in this paper. SS has supported this analysis and contributed to the discussion. CT, DS, TT, and MI contributed to the discussion as experts of ionosphere and space weather. All authors read and approved the final manuscript.

Acknowledgements

GPS data of GEONET were provided by the Geospatial Information Authority of Japan. MN thanks to ionograms scalers of NICT and forerunner of NICT for valuable manually scaled data.

References

1. Bilitza D (2018) IRI the International Standard for the Ionosphere. *Adv Radio Sci* 16:1–11. <https://doi.org/10.5194/ars-16-1-2018>
2. Cherniak I, Zakharenkova I, Redmon RJ (2015) Dynamics of the high-latitude ionospheric irregularities during the 17 March 2015 St. Patrick's Day storm: Ground-based GPS measurements. *Space Weather* 13:585–597. doi:10.1002/2015SW001237
3. Foster JC, Rideout W, Sandel B, Forrester WT, Rich FJ (2007) On the relationship of SAPS to storm-enhanced density. *J Atmos Sol Terr Phys* 69:303–313
4. Greer KR, Immel T, Ridley A (2017) On the variation in the ionospheric response to geomagnetic storms with time of onset. *J Geophys Res Space Phys* 122:4512–4525. doi:10.1002/2016JA02345
5. Huang H, Liu L, Chen Y, Le H, Wan W (2016) A global picture of ionospheric slab thickness derived from GIM TEC and COSMIC radio occultation observations. *J Geophys Res Space Phys* 121:867–880. doi:10.1002/2015JA021964
6. Immel TJ, Mannucci AJ (2013) Ionospheric redistribution during geomagnetic storms. *J Geophys Res Space Phys* 118:7928–7939. doi:10.1002/2013JA018919
7. Jin S, Cho J-H, Park J-U (2007) Ionospheric slab thickness and its seasonal variations observed by GPS. *J Atmos Sol Terr Phys* 69(15):1864–1870. doi:10.1016/j.jastp.2007.07.008

8. Mannucci AJ, Tsurutani BT, Iijima BA, Komjathy A, Saito, W A, Gonzalez D, Guarneri FL, Kozyra JU, Skoug R (2005) Dayside global ionospheric response to the major interplanetary events of October 29, 30, 2003 Halloween Storm. *Geophys Res Lett* 32:L12S02. doi:10.1029/2004GL021467
9. Mannucci AJ, Tsurutani BT, Abdu MA, Gonzalez WD, Komjathy A, Echer E, Iijima BA, Crowley G, Anderson D (2008) Superposed epoch analysis of the dayside ionospheric response to four intense geomagnetic storms. *J Geophys Res* 113:A00A02, doi:10.1029/2007JA012732
10. Maruyama T (2011) Modified solar flux index for upper atmospheric applications. *J Geophys Res* 116:A08303. doi:10.1029/2010JA016322
11. Maruyama T, Ma G, Tsugawa T (2013) Storm-induced plasma stream in the low-latitude to midlatitude ionosphere. *J Geophys Res Space Phys* 118:5931–5941. doi:10.1002/jgra.50541
12. Nishioka M, Saito A, Tsugawa T (2009) Super-medium-scale traveling ionospheric disturbance observed at midlatitude during the geomagnetic storm on 10 November 2004. *J Geophys Res* 114:A07310. doi:10.1029/2008JA013581
13. Otsuka Y, Ogawa T, Saito A, Tsugawa T, Fukao S, Miyazaki S (2002) A new technique for mapping of total electron content using GPS network in Japan. *Earth Planets Space* 54:63–70
14. Riley P (2012) On the probability of occurrence of extreme space weather events. *Space Weather* 10:S02012. doi:10.1029/2011SW000734
15. Saito A, Fukao S, Miyazaki S (1998) High resolution mapping of TEC perturbations with the GSI GPS network over Japan. *Geophys Res Lett* 25:3079–3082. doi:10.1029/98GL52361
16. Stankov SM, Warnant R (2009) Ionospheric slab thickness. Analysis, modelling and monitoring. *Adv Space Res* 44:1295–1303
17. Tao C, Nishioka M, Tsugawa T, Saito S, Shiota D, Watanabe K, Ishii M (2020) Statistical analysis of short-wave fadeout for extreme event estimation. *Earth Planets Space*, submitted to this issue
18. Tsubouchi K, Omura Y (2007) Long-term occurrence probabilities of intense geomagnetic storm events. *Space Weather* 5:S12003. doi:10.1029/2007SW000329
19. Tsurutani BT et al (2004) Global dayside ionospheric uplift and enhancement associated with interplanetary electric fields. *J Geophys Res* 109:A08302. doi:10.1029/2003JA010342
20. US White House (2018) Space Weather Phase 1 Benchmarks. Space Weather Operations, Research, and Mitigation Subcommittee Committee on Homeland and National Security, National Science & Technology Council. <https://www.whitehouse.gov/wp-content/uploads/2018/06/Space-Weather-Phase-1-Benchmarks-Report.pdf>
21. Xu Z-W, Wu J, Igarashi K, Kato H, Wu Z-S (2004) Long-term ionospheric trends based on ground-based ionosonde observations at Kokubunji, Japan. *J Geophys Res* 109:A09307. doi:10.1029/2004JA010572
22. Yue X, Wan W, Liu L, Ning B, Zhao B (2006) Applying artificial neural network to derive long-term foF2 trends in the Asia/Pacific sector from ionosonde observations. *J Geophys Res* 111:A10303. doi:10.1029/2005JA01157

Tables

Table 1.

Estimated TEC of once per one, ten, and hundred years in Tokyo, Kagoshima, and Hokkaido.

	Once-in-one-year TEC	Once-in-ten-year TEC	Once-in-hundred-year TEC
Tokyo	~90 TECU	130 TECU	190 TECU
Kagoshima	~110 TECU	160 TECU	230 TECU
Hokkaido	~70 TECU	105 TECU	150 TECU

Figures




Figure 1

Cumulative distribution function (CDF) of daily TEC occurrence at Tokyo from 1997 to 2018. The occurrence rate, which is the number of days per hundred years, and the occurrence percentage are shown on the left and right axes, respectively. Red and pink (blue and light blue) colors represent days when solar activity is high (low). Red and blue (pink and light blue) colors represent days when geomagnetic activity is high (low). The solid, dotted, and dashed horizontal lines represent occurrence rates of 0.3%, 0.03%, and 0.003%, which correspond to occurrence frequencies of once per year, ten years, and hundred years, respectively.




Figure 2

CDF of the daily foF2 occurrence from 1997 to 2018 at Kokubunji station, Tokyo. The occurrence rate, which is the number of days per hundred years, and the occurrence rate in percentage are shown on the left- and right-hand axes of the ordinate, respectively. Red and pink (blue and light blue) colors represent days when solar activity is high (low). Magenta and blue (pink and light blue) colors represent days when geomagnetic activity is high (low). The solid, dotted, and dashed horizontal lines represent occurrence rates of 0.3%, 0.03%, and 0.003%, which correspond to frequencies of once per year, ten years, and hundred years, respectively.




Figure 3

CDF of the daily foF2 from 1957 to 2018. The plotting format is the same as that of Figure 2.




Figure 4

Scatter plot of daily TEC and corresponding daily NmF2 from 1997 to 2018. The red line represents a linear fitting to the data points.




Figure 5

Slab thickness against day of year: The red polyline is the monthly mean value of slab thickness. Red and blue vertical bars represent $\pm 3\sigma$ and $\pm 4.2\sigma$, respectively.




Figure 6

CDFs of the daily TEC occurrence estimated from daily foF2 from 1957 to 2018 and slab thickness. The occurrence rate, which is the number of days per hundred years, and the occurrence rate in percentage are shown on the left- and right-hand axes of the ordinate, respectively. The black histograms are derived from the average slab thickness shown in Figure 5. The blue and red histograms are derived with slab thicknesses of average+ 3σ and + 4.2σ , which are shown with blue and red vertical lines, respectively. The solid, dotted, and dashed horizontal lines represent occurrence rates of 0.3%, 0.03%, and 0.003%, which correspond to frequencies of once per year, ten years, and hundred years, respectively.




Figure 7

Correlation of daily TEC between (a)Tokyo and Kagoshima and (b)Tokyo and Hokkaido from 1997 to 2018. The red line represents the linear approximation of each set of data.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [abstfig.JPG](#)