Morphological Neurite Changes Induced by Porcupine Inhibition Are Rescued by Wnt Ligands

Juan A. Godoy

Pontificia Universidad Catolica de Chile

Jasson Espinoza-Caicedo

Pontificia Universidad Catolica de Chile

Nibaldo C Inestrosa (✉ ninestrosa@bio.puc.cl)

Pontificia Universidad Catolica de Chile

Research

Keywords: Wnt signaling, porcupine, neurites, dendritic arbor complexity, young hippocampal neurons

Posted Date: August 6th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-42321/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Cell Communication and Signaling on August 16th, 2021. See the published version at <https://doi.org/10.1186/s12964-021-00709-y>.

Abstract

Background: Wnt signaling plays key roles in cellular and physiological processes, including cell proliferation, differentiation and migration during development and tissue homeostasis in adults. This pathway can be defined as Wnt/ β -catenin-dependent or β -catenin-independent or "non-canonical", both signaling are involved in neurite and synapse development/maintenance. Porcupine (PORCN), an acylase that o-acylates Wnt ligands, a major modification in secretion and interaction with its receptors. We use Wnt-C59, a specific PORCN inhibitor, to block the secretion of endogenous Wnts in young hippocampal neurons (DIV 4). Under these conditions, the morphology and length of the neurites and the complexity of the dendritic tree and axonal polarity were evaluated.

Methods: Cultured primary young hippocampal neurons obtained from Sprague-Dawley rat fetuses (E18), were cultured until day *in vitro* (DIV) 4 (according to Banker's protocol) and treated with Wnt-C59 for 24h, Wnt ligands were added to the cultures on DIV 3 for 24h. Dendritic arbors and neurites were analysis by fluorescence microscopy. Transfection with Lipofectamine 2000 on DIV 2 of plasmid expressing eGFP and KIF5-Cherry and neurons were fixed on DIV 4. Immunostaining was performed with MAP1B and Tau protein. Immunoblot analysis was carried out with Wnt3a, b-catenin and GSK-3b (p-Ser9). Quantitative analysis of dendrite morphology was carried out with ImageJ (NIH) software with Neuron J Plugin.

Results: We report, here, that Wnt-C59 treatment changed the morphology of the dendritic arbors and neurites of young hippocampal neurons, with decreases β -catenin and Wnt3a and an increase in GSK-3 β (p-Ser9) levels. No effect was observed on axonal polarity. In sister cultures, addition of exogenous Wnt3a, 5a and 7a ligands rescued the changes in neuronal morphology. Wnt3a restored the length of neurites to near that of the control, but Wnt7a increased the neurite length beyond that of the control. Wnt5a also restored the length of neurites relative Wnt concentrations.

Conclusions: Our results indicated that all 3 Wnt ligands restored dendritic arbor complexity with recovery of secondary and tertiary projections in young hippocampal neurons. We proposed that PORCN is an emerging molecular target of interest in the search for preclinical options to study and treat neurological diseases.

Background

Proper functioning of the Wnt signaling pathway is critical to the regulation of numerous processes, including cellular polarity, cell proliferation and cellular differentiation (1, 2). Wnt ligands are growth factors that can influence the cell cycle and affect cell proliferation; however, they also contribute to cytoskeleton arrangement and therefore give directionality to cell proliferation, regulate spatial growth and provide signals to orient asymmetric stem cell division *in vitro* (3, 4, 5). The Wnt signaling pathway is active in various types of cells in the central nervous system (CNS), including fibroblast-like cells, radial glia, oligodendrocytes, microglia/macrophages, astrocytes and neurons (6, 7, 8, 9). In the CNS, Wnt signaling, also participates in the formation of synapses and neuronal circuits, and in the adult brain, Wnt

signaling modulates synaptic transmission, plasticity and neurogenesis (6, 10, 11, 12). Wnt ligands are secreted short-range molecules that can trigger two intracellular pathways, the Wnt- β -catenin-dependent and the "noncanonical" β -catenin-independent signaling (13, 14). Nineteen Wnt proteins have been described in humans and recognized by the Frizzled (Fzd) family of receptors (2).

Neurons are highly polarized cells with two distinct morphological structures, axons and dendrites. Synapses form between the contacts of these two specialized regions. In general, the axon is a long process that communicates with other neurons through the release of neurotransmitters (15, 16). Dendrites are more complex structures formed by multiple branching processes and dendritic spines, which are small protuberance of the plasma membrane where the postsynaptic excitatory apparatus is assembled and are essential for the regulation of postsynaptic receptor levels (17, 18). The formation and complexity of these processes is critical to the directional flow of information between neurons in the CNS. Granular cerebellar, cortical, and hippocampal neurons become polarized as they differentiate, and a cell culture system was established to study rodent hippocampal neurons *in vitro* more than four decades ago. This *in vitro* system has been fundamental in research on the structuring of neuronal shape and cellular and molecular mechanisms during neuronal polarization (19, 20). In that *in vitro* model, hippocampal neurons spontaneously polarize, spreading filopodia, and lamellipodia appear with small neurites, then one of the neurites grows faster to become an axon, and the remaining neurites develop into dendrites. Finally, in later stages of culture, axons and dendrites undergo further development to form functional synapses (21, 22).

The aim of our work was to evaluate the effects of Wnt-C59 (23) a high affinity specific inhibitor of the activity of PORCN, a membrane-bound O-acyltransferase required for the palmitoylation, secretion and biological activity of Wnts (24, 25, 26, 27). We expected Wnt-C59 to inhibit the action of both canonical and noncanonical endogenous ligands. Rescue experiments with specific Wnt ligands were carried out to determine the contribution of individual Wnt ligands to dendritic arbor formation and the formation/maintenance of dendritic spines.

Methods

Animals and ethical standards

Experiments were performed on Sprague-Dawley rat fetuses (E18). All rats were housed at the Animal Warehouse Facility of the Pontificia Universidad Católica de Chile. The Bioethical and Biosafety Committee of the Faculty of Biological Sciences of our university approved the experimental procedures.

Primary rat hippocampal neuron cultures

Hippocampal neurons were obtained from 18-day-old Sprague-Dawley rat embryos (19). Hippocampi were aseptically dissected and trypsinized for 16 min. After centrifugation for 1 min, cells were seeded in phenol red-free Dulbecco's modified Eagle's medium plus 10% horse serum into 1% poly-L-lysine-coated plates. After 120 min, the medium was removed, and neurobasal medium containing 1% B27 supplement

(Invitrogen), streptomycin and penicillin was added. To obtain young hippocampal neurons, neurons were cultured until day *in vitro* (DIV) 4 and treated with different concentrations of Wnt-C59 for 24 h, and different Wnt ligands were added to the cultures on DIV 3 for 24 h.

Reagents

Antibodies against ATF2, MAP1B, Tau, Ankin G, β -catenin, GSK-3 β (ser9), β -actin, MAP2 and phalloidin were obtained from Cell Signaling Technology, Inc. (Trask Lane, Danvers, MA). Piccolo was a gift from Craig Garner's laboratory. Neurobasal growth medium was obtained from Life Technologies (Carlsbad, CA). Other reagents were purchased from Sigma (St. Louis, MO). Wnt-C59 inhibitor was dissolved in DMSO and added to neuronal cultures at a final concentration of 0.01% DMSO or less (Cellagen Technology, San Diego, CA). Recombinant Wnt3a, Wnt5a, and Wnt7a (R&D Systems, Abingdon, UK) were used at 50 ng/ml, 100 ng/ml, and 200 ng/ml, respectively with corresponding dilutions in vehicle.

Fluorescence microscopy

After fixation, neurons were permeabilized with 0.5% (v/v) Triton X-100 in 1x PBS for 5 min and washed in 1x in PBS. Then, the cells were incubated for 30 min in blocking solution (2% glycine, 2% BSA, 5% FBS). To study dendrite morphology neurons were stained with MAP2 and Tau antibodies. Highly cross-absorbed Alexa-conjugated antibodies were used as secondary labels. Coverslips were mounted on slides with Vectashield (Vector Labs) medium and images were captured in a Carl Zeiss LSM microscope.

Transfection of neuronal hippocampal cultures

Primary hippocampal neuronal cultures were transfected with Lipofectamine 2000 on DIV 2 with plasmid expressing eGFP or KIF5-cherry, and cells were fixed on DIV 4. Immunostaining was performed for MAP1B and Tau protein. Neuronal cultures were fixed at the times indicated in the figure legends with LF fixative solution (60 mM PIPES, 25 mM HEPES, 10 mM EDTA, 2 mM MgCl₂, 0.12M sucrose, 4% paraformaldehyde) for 5 min and then washed once with phosphate buffered saline (PBS) before processing for immunostaining.

Immunoblot analysis.

Total protein extraction from cultured hippocampal neurons and immunoblot analysis were performed as previously described (28). At the designated times, cells were lysed in RIPA buffer (10 mM Tris-HCl, pH 7.4, 5 mM EDTA, 1% NP-40, 1% sodium deoxycholate, and 1% SDS) supplemented with a protease inhibitor mixture (1 mM PMSF, 2 μ g/mL aprotinin, 1 μ g/mL pepstatin and 10 μ g/mL benzamidine) and phosphatase inhibitors (25 mM NaF, 100 mM Na₃VO₄, 1 mM EDTA and 30 μ M Na₄P₂O₇), homogenized by scraping, and then passed sequentially through different caliber syringes. Protein samples were centrifuged twice at 10,000 rpm and 4 °C for 5 min. Protein concentration was determined using the BCA Protein Assay Kit (Pierce). Samples were resolved by 10% SDS-PAGE and transferred to a PVDF membrane. The membranes were incubated with antibodies against β -catenin, GSK-3 β (p-ser9), Wnt3a (Thermo Scientific, Rockford, IL), (Cell Signaling Technology, Danvers, MA) and β -actin (Sigma-Aldrich, St. Louis, MO), followed by anti-mouse, anti-goat or anti-rabbit IgG peroxidase-conjugated antibodies (Pierce,

Rockford, IL). The membranes were developed using an ECL Kit (Western Lightning Plus ECL, PerkinElmer).

Image and statistical analysis

The quantitative analysis of dendrite morphology was carried out with ImageJ (NIH) software with the assistance of the Neuron J Plugin. All dendrites from each neuron were quantified for parameters such as length and primary, secondary and tertiary dendrite numbers. All data are expressed as the mean $\pm$ SEM, and the number of experiments is indicated in the corresponding figures. Differences between groups were determined by ANOVA, one-way post hoc Bonferroni of multiple comparisons to establish significant differences. A value of $p < 0.05$ was considered significant.

Results

Inhibition of porcupine by Wnt-C59 triggers the loss of neurites in young cultures of hippocampal neurons

Neurons construct a characteristic polarized dendrite axis, which is physiologically important for receiving and processing axon synaptic signals. In low-density cultures, polarization and active axon growth occur between DIV 2–6; therefore, this time is ideal to study the role of endogenous Wnt signaling in dendritic tree formation and axonal processes. For this purpose, we blocked Wnt signaling with the acylase inhibitor Wnt- C59 which is specific for the acylase PORCN. Rat hippocampal neurons DIV 4, were treated with different Wnt-C59 concentrations (0, 0.1, 1, 10, and 100 nM) for 36 h, and stained for microtubule-associated protein 1B (MAP1B) which is abundantly expressed in actively spreading neurons (29, 30). Neurites were visualized with phalloidin which labels F-actin and nuclei were stained with ATF-2. Our results showed that PORCN inhibition significantly decreased both the number of neurites/neurons (Fig. 1A; see representative micrographs, $**p < 0.01$) and the length of the neurites (Fig. 1B, $**p < 0.01$). Interestingly, a large amount of terminal filopodia/lamellipodia was observed with Wnt-C59 treatment (Fig. 1A, see white arrow).

Inhibition of porcupine by Wnt-C59 results in loss of contact with other cells in young cultures of hippocampal neurons

Another phenomenon that was verified in our experiments is the loss of contacts with other types of cells present in primary cultures. This phenomenon was evaluated by counting the number of neuronal processes in contact with another type of nonneuronal cell. The number of contacts with another cell type decreases by half when the neuronal cultures are treated with the Wnt-C59 inhibitor (Fig. 2A, see representative micrographs and lower graph, $**p < 0.01$). The treatment of hippocampal neuron cultures with the inhibitor also results in a significant decrease in β -catenin, a key protein in the Wnt signaling pathway, with an increase in the inactive form of GSK-3 β kinase (phosphorylated in serine 9). Both results indicated a reduced Wnt signaling activity in the presence of the PORNC inhibitor (27). The Wnt3a ligand also decreased after Wnt-C59 treatment (Fig. 2B, see representative blot and quantification of different proteins, $*p < 0.05$).

Inhibition of porcupine by Wnt-C59 treatment does not change axonal polarity in young hippocampal neurons

The polarized sorting and trafficking of newly synthesized proteins to the somato-dendritic and axonal domains of neurons occurs by selective incorporation into distinct populations of vesicular transport carriers. Voltage-gated Na^+ and K^+ channels are crucial for the efficient initiation and saltatory propagation of action potentials along myelinated axons of vertebrates, are clustered at the axon initial segment (AIS) and nodes of Ranvier. The clustering of these channels is mediated by ankyrin-G (AnkG) (31). Previous studies concluded that AIS, an actin-based filter, selectively prevents the passage of somato-dendritic vesicles into the axon (32). Conventional kinesin-1 is a major anterograde motor that operates in axons and consists of a heavy-chain (KIF5A, KIF5B, or KIF5C) dimer and two light chains (KLC) binding to the C termini of the dimer (33). One of the questions that arises from our experiments is whether the inhibition of PORCN, with the consequent lack of ligands and activation of the Wnt pathway, might alter neuronal polarity in some way. For this purpose, young neurons were treated with the inhibitor Wnt-C59 and two proteins of the AIS were studied, AnkG and KIF5 (32, 34). Our results showed that both proteins were correctly located in the neuronal axon of the neuron (Fig. 3). Young neurons were treated with different concentrations of Wnt-C59 (0.1, 1, 10 nM) and stained for AnkG and Piccolo, a protein present in the Piccolo-Bassoon transport vesicle (PTV), which are dense particles that form the active zone of the neurons (15). Our experiments show that both proteins are correctly located, together with actin activity in the same place (Fig. 3A, see representative micrograph, white arrow shows AIS). Young hippocampal neurons transfected with KIF5 and treated with 10 nM Wnt-C59, showed a similar localization to control cells together with specific proteins of the neuronal axon (Fig. 3B, see representative micrographs). We conclude that inhibition of porcupine with Wnt-C59 did not affect the neuronal polarity of the axon.

Different exogenous Wnt ligands recover the morphology of young hippocampal neurons in culture

To determine the specific role of specific Wnt ligands, young primary hippocampal neurons were treated with Wnt-C59 to shut down Wnt signaling activity. We evaluated the individual activity of Wnt ligands with canonical activity, such as Wnt 3a and 7a, and a known non canonical Wnt5a ligand.

A. Treatment with exogenous canonical Wnt3a ligand on neuronal morphology

Young hippocampal neurons (DIV 4), were treated with 10 nM Wnt-C59 from DIV 2 to DIV 4. These cultures were treated with different concentrations of the Wnt3a ligand for 24 h (50, 100, 200 ng/ml) and stained with MAP-1B (red), ATF-2 a nuclear transcription factor (blue) and phalloidin to observe actin (green), which can be seen at the tips of neuronal processes and to a lesser extent in neurons treated with Wnt-C59 (see white arrows). Young neurons treated with Wnt3a showed a recovery of the length of

neurites similar to the control (Fig. 4A, see representative micrographs). Another parameter that was recovered was the dendritic arbor complexity, concomitant with the recovery of secondary and tertiary projections (Fig. 4B, see graphs).

B. Treatment with exogenous noncanonical Wnt5a ligand on neuronal morphology

Sister young hippocampal neurons (DIV 4), were treated with 10 nM Wnt-C59 from DIV 2 to DIV 4. These cultures were treated with different concentrations of Wnt5a for 24 h (50, 100, 200 ng/ml) and stained with Tau (blue) and phalloidin to visualize actin (green), which can be observed at the tips of neuronal processes and to a lesser extent in neurons treated with Wnt-C59 (Fig. 5; representative micrographs). Our results showed that young neurons treated with different concentrations of Wnt5a showed a recovery of the length of neurites to a longer extension than control neurons, by 40–50% of the control depending on Wnt5a concentration (Fig. 5A, see representative micrographs) F-actin activity was marked with phalloidin (see white arrow). The complexity of the dendritic arbor was also recovered, together with the secondary and tertiary projections (Fig. 5B, see graph).

C. Effect of treatment with exogenous canonical Wnt7a ligand on neuronal morphology

Young hippocampal neurons (DIV 4), were treated with 10 nM Wnt-C59 from DIV 2 to DIV 4. These cultures were treated with different concentrations of the Wnt7a ligand for 24 h (50, 100, 200 ng/ml) and stained with MAP-1B (red), ATF-2 a nuclear transcription factor (blue) and phalloidin to visualize actin (green), which can be observed at the tips of neuronal processes (see white arrow). The young neurons treated with Wnt7a ligand showed a recovery of the length of neurites to a much longer extension than the control, but at a much lower concentration of Wnt7a than of Wnt5a (Fig. 6A, see representative micrographs; B, left graph). According to our results, 24-hour treatment of young neurons restores dendritic tree complexity, to an extent similar to that observed with the control (Fig. 6B, see right graph).

Discussion

Wnt signaling acts at various stages during neural development, including neuronal migration, axon and dendrite development, synapse formation/maturation, and neuronal plasticity (2, 6, 35). Wnt pathways are classified into canonical Wnt/ β -catenin or noncanonical (β -catenin-independent) pathways. Canonical Wnt/ β -catenin signaling is mediated by nuclear translocation of its central effector β -catenin. Noncanonical Wnt signaling occurs independently of β -catenin and is stimulated by Wnt ligands that bind to a receptor complex of Fzd, Ror1/2 or Ryk (1, 13, 36).

A series of small molecule inhibitors specifically target several of the proteins in the Wnt signaling pathway, such as Fzd, DVL, PORCN or Tankyrase, and could also be used as chemical probes to dissect the mechanism(s) of these signaling pathways (37, 38). We therefore used Wnt-C59, a PORCN inhibitor, in

young hippocampal neurons, to research polarization, the development of dendritic trees, and maintenance and/or maturation of dendritic spines; these inhibitory molecules have the potential to be developed both as study reagents in cell biology as therapeutic agents (39, 40). Our present results show that the early inhibition of PORCN in young hippocampal neurons, starting at picomolar Wnt-C59 concentration, drastically reduces the length of the neurites by up to 80%, and affects the dendritic arbor complexity and the signaling and contact that could be established between the different cells in culture. Nonneuronal cells are known to show no contact inhibition when in contact with the growth cones or neurites of neurons. These observations on the contact behavior of nonneuronal cells imply a series of signals from both types of cells, which in our experiments are lost with the inhibition of the Wnt pathway (41). Together, these results, show that the endogenous production of the different Wnt ligands plays an early and important role in the growth and architecture of neurites and the signaling between the different cells of the *in vitro* neuronal culture.

To evaluate the individual activity of Wnt ligands, sister cultures of hippocampal neurons previously treated with Wnt-C59 to shut down Wnt signaling activity were incubated with exogenous Wnts. First, we tested Wnt3a, a ligand that has canonical activity but has also shown some Ca^{2+} modulation (42, 43). Then, tested a noncanonical ligand Wnt5a which stimulates PSD-95 clustering (44) and finally, Wnt7a, a classic ligand of the Wnt- β -catenin-dependent pathway, which promotes the presynaptic colocalization of several neural receptors (35, 45). Our results indicated that all exogenous Wnt ligands (Wnt3a, 5a and 7a) rescued the changes in neuronal morphology. Specifically, Wnt3a restores the length of neurites to a value similar to that of the control; however, Wnt7a increased the neurite length beyond that of the control. Wnt5a had the same effect but at higher relative Wnt concentration. In addition, all 3 Wnts ligands restored dendritic arbor complexity with the recovery of secondary and tertiary projections.

Previous studies, have indicated that specific Wnt ligands induce neurite extension in various neuronal types: Wnt3 and Wnt3a in cultures of neural precursors of spinal cord cells (46), Wnt7a in cultured mouse cerebellar granule cells (47), Wnt5a and Wnt3b in chick retinal ganglion cell axons (48) and Wnt7b in mouse hippocampal neurons (49). More interestingly, different Wnt ligands have distinct effects on growing axons, and canonical and noncanonical pathways can have opposite effects (2). Adding to the complexity, the canonical ligand Wnt3a can activate both canonical and noncanonical signaling pathways in the same cell type (13, 50). Although the interaction between canonical and noncanonical Wnt signaling is complex, these signaling pathways generally inhibit each, as demonstrated during development (51).

Recent studies in our laboratory with Wnt-C59, a specific PORCN inhibitor in SH-SY5Y cells, indicated that both the canonical Wnt3a ligand and a noncanonical Wnt5a ligand increased the basal expression of Teneurin (52), a new trans-synaptic signal in both the peripheral nervous system and the CNS (53, 54). This work not only demonstrate the involvement of Wnt signaling in regulating Ten-3 expression but also revealed that Wnt3a (a canonical Wnt ligand) increases the expression of Ten-3 through a mechanism dependent on the secretion and activity of Wnt5a (a noncanonical ligand). Although the work raises new questions, the results seem to demonstrate the upregulation of Ten-3 by Wnt signaling and suggest that

Ten-3 modulation is possible because of crosstalk between the canonical and noncanonical Wnt pathways (52).

An alternative derived from our results is the possibility of modulating the neuronal morphology in the neurogenic niche and in a determined time frame, after PORCN inhibition by Wnt-C59, using the "replacement" of specific Wnt ligands, and thereby avoiding other secreted endogenous Wnt inhibitors, such as Dickkopf-1 (Dkk-1) or the secreted-frizzled-related protein (sFRP), which would allow the formation of new connections and therefore new neural networks (39, 55, 56). An interesting example is Wnt signaling inhibition by Wnt-C59 which prevents the activity of Wnt3a and Wnt8B, in canonical and noncanonical Wnt-mediated signaling to induce cortical motor neurons or their progenitors from iPSCs in humans (26, 57). This strategy might offer a proof of concept on the preclinical side, where the inhibition of Wnts signaling in mammals can be achieved by inhibiting PORCN with small molecule inhibitors such as Wnt-C59, providing a safe and feasible strategy *in vivo*. Our results increase the possibility that a specific PORCN inhibitor contributes to regulating the Wnt pathway so that it become a safe and plausible methodology for cell replacement therapy in neurological diseases.

Conclusions

In summary, we use a Wnt-C59 a specific PORCN inhibitor, to block secretion of endogenous Wnts in young hippocampal neurons. Under these conditions, the morphology and length of the neurites and the complexity of the dendritic tree were evaluated. Our results indicated shows that Wnt3a, Wnt7a and Wnt5a ligands restored dendritic arbor complexity with recovery of secondary and tertiary projections in hippocampal neurons. We proposed that PORCN in Wnt signaling is an emerging molecular target of interest in the search for preclinical options to study and treat neurological diseases.

Abbreviations

AIS: Axon initial segment

ANOVA: Analysis of variance

ATF2: Activating transcription factor 2

CNS: Central nervous system

Dkk-1: Dickkopf-1

DIV: days *in vitro*

DVL: Dishevelled

ECL: Enhanced chemiluminescence

EDTA: Ethylenediaminetetraacetic acid

eGFP: enhanced Green Fluorescent Protein

Fzd: Frizzled

GSK-3b: Glycogen synthase kinase-3 b

MAP1B: Microtubule-associated protein 1B

NaF: Sodium fluoride

PSD-95: Postsynaptic density protein 95

PTV: Piccolo-Bassoon transport vesicle

PVDF: Polyvinylidene difluoride

PORCN: Porcupine

sFRP: Secreted-frizzled-related protein

SDS-PAGE: Sodium dodecyl sulfate polyacrylamide gel electrophoresis

SEM: Standard error of the mean

RIPA: Radioimmunoprecipitation assay buffer

Wnt: Wingless

Declarations

Ethical Approval and Consent to participate

Rats were housed at the Animal Warehouse Facility of P. Catholic University of Chile. The experimental procedure was approved by the Bioethical and Biosafety Committee of the Faculty of Biological Sciences of our university.

Consent for publication

The experiments were carried out in the Department of Cellular and Molecular Biology of the P. Catholic University of Chile, Santiago of Chile.

Availability of supporting data

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Basal Center of Excellence of Aging and Regeneration CONICYT-ANID (AFB 170005) and a special grant “Lithium in Health and Disease” from the Sociedad Química y Minera de Chile (SQM).

Authors' contributions

JAG and NIC designed the study, together with JE-C performed the experiments and analyzed the data. NCI obtained resources. JAG and NIC wrote the paper. All authors read and approved the final manuscript.

Acknowledgements

Not applicable

References

1. Nusse R, Clevers H. Wnt/ β -Catenin Signaling, Disease, and Emerging Therapeutic Cell 2017; 169:985-999.
2. Oliva CA, Montecinos-Oliva C, Inestrosa NC. Wnt Signaling in the Central Nervous System: New Insights in Health and Disease. Prog Mol Biol Transl Sci. 2018; 153:81-130.
3. Habib SJ, Chen BC, Tsai FC, Anastassiadis K, Meyer T, Betzig E, Nusse R. A localized Wnt signal orients asymmetric stem cell division in vitro. Science 2013; 339:1445-8.
4. Loh KM, van Amerongen R, Nusse R. Generating Cellular Diversity and Spatial Form: Wnt Signaling and the Evolution of Multicellular Animals. Dev Cell 2016; 38: 643-55.
5. Cerpa W, Ramos-Fernandez E, Inestrosa NC. Modulation of the NMDA Receptor Through Secreted Soluble Factor. Mol. Neurobiol. 2016; 53: 299-309.
6. Inestrosa NC, Arenas E. Emerging roles of Wnts in the adult nervous system. Nature Rev. Neurosci. 2010; 11: 77–86.
7. Briona LK, Poulain FE, Mosimann C, Dorsky RI. Wnt/ β -catenin signaling is required for radial glial neurogenesis following spinal cord injury. Dev Biol. 2015; 403:15-21.
8. Guo F, Lang J, Sohn J, Hammond E, Chang M, Pleasure D. Canonical Wnt signaling in the oligodendroglial lineage—puzzles remain. Glia 2015; 63:1671-93.

9. Lambert C, Cisternas P, Inestrosa NC. Role of Wnt signaling in central nervous system injury. *Mol Neurobiol.* 2016; 53:2297–2311.
10. Varela-Nallar L, Alfaro IE, Serrano FG, Parodi J, Inestrosa NC. Wingless-type family member 5A (Wnt-5a) stimulates synaptic differentiation and function of glutamatergic synapses. *Proc Natl Acad Sci U S A.* 2010; 107: 21164-9.
11. Oliva, C.A., Inestrosa, N.C. A novel function for Wnt signaling modulating neuronal firing activity and the temporal structure of spontaneous oscillation in the entorhinal-hippocampal circuit. *Exp. Neurol.* 2015; 269: 43-55.
12. Cheng J, Shen W, Jin L, Pan J, Zhou Y, Pan G, Xie Q, Hu Q, Wu S, Zhang H, Chen X. Treadmill exercise promotes neurogenesis and myelin repair via upregulating Wnt/ β catenin signaling pathways in the juvenile brain following focal cerebral ischemia/reperfusion. *Int J Mol Med.* 2020; 45:1447-63.
13. Gordon MD, Nusse R. Wnt Signaling: Multiple Pathways, Multiple Receptors, and Multiple Transcription Factors. *J Biol Chem.* 2006; 281: 22429–33.
14. Lai KKY, Nguyen C, Lee KS, Lee A, Lin DP, Teo JL, Kahn M. Convergence of Canonical and Non-Canonical Wnt Signal: Differential Kat3 Coactivator Usage. *Curr Mol Pharmacol.* 2019; 12:167-183.
15. Torres VI, Inestrosa NC. Vertebrate Presynaptic Active Zone Assembly: a Role Accomplished by Diverse Molecular and Cellular Mechanisms. *Mol Neurobiol.* 2018; 55: 4513-28.
16. Zuber B, Lučić V. Molecular architecture of the presynaptic terminal. *Curr Opin Struct Biol.* 2019; 54:129-138.
17. Südhof TC. Towards an Understanding of Synapse Formation. *Neuron.* 2018; 100:276-293.
18. Xiao S, McKeever PM, Lau A, Robertson J. Synaptic localization of C9orf72 regulates post-synaptic glutamate receptor 1 levels. *Acta Neuropathol Commun.* 2019; 7:161.
19. Banker GA, Cowan WM. Rat hippocampal neurons in dispersed cell culture. *Brain Res* 1977; 126:397-442.
20. Kaech S, Banker G. Culturing hippocampal neurons. *Nat Protoc.* 2006; 1: 2406-15.
21. Dotti CG, Sullivan CA, Banker GA. The establishment of polarity by hippocampal neurons in culture. *J Neurosci.* 1988; 8:1454-68.
22. Craig AM, Wyborski RJ, Banker G. Preferential addition of newly synthesized membrane protein at axonal growth cones. *Nature.* 1995; 375:592-4.
23. Motono M, Ioroi Y, Ogura T, Takahashi J. WNT-C59, a Small-Molecule WNT Inhibitor, Efficiently Induces Anterior Cortex That Includes Cortical Motor Neurons From Human Pluripotent Stem Cells.

Stem Cells Transl Med. 2016; 5:552-60.

24. Hofmann, K. A superfamily of membrane-bound O-acyltransferases with implications for Wnt signaling. *Trends in Biochem. Sci.*, 2000; 25:111–112.
25. Takada, R., Satomi, Y., Kurata, T., Ueno, N., Norioka, S., Kondoh, H., Takao, T., and Takada, S. Monounsaturated fatty acid modification of Wnt protein: its role in Wnt secretion. *Dev. Cell* 2006; 11: 791–801.
26. Clements, S.E. Importance of PORCN and Wnt signaling pathways in embryogenesis. *Amer. J. Med. Gen.* 2009; 149A: 2050-2051.
27. Torres VI, Godoy JA, Inestrosa NC. Modulating Wnt signaling at the root: Porcupine and Wnt acylation. *Pharmacol Ther.* 2019; 198: 34-45.
28. Varella-Nallar L, Grabowski CP, Alfaro IE, Alvarez AR, Inestrosa NC. Role of the Wnt receptor Frizzled-1 in presynaptic differentiation and function. *Neural Dev.* 2009; 4: 41.
29. Calvert R, Anderton BH. A Microtubule-Associated Protein (MAP1) Which Is Expressed at Elevated Levels During Development of the Rat Cerebellum. *EMBO J.* 1985; 4:1171-6.
30. Calvert R, Anderton BH. A Microtubule-Associated Protein (MAP1) Which Is Expressed at Elevated Levels During Development of the Rat Cerebellum. *EMBO J.*; 1985; 4:1171-6.
31. Barry J, Gu Y, Jukkola P, O'Neill B, Gu H, Mohler PJ, Rajamani KT, Gu C. Ankyrin-G directly binds to kinesin-1 to transport voltage-gated Na⁺ channels into axons. *Dev Cell* 2014; 28:117-31.
32. Farías GG, Guardia CM, Britt DJ, Guo X, Bonifacino JS. Sorting of Dendritic and Axonal Vesicles at the Pre-axonal Exclusion Zone. *Cell Rep.* 2015; 13:1221-1232.
33. Hirokawa N, Niwa S, Tanaka Y. Molecular motors in neurons: transport mechanisms and roles in brain function, development, and disease. *Neuron* 2010; 68:610-38.
34. Norris SR, Soppina V, Dizaji AS, Schimert KI, Sept D, Cai D, Sivaramakrishnan S, Verhey KJ. A method for multiprotein assembly in cells reveals independent action of kinesins in complex. *J Cell Biol.* 2014; 207:393-406.
35. Ciani L, Salinas PC. WNTs in the vertebrate nervous system: from patterning to neuronal connectivity. *Nat Rev Neurosci.* 2005; 6:351-62.
36. Gómez-Orte E, Sáenz-Narciso B, Moreno S, Cabello J. Multiple functions of the noncanonical Wnt pathway. *Trends Genet.* 2013; 29:545-53.
37. Tapia-Rojas C, Burgos PV, Inestrosa NC. Inhibition of Wnt signaling induces amyloidogenic processing of amyloid precursor protein and the production and aggregation of Amyloid- β (Ab)₄₂

- peptides. J Neurochem. 2016; 139:1175-91.
38. Yamaguchi K, Nagatoishi S, Tsumoto K, Furukawa Y. Discovery of chemical probes that suppress Wnt/ β -catenin signaling through high-throughput screening. Cancer Sci. 2020; 111:783-794.
 39. Voronkov A and Krauss S. Wnt/b-Catenin Signaling and Small Molecule Inhibitors. Curr. Pharmacol. Design. 2013; 19: 634-664.
 40. Tran FH, Zheng JJ. Modulating the Wnt signaling pathway with small molecules. Protein Sci. 2017; 26:650-661.
 41. Wessells NK, Letourneau PC, Nuttall RP, Ludueña-Anderson M, Geiduschek JM. Responses to cell contacts between growth cones, neurites and ganglionic non-neuronal cells. J Neurocytol. 1980; 9:647-64.
 42. Cerpa W, Godoy JA, Alfaro I, Farías GG, Metcalfe MJ, Fuentealba R, Bonansco C, Inestrosa NC. [Wnt-7a modulates the synaptic vesicle cycle and synaptic transmission in hippocampal neurons](#). J Biol Chem. 2008; 283:5918-27.
 43. Avila ME, Sepúlveda FJ, Burgos CF, Moraga-Cid G, Parodi J, Moon RT, Aguayo LG, Opazo C, De Ferrari GV. Canonical Wnt3a modulates intracellular calcium and enhances excitatory neurotransmission in hippocampal neurons. J Biol Chem. 2010; 285:18939-47.
 44. Farías GG, Alfaro IE, Cerpa W, Grabowski CP, Godoy JA, Bonansco C, Inestrosa NC. Wnt-5a/JNK signaling promotes the clustering of PSD-95 in hippocampal neurons. J Biol Chem. 2009; 284:15857-66.
 45. Palomer E, Buechler J, Salinas PC. Wnt Signaling Deregulation in the Aging and Alzheimer's Brain. Front Cell Neurosci 2019; 13:227.
 46. David MD, Cantí C, Herreros J. Wnt-3a and Wnt-3 differently stimulate proliferation and neurogenesis of spinal neural precursors and promote neurite outgrowth by canonical signaling. J Neurosci Res. 2010; 88:3011-23.
 47. Hall AC, Lucas FR, Salinas PC. Axonal remodeling and synaptic differentiation in the cerebellum is regulated by WNT-7a signaling. Cell 2000; 100:525-35.
 48. Rodriguez J, Esteve P, Weinl C, Ruiz JM, Fermin Y, Trousse F, Dwivedy A, Holt C, Bovolenta P. SFRP1 regulates the growth of retinal ganglion cell axons through the Fz2 receptor. Nat Neurosci. 2005; 8:1301-9.
 49. Rosso SB, Sussman D, Wynshaw-Boris A, Salinas PC. Wnt signaling through Dishevelled, Rac and JNK regulates dendritic development. Nat Neurosci. 2005; 8: 34-42.

50. Nalesso G, Sherwood J, Bertrand J, Pap T, Ramachandran M, De Bari C, Pitzalis C, Dell'accio F. WNT-3A modulates articular chondrocyte phenotype by activating both canonical and noncanonical pathways. *J Cell Biol.* 2011; 193:551–564.
51. Topol L, Jiang X, Choi H, Garrett-Beal L, Carolan PJ, Yang Y. Wnt-5a inhibits the canonical Wnt pathway by promoting GSK-3-independent b-catenin degradation. *J Cell Biol* 2003; 162:899-908.
52. Bastías-Candia S, Martínez M, Zolezzi JM, Inestrosa NC. Wnt Signaling Upregulates Teneurin-3 Expression via Canonical and Noncanonical Wnt Pathway Crosstalk. *Front Neurosci.* 2019; 13:505.
53. Bastías-Candia S, Braidy N, Zolezzi JM, Inestrosa NC. Teneurins and Alzheimer's disease: a suggestive role for a unique family of proteins. *Med Hypotheses* 2015; 84:402-7.
54. Mosca TJ. On the Teneurin track: a new synaptic organization molecule emerges. *Front. Cell. Neurosci.* 2015; 9:204.
55. Yan M, Li G, Jing An J. Discovery of Small Molecule Inhibitors of the Wnt/ β -catenin Signaling Pathway by Targeting β -catenin/Tcf4 Interactions. *Exp Biol Med* 2017; 242:1185-1197.
56. Shin SH, Lim DY, Reddy K, Malakhova M, Liu F, Wang T, Song M, Chen H, Bae KB, Ryu J, Liu K, Lee MH, Bode AM, Dong Z. A Small Molecule Inhibitor of the β -Catenin-TCF4 Interaction Suppresses Colorectal Cancer Growth *In Vitro* and *In Vivo*. *EBioMedicine* 2017; 25:22-31.
57. Munji RN, Choe Y, Li G, Siegenthaler JA, Pleasure SJ. Wnt signaling regulates neuronal differentiation of cortical intermediate progenitors. *J Neurosci.* 2011, 31:1676-87.

Figures

Figure 1

A Porcupine inhibitor, Wnt-C59 induces morphological change in dendritic arbor. Hippocampal neurons were cultured until 4DIV and treated with different concentration of Wnt-C59 (0.01, 0.1, 1, 10, 100 nM) for 36 h, and stained with MAP1B (red), ATF-2 (blue) and F-actin marker phalloidin (green). Neurons were imaged using a confocal microscope to evaluate neuronal morphology. A. Representative Micrographs are showed, Wnt-C59 produces a decrease of the dendritic arbor in these young neurons; B. The quantification of these processes is shows as representative data from young cultured neurons. The results are presented as the mean $\pm$ SEM for n: 10-12 experiments), $**p < 0.01$. Bonferroni's test. Bar represents 10 μm .

Figure 2

A Porcupine inhibitor, Wnt-C59, produces a loss of contacts with other types of cells in culture. A. Sister cultures of hippocampal neurons were used to quantify the contact of neurons with other cell types, without specifying the cell type; these contacts of neuronal processes decrease by half, depending also on the concentration of the inhibitor. See representative micrographs of control neurons and treated with 10 nM Wnt-C59 for 24 h. and zoom. B. The figure shows a representative blot of some proteins of the Wnt

pathway, under the inhibitory activity of Wnt-C59 in hippocampal neurons. Graph shows quantification of these contacts. The results are presented as the mean $\pm$ SEM for n= 10 experiments), *p<0.05, **p<0.01. Bonferroni's test. Bar represents 10 μ m.

Figure 3

Porcupine inhibition by Wnt-C59 treatments does not change the axon polarity in young hippocampal neurons. Young neurons (DIV 4) were treated with different concentration of Wnt-C59 (0.1, 1, 10 nM) for

24 h. A. AnkG protein (red), Piccolo a protein present in the Piccolo-Bassoon transport vesicle (PTV) (blue) and actin (green) were stained to locate AIS (see white arrow). B. Sister cultured were transfected with KIF 5-Cherry and with MAP1b and Tau protein, to locate AIS in axonal processes (see zoom of label region in axonal region). Bar represents 10 μ m.

Figure 4

Exogenous Wnt3a ligand treatment recovery dendritic arbor after inhibition of PORCN. Sister cultures of young hippocampal neurons were used to quantify the recovery of dendritic arbor after treatment with different concentrations of the canonical Wnt3a ligand. A. See representative micrographs of control

neurons and treated with Wnt3a (50ng/ml), 10 nM Wnt-C59 or plus Wnt3a, (50, 100, 200 ng/ml) for 24 h, White arrowhead show actin on the tip of different neurites. B. Graph shows quantification of average of neurites length (see left graph) and recovery of dendritic complexity (see right graph). The results are presented as the mean $\pm$ SEM for n= 5 experiments, **p<0.01, ***p<0.001. Bonferroni's test. Bar represents 10 μ m.

Figure 5

Exogenous Wnt5a ligand treatment recovers dendritic arbor after inhibition of PORCN. Sister cultures of young hippocampal neurons were used to quantify the recovery of dendritic arbor, after treatment with

different concentrations of the non-canonical Wnt5a ligand. A. See representative micrographs of control neurons and treated with Wnt5a (50ng/ml), 10 nM Wnt-C59 or plus Wnt5a, (50, 100, 200 ng/ml) for 24 h, White arrowhead show actin on the tip of different neurites. B. Graph shows quantification of average of neurite length (see left graph) and recovery of dendritic complexity (see right graph). The results are presented as the mean $\pm$ SEM for n= 5 experiments, **p<0.01. Bonferroni's test. Bar represents 10 μ m.

Figure 6

Exogenous Wnt7a ligand treatment recovery dendritic arbor after inhibition of PORCN. Sister cultures of young hippocampal neurons were used to quantify the recovery of dendritic arbor after treatment with

different concentrations of the canonical Wnt7a ligand, and stained with MAP1B (red), ATF-2 (blue) and phalloidin (green). Neurons were imaged using a confocal microscope to evaluate neuronal morphology. A. See representative micrographs of control neurons and treated with Wnt7a (50ng/ml), 10 nM Wnt-C59 or plus Wnt7a (50, 100, 200 ng/ml) for 24 h, White arrowhead show actin on the tip of different neurites. B. Graph shows quantification of average of neurites length (see left graph) and recovery of dendritic complexity (see right graph). The results are presented as the mean $\pm$ SEM for n= 5 experiments, **p<0.01, ***p<0.001. Bonferroni's test. Bar represents 10 μ m.