

Triglyceride-glucose index linked to hospital mortality in critically ill stroke patients: an observational multicentre study on eICU database

Bingjun Zhang

Third Affiliated Hospital of Sun Yat-Sen University

Lingling Liu

Third Affiliated Hospital of Sun Yat-Sen University

Hengfang Ruan

Third Affiliated Hospital of Sun Yat-Sen University

Qiang Zhu

Third Affiliated Hospital of Sun Yat-Sen University

Dafan Yu

Third Affiliated Hospital of Sun Yat-Sen University

Yu Yang

Third Affiliated Hospital of Sun Yat-Sen University

Zhengqi Lu (✉ lzq1828@outlook.com)

Third Affiliated Hospital of Sun Yat-Sen University

Original investigation

Keywords: Triglyceride-glucose index, Insulin resistance, Stroke, Mortality, ICU

Posted Date: July 28th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-42390/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: The triglyceride-glucose (TyG) index is a reliable surrogate of insulin resistance and a marker for ischemic stroke (IS) incident. Whether the TyG index predicts stroke outcome remains uncertain. This study investigated the prognostic value of the TyG index in critically ill stroke patients.

Methods: This was a retrospective observational study that included stroke patients, and all data were extracted from the eICU Collaborative Research Database. The TyG index was calculated as the $\ln(\text{fasting glucose level [mg/dL]} \times \text{triglyceride level [mg/dL]}/2)$. The outcomes included the hospital and intensive care unit (ICU) death. Multivariate logistic regression was used to determine independent risk factors. The smoothing curves and forest plots were illustrated.

Results: A total of 4570 eligible subjects were enrolled. The mean level of TyG index was 9.1 ± 0.7 . The hospital and ICU mortality rate were 10.3% and 5.0% respectively. TyG index as a continuous variable was associated hospital mortality in univariate analysis (OR 1.723, 95% CI 1.524-1.948, $P < 0.001$), adjusted model 1 (OR 1.861, 95% CI 1.637-2.116, $P < 0.001$) and adjusted model 2 (OR 2.543, 95% CI 1.588-4.073, $P < 0.001$). TyG was also associated ICU mortality in univariate analysis (OR 2.146, 95% CI 1.826-2.523, $P < 0.001$), adjusted model 1 (OR 2.183, 95% CI 1.847-2.580, $P < 0.001$), and adjusted model 2 (OR 2.672, 95% CI 1.376-5.188, $P < 0.001$). The smoothing curves observed a continuous linear association after adjusting all covariates both in hospital and ICU mortality. Subgroup analysis demonstrated TyG index was associated with increased risk of hospital and ICU death in critically ill IS ($P < 0.05$), but not in hemorrhage stroke ($P > 0.05$).

Conclusion: The TyG index is a potential predictor for hospital and ICU mortality in critically ill stroke patients, especially in IS patients.

Background

Stroke is a leading cause of mortality and disability worldwide. According to global burden of disease (GBD) 2016 stroke surveys, it was estimated that there were 5.5 million deaths due to stroke, and the global lifetime risk of stroke from the age of 25 years onward was approximately 25%. [1, 2] The economic costs of treatment and post-stroke care are substantial, while the global burden of stroke has been increasing. Furthermore, an increasing number of stroke patients, who have higher hospital mortality, are being admitted to an intensive care unit (ICU) for management of severe neurological impairment and post-stroke complications. [3, 4] Therefore, it is still important to identify the controllable risk factors targeted by preventive strategies and health-care management that lead to a larger decrease in stroke mortality.

The triglyceride-glucose (TyG) index combines both levels of fasting plasma glucose and triglyceride, and it has been reported to be significantly correlated with insulin resistance (IR) and to be a reliable surrogate marker of IR. [5] Previous studies have indicated that the TyG index is associated with cardiovascular disease morbidity and mortality in the general population and different types of patient cohorts. [6-9]

Lately, the TyG index has been proposed as a direct marker for the risk of incident ischemic stroke (IS) in general population.[10] To date, no relevant study has focused on the impact of the TyG index on mortality in patients with critically ill stroke including IS and hemorrhage stroke (HS). Therefore, this study aimed to specifically investigate the association between TyG index and mortality of stroke in a large observational multicentre study on eICU database.

Methods

Data source

Data were collected from the eICU Collaborative Research Database v2.0. The database was a multi-center ICU database for over 200,000 admissions in 2014 and 2015 at 208 United States hospitals.[11] The eICU database included hourly physiological readings from bedside monitors, records of demographic characteristics, severity of illness measures, diagnoses, treatment, and other clinical data collected during routine medical care. The use of this database has been approved by the institutional review boards of Massachusetts Institute of Technology (Cambridge, MA, USA). One author (Bingjun Zhang) obtained the access and was responsible for the data extraction (certification number: 36492956).

Participants

Patients with a primary diagnosis of brain stroke, recorded on patient dataset, were potentially eligible. The stroke patients can be grouped into 4 categories: IS group, HS group, IS with HS group, and unknown/others group.

Demographic and laboratory variables

The following data were extracted from the eICU database: gender, age, ethnicity, admission height, admission weight, stroke types, plasma glucose values, blood lipids values, acute physiology and chronic health evaluation (APACHE) IV score, the status of discharge hospital or ICU, in-hospital length of stay (LOS), ICU LOS, comorbidities, mechanical ventilation. APACHE IV score was calculated within the first 24 hours after the ICU admission.

If glucose, lipids, and some variables were measured for several times after the ICU entry, data of the first time were used. The TyG index was calculated as the $\ln(\text{fasting glucose level [mg/dL]} \times \text{triglyceride level [mg/dL]}/2)$.

The primary outcome was hospital mortality, with ICU mortality as a secondary outcome.

Variables with more than 10% missing values were excluded from the analysis. Multiple imputation was performed for variables with missing values of less than 10%.

Statistical analyses

Categorical variables were expressed in absolute number with percentage and analyzed by chi-square or Fisher exact test. Continuous variables were first assessed for normality. Normal data were expressed in mean with standard deviation (SD) and compared using Student T test or one-way ANOVA. Non-normal data were expressed in median with interquartile range (IQR) and were compared using Wilcoxon rank-sum test. Variables with two-tailed p value less than 0.05 were considered to be statistically significant and were included in the regression model. Multivariate-adjusted odds ratios (OR) and 95% confidence intervals (CI) for the study outcomes and TyG index (1 unit and quartile) were calculated by logistic regression analysis. The multivariate model 1 included gender, age, ethnicity, while model 2 included model 1 plus stroke types, coronary heart disease, diabetes mellitus, heart failure, plasma glucose, total cholesterol (TC), low-density lipoprotein-cholesterol (LDL-C), triglyceride, mechanical ventilation, APACHE IV score, LOS. All of the analyses were performed using the SPSS version 26 (Chicago, Illinois). All p values were two-tailed and a p value < 0.05 was considered statistically significant. The smoothing curves and forest plots were illustrated by EmpowerStats (X&Y Solutions, Inc., Boston, MA).

Results

The initial search identified 200,859 ICU admissions from the eICU database. A total of 6849 subjects with a primary diagnosis of brain stroke were identified. A total of 2279 subjects were excluded because they did not have mortality values. The final cohort included 4570 patients, including 4,101 (89.7%) survivors before hospital discharge and 4341 (95.0%) survivors before ICU discharge. The study patients had an average age of 66.3 ± 14.2 years and 2358 (51.6%) patients were male. The mean level of TyG index was 9.1 ± 0.7 . The hospital and ICU mortality rate were 10.3% and 5.0% respectively.

Baseline characteristics

The baseline characteristics between survivors and non-survivors groups are described in Table 1. Hospital non-survivor group had higher age, plasma glucose, triglyceride, ICU LOS than hospital survivor group ($P < 0.05$), while lower TC and LDL-C ($P < 0.05$). There was no significant difference on sex, ethnicity, height, weight, HDL-C, hypertension, atrial fibrillation, hospital LOS ($P > 0.05$). Furthermore, hospital non-survivor group had more frequent diabetes mellitus, coronary heart disease, heart failure, mechanical ventilation than hospital survivor group ($P < 0.05$). Stroke type was significantly different between the two groups ($P < 0.05$). Patients in the hospital non-survivor group had higher APACHE IV score (70.8 ± 25.0 vs 47.7 ± 19.0 , $P < 0.001$) and TyG index (9.3 ± 0.7 vs 9.0 ± 0.7 , $P < 0.001$) than those survivor group.

In addition, there was no significant difference on age and TC between ICU survivors and ICU non-survivors groups ($P > 0.05$). ICU non-survivor group had lower hospital LOS than ICU survivor group ($P < 0.001$). Patients in the ICU non-survivor group had higher APACHE IV score (70.8 ± 25.0 vs 47.7 ± 19.0 , $P < 0.001$) and TyG index (9.3 ± 0.7 vs 9.0 ± 0.7 , $P < 0.001$) than those survivor group. Other results in ICU groups were consistent with those in-hospital groups.

TyG index and mortality of critically ill stroke

Univariate and multivariate logistic regression revealed the association between TyG index and critically ill stroke mortality (Table 2). In univariate analysis, TyG index as a continuous variable was associated hospital mortality (OR 1.723, 95% CI 1.524-1.948, $P < 0.001$). Several risk factors including important clinical and significant variables in the univariate model were included in the multivariate model for adjustment, and the TyG index also remained to be an independent predictor of hospital mortality in adjusted model 1 (OR 1.861, 95% CI 1.637-2.116, $P < 0.001$) and model 2 (OR 2.543, 95% CI 1.588-4.073, $P < 0.001$). Furthermore, TyG index was also associated ICU mortality in univariate analysis (OR 2.146, 95% CI 1.826-2.523, $P < 0.001$), adjusted model 1 (OR 2.183, 95% CI 1.847-2.580, $P < 0.001$), and adjusted model 2 (OR 2.672, 95% CI 1.376-5.188, $P < 0.001$).

In addition, all participants were stratified into four groups based on the quartile of TyG index. The mean levels of TyG index were 8.3 ± 0.3 , 8.8 ± 0.1 , 9.2 ± 0.1 , and 10.0 ± 0.1 , respectively. The hospital and ICU mortality significantly increased with increasing quartiles of the TyG index ($P < 0.001$) (Fig. 1). When dividing TyG index into quartiles, we observed a quartile increment in TyG index was associated with the increased hospital and ICU mortality of critically ill stroke in univariate, adjusted model 1 and adjusted model 2 logistic regression analysis, with a significant trend across the quartiles (P for trend < 0.001 , Table 2).

To further explore the relationship between TyG index and the mortality of critically ill stroke, we plotted the smoothing curves of TyG index against the hospital and ICU mortality of critically ill stroke (Fig. 2). In this plot, we observed a continuous linear association after adjusting all covariates both in hospital (Fig. 2A) and ICU (Fig. 2B) mortality. This finding was consistent with the stepwise increased OR in the analysis of multivariate logistic regression.

Relationship of TyG index to mortality according to stroke types

The stroke patients were grouped into 4 types: IS group, HS group, IS with HS group, and unknown/others group. Multivariate logistic regression models were analyzed to identify the association between TyG index and mortality according to the stroke type and TyG index. We found that a higher TyG index was significantly associated with the increased hospital and ICU mortality in IS group and unknown/others group (Fig. 3 and Fig. 4, adjusted $P < 0.05$). However, the similar result did not occur in HS group and IS with HS group (Fig. 3 and Fig. 4, adjusted $P > 0.05$).

Discussion

To the best of our knowledge, this is the first study to evaluate the association of the TyG index with hospital and ICU mortality in critically ill stroke patients. The main findings are as follows: (1) the TyG index is an independent predictor for hospital and ICU mortality in patients with critically ill stroke; (2) the hospital and ICU mortality correlated proportionally with the increment of TyG index, implicating the linearity of TyG index as an indicator of critically ill stroke; (3) subgroup analysis demonstrated TyG index was associated with increased risk of hospital and ICU death in critically ill IS, but not in HS.

The TyG index, as the product of fasting plasma glucose and triglyceride, is a novel index that has been well recognized as a simple and reliable surrogate of IR.[12] The homeostatic model assessment of IR (HOMA-IR) has been traditionally used to estimate IR.[13] However, insulin levels must be required to calculate the HOMA-IR index. Compared with the inconvenient HOMA-IR, the TyG index does not require levels of insulin and may apply to all of the patients and healthy population. Recent studies indicated that the TyG index has been shown to be superior to HOMA-IR in predicting IR.[14] Furthermore, several studies conducted in Asia and Europe validated the strong association between TyG and incidence of diabetes mellitus.[12, 15] Won et al. reported the TyG index was independently associated with arterial stiffness in a relatively healthy Korean population.[16] Other studies demonstrated the TyG index was an independent predictor of coronary artery calcification progression and risk of cardiovascular diseases.[8, 17] In addition, previous studies suggested the TyG index predicted severity and outcomes in patients with acute coronary syndrome.[6, 18, 19] Wang et al. reported the TyG index predicted future cardiovascular events in patients with diabetes and acute coronary syndrome independently of known cardiovascular risk factors.[7] However, in a previous study based on a Caucasian population, the TyG index displayed an insignificant association with stroke.[20] Later, another study based on the same Caucasian cohort, identified metabolic health and obesity states groups based on TyG index was significantly associated with the risk of IS.[21] Additionally, a recently published epidemiological investigation expanded the use of TyG index as a direct marker for the risk of IS.[10] However, no relevant study focus on the association between TyG index and outcome of stroke. The present study investigated the relationship between the TyG index and the hospital mortality in critically ill stroke on eICU database. We found that the TyG index was significantly associated with mortality in ICU stroke after adjusting for confounding factors. Furthermore, the TyG index had a continuous linear correlation with the mortality of ICU stroke.

Although the mechanism underlying the relationship between the TyG index and stroke is not fully elucidated, the TyG index has been deemed as a useful atherogenic indicator linked to IR and metabolic syndrome. IR may be the mechanism in increasing mortality in stroke. Firstly, IR may increase proinflammatory cytokines and enhance prothrombotic responses, thus exacerbating damage in the brain after stroke.[22, 23] Secondly, IR may cause sympathetic activity and catabolism in muscles, thus enhancing muscle loss and leading to poor functional outcomes.[24] Thirdly, IR may increase platelet adhesion, enhance atherosclerosis progression, facilitate plaque instability, and therefore contribute to severity of stroke, via promoting apoptosis of vascular smooth muscle cells macrophages, and endothelial cells.[25, 26] Fourthly, recent evidence has proved that triglyceride and glucose disorder are the risk factors of stroke, while the formula of TyG index is composed of triglyceride and glucose.[27, 28] Lastly, IR may augment the role of modifiable risk factors of stroke, such as hypertension, atrial fibrillation.[10] Although the present study had showed the association between TyG index and poststroke outcomes, the underlying molecular mechanisms involved in this association should be further investigated in the future study.

Stroke accounts for almost 5% of all disability-adjusted life-years and 10% of all deaths worldwide.[2] From different epidemiological surveys, stroke mortality in different periods varies from 5%-72 %.[29, 30]

Stroke mortality has been declining since the early 20th century.[31] The reasons for this are not completely understood, although the decline is welcome.[31] However, due to the increasing complexity of stroke treatment and severe conditions, an increasing proportion of acute stroke patients are being admitted to an ICU.[32, 33] A study including 4958 consecutive stroke patients reported 347 (7.0%) patients required ICU admission at any time point during their index hospitalization.[4] In-hospital mortality of ICU stroke reported in the literature varies widely, highly depending on the patient characteristics. A previous small study found the mortality was 38.7% (43/111) in stroke patients requiring ICU admission.[32] An United States study including 448 ICU stroke patients provided hospital mortality was 30%.[34] Recently, a prospective observation trial reported the mortality was 7.5% in critically ill stroke patients.[35] These patients with high hospital mortality mainly associated with older age, poor neurological severity at admission, high APACHE score, impaired consciousness, intracranial hemorrhage and need for mechanical ventilation.[32, 33] Moreover, functional outcomes in survivors appear to be poor.[36] The current study based on eICU database, indicated the hospital and ICU mortality were 10.3% and 5.0%, respectively. Furthermore, we found a new risk factor, the TyG index, for the mortality of critically ill stroke.

In subgroup analysis, we demonstrated TyG index was associated with increased risk of hospital and ICU death in critically ill IS, but not in HS. In agreement with previous GBD 2016 report, we did not estimate the mortality due to subarachnoid hemorrhage and intracerebral hemorrhage separately.[1] Of the total number of prevalent strokes, over 80% were IS. However, the number of global deaths due to IS was slightly lower than the number due to HS deaths.[1] In contrast, a previous neurology ICU study demonstrated HS was more frequent than IS (71.9% vs 28.1%).[37] Several studies, including our present study, described TyG index was a useful marker in IS. However, to our knowledge, no relevant study has evaluated the relationship between the TyG index and HS. Although previous studies had focusing on the relationship between IR and HS, the results had been discrepancy. The Rotterdam study and the Uppsala study had previously examined the association between IR and risk of HS, finding virtually no evidence of an association.[38, 39] Later, a large United States stroke cohort showed IR may be a protective effect on HS.[40] In the current study, we did not observe a significant association of the TyG index with all-cause death in critically ill HS in either unadjusted or adjusted analysis.

Although our study based on a large multicentre critical care database, it still has some limitations. First, this was a retrospective analysis derived from an observational study, which could not definitively establish causality. Second, the eICU v2.0 did not contain the data on head imaging, neurological severity scores, and follow-up outcomes after discharge. Residual confounding could exist. Third, the baseline levels of plasma glucose and triglyceride could be affected by the use of antidiabetic and lipid-lowering drugs before ICU admission. The TyG index might have changed during hospital; therefore, it is unknown whether the change in the TyG index could have predicted the mortality. Fourth, we did not measure HOMA-IR because the examination of insulin levels is not included in the eICU v2.0. Last, the data were from the United States, and thus the results may not apply fully to ICUs elsewhere with different practices or resources.

Conclusions

In this multicentre critically ill stroke cohort, we observed the association of the TyG index with hospital and ICU mortality in stroke patients. For the first time, this study demonstrated that the TyG index is a potential predictor for hospital and ICU mortality in critically ill stroke patients, especially in IS patients. Furthermore, the TyG index has a linear correlation with the mortality of ICU stroke. Most importantly, these findings suggested that the TyG index may be a useful indicator for risk stratification and prognosis in patients with critically ill stroke. Further prospective studies are required to confirm our findings.

Abbreviations

APACHE: acute physiology and chronic health evaluation; CI: confidence intervals; GBD: global burden of disease; HS: hemorrhage stroke; HOMA-IR: homeostatic model assessment of IR; ICU: intensive care unit; IQR: interquartile range; IR: insulin resistance; IS: ischemic stroke; LDL-C: low-density lipoprotein-cholesterol; LOS: length of stay; OR: odds ratios; SD: standard deviation; TC: total cholesterol; TyG: triglyceride-glucose.

Declarations

Acknowledgements

We thank all the investigators and subjects who participated in this project.

Authors' contributions

BJZ, LLL, HFR, and ZQL designed research. QZ and DFY performed experiments and analyzed data. BJZ and YY wrote the main manuscript text and prepared figures. BJZ edited and revised manuscript. All authors reviewed and approved the manuscript.

Funding

This study was supported by the National Natural Science Foundation of China (81971110, 81671178); the Guangdong Basic and Applied Basic Research Foundation (2020A1515010056); the Third Affiliated Hospital of Sun Yat-Sen University, Clinical Research Program (QHJH201907).

Availability of data and materials

Data were fully available at <https://eicu-crd.mit.edu/>.

Ethics approval and consent to participate

The establishment of this database was approved by the Massachusetts Institute of Technology (Cambridge, MA), and consent was obtained for the original data collection. Therefore, the ethical approval statement and the need for informed consent were waived for this manuscript.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Collaborators GBDS: **Global, regional, and national burden of stroke, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016.** *Lancet Neurol* 2019, **18**(5):439-458.
2. Feigin VL, Nguyen G, Cercy K, Johnson CO, Alam T, Parmar PG, Abajobir AA, Abate KH, Abd-Allah F, Abejie AN *et al*: **Global, Regional, and Country-Specific Lifetime Risks of Stroke, 1990 and 2016.** *New Engl J Med* 2018, **379**(25):2429-2437.
3. Smith M, Citerio G: **Focus on clinical neuroscience.** *Intensive care medicine* 2016, **42**(3):460-462.
4. Alonso A, Ebert AD, Kern R, Rapp S, Hennerici MG, Fatar M: **Outcome Predictors of Acute Stroke Patients in Need of Intensive Care Treatment.** *Cerebrovasc Dis* 2015, **40**(1-2):10-17.
5. Unger G, Benozzi SF, Perruzza F, Pennacchiotti GL: **Triglycerides and glucose index: a useful indicator of insulin resistance.** *Endocrinologia y nutricion : organo de la Sociedad Espanola de Endocrinologia y Nutricion* 2014, **61**(10):533-540.
6. Ma X, Dong L, Shao Q, Cheng Y, Lv S, Sun Y, Shen H, Wang Z, Zhou Y, Liu X: **Triglyceride glucose index for predicting cardiovascular outcomes after percutaneous coronary intervention in patients with type 2 diabetes mellitus and acute coronary syndrome.** *Cardiovascular diabetology* 2020, **19**(1):31.
7. Wang L, Cong HL, Zhang JX, Hu YC, Wei A, Zhang YY, Yang H, Ren LB, Qi W, Li WY *et al*: **Triglyceride-glucose index predicts adverse cardiovascular events in patients with diabetes and acute coronary syndrome.** *Cardiovascular diabetology* 2020, **19**(1):80.
8. Won KB, Park EJ, Han D, Lee JH, Choi SY, Chun EJ, Park SH, Han HW, Sung J, Jung HO *et al*: **Triglyceride glucose index is an independent predictor for the progression of coronary artery calcification in the absence of heavy coronary artery calcification at baseline.** *Cardiovascular diabetology* 2020, **19**(1):34.
9. Li S, Guo B, Chen H, Shi Z, Li Y, Tian Q, Shi S: **The role of the triglyceride (triacylglycerol) glucose index in the development of cardiovascular events: a retrospective cohort analysis.** *Sci Rep* 2019, **9**(1):7320.
10. Shi W, Xing L, Jing L, Tian Y, Yan H, Sun Q, Dai D, Shi L, Liu S: **Value of triglyceride-glucose index for the estimation of ischemic stroke risk: Insights from a general population.** *Nutrition, metabolism, and cardiovascular diseases : NMCD* 2020, **30**(2):245-253.

11. Pollard TJ, Johnson AEW, Raffa JD, Celi LA, Mark RG, Badawi O: **The eICU Collaborative Research Database, a freely available multi-center database for critical care research.** *Scientific data* 2018, **5**:180178.
12. Du T, Yuan G, Zhang M, Zhou X, Sun X, Yu X: **Clinical usefulness of lipid ratios, visceral adiposity indicators, and the triglycerides and glucose index as risk markers of insulin resistance.** *Cardiovascular diabetology* 2014, **13**:146.
13. Wallace TM, Matthews DR: **The assessment of insulin resistance in man.** *Diabetic medicine : a journal of the British Diabetic Association* 2002, **19**(7):527-534.
14. Vasques AC, Novaes FS, de Oliveira Mda S, Souza JR, Yamanaka A, Pareja JC, Tambascia MA, Saad MJ, Geloneze B: **TyG index performs better than HOMA in a Brazilian population: a hyperglycemic clamp validated study.** *Diabetes Res Clin Pract* 2011, **93**(3):e98-e100.
15. Lee JW, Lim NK, Park HY: **The product of fasting plasma glucose and triglycerides improves risk prediction of type 2 diabetes in middle-aged Koreans.** *BMC endocrine disorders* 2018, **18**(1):33.
16. Won KB, Park GM, Lee SE, Cho IJ, Kim HC, Lee BK, Chang HJ: **Relationship of insulin resistance estimated by triglyceride glucose index to arterial stiffness.** *Lipids in health and disease* 2018, **17**(1):268.
17. Park K, Ahn CW, Lee SB, Kang S, Nam JS, Lee BK, Kim JH, Park JS: **Elevated TyG Index Predicts Progression of Coronary Artery Calcification.** *Diabetes Care* 2019, **42**(8):1569-1573.
18. Mao Q, Zhou D, Li Y, Wang Y, Xu SC, Zhao XH: **The Triglyceride-Glucose Index Predicts Coronary Artery Disease Severity and Cardiovascular Outcomes in Patients with Non-ST-Segment Elevation Acute Coronary Syndrome.** *Disease markers* 2019, **2019**:6891537.
19. Jin JL, Cao YX, Wu LG, You XD, Guo YL, Wu NQ, Zhu CG, Gao Y, Dong QT, Zhang HW *et al*: **Triglyceride glucose index for predicting cardiovascular outcomes in patients with coronary artery disease.** *Journal of thoracic disease* 2018, **10**(11):6137-6146.
20. Sánchez-Iñigo L, Navarro-González D, Fernández-Montero A, Pastrana-Delgado J, Martínez JA: **The TyG index may predict the development of cardiovascular events.** *European journal of clinical investigation* 2016, **46**(2):189-197.
21. Sánchez-Iñigo L, Navarro-González D, Fernández-Montero A, Pastrana-Delgado J, Martínez JA: **Risk of incident ischemic stroke according to the metabolic health and obesity states in the Vascular-Metabolic CUN cohort.** *Int J Stroke* 2017, **12**(2):187-191.
22. Ozkul A, Turgut ET, Akyol A, Yenisey C, Kadikoylu G, Tataroglu C, Kiylioglu N: **The relationship between insulin resistance and hypercoagulability in acute ischemic stroke.** *European neurology* 2010, **64**(4):201-206.
23. Harada S, Fujita-Hamabe W, Tokuyama S: **Ischemic stroke and glucose intolerance: a review of the evidence and exploration of novel therapeutic targets.** *Journal of pharmacological sciences* 2012, **118**(1):1-13.
24. Narici MV, Maffulli N: **Sarcopenia: characteristics, mechanisms and functional significance.** *British medical bulletin* 2010, **95**:139-159.

25. Bornfeldt KE, Tabas I: **Insulin resistance, hyperglycemia, and atherosclerosis.** *Cell metabolism* 2011, **14**(5):575-585.
26. Sung KC, Wild SH, Kwag HJ, Byrne CD: **Fatty liver, insulin resistance, and features of metabolic syndrome: relationships with coronary artery calcium in 10,153 people.** *Diabetes Care* 2012, **35**(11):2359-2364.
27. Budoff M: **Triglycerides and Triglyceride-Rich Lipoproteins in the Causal Pathway of Cardiovascular Disease.** *Am J Cardiol* 2016, **118**(1):138-145.
28. Yao M, Ni J, Zhou L, Peng B, Zhu Y, Cui L: **Elevated Fasting Blood Glucose Is Predictive of Poor Outcome in Non-Diabetic Stroke Patients: A Sub-Group Analysis of SMART.** *PLoS One* 2016, **11**(8):e0160674.
29. García-Berrocoso T, Giralt D, Bustamante A, Etgen T, Jensen JK, Sharma JC, Shibazaki K, Saritas A, Chen X, Whiteley WN *et al.*: **B-type natriuretic peptides and mortality after stroke: a systematic review and meta-analysis.** *Neurology* 2013, **81**(23):1976-1985.
30. Singh RJ, Chen S, Ganesh A, Hill MD: **Long-term neurological, vascular, and mortality outcomes after stroke.** *Int J Stroke* 2018, **13**(8):787-796.
31. Seminog OO, Scarborough P, Wright FL, Rayner M, Goldacre MJ: **Determinants of the decline in mortality from acute stroke in England: linked national database study of 795 869 adults.** *Bmj* 2019, **365**:l1778.
32. Sonnevile R, Gimenez L, Labreuche J, Smonig R, Magalhaes E, Bouadma L, Timsit JF, Mazighi M: **What is the prognosis of acute stroke patients requiring ICU admission?** *Intensive care medicine* 2017, **43**(2):271-272.
33. van Valburg MK, Arbous MS, Georgieva M, Brealey DA, Singer M, Geerts BF: **Clinical Predictors of Survival and Functional Outcome of Stroke Patients Admitted to Critical Care.** *Crit Care Med* 2018, **46**(7):1085-1092.
34. Rincon F, Morino T, Behrens D, Akbar U, Schorr C, Lee E, Gerber D, Parrillo J, Mirsen T: **Association between out-of-hospital emergency department transfer and poor hospital outcome in critically ill stroke patients.** *Journal of critical care* 2011, **26**(6):620-625.
35. Suntrup-Krueger S, Schmidt S, Warnecke T, Steidl C, Muhle P, Schroeder JB, Labeit B, Minnerup J, Dziawas R: **Extubation Readiness in Critically Ill Stroke Patients.** *Stroke* 2019, **50**(8):1981-1988.
36. Kirkman MA, Citerio G, Smith M: **The intensive care management of acute ischemic stroke: an overview.** *Intensive care medicine* 2014, **40**(5):640-653.
37. Wang D, Guo Y, Zhang Y, Li Z, Li A, Luo Y: **Epidemiology of acute kidney injury in patients with stroke: a retrospective analysis from the neurology ICU.** *Internal and emergency medicine* 2018, **13**(1):17-25.
38. Wiberg B, Sundström J, Zethelius B, Lind L: **Insulin sensitivity measured by the euglycaemic insulin clamp and proinsulin levels as predictors of stroke in elderly men.** *Diabetologia* 2009, **52**(1):90-96.
39. Wieberdink RG, Koudstaal PJ, Hofman A, Witteman JC, Breteler MM, Ikram MA: **Insulin resistance and the risk of stroke and stroke subtypes in the nondiabetic elderly.** *American journal of epidemiology* 2012, **176**(8):699-707.

40. Howard G, Wagenknecht LE, Kernan WN, Cushman M, Thacker EL, Judd SE, Howard VJ, Kissela BM: **Racial differences in the association of insulin resistance with stroke risk: the REasons for Geographic And Racial Differences in Stroke (REGARDS) study.** *Stroke* 2014, **45**(8):2257-2262.

Tables

Table 1 Baseline clinical and laboratory characteristics of the study patients

Variables	Hospital mortality		P	ICU mortality		P
	Non-survivors n = 469	Survivors n = 4101		Non-survivors n = 229	Survivors n = 4341	
Age, years	69.2 ± 13.3	66.3 ± 14.0	< 0.001	65.8 ± 14.1	66.7 ± 14.0	0.336
Male, n (%)	237(50.5)	2121(51.7)	0.837	116(50.7)	2242(51.6)	0.932
Ethnicity, n (%)			0.172			0.868
Caucasian	341(72.7)	3059(74.6)		170(74.2)	3230(74.4)	
African American	49(10.4)	452(11.0)		25(10.9)	476(11.0)	
Native American	1(0.2)	24(0.6)		0	25(0.6)	
Hispanic	40(8.5)	229(5.6)		14(6.1)	255(5.9)	
Asian	9(1.9)	89(2.2)		4(1.7)	94(2.2)	
Other/unknown	29(6.2)	248(6.0)		16(7.0)	261(6.0)	
Admission height (cm)	168.4 ± 13.3	168.9 ± 15.6	0.514	168.8 ± 11.2	168.8 ± 15.6	0.966
Admission weight (kg)	81.0 ± 21.9	82.7 ± 22.4	0.108	82.4 ± 24.1	82.6 ± 22.3	0.9
Stroke types, n (%)			< 0.001			< 0.001
IS	201 (42.9)	1184 (45.9)		92 (40.2)	1993 (45.9)	
HS	174 (37.1)	907 (22.1)		89 (38.9)	992 (22.9)	
IS with HS	27 (5.8)	109 (2.7)		17 (7.4)	119 (2.7)	
Unknown/others	67 (14.3)	1201 (29.3)		31 (13.5)	1237 (28.5)	
Hypertension, n (%)	169(36.0)	1377(33.6)	0.287	75(32.8)	1471(33.9)	0.723
Diabetes mellitus, n (%)	75(16.0)	480(11.7)	0.007	37(16.2)	518(11.9)	0.056
Coronary heart disease, n (%)	42(9.0)	214(5.2)	0.001	23(10.0)	233(5.4)	0.003
Atrial fibrillation, n (%)	62(13.2)	463(11.3)	0.214	31(13.5)	494(11.4)	0.318
Heart failure, n (%)	24(5.1)	120(2.9)	0.01	15(6.6)	129(3.0)	0.003
Plasma glucose (mg/dl)	211 ± 95.8	164.6 ± 80.7	< 0.001	230.7 ± 112	166.2 ± 80.5	< 0.001
TC (mg/dL)	156.2 ± 47.1	163.2 ± 45.7	0.002	159.4 ± 48.1	162.7 ± 45.8	0.291
LDL-C (mg/dL)	84.3 ± 37.9	92.4 ± 38.3	< 0.001	86.4 ± 38.6	91.8 ± 38.3	0.037
HDL-C (mg/dL)	45.2 ± 17.9	45.5 ± 15.7	0.775	44.2 ± 19.3	45.5 ± 15.8	0.216
Triglyceride (mg/dL)	138.5 ± 95.5	129.9 ± 94.7	0.064	151.9 ± 102.1	129.7 ± 94.3	0.001
Mechanical ventilation, n (%)	387 (82.5)	2217 (54.1)	< 0.001	202 (88.2)	2402 (55.3)	< 0.001
APACHE IV score	70.8 ± 25.0	47.7 ± 19.0	< 0.001	70.8 ± 25.0	47.7 ± 19.0	< 0.001
Hospital LOS, days	5.8 (3.1, 10.2)	5.8 (3.3, 9.0)	0.766	4.6 (2.6, 8.7)	5.8 (3.4, 9.1)	< 0.001
ICU LOS, days	3.7 (2.0, 6.8)	2.0 (1.2, 3.7)	< 0.001	3.9 (2.3, 7.1)	2.0 (1.2, 3.8)	< 0.001
TyG index	9.3 ± 0.7	9.0 ± 0.7	< 0.001	9.3 ± 0.7	9.0 ± 0.7	< 0.001

ICU intensive care unit, IS ischemic stroke, HS hemorrhage stroke, TC total cholesterol, LDL-C low-density lipoprotein-cholesterol, HDL-C high-density lipoprotein-cholesterol, APACHE acute physiology and chronic health evaluation, LOS length of stay, TyG triglyceride-glucose

Table 2 Logistic regression of TyG index for mortality in all study groups.

		Univariate		Model 1		Model 2	
ospital mortality	TyG index	OR (95% CI)	p	OR (95% CI)	p	OR (95% CI)	p
	Per 1-unit increase	1.723 (1.524-1.948)	< 0.001	1.861 (1.637-2.116)	< 0.001	2.543 (1.588-4.073)	< 0.001
	Quartile 1	Reference		Reference		Reference	
	Quartile 2	1.635 (1.170-2.283)	0.004	1.664 (1.190-2.326)	0.003	1.483 (1.030-2.137)	0.034
	Quartile 3	2.294 (1.669-3.153)	< 0.001	2.373 (1.723-3.268)	< 0.001	2.018 (1.383-2.946)	< 0.001
	Quartile 4	3.486 (2.572-4.724)	< 0.001	3.838 (2.819-5.224)	< 0.001	2.546 (1.547-4.191)	< 0.001
	P for trend	< 0.001		< 0.001		< 0.001	
U mortality	TyG index	OR (95% CI)	P	OR (95% CI)		OR (95% CI)	P
	Per 1-unit increase	2.146 (1.826-2.523)	< 0.001	2.183 (1.847-2.580)	< 0.001	2.672 (1.376-5.188)	< 0.001
	Quartile 1	Reference		Reference		Reference	
	Quartile 2	1.977 (1.130-3.459)	0.017	1.973 (1.127-3.452)	0.017	1.521 (0.827-2.797)	0.177
	Quartile 3	3.564 (2.123-5.982)	< 0.001	3.573 (2.127-6.002)	< 0.001	2.513 (1.377-4.586)	0.003
	Quartile 4	6.173 (3.764-10.126)	< 0.001	6.193 (3.767-10.180)	< 0.001	2.826 (1.341-5.956)	0.006
	P for trend	< 0.001		< 0.001		< 0.001	

Model 1 adjusted for gender, age, ethnicity.

Model 2 adjusted for model 1 plus stroke types, coronary heart disease, diabetes mellitus, heart failure, plasma glucose, TC, LDL-C, triglyceride, mechanical ventilation, APACHE IV score, LOS.

ICU intensive care unit, TyG triglyceride-glucose, OR odds ratios, CI confidence intervals, TC total cholesterol, LDL-C low-density lipoprotein-cholesterol, APACHE acute physiology and chronic health evaluation, LOS length of stay

Figures




Figure 1

The hospital and ICU mortality according to TyG index quartiles. ICU intensive care unit, TyG triglyceride-glucose




Figure 2

The smoothing curves of the hospital (A) and ICU (B) mortality of critically ill stroke against TyG index. ICU intensive care unit, TyG triglyceride-glucose




Figure 3

The forest plot of the hospital mortality of critically ill stroke according to stroke types and TyG index. TyG triglyceride-glucose




Figure 4

The forest plot of the ICU mortality of critically ill stroke according to stroke types and TyG index. ICU intensive care unit, TyG triglyceride-glucose